

# Genome Sequence of *Paraburkholderia nodosa* Strain CNPSO 1341, a N<sub>2</sub>-Fixing Symbiont of the Promiscuous Legume *Phaseolus vulgaris*

Rebeca Fuzinato Dall'Agnol,<sup>a,b</sup> Maira Rejane Costa,<sup>a,c</sup> Renan Augusto Ribeiro,<sup>d</sup> Jakeline Renata Marçon Delamuta,<sup>a,e</sup> Ligia Maria Oliveira Chueire,<sup>d</sup>  Mariangela Hungria<sup>a,b,c,d</sup>

Embrapa Soja, Soil Biotechnology Laboratory, Londrina, Paraná, Brazil<sup>a</sup>; Department of Biochemistry and Biotechnology, Universidade Estadual de Londrina, Londrina, Paraná, Brazil<sup>b</sup>; Department of Microbiology, Universidade Estadual de Londrina, Londrina, Paraná, Brazil<sup>c</sup>; CNPq, SHIS QI 1 Conjunto B, Brasília, Federal District, Brazil<sup>d</sup>; CAPES, SBN, Quadra 2, Brasília, Federal District, Brazil<sup>e</sup>

***Paraburkholderia nodosa* CNPSO 1341 is a N<sub>2</sub>-fixing symbiont of *Phaseolus vulgaris* isolated from an undisturbed soil of the Brazilian Cerrado. Its draft genome contains 8,614,032 bp and 8,068 coding sequences (CDSs). Nodulation and N<sub>2</sub>-fixation genes were clustered in the genome that also contains several genes of secretion systems and quorum sensing.**

Received 8 August 2016 Accepted 12 September 2016 Published 3 November 2016

**Citation** Dall'Agnol RF, Costa MR, Ribeiro RA, Delamuta JRM, Chueire LMO, Hungria M. 2016. Genome sequence of *Paraburkholderia nodosa* strain CNPSO 1341, a N<sub>2</sub>-fixing symbiont of the promiscuous legume *Phaseolus vulgaris*. *Genome Announc* 4(6):e01073-16 doi:10.1128/genomeA.01073-16.

**Copyright** © 2016 Dall'Agnol et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Mariangela Hungria, mariangela.hungria@embrapa.br.

*Paraburkholderia* has been recently validated as a new genus (1–3) and encompasses several environmental *Burkholderia* (1, 4, 5). The N<sub>2</sub>-fixing *Paraburkholderia* that nodulate legumes are also called beta-rhizobia (6). *Paraburkholderia nodosa* was first isolated from nodules of *Mimosa* (*M. bimucronata* and *M. scabrella*) (7), and later from other leguminous plants from the “*Piptadenia* group” (8), all belonging to the *Mimosoideae* subfamily; more rarely, it nodulates members of the *Papilionoidea* subfamily, as *Phaseolus vulgaris* (9). *P. nodosa* strain CNPSO 1341 was trapped by the promiscuous common bean (*Phaseolus vulgaris*) from an undisturbed soil of the Brazilian Cerrado (9), and here we present its draft genome.

To extract the total bacterial DNA, we used the DNeasy Blood and Tissue Kit (Qiagen) and processed genome sequencing on the MiSeq platform (Illumina) at Embrapa Soja, Londrina, Brazil. Shotgun sequencing generated 2,081,314 paired-end reads (2 × 300 bp), corresponding to approximately a 73.45-fold coverage. The FASTQ files were assembled by the A5-miseq pipeline (*de novo* assembly) (10). The genome of strain CNPSO 1341 was estimated at 8,614,032 bp, with a G+C content of 64.2 mol%, and assembled in 86 contigs, with 8,068 predicted coding sequences (CDSs).

Sequences were submitted to RAST (11) and by analyzing the sequences in the SEED system (11), we determined that 47% of CDSs had coverage in 530 subsystems, the majority in the carbohydrates and amino acids and derivatives categories. The highest genome scores were with *Paraburkholderia* sp. Ch1-1 and *Paraburkholderia xenovorans* LB400, two strains well known for their capacity of degradation of xenobiotics and tolerance of stresses. Indeed, *P. nodosa* CNPSO 1341 carries 249 CDSs related to stress response, 45% of them of oxidative stress, in addition to 204 CDSs of the metabolism of aromatic compounds.

In relation to the symbiosis, the nodulation (*nod* genes) operons were followed by the nitrogen fixation (*nif*) operons. We found one copy of the regulatory *nodD* gene (LysR family) that

orchestrates the nodulation process (12), and interestingly, adjacent to the *nodD* there is a gene related to nikkomycin biosynthesis. The genome of *P. nodosa* CNPSO 1341 carries several other CDSs of resistance to and biosynthesis of antibiotics. Preceding the *nifA* there is another regulatory gene of the LysR family that is also present in a plasmid of *Burkholderia* CCGE 1001 and deserves further investigation.

Noteworthy is the variety of secretion systems found in the genome of *P. nodosa* CNPSO 1341. Among them, the complete operon of the secretion system X, which shows greater similarity to the plant pathogen *Ralstonia solanacearum*. In relation to quorum sensing, there is a pair of genes of quorum-sensing LuxR-LuxI, of homoserine lactone, but in addition there are nine other CDSs of transcription regulators of the LuxR family, which might be related to the perception of several signals.

**Accession number(s).** This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession number SUBID SUB1743636 and BioProject PRJNA337952, BioSample SAMN05514346, and accession number [MCNV00000000](https://ncbi.nlm.nih.gov/sra/NCN00000000). This paper describes the first version.

## FUNDING INFORMATION

This work was funded by Embrapa (02.13.08.001.00.00) and MCTI/CNPq/CAPES/FAPs (INCT-MPCPAgro).

## REFERENCES

1. Sawana A, Adeolu M, Gupta RS. 2014. Molecular signatures and phylogenomic analysis of the genus *Burkholderia*: proposal for division of this genus into the emended genus *Burkholderia* containing pathogenic organisms and a new genus *Paraburkholderia* gen. nov. harboring environmental species. *Front Genet* 5:1–22. <http://dx.doi.org/10.3389/fgene.2014.00429>.
2. Oren A, Garrity GM. 2015. List of new names and new combinations previously effectively, but not validly, published. *Int J Syst Evol Microbiol* 65:2017–2025. <http://dx.doi.org/10.1099/ijs.0.000317>.
3. Dobritsa AP, Samadpour M. 2016. Transfer of eleven *Burkholderia* species to the genus *Paraburkholderia* and proposal of *Caballeronia* gen. nov.,

- a new genus to accommodate twelve species of *Burkholderia* and *Paraburkholderia*. *Int J Syst Evol Microbiol* 66:2836–2846. <http://dx.doi.org/10.1099/ijsem.0.001065>.
4. Estrada-de los Santos P, Vinuesa P, Martínez-Aguilar L, Hirsch AM, Caballero-Mellado J. 2013. Phylogenetic analysis of *Burkholderia* species by multilocus sequence analysis. *Curr Microbiol* 67:51–60. <http://dx.doi.org/10.1007/s00284-013-0330-9>.
  5. Zuleta LF, Cunha Cde O, de Carvalho FM, Ciapina LP, Souza RC, Mercante FM, de Faria SM, Baldani JI, Stralioetto R, Hungria M, de Vasconcelos AT. 2014. The complete genome of *Burkholderia phenoliruptrix* strain BR3459a, a symbiont of *Mimosa flocculosa*: highlighting the coexistence of symbiotic and pathogenic genes. *BMC Genomics* 15:535. <http://dx.doi.org/10.1186/1471-2164-15-535>.
  6. Moulin L, Munive A, Dreyfus B, Boivin-Masson C. 2001. Nodulation of legumes by members of the beta-subclass of *Proteobacteria*. *Nature* 411:948–950. <http://dx.doi.org/10.1038/35082070>.
  7. Chen WM, De Faria SM, James EK, Elliott GN, Lin KY, Chou JH, Sheu SY, Cnockaert M, Sprent JI, Vandamme P. 2007. *Burkholderia nodosa* sp. nov., isolated from root nodules of the woody Brazilian legumes *Mimosa bimucronata* and *Mimosa scabrella*. *Int J Syst Evol Microbiol* 57:1055–1059. <http://dx.doi.org/10.1099/ijms.0.64873-0>.
  8. Bournaud C, da Faria SM, dos Santos JM, Tisseyre P, Silva M, Chaintréuil C, Gross E, James EK, Prin Y, Moulin L. 2013. *Burkholderia* species are the most common and preferred nodulating symbionts of the Piptadenia group (tribe Mimoseae). *PLoS One* 8:e63478. <http://dx.doi.org/10.1371/journal.pone.0063478>.
  9. Dall'Agnol RF, Plotegher F, Souza RC, Mendes IC, Dos Reis Junior FB, Béna G, Moulin L, Hungria M. 2016. *Paraburkholderia nodosa* is the main N<sub>2</sub>-fixing species trapped by promiscuous common bean (*Phaseolus vulgaris* L.) in the Brazilian “Cerradão”. *FEMS Microbiol Ecol* 92:fiw108. <http://dx.doi.org/10.1093/femsec/fiw108>.
  10. Coil D, Jospin G, Darling AE. 2015. A5-miseq: an updated pipeline to assemble microbial genomes from Illumina MiSeq data. *Bioinformatics* 31:587–589. <http://dx.doi.org/10.1093/bioinformatics/btu661>.
  11. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST Server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
  12. Del Cerro P, Rolla-Santos AA, Gomes DF, Marks BB, Pérez-Montaña F, Rodríguez-Carvajal MÁ, Nakatani AS, Gil-Serrano A, Megias M, Ollero FJ, Hungria M. 2015. Regulatory *nodD1* and *nodD2* genes of *Rhizobium tropici* strain CIAT 899 and their roles in the early stages of molecular signaling and host-legume nodulation. *BMC Genomics* 16:251. <http://dx.doi.org/10.1186/s12864-015-1458-8>.