



Pharmacognosy Research

Publication of Pharmacognosy Network Worldwide

www.phcogres.com



Medknow

 Wolters Kluwer

Phcog.Net - Bringing Medicinal Plant Researchers Together

Synergisms in Alpha-glucosidase Inhibition and Antioxidant Activity of *Camellia sinensis* L. Kuntze and *Eugenia uniflora* L. Ethanolic Extracts

Juliana Vinholes, Márcia Vizzotto

Embrapa Clima Temperado, Rodovia BR-392, Km 78, 9º Distrito, Monte Bonito Caixa Postal 403, CEP: 96010-971 - Pelotas, RS, Brazil

ABSTRACT


Background: *Camellia sinensis*, the most consumed and popular beverages worldwide, and *Eugenia uniflora*, a Brazilian native species, have been already confirmed to have beneficial effects in the treatment of diabetes mellitus. However, their potential acting together against an enzyme linked to this pathology has never been exploited. **Objective:** The aim of this study was to evaluate the inhibitory properties of individual and combined ethanolic extracts of the leaves of *C. sinensis* and *E. uniflora* over alpha-glucosidase, a key digestive enzyme used on the Type 2 diabetes mellitus (T2DM) control. In addition, their inhibitory activity against 2,2-diphenyl-1-picrylhydrazyl radical (DPPH[•]) and peroxy radicals was also assayed. **Materials and Methods:** Enzyme inhibition and antioxidant potential were assessed based on *in vitro* assays. Total phenolic compounds, carotenoids, and chlorophylls A and B were achieved using spectrophotometric methods. **Results:** *E. uniflora* was almost 40 times more active on alpha-glucosidase than *C. sinensis* and combined extracts showed a significant synergistic effect with an obtained IC₅₀ value almost 5 times lower than the theoretical value. *C. sinensis* extract was twice more active than *E. uniflora* concerning DPPH[•], in contrast, *E. uniflora* was almost 10 times more effective than *C. sinensis* on inhibition of peroxy radicals with a significant synergistic effect for combined extracts. The extracts activities may be related with their phytochemicals, mainly phenolic compounds, and chlorophylls. **Conclusion:** Combined *C. sinensis* and *E. uniflora* ethanolic extracts showed synergistic effect against alpha-glucosidase and lipid peroxidation. These herbal combinations can be used to control postprandial hyperglycemia and can also provide antioxidant defenses to patients with T2DM.

Key words: Additive effect, antihyperglycemic effect, antiradical activity, diabetes, phytochemicals, synergistic effect

SUMMARY

- Alpha-glucosidase and antioxidant Interaction between *Camellia sinensis* L. Kuntze and *Eugenia uniflora* L. ethanolic extracts was investigated.

- Extracts showed synergistic effect over alpha-glucosidase and peroxy radicals.
- Total phenolic, carotenoids and chlorophylls A and B can be responsible by the observed activities.
- Extracts could be used as alternative to control postprandial hyperglycemia.
- Extracts could increase antioxidant defenses to patients with T2DM.



Abbreviations Used: T2DM: Type 2 diabetes mellitus; DPPH[•]: 2,2-diphenyl-1-picrylhydrazyl radical; PNPG: 4-Nitrophenyl β-D-glucuronide; LOO[•]: Lipid peroxidation; SEM: Standard error of the mean; CAE: Chlorogenic acid equivalent

Correspondence:

Dr. Juliana Vinholes,
Embrapa Clima Temperado, BR 392, KM 78,
C. P. 403, CEP 96010-971, Pelotas-RS, Brazil.
E-mail: julianarochavinholes@gmail.com
DOI: 10.4103/0974-8490.197797

Access this article online

Website: www.phcogres.com

Quick Response Code:



INTRODUCTION

Diabetes mellitus is becoming a major public health concern, with high social and health-care costs since it affects over 387 million people worldwide causing 4.9 million deaths in 2014 (1 death each 7 s) according to the International Diabetes Federation. In Brazil, this pathology is present in around 13 million people between 20 and 79 years old.^[1] Type 2 diabetes mellitus (T2DM) accounts for 90% of cases of diabetes and is characterized by individuals with postprandial hyperglycemia associated with low production of insulin, resistance to insulin, or both. One strategy used to control T2DM is the use of inhibitors of digestive enzymes such as alpha-glucosidase that is present in the intestine which catalyzes the digestion of complex carbohydrates, converting them into easily digestible monosaccharides.^[2] Inhibitors of this enzyme are used by individuals with T2DM to promote a decrease in glucose uptake and consequently a reduction in blood sugar levels. Different glucosidase inhibitors are currently used in patients with T2DM, namely acarbose, the first alpha-glucosidase inhibitor that is produced by fermentation of actinomycetes called *Actinoplanes* sp. and miglitol which is

synthesized starting from the naturally occurring 1-deoxynojirimycin as a lead structure.^[3] Since then, several studies have been performed in different species aiming to find new sources of inhibitors of this enzyme due to increase cases of T2DM in the world. Researchers have proposed different species as natural sources of alpha-glucosidase inhibitors including *Camellia sinensis* L. Kuntze (green tea), the most consumed and popular beverages worldwide,^[4] and *Eugenia uniflora* L. (Brazilian Pitanga), a Brazilian native species that, shows in recent years an interesting potential as source of bioactive compounds. Both

This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Cite this article as: Vinholes J, Vizzotto M. Synergisms in alpha-glucosidase inhibition and antioxidant activity of *Camellia sinensis* L. Kuntze and *Eugenia uniflora* L. Ethanolic Extracts. Phcog Res 2017;9:101-7.

species have been already confirmed to have beneficial effects in the treatment of diabetes mellitus. *C. sinensis* tea consumption was effective against type 2 diabetes in a retrospective cohort studies in Japan and Taiwan.^[5,6] Moreover, different studies also showed the positive effect of *C. sinensis* on T2DM prevention and treatment that was related with their phytochemicals mainly flavonols,^[7] methylxanthine alkaloids,^[8] and polysaccharides^[9] by different mechanisms including inhibition of glucosidases.^[10] *E. uniflora* leaves have been empirically used in the T2DM treatment and also showed inhibitory properties against alpha-glucosidase.^[11-15] *E. uniflora* leaves are source of macrocyclic hydrolysable tannin dimers (eugeniflorins D1 and D2), oenothien B, 1,2,4,6-tetra-*O*-galloyl-fl-*o*-glucose, galocatechin and myricitrin, compounds that may be responsible for the species activity.^[16] Besides, it is well known that leaves also contain chlorophylls and carotenoids pigments, and these phytochemicals compounds can also confer good antioxidant activities for both species, which is an additional feature for the treatment of patients with T2DM which have their antioxidants defenses altered.

It is well established that when compounds with different properties are combined, numerous interactions can occur toward each other which can result in effects different from the formers. These effects can be classified as synergistic, antagonistic, or additive. The combination of extracts with different composition and the presence of synergistic effect is of great interest from the pharmacological point of view since the biological effect of the combined product is greater than the sum of individual agents.^[17] Thus, smaller quantities of extracts are required to achieve the desirable effect which may improve the health-promoting properties of both products. Thus, the aim of this work was to determine the alpha-glucosidase inhibitory activity, and antioxidant effect of combined ethanolic extracts of *C. sinensis* L. Kuntze and *E. uniflora* L. leaves to assess their potential use in the T2DM treatment. In addition, the phytochemicals present in both extract were also determined and their relation with the observed biological effects discussed.

MATERIALS AND METHODS

Chemicals

Reference compounds and reagents were purchased from different suppliers. 2,2-diphenyl-1-picrylhydrazyl radical (DPPH[•]), iron sulfate heptahydrate, linoleic acid, Tris-HCl, phosphate buffer, alpha-glucosidase (type I from baker's yeast), and 4-nitrophenyl α -D-glucopyranoside (PNP-G) were obtained from Sigma-Aldrich (St. Louis, MO, USA). Ethanol was purchased from Synth (Diadema, SP, Brazil), ethyl ether, dichloromethane, ascorbic acid, phosphoric acid hydrogen peroxide solution (30% w/w) was from Impex (Diadema, SP, Brazil) and Acarbose (Glucobay[®]) was from Bayer Pharma AG (Leverkusen, Germany).

Samples

C. Sinensis (green tea), from a commercial brand, was purchased from local market in Pelotas and deposited in the laboratory of Food Science and Technology under the number: CS-01. *E. uniflora* leaves from purple genotype (plant identification PIT102, deposit number: ECT450) were collected from the Active Germplasm Bank of native fruits at Embrapa Clima Temperado (Brazil, 31°40'47"S, 52°26'24"W), on November 21, 2014 and identified by the PhD Gustavo Heiden Curator of the Embrapa Clima Temperado Herbarium. After collection, the sample was transported immediately to the laboratory, where it was placed in oven at 37°C until constant weight. Samples were powdered and sieved (<1 mm) for further extraction.

Preparation of extracts

Samples were extracted with ethanol 95% using 1 g of sample/100 ml of solvent, at 200 rpm, during 60 min. Extracts were filtered through paper filter (Whatman n°4) and further evaporated under pressure at 40°C. Ethanolic extracts yields were 14.47% \pm 2.47% and 6.87% \pm 1.08% for *C. sinensis* and *E. uniflora*, respectively. Samples were redissolved in ethanol leading to extracts with concentration of 3.5 mg/ml, which were further stored at -20°C until analysis.

Alpha-glucosidase inhibition and antioxidant potential

General

To evaluate the antihyperglycemic and antioxidant potential of leaves extracts *in vitro* assays were performed by spectrophotometric methods using an Amersham, ultraviolet-visible Ultrospec-3100 Pro Amersham Bioscience spectrophotometer. The IC₅₀ values were calculated using at least 5 concentrations for each extract, and combined extracts (1:1). Three extracts were prepared for each sample and assays were performed in triplicate (*n* = 9) and expressed as \pm standard error of the mean (SEM).

Alpha-glucosidase inhibitory activity

The effect on alpha-glucosidase was assessed using a procedure previously reported with a slightly modification.^[18] Briefly, 20 μ l of extract or ethanol (control) was added to a vial with 100 μ l of PNP-G (3.25 mM) in phosphate buffer (pH 7.0). The reaction was initiated by the addition of 100 μ l of enzyme (9.37 U/ml in phosphate buffer, pH 7.0), and vials were incubated at 37°C for 10 min. The reaction was stopped by adding 0.600 ml of Na₂CO₃ (1M), and the absorbance at 405 nm was measured. Acarbose was used as positive control (85-1360 μ g/ml).

2,2-diphenyl-1-picrylhydrazyl radical scavenging activity

The hydrogen atoms or electrons donation ability of the extracts was measured from the bleaching of purple-colored methanol solution of DPPH[•] by adaptation of the methods reported in the literature.^[19,20] Briefly, 50 μ l of each extract or ethanol (control) were added to 200 μ l of a 0.6 mM DPPH[•] methanol solution. The reaction was mixed and incubated in the dark for 30 min at room temperature; samples were read at 515 nm.

Lipid peroxidation inhibition

Lipid peroxidation (LOO[•]) was measured according to the method described in the literature.^[21] Briefly, the reaction mixture contained 50 μ l of extract or ethanol (control), 250 μ l linoleic acid (20 mM), 150 μ l Tris-HCl (100 mM, pH 7.5), and 50 μ l FeSO₄·7H₂O (4 mM). Linoleic acid peroxidation was initiated by the addition of 50 μ l of ascorbic acid (5 mM) followed by incubation for 60 min at 37°C. The addition of 1.5 ml of ethanol-ether (3:1, v/v) and vortexing for 1 min allowed the separation of conjugated dienes in the organic layer that was spectrophotometrically measured at 233 nm.

Calculation of effects

Theoretical effects values for alpha-glucosidase and antioxidant activities of the mixtures were calculate as weighted mean experimental IC₅₀ values [Table 1] and considering the additive contributions of 50% individual extracts as follows:

$$\text{Theoretical IC}_{50} = \text{IC}_{50} \text{ } C. \text{ sinensis} \times 0.50 + \text{IC}_{50} \text{ } E. \text{ uniflora} \times 0.50$$

The classification in additive, synergistic, or antagonistic effects was performed by comparison of obtained IC₅₀ values with the theoretical IC₅₀ value according to literature.^[22,23] The interaction was considered additive when theoretical IC₅₀ and experimental IC₅₀ values show

Table 1: IC₅₀ values for alpha-glucosidase, 2,2-diphenyl-1-picrylhydrazyl radical, and lipid peroxidation inhibition of individual and combined ethanolic extracts of *Camellia sinensis* L. Kuntze and *Eugenia uniflora* L.

	Acarbose	<i>Camellia sinensis</i> L.	<i>Eugenia uniflora</i> L.	<i>Camellia sinensis</i> L. + <i>Eugenia uniflora</i> L.		Effect
				Experimental	Theoretical	
Alpha-glucosidase, IC ₅₀ (µg/ml)	413.6±20.2	10.68±2.02	0.26±0.05	1.04±0.20***	5.47±1.02	SN
DPPH, IC ₅₀ (µg/ml)		8.28±0.27	15.45±0.70	12.72±0.04	11.87±0.36	AN
Lipid peroxidation, IC ₅₀ (µg/ml)		10.42±0.09	0.96±0.18	3.87±0.54*	5.69±0.23	SN

Results are expressed as mean values±SEM of three experiments performed in triplicate (n=9). Significance differences were compared between obtained and theoretical values (*P<0.05, ***P<0.0001). SN: Synergistic effect; AN: Antagonistic effect; DPPH: 2,2-diphenyl-1-picrylhydrazyl radical; SEM: Standard error of the mean

differences lower than 5%; for synergistic effect, the experimental IC₅₀ values are more than 5% lower than theoretical values, and antagonistic effect when experimental IC₅₀ values was more than 5% higher when compared with theoretical values as can be seen in Figure 1.^[24]

Phytochemical analysis

Total phenolic compounds

Total phenolic content was measured according to the Folin-Ciocalteu method adapted from Swain and Hillis.^[25] Briefly, 50 µl aliquot of the extract and the control (50 µl of ethanol) were each combined with 250 µL of 0.25 N Folin-Ciocalteu reagent. After 3 min reaction, 500 µl of Na₂CO₃ (1N) was added, the mixtures were incubated for 2 h at room temperature, the absorbance was measured at 725 nm, and the results were expressed as mg of chlorogenic acid equivalents per 100 g of sample (CAE mg/100 g of sample) using a chlorogenic acid (0–0.4 mg/ml) standard curve.

Total carotenoids and chlorophylls content

Carotenoid, chlorophyll A, and chlorophyll B contents were assessed according to literature.^[26] The ethanolic extracts were analyzed at different wavelengths and quantified according to the following equations:

$$\text{Chlorophyll A} = 13.36A_{664} - 5.19 A_{649}$$

$$\text{Chlorophyll B} = 27.43A_{649} - 8.12 A_{664}$$

Carotenoids = (1000A₄₇₀ - 2.13Chlorophyll A - 97.63Chlorophyll B)/209 where A₄₇₀, A₆₄₉, and A₆₆₄ are the absorbance's read at 470 nm, 649 nm, and 664 nm, respectively. Results are expressed as µg/g of sample.

Statistical analysis

Data are reported as mean ± SEM of three extracts analyzed in triplicate (n = 9). One-way analysis of variance was employed to compare the means related to the evaluated parameters. Significant differences were considered when P < 0.05.

RESULTS

Alpha-glucosidase inhibition

A dose response inhibitory effect over alpha-glucosidase was observed for both extracts and their combination as can be observed in Figure 2. The inhibition of 50% of the enzyme (IC₅₀) was lower for *E. uniflora*, followed by combined extracts and *C. sinensis* [Table 1]. The combination was found to be synergistic since the experimental IC₅₀ values were more than 5% lower than theoretical value [Table 1]. Both extracts and their combination showed IC₅₀ lower than acarbose (positive control).

2,2-diphenyl-1-picrylhydrazyl radical scavenging activity

All ethanolic extracts, under the assays conditions, inhibit the free radical DPPH[•] in a concentration-dependent way [Figure 3a] with IC₅₀ values ranging from 8.28 to 15.45 µg/ml [Table 1]. *C. sinensis* was almost




Figure 1: Classification in additive, synergistic, or antagonistic effects using the inhibition of alpha-glucosidase as example. TIC₅₀ = Theoretical IC₅₀ and EIC₅₀ = Experimental IC₅₀

2 times more effective than *E. uniflora*. The extracts combination showed an antagonistic effect on DPPH[•] since the experimental value of IC₅₀ was 6.68%, which was higher than 5%, when compared with the theoretical IC₅₀ value [Table 1].

Lipid peroxidation inhibition

C. sinensis and *E. uniflora* ethanolic leaves extracts and their combination were evaluated as inhibitors of LOO[•]. Figure 3b displays the dose-dependent behavior observed for all extracts. *E. uniflora* was 10 times more effective than *C. sinensis* with IC₅₀ values of 0.96 and 10.42 µg/ml [Table 1], respectively. A synergistic effect was observed when combining both extracts on the LOO[•] inhibition, with an experimental value of IC₅₀ 3.87 µg/ml, which is lower than the theoretical value and lower than the IC₅₀ found for *C. sinensis* alone.

Phytochemicals

Total phenolic compounds

The total phenolic content of the *C. sinensis* and *E. uniflora* extracts was measured, and the results were expressed as milligram of CAE per 100 g dried weight sample. As can be observed on Table 2, the total phenolic compounds in *C. sinensis* (812.58 ± 117.15 mg of CAE/100 g of sample) and *E. uniflora* (663.82 ± 107.58 mg of CAE/100 g of sample) were not significantly different.

Total carotenoids

The total content of carotenoids in ethanolic extracts of *C. sinensis* and *E. uniflora* is presented in Table 2. *C. sinensis* ($0.13 \pm 0.03 \mu\text{g/g}$ of sample) was 2-fold richer in these compounds than *E. uniflora* ($0.07 \pm 0.02 \mu\text{g/g}$ of sample).

Chlorophylls

The contents of chlorophylls in the extracts are presented in Table 2. *C. sinensis* also showed higher concentrations of chlorophyll A ($0.96 \pm 0.16 \text{ mg/g}$ of sample) than *E. uniflora* ($0.60 \pm 0.09 \text{ mg/g}$ of sample), conversely, *E. uniflora* had higher amounts of chlorophyll B ($0.46 \pm 0.11 \text{ mg/g}$ of extract), almost 2 times, when compared to *C. sinensis* ($0.18 \pm 0.02 \text{ mg/g}$ of sample).

DISCUSSION

Hyperglycemia, typical in patients with T2DM, is characterized by the abnormal increase in the level of fasting and postprandial blood glucose. Thus, controlling postprandial hyperglycemia is a major therapeutic approach for T2DM management. Different strategies can be used to decrease the high levels of glucose after a carbohydrate-rich meal, inhibit the enzymes responsible for their digestion is an example. Thus, in this study, we evaluated the potential of *C. sinensis* and *E. uniflora* and their combination as inhibitors of alpha-glucosidase, an enzyme present in the human body responsible for the carbohydrate breakdown.

Both extracts were tested at same concentration; however, *E. uniflora* achieved almost 100% of alpha-glucosidase inhibition with lower amounts. As consequence, the IC_{50} found for *C. sinensis* was almost 40 times higher than the IC_{50} of *E. uniflora*. *C. sinensis* and *E. uniflora* ethanolic extracts were 40 and 1500 times more effective than the positive control acarbose, respectively. *C. sinensis* showed an IC_{50} ($10.68 \mu\text{g/ml}$) almost 30 times lower than reported for a hydro-alcoholic extract ($IC_{50} = 299 \mu\text{g/ml}$).^[27] In addition, this value is slightly higher than the IC_{50} reported in the literature for *C. sinensis* water extracts $4.42 \mu\text{g/ml}$, but lower than $2040 \mu\text{g/ml}$ reported by other authors.^[28,29] This variation was expected since samples, extraction procedure and enzymatic protocols used were not the same. The alpha-glucosidase inhibitory value for *E. uniflora* ($0.26 \mu\text{g/ml}$) [Table 1] was lower than the value described in the literature.^[14] These authors have studied different fractions of leaves ethanolic extract, instead of whole extract, and have

found that 4 fractions were capable of inhibiting almost 50% of the enzyme at a concentration of $100 \mu\text{g/ml}$. Nevertheless, both extracts were more effective on alpha-glucosidase inhibition than extracts of *Bauhinia* species, commonly used to treat T2DM.^[30]

Concerning the interaction between *C. sinensis* and *E. uniflora* ethanolic leaves extracts on the inhibitory activity of alpha-glucosidase, the value obtained was almost 400 times more effective than acarbose. A similar result was observed for the herbal mixture of *Allium sativum* plus *Lagerstroemia speciosa* on the inhibition of alpha-glucosidase, where their combination was also more effective than the positive control miglitol.^[31]

Since the alpha-glucosidase inhibition by combined extracts was greater than the individual extracts, lower amounts of extract were needed to achieve the biological effect. This fact can be very beneficial to human health, due to the increased therapeutic effect and the reduced toxicity. As far as we are concerned, this is the first report about the interaction of *C. sinensis* and *E. uniflora* ethanolic leaves extracts and their inhibitory effect against the alpha-glucosidase enzyme.

As recent studies shows that T2DM patients have an increased free-radical production and reduced antioxidant defense, which leads to different health complications, supplementation with antioxidants agents can be useful in their treatment. Therefore, the antiradical potential of *C. sinensis*, *E. uniflora*, and combined extracts was evaluated.

Different *in vitro* methods can be used to measure the efficiency of natural antioxidant compounds either as pure or as plant mixtures. Owing to the complex nature of extracts and their mechanisms of




Figure 2: Effect of different concentrations of individual and combined extracts of *Camellia sinensis* and *Eugenia uniflora* on alpha-glucosidase




Figure 3: Effect of different concentrations of individual and combined extracts of *Camellia sinensis* and *Eugenia uniflora* on 2,2-diphenyl-1-picrylhydrazyl radical (a) and lipid peroxidation (b)

Table 2: Total phenolic, carotenoids, chlorophylls (A and B) contents in ethanolic extracts of *Camellia sinensis* L. Kuntze and *Eugenia uniflora* L

Sample	Phenolics ^a	Carotenoids ^b	Chlorophyll A ^b	Chlorophyll B ^b
<i>Camellia sinensis</i>	812.58±117.15	0.13±0.02	0.96±0.16	0.18±0.02
<i>Eugenia uniflora</i>	663.82±107.58	0.07±0.01	0.60±0.09	0.46±0.11

^amg of CAE/100 g of sample, ^bmg/g of sample. Results are expressed as mean values±SEM of three experiments performed in triplicate (n=9). CAE: Chlorogenic acid equivalent; SEM: Standard error of the mean

action, a single method is not capable to provide a comprehensive view of their antioxidant profile. In addition, the comparison of results with those found in the literature is very difficult since the assays conditions cannot be exactly the same and the relative effectiveness of antiradical compound is highly dependent on their concentration, test system, time, and selected assay. Thus, the radical-scavenging activity of *C. sinensis* and *E. uniflora* and their combination was assessed by means of two assays: DPPH[•] and peroxy radicals.

The hydrogen atoms or electrons donation ability of *C. sinensis* and *E. uniflora* ethanolic leaves extracts and their combination was measured from the bleaching of purple colored methanol solution of DPPH[•]. This assay is frequently used for the screening of the antiradical activity of different matrix and isolated compounds, due to its reproducibility, simplicity, and fastness. The present study showed better results on the inhibitory effect over the free radical DPPH[•] when comparing the IC₅₀ found for *C. sinensis* with those reported in the literature for ethanolic extract (IC₅₀ = 10.35 ± 0.14 µg/ml), water extracts (IC₅₀ = 15.63 and 60.00 µg/ml), and hydroalcoholic extracts (IC₅₀ = 201.3 µg/ml).^[29,32,33] Antagonistic effect occurs when the combination of extracts reduces the observed activity relatively to extracts tested alone, indicating the *E. uniflora* and *C. sinensis* combined extracts had a negative effect over DPPH[•] inhibition. Nevertheless, this is the first report of the DPPH[•] inhibitory effect of *C. sinensis* and *E. uniflora* combined extracts, as far as we known.

Peroxy radicals (LOO[•]) are the product of oxidation of lipids after the attack of reactive oxygen species. In individuals with diabetes, this product is present in higher amounts when compared to nondiabetic ones.^[34] This radical species can cause alterations on cell membrane lipids, process that can result in cell damage, death, and neoplasia which are probably involved in the complications of diabetes and also in the incidence of several chronic and degenerative diseases. Concerning *E. uniflora* extracts, results found under the assay conditions tested in the present study showed inhibitory properties 6–60 times lower than reported in the literature.^[32,35] *E. uniflora* ethanolic extracts inhibited the LOO[•] in rat's brain and liver homogenates at concentrations ranging from 6.3 to 50 µg/ml.^[35] In addition, the administration of an aqueous extract (IC₅₀ = 60.00 µg/ml) to Type 1 non obese diabetic mice reduces in 76% the serum LOO[•] when compared to the untreated and by 69% when compared with acute diabetic animals.^[32] In respect to *C. sinensis*, the IC₅₀ value found was 30 times lower than that reported for an ethanolic extract (IC₅₀ = 333.29 ± 17.90 µg/ml).^[36] Moreover, a statistically significant decrease on LOO[•] markers in diabetic patients treated with green tea extract capsule was also reported (200 mg of standardized extract of *C. sinensis* L. leaves, adjusted to 70% polyphenols) after 9 months or after 18 months on a follow-up study.^[37]

A synergistic effect was observed when combining both extracts on the LOO[•] inhibition. The combination of both extracts had a positive effect on the studied biological property. Thus, inhibition of LOO[•] can be achieved at low concentrations reducing the amounts of individual extracts needed, which can improve, for instance, the protection of membrane cells against oxidative injuries.

Different biological activities have been attributed to phenolic compounds, including the alpha-glucosidase and antioxidant activities studied here in. It is well established by different studies that the total phenolic composition of a matrix gives an idea of how rich this product is in antioxidants since these parameters are closely related. The total phenolic composition found for *C. sinensis* was similar to those found by other authors while for *E. uniflora* higher values were reported.^[38,39]

Although different studies suggest a correlation between the total amounts of phenolic compounds and the biological activity, this was not observed in the present study. Our result indicates that

probably, the type and amounts of individual phenolic compounds present in these matrices drives the inhibitory effect. *C. sinensis* is known to have high flavonoid content, primarily catechins such as (–)-epigallocatechin gallate, (–)-epigallocatechin, (–)-gallocatechin, and (+)-catechin.^[40] Conversely, *E. uniflora* leaves extract is poorly characterized with only two studies reporting its composition. One report of identification of myricetin and quercetin derivatives in a fraction obtained from ethanolic extract and other about macrocyclic hydrolysable tannin dimers (eugeniflorins D1 and D2), oenothin B, 1,2,4,6-tetra-O-galloyl-fl-o-glucose, gallocatechin, and myricitrin isolated from the leaves.^[16,41] Thus, further study is needed, such as bio-guided fractionation, to determine the phenolic compound or group of compounds, responsible for the observed biological activity. Nevertheless, the amounts of phenolic compounds provided by both extracts can be partially responsible for the alpha-glucosidase inhibition and can also offer good antioxidant protection for patients with T2DM.

Other important phytochemicals are carotenoids and chlorophylls; these compounds are colorful pigments abundant in fruits and vegetables. Carotenoids are considered important bioactive compounds for human's health as scientific studies demonstrate their important role in reducing the risk of degenerative diseases.^[42] Different studies have demonstrated significant decrease of plasma antioxidants by carotenoids (α- and γ-tocopherol, α and β-carotene, lycopene, β-cryptoxanthin, lutein, and zeaxanthin) in the progression of diabetes and its associated complications such as endothelial dysfunction and atherosclerosis.^[43–45] The amounts of carotenoids in *C. sinensis* was similar to those found in literature, and these are the first report about the amounts of carotenoids in *E. uniflora* leaves extracts.^[46] As far as we are concerned carotenoids have never been studied as inhibitors of alpha-glucosidase; however, as previously stated, they are important phytochemicals that could prevent the development of degenerative diseases. This fact is mainly because carotenoids are important antioxidant compounds, they have been reported as active toward peroxy radicals, a deleterious radical species that can be in the origin of different degenerative diseases, but with no activity against DPPH[•] radicals.^[47]

Regarding chlorophylls and their related compounds, they are among the best candidates for the chemicals responsible for the general protection afforded by vegetables. The main benefits of chlorophylls are their anticarcinogenic activity, related to their antioxidant effects; and their contribution to a positive hematological status due to the similarity between chlorophyll structure and hemoglobin. The total content of chlorophylls was proposed as quality parameter for *C. sinensis* and the concentration found among 14 samples ranged from 1.18 ± 0.16 to 1.98 ± 0.11 mg/g of sample.^[48] Thus, the value found in the present study was in agreement with the reported value. No report was found about the chlorophyll content in *E. uniflora* as far as we known. Although there are no studies reporting the inhibition of alpha-glucosidase by chlorophylls, different studies showed that natural matrices rich in these types of compounds have an important role in diabetes.^[49] The chlorophylls effect over DPPH[•] and LOO[•] have been reported.^[50] Chlorophyll A and B inhibited 40% of DPPH[•] at a concentration of 0.18 mg for both compounds. This value is lower than the concentrations of chlorophylls in both extracts [Table 2], indicating that these compounds can partially explain their DPPH[•] inhibition. The same behavior was also observed in the reduction on LOO[•] for both compounds, where chlorophyll A and B showed IC₅₀ values of 4.40 and 23.59 µg/g, respectively.^[50]

Taken together, these results indicates that the inhibitory effect on alpha-glucosidase and LOO[•] activities is likely mediated by the contribution of several or multiple bioactive compounds of the extracts, which is confirmed when extracts are combined, and a synergistic effect was observed.

CONCLUSION

The ethanolic extracts of *E. uniflora* and *C. sinensis* and their combination showed remarkable inhibitory activity over alpha-glucosidase, being 40–1500 times more effective than acarbose. A synergistic effect was found in the interaction of *E. uniflora* and *C. sinensis* over alpha-glucosidase and LOO[•]. These results may be partially explained by the presence and combination of phenolic compounds, chlorophylls A and B, and carotenoids. This study indicates that individual and combined extracts may be used for the treatment of diabetes mellitus, by inhibiting the enzyme, and also protecting the T2DM patients by improving their antioxidant status.

Financial support and sponsorship

Financial support of CNPq/ Science Without Borders Program project “Frutas Nativas do Brasil: potencial anti-hiperglicemiante e antioxidante. Juliana Vinholes thanks the Science without Borders Program (CNPq) for the Young Talent attraction fellowship.


Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Brazilian Diabetes Society (SBD). There are 13.4 million people with diabetes in Brazil; 2013. Available from: <http://www.diabetes.org.br/sala-noticias/2364-sao-134-milhoes-de-pessoas-portadoras-de-diabetes-no-brasilbet-es-no-Brasil>. [Last retrieved on 2013 Aug 10].
- Li DQ, Qian ZM, Li SP. Inhibition of three selected beverage extracts on alpha-glucosidase and rapid identification of their active compounds using HPLC-DAD-MS/MS and biochemical detection. *J Agric Food Chem* 2010;58:6608-13.
- Grabley S, Thiericke R. Bioactive agents from natural sources: Trends in discovery and application. *Adv Biochem Eng Biotechnol* 1999;64:101-54.
- Tadesse A, Hymete A, Bekhit AA, Mohammed SF. Quantification of total polyphenols, catechin, caffeine, L-theanine, determination of antioxidant activity and effect on antileishmanial drugs of ethiopian tea leaves extracts. *Pharmacognosy Res* 2015;7 Suppl 1:S7-14.
- Iso H, Date C, Wakai K, Fukui M, Tamakoshi A; JACC Study Group. The relationship between green tea and total caffeine intake and risk for self-reported type 2 diabetes among Japanese adults. *Ann Intern Med* 2006;144:554-62.
- Wu CH, Lu FH, Chang CS, Chang TC, Wang RH, Chang CJ. Relationship among habitual tea consumption, percent body fat, and body fat distribution. *Obes Res* 2003;11:1088-95.
- Bahmani M, Golshahi H, Saki K, Rafieian-Kopaei M, Delfan B, Mohammadi T. Medicinal plants and secondary metabolites for diabetes mellitus control. *Asian Pac J Trop Dis* 2014;4:S687-92.
- Sato T, Igarashi M, Yamada S, Takahashi N, Watanabe K. Inhibitory effect of black tea and its combination with acarbose on small intestinal a-glucosidase activity. *J Ethnopharmacol* 2015;161:147-55.
- Boros K, Jedlinszki N, Csupor D. Theanine and caffeine content of infusions prepared from commercial tea samples. *Pharmacogn Mag* 2016;12:75-9.
- Wang Y, Yang Z, Wei X. Sugar compositions, a-glucosidase inhibitory and amylase inhibitory activities of polysaccharides from leaves and flowers of *Camellia sinensis* obtained by different extraction methods. *Int J Biol Macromol* 2010;47:534-9.
- de Souza GC, Haas AP, von Poser GL, Schapoval EE, Elisabetsky E. Ethnopharmacological studies of antimicrobial remedies in the South of Brazil. *J Ethnopharmacol* 2004;90:135-43.
- Santos MM, Nunes MG, Martins RD. Empirical use of medicinal plants for diabetes treatment. *Rev Bras Planta Med* 2012;14:327-34.
- Trojan-Rodrigues M, Alves TL, Soares GL, Ritter MR. Plants used as antidiabetics in popular medicine in Rio Grande do Sul, Southern Brazil. *J Ethnopharmacol* 2012;139:155-63.
- Arai I, Amagaya S, Komatsu Y, Okada M, Hayashi T, Kasai M, et al. Improving effects of the extracts from *Eugenia uniflora* on hyperglycemia and hypertriglyceridemia in mice. *J Ethnopharmacol* 1999;68:307-14.
- Matsumura T, Kasai M, Hayashi T, Arisawa M, Momose Y, Arai I, et al. a-glucosidase inhibitors from paraguay natural medicine, ñangapiry, the leaves of *Eugenia uniflora*. *Pharm Biol* 2000;38:302-7.
- Mei-Hsien L, Saori N, Ling-Ling Y, Kun-Ying Y, Tsutomu H, Takashi Y, et al. Two macrocyclic hydrolysable tannin dimers from *Eugenia uniflora*. *Phytochemistry* 1997;44:1343-9.
- Gallucci MN, Oliva M, Asero C, Dambolena J, Luna A, Zygadlo J, et al. Antimicrobial combined action of terpenes against the food-borne microorganisms *Escherichia coli*, *Staphylococcus aureus* and *Bacillus cereus*. *Flavour Frag J* 2009;24:348-54.
- Ferreres F, Vinholes J, Gil-Izquierdo A, Valentão P, Gonçalves RF, Andrade PB. *In vitro* studies of a-glucosidase inhibitors and antiradical constituents of *Glandora diffusa* (Lag.) D.C. Thomas infusion. *Food Chem* 2013;136:1390-8.
- Vinholes J, Grosso C, Andrade PB, Gil-Izquierdo A, Valentão P, Pinho PG, et al. *In vitro* studies to assess the antidiabetic, anti-cholinesterase and antioxidant potential of *Spergularia rubra*. *Food Chem* 2011;129:454-62.
- Vinholes J, Gonçalves P, Martel F, Coimbra MA, Rocha SM. Assessment of the antioxidant and antiproliferative effects of sesquiterpenic compounds in *in vitro* Caco-2 cell models. *Food Chem* 2014;156:204-11.
- Grosso C, Vinholes J, Silva LR, de Pinho PG, Gonçalves RF, Valentão P, et al. Chemical composition and biological screening of *Capsella bursa-pastoris*. *Rev Bras Farmacogn* 2011;21:635-43.
- Pereira C, Calhelha RC, Barros L, Queiroz MJ, Ferreira IC. Synergisms in antioxidant and anti-hepatocellular carcinoma activities of artichoke, milk thistle and borututu syrups. *Ind Crops Prod* 2014;52:709-13.
- Nedamani ER, Mahoonak AS, Ghorbani M, Kashaninejad M. Antioxidant properties of individual vs. combined extracts of rosemary leaves and oak fruit. *J Agric Sci Technol* 2014;16:1575-86.
- Yeh P, Tschumi AI, Kishony R. Functional classification of drugs by properties of their pairwise interactions. *Nat Genet* 2006;38:489-94.
- Swain T, Hillis WE. The phenolic constituents of *Prunus domestica*. I. – The quantitative analysis of phenolic constituents. *J Sci Food Agric* 1959;10:63-8.
- Sumanta N, Haque CI, Nishika J, Suprakash R. Spectrophotometric analysis of chlorophylls and carotenoids from commonly grown fern species by using various extracting solvents. *Res J Chem Sci* 2014;4:63-9.
- Ramírez G, Zavala M, Pérez J, Zamilpa A. *In vitro* screening of medicinal plants used in Mexico as antidiabetics with glucosidase and lipase inhibitory activities. *Evid Based Complement Alternat Med* 2012;2012:701261.
- Gao J, Xu P, Wang Y, Wang Y, Hochstetter D. Combined effects of green tea extracts, green tea polyphenols or epigallocatechin gallate with acarbose on inhibition against a-amylase and a-glucosidase *in vitro*. *Molecules* 2013;18:11614-23.
- Oh J, Jo SH, Kim JS, Ha KS, Lee JY, Choi HY, et al. Selected tea and tea pomace extracts inhibit intestinal a-glucosidase activity *in vitro* and postprandial hyperglycemia *in vivo*. *Int J Mol Sci* 2015;16:8811-25.
- Ferreres F, Gil-Izquierdo A, Vinholes J, Silva ST, Valentão P, Andrade PB. *Bauhinia forficata* Link authenticity using flavonoids profile: Relation with their biological properties. *Food Chem* 2012;134:894-904.
- Kesavanarayanan KS, Sathiya S, Ranju V, Sunil AG, Ilavarasan R, Saravana Babu C, et al. *In vitro* cytotoxic, antioxidative and alpha-glucosidase inhibitory potential of a herbal mixture comprised of *Allium sativum* and *Lagerstroemia speciosa*. *Eur Rev Med Pharmacol Sci* 2012;16 Suppl 3:58-68.
- Schumacher NS, Colomeu TC, de Figueiredo D, Carvalho Vde C, Cazarin CB, Prado MA, et al. Identification and antioxidant activity of the extracts of *Eugenia uniflora* leaves. characterization of the anti-inflammatory properties of aqueous extract on diabetes expression in an experimental model of spontaneous type 1 diabetes (NOD Mice). *Antioxidants (Basel)* 2015;4:662-80.
- Tewari I, Sharma L, Lal Gupta G. Synergistic antioxidant activity of three medicinal plants *Hypericum perforatum*, *Bacopa monnieri*, and *Camellia Sinensis*. *Indo Am J Pharm Sci Res* 2014;4:2563-8.
- Seghrouchni I, Draï J, Bannier E, Rivière J, Calmard P, Garcia I, et al. Oxidative stress parameters in type I, type II and insulin-treated type 2 diabetes mellitus; insulin treatment efficiency. *Clin Chim Acta* 2002;321:89-96.
- Kade IJ, Ibukun EO, Nogueira CW, da Rocha JB. Sun-drying diminishes the antioxidative potentials of leaves of *Eugenia uniflora* against formation of thiobarbituric acid reactive substances induced in homogenates of rat brain and liver. *Exp Toxicol Pathol* 2008;60:365-71.
- Lassed S, Amrani A, Altun M, Zama D, Demirtas I, Benayache F, et al. *In vitro* antioxidant, inhibition of oxidative DNA damage and antiproliferative activities of ethanolic green tea (*Camellia sinensis*) extract. *Int J Pharm Sci Res* 2015;35:36-42.
- Spadiene A, Savickiene N, Ivanauskas L, Jakstas V, Skesters A, Silova A, et al. Antioxidant effects of *Camellia sinensis* L. extract in patients with type 2 diabetes. *J Food Drug Anal* 2014;22:505-11.
- Pereira VP, Knor FJ, Velloso JC, Beltrame FL. Determination of phenolic compounds and antioxidant activity of green, black and white teas of *Camellia sinensis* (L.) Kuntze, Theaceae. *Rev Bras Planta Med* 2014;16:490-8.

39. Santos RM, Oliveira MS, Ferri PH, Santos SC. Seasonal variation in the phenol content of *Eugenia uniflora* L. leaves. *Rev Bras Planta Med* 2011;13:85-9.
40. Khan N, Mukhtar H. Tea polyphenols for health promotion. *Life Sci* 2007;81:519-33.
41. Rattmann YD, De Souza LM, Malquevicz-Paiva SM, Dartora N, Sassaki GL, Gorin PA, *et al.* Analysis of flavonoids from *Eugenia uniflora* leaves and its protective effect against murine sepsis. *J Evid Based Complement Altern Med* 2012;2012:9.
42. Rao AV, Rao LG. Carotenoids and human health. *Pharmacol Res* 2007;55:207-16.
43. Polidori MC, Mecocci P, Stahl W, Parente B, Cecchetti R, Cherubini A, *et al.* Plasma levels of lipophilic antioxidants in very old patients with type 2 diabetes. *Diabetes Metab Res Rev* 2000;16:15-9.
44. Polidori MC, Stahl W, Eichler O, Niestroj I, Sies H. Profiles of antioxidants in human plasma. *Free Radic Biol Med* 2001;30:456-62.
45. Valabhji J, McColl AJ, Richmond W, Schachter M, Rubens MB, Elkeles RS. Total antioxidant status and coronary artery calcification in type 1 diabetes. *Diabetes Care* 2001;24:1608-13.
46. Kumar N, Rai R, Bera B. Impact of organic farming on the biochemical constituents and quality parameters of Darjeeling tea (*Camellia sinensis* (L) O Kuntze). *Two and a Bud* 2012;59:50-5.
47. Müller L, Fröhlich K, Böhm V. Comparative antioxidant activities of carotenoids measured by ferric reducing antioxidant power (FRAP), ABTS bleaching assay (α TEAC), DPPH assay and peroxy radical scavenging assay. *Food Chem* 2011;129:139-48.
48. Ošťádalová M, Tremlová B, Pokorná J, Král M. Chlorophyll as an indicator of green tea quality. *Acta Vet Brno* 2014;83:S103-9.
49. Bar-Sela G, Cohen M, Ben-Arye E, Epelbaum R. The medical use of wheatgrass: Review of the gap between basic and clinical applications. *Mini Rev Med Chem* 2015;15:1002-10.
50. Hsu C, Chao P, Hu S, Yang C. The Antioxidant and free radical scavenging activities of chlorophylls and pheophytins. *Food Nutr Sci* 2013;4:1-8.



Juliana Vinholes

ABOUT AUTHOR

Juliana Vinholes, Graduated in Chemistry from the Federal University of Pelotas (2002) and PhD in Chemistry from the University of Aveiro (2013). I have experience in analytical chemistry, focused on the development of chromatographic methods for primary and secondary metabolites analysis. In addition, I am also involved in the evaluation of biological potential of isolated compounds and natural matrices.