

Chemical characterization and antioxidant capacity of 'Cambui' (*Myrciaria tenella*) products

Ana Cecília Poloni Rybka¹, Aline Camarão Telles Biasoto Marques¹,
Ana Júlia Araújo², Rita de Cassia Mirela Resende Nassur^{3*}

¹Brazilian Agricultural Research Corporation, Petrolina, Brazil

²Federal Institute of Education, Science and Technology of the Sertão, Petrolina, Brazil

³University of the State of Bahia, Salvador, Brazil.

*Corresponding author, e-mail: ritarnassur@hotmail.com

Abstract

Very little information is found about the 'Cambui' products and the fruit is almost unexplored by the Brazilian agriculture or extractive activities. The present study aimed to characterize the physicochemical composition and antioxidant capacity of the juice, jam and fermented beverage obtained from 'Cambuí' fruits aiming to add value to the fruit and generate alternatives to Brazilian extractive population. These products were elaborated in small scale of production. The 'Cambuí' juice was prepared using the method of steam distillation and the jam was obtained from the residue of the juice with addition of sugar and citric acid. The 'Cambuí' fermented beverage was elaborated using procedures of winemaking. For the evaluation of the products quality, chemical analysis and evaluation of the antioxidant capacity were performed. All products presented high concentrations of the studied variables, when comparing to grape wines, fruit jams and grape juices. The 'Cambui' products can be considered sources of anthocyanins and other phenolic compounds, with high antioxidant potential. The use of 'Cambui' has potential for the development of juice, jam and fermented beverage, and these products can be an alternative to add value to fruit and generate income for populations that work with which income depend on extractive activities.

Keywords: fruit juice, fruit wine, jam

Introduction

The 'Cambui' tree (*Myrciaria tenella*) belongs to the *Myrtaceae* family, originating from South America (Sessa et al., 2016), South (Belem et al. 2016; Tonetti & Biondi, 2016; Cordero et al. 2016) and Southeast states of Brazil, it can also be found in the Northeast of the country (Silva et al., 2012), as in areas of the 'Chapada Diamantina', Bahia State. It is mainly used as an ornamental plant and its wood is used for fence posts, tools and firewood (Lorenzi 2000). Its fruit is named 'Cambui' in Brazil, with a rounded shape, with one to three seeds and color ranging from yellow, deep red and dark violet, depending on the maturity stage. Its fruits are glabrous and

bright globose berries, red or dark purplish when ripe (Lédo et al. 2014).

According to the few studies that evaluated the chemical composition 'Cambuí' it is possible to conclude that this fruit has a diameter between 0.5-1.0 cm, an average flesh weight of 0.60 to 0.15 g, pH of 3.3-4.8, total soluble solids around 10.5°Brix, ascorbic acid content of 110 mg.100g⁻¹, with significant amounts of lipids (>33%) and proteins (>7.5%) based on dried fruit (Pinheiro et al. 2011).

This fruit is able to produce essential oil rich in volatile compounds and Apel et al. (2010) identified 34 volatiles in the 'Cambuí' essential oil, with higher contents of β-caryophyllene

Received: 31 July 2017
Accepted: 11 June 2018

terpenes (25%) and spathulenol (10%). The anti-inflammatory effect of 'Cambui' essential oil was investigated by the authors 'in vitro' and 'in vivo' and the use of the essential oil significantly reduced the chemotaxis of neutrophils treated, with 93% of inhibition rate.

The evaluation regarding the chemical composition of fresh 'Cambui' reveal that it is an attractive option for human consumption, being necessary further studies about the species. This result probably shows potential for use in the agro-industry, both for the development of products by industrial or handmade scales, being an interesting alternative for familiar agriculture or small extractive growers and farmers. Currently, very little information is found about the 'Cambui' processing and this fruit is almost unexplored by the Brazilian agriculture or extractivism. In this sense, this study aimed to carry out the chemical characterization and evaluation of the antioxidant capacity of 'Cambui's' juice, jam and fermented alcoholic beverage.

Material and methods

Ninety kilograms (approximately 198 pounds) of 'Cambui' were harvest in the municipality of Morro do Chapéu - BA, Brazil, when the soluble solids content reached 11°Brix. The fruits were immediately transported to the Embrapa Semi-Arid, Petrolina - PE, Brazil, to proceed the development of the products. Two products were elaborated- the juice and the fermented beverage and a subproduct- the jam, which was obtained from the juice residue. For the products preparation, the methodology suggested and described by Marques et al. (2013) were followed. All elaborations were conducted in triplicate.

For the preparation of the fermented alcoholic beverage (wine), glass bottles with 20L of capacity adapted with Müller valve to control the entry of oxygen and allow the carbon dioxide output were used, by adding 0.01 g L⁻¹ of the preservative potassium metabisulfite and 30 mg.L⁻¹ of *Saccharomyces cerevisiae* commercial yeast, which was previously activated at 35°C for 40 minutes, and sucrose (36 g.L⁻¹) in the form of crystallized sugar to increase the alcohol content of the beverage.

Must preparation: for the obtainment of 10L of fermented beverage, 20Kg of healthy and mature cambui fruits were used, stored in a cold room at 8 ± 2 °C since harvest. Fruit were weighted, mashed and placed in a 20L container, adding 0,1g/L of potassium metabisulfite, avoiding enzymatic reactions and not desirable fermentation. For the alcoholic fermentation, the *Saccharomyces cerevisiae* yeast was added (0.35 g/L of the yeast in 10mL of water previously heated at 35 °C). Fermentation was carried out at a temperature of 22 ± 2°C during 20 days. At the same time with the alcoholic fermentation, the maceration was carried out, remaining fruits for 10 days in contact with the fermenting must to promote a better extraction of film compounds, seed and pulp. After fermentation, stabilization was performed in a cold (0°C) room for 30 days to improve the clarity and stability of the beverage. After this period, the fermented beverage was bottled and stored horizontally in a room with control of temperature (18°C), until analysis.

For the juice preparation, 10 kg (approximately 22 pounds) of fruits were used for each replicate, being previously sanitized with 200 ppm of active chlorine solution by immersion during 20 minutes. For extraction of the 'Cambui' juice, the stem extraction was used with the aid of a stainless steel juicer, being a simple technique with low cost and that can provide the preparation of pasteurized juice with very similar characteristics when compared to the fresh fruit. The extraction of juice was performed for 60 minutes at 75±5°C, then previously sterilized glass bottles (75°C) was filled by the hot juice. The product was cooled at a room with controlled temperature and stored in a horizontal position at the same condition (18°C), until analysis.

To obtain the jam, the residue of the juice preparation was used, with addition of 0.6 kg of sugar and 1.3 g of citric acid per kg (2.2 pounds) of the 'Cambui' residue. The mixture was heated at 90°C for 60 minutes and mixed constantly to prevent fouling. Then the product was passed through a sieve. The obtained jam was then bottled hot (85°C) in previously sterilized glass vials and cooled in a room with temperature control.

The physico-chemical analysis were performed one month after the preparation

of the products. For the evaluation of the titratable acidity ($\text{g}\cdot\text{L}^{-1}$) and the relative density, the procedures described by the Association of Official Analytical Chemists (AOAC 2016) were used. The hydrogenionic potential (pH) was directly determined in the sample with the aid of a pH meter (Tecnal, Tec-3MP model), previously calibrated. The alcohol content ($\%v/v$) was determined by hydrostatic balance (Model Super Gilbertini Alcomat), after distillation of the sample by an automatic Oenochemical Electronic Distilling (Gilbertini model Super E. D. E.). In the same hydrostatic scale, the dry matter content, density and soluble solids ($^{\circ}\text{Brix}$) were evaluated. The total monomeric anthocyanins were determined by the method described by Rizzon (2010), using a spectrophotometer Biomol, model SP-220. In the same spectrophotometer, it was determined the total polyphenol index (TPI) for the fermented beverage and the juice,

performing the direct reading of the samples at 280nm of absorbance (Harbertson & Spayd 2006), and the color intensity, adding up readings obtained at wavelengths of 420, 520 and 620 nm (Ough & Amerine 1988). The antioxidant activity was evaluated using the reagent 2,2-diphenyl-1-picrylhydrazyl (DPPH), based on the method of Brand-Williams et al. (1995) and adapted by Rybka (2010).

The test results were submitted to ANOVA and Tukey test ($p \leq 0.05$) using the statistical software Statistical Analysis System - SAS[®], version 9.3.1.

Results and Discussion

The obtained results for the chemical analysis and for the evaluation of the antioxidant capacity of the fermented beverage, juice and jam are presented on Tables 1 and 2, respectively.

The fermented beverage of 'Cambui'

Table 1. Results for the physico-chemical analysis performed on 'Cambui' products.

Product ¹	pH	Density g/cm^3	Total acidity ²	Total polyphenol index (IPT)	Total anthocyanins ³	Volatile acidity ⁴	Soluble solids ⁵	Color index ⁶
Beverage	3.3 ^b	1.032	31.6 ^a	166.3 ^b	250.9 ^b	1.1 ^a	-	61.5 ^c
Juice	3.4 ^a	-	17.6 ^b	175.3 ^a	311.7 ^a	0.5 ^b	10.9 ^b	79.6 ^a
Jam	3.2 ^c	-	12.4 ^c	175.0 ^a	120.3 ^c	-	79 ^a	63.1 ^b

¹ Products with a common letter in the same column are not statistically different, according to Tukey's test ($p \leq 0.05$). The variability coefficients (CV%) between all replicates for all analysis were lower than 5%. ² expressed as tartaric acid in $\text{g}\cdot 100\text{mL}^{-1}$ for the fermented beverage and the juice, and $\text{g}\cdot 100\text{g}^{-1}$ for the jam. ³ Monomeric anthocyanins expressed as malvidin-3-O-glucoside in $\text{mg}\cdot 100\text{mL}^{-1}$ for the fermented beverage and the juice, and $\text{mg}\cdot 100\text{g}^{-1}$ for the jam. ⁴ expressed as acetic acid in $\text{g}\cdot 100\text{mL}^{-1}$ for the fermented beverage and the juice and $\text{g}\cdot 100\text{g}^{-1}$ for the jam. ⁵ $^{\circ}\text{Brix}$. ⁶ sum for the lectures in the spectrophotometer at the 420, 520 e 620nm wavelengths.

Table 2. Results for the antioxidant capacity using the DPPH methodology for the 'Cambui' products.

Product ¹	Antioxidant Capacity ($\mu\text{g EAA}\cdot 100\text{mL}^{-1}$)
Fermented Beverage	1.055,15 ^c
Juice	1.083,54 ^a
Jam	1.081,18 ^b

¹ Products with the same letter in the column are not statistically different according to Tukey's test ($p \leq 0.05$). The variability coefficients (CV%) between all replicates for all analysis were lower than 5%. The antioxidant capacity is expressed in μmol of ascorbic acid equivalent (AAE) for 100mL of the fermented beverage and the juice and $\mu\text{mol EAA}\cdot 100\text{g}^{-1}$ for the jam.

presented reasonable alcohol content (11.3 $\%v/v$) and low pH (3.3), factors that contribute to a good chemical and microbiological stability of an alcoholic beverage. The drink also showed high anthocyanin (250.9 $\text{mg}\cdot 100\text{mL}^{-1}$) and total polyphenol (166.3) contents, and intense color (Table 1). Torres et al. (2013) found higher anthocyanin content in wines from Syrah, Cabernet Sauvignon (*Vitis vinifera* L.) and Petit Verdot (*Vitis vinifera* L.) varieties of 118, 106 and 123 $\text{mg}\cdot 100\text{mL}^{-1}$, respectively, values lower

than the observed for the 'Cambui' fermented beverage. Analyzed by the DPPH method, the fermented beverage also showed high antioxidant potential (1055.15 $\mu\text{mol EAA}\cdot 100\text{mL}^{-1}$), as can be seen in Table 1, possibly due to the high content of anthocyanins and polyphenols of the beverage. Picinelli Lobo et al. (2009) found in ciders an average of 290 $\mu\text{mol EAA}\cdot 100\text{mL}^{-1}$, being possible to conclude that the antioxidant capacity of the 'Cambui' fermented beverage is about three times higher than the cider.

The 'Cambui' juice presented pH value of 3.4, similar to that observed in grape juices evaluated by Lima et al. (2014) and Rizzon & Link (2006). The soluble solids content of the 'Cambui' juice was 10.9 °Brix, content that can also be compared to the value related by Rizzon & Link (2006), evaluating grape juices, that ranged between 12.2 and 13.1 °Brix. The color of the Cambuí juice was much more intense than the color of all grape juices evaluated by Lima et al. (2014), whose result is the sum of the reading absorbances at wavelengths of 420, 520 and 620 nm, ranged from 2.78 to 11.15, while the intensity of the 'Cambui' juice color was 79.6.

The 'Cambui' juice, similar to the fermented beverage, also presented high antioxidant potential, being observed values of 1083.54 $\mu\text{mol EAA.100mL}^{-1}$ (Table 2), standing out significantly ($p=0.05$) when compared to the content observed for the fermented beverage and the jam, possibly due to the higher concentration of total polyphenols (175.3) and anthocyanins (311.7mg.100mL⁻¹) of this product (Table 1). Zdunic et al. (2016) observed an increase in the phenolic content comparing pumpkin juice to the 'in natura' fruit, from 93.0 $\mu\text{g GAE/g}$ of pumpkin juice to 905.9 $\mu\text{g GAE/g}$ of fresh vegetable, demonstrating that the process can decrease the phenolic content in fruits and vegetables. The anthocyanins content of 'Cambui' juice was higher than that found by Lima et al. (2014) for grape juices produced with different grape varieties (BRS Cora, Magna BRS and BRS Violeta), whose maximum value observed was 46.4 mg.100mL⁻¹.

The 'Cambui' jam, although processed from the juice residue, presented antioxidant capacity of (1081.18 $\mu\text{mol EAA.100mL}^{-1}$), high content of total polyphenols (175.0) and high concentration of anthocyanins (120,3mg.100g⁻¹). The anthocyanins content of the 'Cambui' jam was similar to anthocyanin content quantified by Mota (2006) in jams elaborated with seven blackberry cultivars, which contents varied between 98.58 and 170.66 mg.100g⁻¹, being considered a product rich in phenolic compounds by the author.

Compared to other fruits, the anthocyanins content observed in 'Cambui'

juice and fermented beverage are similar to the contents of fresh 'jabuticaba' (314 mg.100g⁻¹) and superior to the anthocyanins content observed by Bagetti et al. (2011) in fresh purple, red and orange 'pitangas', which were respectively 136, 69 and 25 mg.100g⁻¹. Teixeira et al (2008) found a mean of 20 mg.100g⁻¹ of anthocyanins in strawberries, 'açai' and 'maria pretinha' and 13 mg.100g⁻¹ in fresh pomegranate. According to Araujo et al. (2016), the total anthocyanins content of myrtle fresh fruits can vary from 4.94 to 125.82 mg.100g⁻¹, depending on the maturity stage.

The processing of the 'Cambui' fruit can lead to a reduction in antocyanins content, but in the present study it was observed that the originated products presented contents of this compound similar to those found in fresh fruits and generally superior to many processed food products, such as wine and grape juice, beverages considered sources of phenolics and anthocyanins. Thus, 'Cambuí' products possibly have good nutraceutical potential, since it presented high anthocyanins and high antioxidant contents, being sources of other phenolic compounds that can be beneficial to health, due to the high content of total phenolic compounds found.

Conclusion

'Cambui' presented a good potential for the elaboration of the three studied products.

The 'Cambui' products can be considered sources of antocyanins and other phenolic compounds, with high antioxidant potential.

References

- AOAC International - *Association of official analytical chemists*, 2016. Official methods of analysis (20th Ed.). AOAC International, Rockville, MD, USA.
- Apel, M.A., Lima, M.E., Sobral, M., Young, M.C., Cordeiro, I., Schapoval, E.E., Henriques, A.T, Moreno, P.R. 2010. Anti-inflammatory activity of essential oil from leaves of *Myrciaria tenella* and *Calycorectes sellowianus*. *Pharmaceutical Biology*, 48 (4): 433-438.
- Araujo, D.R.D., Lucena, E.M.P., Gomes, J.P., Figueredo, R.M.F., Silva, C.P.D. 2016.

- Characterization of ripening stages on myrtle fruit. *Revista Brasileira de Fruticultura* 38(2): 1-10.
- Bagetti, M., Facco, E.M.P., Piccolo, J., Hirsch, G.E., Rodriguez-Amaya, D., Kobori, C.N., Vizzotto, M., Emanuell, T. 2011. Physicochemical characterization and antioxidant capacity of pitanga fruits (*Eugenia uniflora* L.) *Ciência e Tecnologia de Alimentos* 31 (1): 147-154.
- Belem, A.L.G., Branco, A.P.R. 2016. Análise florística e da distribuição espacial das espécies arbóreas do eixo de animação Arthur Bernardes, Curitiba-PR. *Revista Geonorte* 3(6): 583-592.
- Brand-Williams, W., Cuvelier M.E.; Berset C. 1995. Use of a free radical method to evaluate antioxidant activity. *Lebensmittel-Wissenschaft und-Technologie*, 28: 25-30.
- Cordero, R.L., Torchelsen, F.P., Overbeck, G.E., Anand, M. 2016. Invasive gorse (*Ulex europaeus*, Fabaceae) changes plant community structure in subtropical forest-grassland mosaics of southern Brazil. *Biological Invasions*. 18 (6): 1629-1643.
- Harbertson, J., Spayd, S. 2006. Measuring phenolics in the winery. *American Journal Enological and Viticultural* 57: 280-288.
- Lédo, A.S., Barin, L.B., Da Silva, A.V.C., de Sa, F.P., Machado, C. de A. 2014. In vitro germination and acclimatization of cambui tree type seedlings. *Ciência Rural* 44(8): 1355-1360
- Lima, M.S., Silani I.M.T., Corrêa, L.C., Biasoto, A.C.T., Pereira, G.E., Bordignon-Luiz, M.T.; Ninow, J.I. 2014. Phenolic compounds, organic acids and antioxidant activity of grape juices produced from new Brazilian varieties planted in the Northeast Region of Brazil. *Food Chemistry* 161: 94-103.
- Lorenzi, H. 2000. *Árvores Brasileiras*. 3. ed. São Paulo: Plantarum, v.1, 264 p.
- Marques, A.T.B., Rybka, A.C.P., ARAÚJO, A.J.B. 2013. *Suco, geleia e bebida alcoólica fermentada derivados de cambuí*. Petrolina-PE: Empresa Brasileira de Pesquisa Agropecuária *Comunicado Técnico* 156.
- Mota, R.V. 2006. Caracterização física e química de geleia de amora-preta. *Ciência e Tecnologia de Alimentos* 26: 539-543.
- Ough, C.S., Amerine, M.A. 1988. *Methods for analysis of musts and wines* (2 ed.). New York: John Wiley and Sons, Inc.
- Picinelli Lobo A., Diniero-Garcia, Y., Mangas-Sanchez, J., Rodriguez-
- Composition and Analysis* 22: 644-648.
- Pinheiro, L.R., Almeida, C.S., Silva, A.V.C. 2011. Diversidade genética de uma população natural de cambuizeiro e avaliação pós-colheita de seus frutos. *Scientia Plena* 7(6): 1-5.
- Rizzon, L. A.; Link, M. 2006. Composição do suco de uva caseiro de diferentes cultivares. *Ciência Rural*, 36: 689-692.
- Rizzon, L.A. *Metodologia para análise de vinho*. Empresa Brasileira de Pesquisa Agropecuária-Embrapa Uva e Vinho. Bento Gonçalves, 2010.
- Rybka, A.C.P. 2010. *Folatos, capacidade antioxidante e trans-2-nonenal em cerveja brasileira*. 184p. (PhD Dissertation). Universidade Estadual de Campinas. Departamento de Ciência de Alimentos, Campinas.
- Sessa, L., Abreo, E., Bettucci, L., Lupo, S. 2016. Botryosphaeriaceae species associated with wood diseases of stone and pome fruits trees: symptoms and virulence across different hosts in Uruguay. *European Journal of Plant Pathology* 1-12.
- Silva, A.V.C. da., Rabbani, A.R.C., Costa, T.S., Clivati, D. 2012. Fruit and seed biometry of cambuí (*Myrciaria tenella* O. Berg) . *Agro@ambiente On-line* 6 (3): 258-262.
- Teixeira L.N., Stringheta P.S., Oliveira F.A. 2008. Comparação de métodos para quantificação de antocianinas. *Revista Ceres* 55 (4): 297-304.
- Tonetti, A., Biondi, D. 2016. Dieta de capivara (*Hydrochoerus hydrochaeris*, L., 1766) em ambiente urbano, parque municipal do Tinguí, Curitiba-PR. *Acta Veterinaria Brasilica* 9(4): 316-326.
- Torres, A.P., Oliveira J.B., Berron L., Vaz J., Gomes O., Menezes T.R., Morais L.R.V., Silva P.F., Biasoto A.C.T., Pereira G.E. 2013. Characterization of wines from a new region of altitude in the Northeast of Brazil. *Ciência e Técnica Vitivinícola* 28: 260-265.
- Zdunić, G. M., Menković, N. R., Jadranin, M. B., Novaković, M. M., Šavikin, K. P., & Živković, J. Č. 2016. Phenolic compounds and carotenoids in pumpkin fruit and related traditional products. *Hemijska industrija* 00: 49-49.