

## 'UENF 506-11': a new maize cultivar for the North and Northwest of Rio de Janeiro State

Messias Gonzaga Pereira<sup>1</sup>, Ana Paula Candido Gabriel Berilli<sup>2</sup>, Roberto dos Santos Trindade<sup>3</sup>, Geovana Cremonini Entringer<sup>1</sup>, Pedro Henrique Araújo Diniz Santos<sup>1</sup>, Julio Cesar Fiorio Vettorazzi<sup>1\*</sup> and Keila Silva da Cunha Galvão<sup>4</sup>

Crop Breeding and Applied Biotechnology  
19: 141-144, 2019  
Brazilian Society of Plant Breeding.  
Printed in Brazil  
<http://dx.doi.org/10.1590/1984-70332019v19n1c20>

**Abstract:** *The intervarietal hybrid 'UENF 506-11' was derived from full-sib maize families in the 11<sup>th</sup> cycle of reciprocal recurrent selection. It produces a grain yield of 7.57 tons ha<sup>-1</sup>, and is recommended for cultivation in the North and Northwest regions of the Rio de Janeiro State.*

**Keywords:** *Zea mays L., reciprocal recurrent selection, cultivar, plant breeding.*


### INTRODUCTION

Maize (*Zea mays* L.) is a cereal of great economic and social importance, cultivated in almost all agricultural regions of the world and used in human and animal nutrition and, more recently, as a source of biofuel. Brazil is the third largest producer in the world, with a production in the year 2017 of 97 million tons, which is surpassed only by the United States and China (FAOSTAT, 2019). Much of the yield potential of this crop in the country is the result of significant advances in breeding programs, which have progressively improved agronomic traits related to grain yield, disease resistance and adaptability to adverse conditions.

For the state of Rio de Janeiro, no agricultural zoning indicates which cultivars are best planted. Maize is grown almost exclusively by small farmers with low technological input, generally without irrigation, for grain and silage production. Mostly, the farmers grow open-pollinated varieties and hybrids developed for other regions of Brazil.

Historically, until the 1980s, the North and Northwest of Rio de Janeiro stood out as one of the largest sugarcane-producing regions of Brazil (Silva 2006). However, dwindling yields caused by a decline in natural soil fertility due to successive cultivation without nutrient replenishment and increased production costs, led to a decline in sugarcane farming in the region (Castro and Ribeiro 2010). In this context, considering the edaphoclimatic conditions of the North and Northwest of Rio de Janeiro, with an undulating plain relief, appropriate for a mechanized crop management, and for maize cultivation destined for grain as well as silage for livestock, the potential of maize is perceived as an important alternative of crop diversification for the region.

In 1996, the Universidade Estadual do Norte Fluminense Darcy Ribeiro (UENF) initiated a maize breeding program based on Full-sib Reciprocal

**\*Corresponding author:**  
E-mail: [juliocesar.f.v@hotmail.com](mailto:juliocesar.f.v@hotmail.com)  
 ORCID: 0000-0002-3887-7735

**Received:** 02 October 2017  
**Accepted:** 05 November 2018

<sup>1</sup> Universidade Estadual do Norte Fluminense Darcy Ribeiro, 28.013-602, Campos dos Goytacazes, RJ, Brazil

<sup>2</sup> Instituto Federal do Espírito Santo, 29.070-910, Colatina, ES, Brazil

<sup>3</sup> Embrapa Milho e Sorgo, Núcleo de Recursos Genéticos e Geração de Cultivares (NRGC), Embrapa Milho e Sorgo, Rodovia MG 424, km 65, CEP: 35.701-970, Sete Lagoas, MG, Brazil.

<sup>4</sup> Instituto de Ensino Superior do Espírito Santo, 29.360-000, Castelo, ES, Brazil

Recurrent Selection (FSRRS). The continuous progress of this breeding program has provided maize producers in the North and Northwest regions of Rio de Janeiro with hybrids with excellent agronomic response and adapted to the regional conditions. In 2000, this program released the inter-populational hybrid 'UENF 506-6' with superior agronomic performance over other cultivars, which has been successfully used by farmers (Tardin et al. 2007, Berilli et al. 2013).

The objective of this study was to describe the main traits of the intervarietal maize hybrid 'UENF 506-11', of the 11<sup>th</sup> cycle of FSRRS.

### Pedigree and Breeding Method

Two maize populations of different heterotic groups were used, one called CIMMYT and the other Piranão, of the heterotic groups "flint" and "dent", respectively. Both populations have the brachytic gene ( $br_2$ ), introduced by backcrossing in a pre-breeding stage, resulting in short plant height. Both populations came from the Universidade Federal de Viçosa (UFV), where they undergone five cycles of FSRRS. As of the sixth cycle, selection was carried out at the UENF.

The FSRRS was described by Hallauer and Eberhart (1970). This methodology promotes the accumulation of favorable alleles within the involved populations and increases the combining ability among them, preserving the genetic variability among and within them (Reis et al. 2009).

### Field trials

Field evaluations were performed from September to February in the growing seasons of 2009/2010, 2010/2011 and 2011/2012 at three locations: two in Campos dos Goytacazes – RJ and one in Itaocara – RJ, totaling nine environments. These nine constitute the environments where the VCU (Value for Cultivation and Use) trials were carried out. In all trials, a randomized complete block design with five replications was used. The experimental plots consisted of a single 4.0 m row, at a plant spacing of 0.2 m between plants and 1.0 m between rows.

Twenty-one genotypes of the FSRRS program of the UENF, seven of which were intervarietal hybrids between the CIMMYT and Piranão populations and the respective parent populations, were evaluated. In addition, seven control genotypes were evaluated: two commercial hybrids ('AG 1051' and 'BM 2202'), an intervarietal hybrid ('Fortaleza' versus CIMMYT 8), three commercial varieties ('BR 106', 'Fortaleza' and 'Sol da Manhã'), and the Composto Caboclo  $br_2$ . For the sake of simplicity, only the results of the registered intervarietal hybrid ('UENF 506-11') and the controls used in the experiment were presented (Table 1).

The crop management at the different locations and in all trials was applied as recommended for maize (Fancelli and Dourado Neto 2000). During the experimental period, seven maize descriptors and nine morpho-agronomic traits were evaluated, according to the minimum requirements for registry in the national cultivar registration system (MAPA 2019). The plots were harvested by hand and the grain yield was weighed considering a grain moisture content of 13%.

**Table 1.** Summary of analysis of variance with the respective mean squares, coefficients of experimental variation (CV), means of the evaluated traits and coefficient of genotypic determination ( $H^2$ ), between 2009 and 2012 in three environments.

SV	df	Mean squares								
		STAND	NLP	DFL	NE	EW	GW	HW	W100	GY
B/E	36	7.33	5.89	5.39	27.93	0.88	0.64	825.97	664.53	3.21
Block (B)	4	12.5	8.18	13.79	44.14	0.90	1.20	1213.34	320.78	5.99
B X E	32	6.68	5.6	4.34	25.91	0.88	0.57	777.55	707.50	2.86
Genotype (G)	4	285.19**	113.49**	155.44**	576.64**	36.5**	33.14**	29211.8**	12463.20**	163.74**
Environment (E)	8	97.47**	76.90**	424.55**	220.75**	13.26**	9.98**	5252.86**	11634.98**	49.34**
G X E	17	85.92**	12.16	13.02	127.56**	1.89	0.83*	2063.12	1616.72**	4.11*
Error	71	14.10	8.8	8.03	39.34	1.29	0.47	1378.66	556.26	2.35
Mean		23.23	2.4	62.99	25.27	3.61	3.02	744.29	304.34	6.72
CV (%)		16.16	123.59	4.49	24.81	31.43	22.77	4.9	7.75	22.78
$H^2$		0.95	0.92	0.95	0.93	0.96	0.98	0.95	0.95	0.98

STAND: Number of plants per experimental unit; NLP: Number of lodged plants; DFL: Days until flowering; NE: Number of ears; EW: Ear weight; GW: Grain weight; HW: Hectoliter weight; W1000: 1000-grain weight; GY: Grain yield (tons ha<sup>-1</sup>). \* and \*\* Significant at 5 and 1% by the F test, respectively.

## Statistical Analyses

An individual ANOVA of the morpho-agronomic traits was performed for each environment (data not shown). The coefficients of variation for all environments were found to be lower than 20%, so the homogeneity of the residual variances was evaluated by the F test (Hartley 1950). The combined analysis of experiments was performed since the ratio between the highest and the lowest residual F for the evaluated environments was less than seven (Table 1). Tukey mean test at 5% probability was applied based on the mean of the tested environments. Software GENES (Cruz 2013) was used for all statistical analyses. For the genotype x environment interaction for the trait grain yield, significance at 5% probability was observed, but since these environments were close together and in view of the excellent performance of hybrid 'UENF 506-11' in the different environments, there was no need to register different hybrids for these regions. Thus, the overall means were considered for the mean test comparisons.

## Agronomic Performance

In three growing seasons at three locations the grain yield of the inter-population hybrid 'UENF 506-11' was only surpassed by hybrid 'AG1051' (Table 2). The Tukey mean test indicated that the intervarietal hybrid 'UENF 506-11' had a similar grain yield to the commercial hybrid 'BM 2202' and higher than that of the commercial varieties 'BR 106' and 'Sol da Manhã', and was only lower than that of the commercial hybrid 'AG 1051'.

**Table 2.** Yield of 'UENF 506-11' compared to the mean of four controls by Tukey test ( $P>0.05$ ) in evaluations between 2009 and 2012 in three environments

Cultivars	Grain yield (tons ha <sup>-1</sup> )
AG 1051	8.86 a
'UENF 506-11'	7.57 b
BM 2202	7.23 b
BR 106	6.14 c
SOL DA MANHÃ	3.78 d
MSD	0.69

MSD = Minimal Significant Difference.

**Table 3.** Descriptive statistics for eight agronomic traits of three maize genotypes evaluated in tests of value for cultivation and use (VCU), in the growing seasons of 2009, 2010 and 2012, at three locations

	Morpho-agronomics traits				
	Plant height	Ear height	Plant density	Ear index	
'UENF 506-11'	Mean	2.07	1.27	24	1.2
	Upper limit	2.63	2.25	30	1.6
	Lower limit	1.44	0.80	17	0.4
		<b>Female flowering</b>	<b>Male flowering</b>	<b>1000-grain weight (g)</b>	<b>Grain yield (t ha<sup>-1</sup>)</b>
	Mean	64	64	314	7.57
	Upper limit	71	75	380	11.56
	Lower limit	59	58	250	1.78
		<b>Plant height</b>	<b>Ear height</b>	<b>Plant density</b>	<b>Ear index</b>
	Mean	1.85	0.99	25	1.1
'BM 2202'	Upper limit	2.31	1.33	41	1.5
	Lower limit	1.41	0.54	18	0.5
		<b>Female flowering</b>	<b>Male flowering</b>	<b>1000-grain weight (g)</b>	<b>Grain yield (t ha<sup>-1</sup>)</b>
	Mean	59	60	296	7.24
	Upper limit	66	67	380	11.78
	Lower limit	53	54	210	0.57
		<b>Plant height</b>	<b>Ear height</b>	<b>Plant density</b>	<b>Ear index</b>
	Mean	1.94	1.07	24	1.1
	'BR 106'	Upper limit	2.38	1.80	29
Lower limit		1.25	0.65	20	0.3
		<b>Female flowering</b>	<b>Male flowering</b>	<b>1000-grain weight (g)</b>	<b>Grain yield (t ha<sup>-1</sup>)</b>
Mean		59	61	304	6.15
Upper limit		62	65	360	11.56
Lower limit		53	54	248	1.08

## Other Traits

The main traits of hybrid 'UENF 506-11' were compared to the other cultivars used in the region and evaluated in the experiment (Table 3). Hybrid 'UENF 506-11' is appropriate for grain or silage production or for fresh consumption. The grain is classified as semi-hard and yellow/orange. The recommended cultivation period for the intervarietal hybrid is between September and February.

## Seed Production and Distribution

The intervarietal hybrid 'UENF 506-11' was registered by the Ministry of Agriculture, Livestock and Supply – MAPA (no. 30595). Seeds are produced and commercialized by Rio Norte Sementes Company in partnership with UENF.

## REFERENCES

- Berilli APCG, Pereira MG, Trindade RS and Costa FR (2013) Response to the selection in the 11<sup>th</sup> cycle of reciprocal recurrent selection among full-sib families of maize. *Acta Scientiarum* **35**: 435-441.
- Castro RAR and Ribeiro AC (2010) Relações de cooperação no aglomerado sucroalcooleiro fluminense: reflexões sobre o papel do recurso intangível no desenvolvimento regional. *Agenda Social* **4**: 59-75.
- Cruz CD (2013) Genes: a software package for analysis in experimental statistics and quantitative genetics. *Acta Scientiarum Agronomy* **35**: 271-276.
- Fancelli AL and Dourado Neto D (2000) *Produção de milho*. Agropecuária, Guaíba, 360p.
- FAOSTAT (2019) Food and Agriculture organization of the United Nations. Production quantities of Maize by country 2017. Available at: <<http://www.fao.org/faostat/en/#data/QC/visualize>>. Access in Marh, 2019.
- Hallauer AR and Eberhart SA (1970) Reciprocal full-sib selection. *Crop Science* **10**: 315-316.
- Hartley HO (1950) The maximum F-ratio as a short-cut teste for heterogeneity of variances. *Biometrika* **37**: 308-312.
- MAPA - Ministério da Agricultura, Pecuária e Abastecimento (2019). Valor de cultivo e uso. Available at: <<http://sistemas.agricultura.gov.br/snpc/cultivarweb/index.php>>. Access in March, 2019.
- Reis MC, Souza JC, Ramalho MAP, Guedes FL and Santos PHAD (2009) Progresso genético com a seleção recorrente recíproca para híbridos interpopulacionais de milho. *Pesquisa Agropecuária Brasileira* **44**: 1667-1672.
- Silva LC (2006) Reconfiguração do Norte Fluminense a partir dos anos 70: a chegada do petróleo e suas consequências na dinâmica de crescimento regional. *Revista Plurais* **1**: 167-187.
- Tardin FD, Pereira MG, Gabriel APC, Amaral Junior AT and Filho GAS (2007) Selection index and molecular markers in reciprocal recurrent selection in maize. *Crop Breeding and Applied Biotechnology* **7**: 225-233.