



Asian Pacific Journal of Tropical Biomedicine

Original Article



doi: 10.4103/2221–1691.259003

www.apjtb.org

Chemical composition, antiparasitic and cytotoxic activities of aqueous extracts of *Ziziphus joazeiro* Mart.

Jacqueline Cosmo Andrade^{1,2}, Ana Raquel Pereira da Silva¹, Antônia Thassy Lucas dos Santos¹, Maria Audilene Freitas¹, Yedda Maria Lobo Soares de Matos¹, Maria Flaviana Bezerra Moraes Braga¹, Camila Fonseca Bezerra¹, Maria Isabeli Pereira Gonçalo², Maria Celeste Vega Gomez³, Míriam Rolóm³, Cathia Coronel³, Paulo Riceli Vasconcelos Ribeiro⁴, Edy Sousa de Brito⁴, Henrique Douglas Melo Coutinho^{1✉}

¹Microbiology and Molecular Biology Laboratory, Biological Chemistry Department, Regional University of Cariri – URCA, Crato (CE), Brazil

²Bioassays Laboratory – LABIO, Federal University of Cariri – UFCA, Brejo Santo University campus (CE), Brazil

³Center for the Development of Scientific Research, Moisés Bertoni Foundation/Díaz Gill Laboratories, Asunción, Paraguay

⁴Natural Products Chemistry Multiuser Laboratory LMQPN, Embrapa Tropical Agroindustry, Fortaleza (CE), Brazil

ARTICLE INFO

Article history:

Received 17 January 2019

Revision 16 February 2019

Accepted 8 May 2019

Available online 28 May 2019

Keywords:

Antiepipmastigote

Antipromastigote

UPLC-MS-ESI-QTOF

ABSTRACT

Objective: To compare the *in vitro* antiparasitic activity of aqueous extracts from *Ziziphus joazeiro* leaves and stem bark against *Trypanosoma cruzi*, *Leishmania braziliensis*, and *Leishmania infantum*, as well as to evaluate its cytotoxicity in mammalian cells, in addition to identifying the chemical composition of the extracts.

Methods: *Ziziphus joazeiro* leaf and stem bark aqueous extracts were prepared by cold extraction maceration and subjected to ultra-efficient liquid chromatography coupled to a quadrupole/time of flight system. The susceptibility assays used *Trypanosoma cruzi* CL-B5 strains and promastigote forms of *Leishmania braziliensis* and *Leishmania infantum* for antiparasitic activity of the extracts. Moreover, mammalian fibroblasts NCTC clone 929 were used for cytotoxicity analysis.

Results: Terpenoid compounds, flavonoids and phenolic acid were identified in extracts. The stem bark aqueous extracts presented more significant results in terms of antiparasitic activity compared with the leaf aqueous extracts, especially against *Leishmania braziliensis* and *Leishmania infantum* promastigote forms with an $IC_{50} < 500 \mu\text{g/mL}$. The cytotoxicity evaluation showed moderate toxicity of the stem bark aqueous extracts, which is relevant information for the rational use of this plant part since it is widely used by the population.

Conclusions: These preliminary results may contribute to the formulation of new therapeutic agents against this group of neglected diseases, so further investigations are required to delineate the mechanisms of action mainly of the aqueous extract of stem bark of *Ziziphus joazeiro*.

1. Introduction

Infectious diseases termed as neglected diseases are caused by parasitic or infectious agents which mainly affect populations living in conditions of poverty and social inequality. The World Health Organization (WHO) emphasizes that housing, food, poor sanitation

and lack of health care are the main causes of these diseases[1,2].

Leishmaniasis, malaria, dengue, Chagas disease, leprosy and tuberculosis can be highlighted as examples of neglected diseases[3].

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2019 Asian Pacific Journal of Tropical Biomedicine Produced by Wolters Kluwer-Medknow. All rights reserved.

How to cite this article: Andrade JC, da Silva ARP, dos Santos ATL, Freitas MA, de Matos YMLS, Braga MFBM, et al. Chemical composition, antiparasitic and cytotoxic activities of aqueous extracts of *Ziziphus joazeiro* Mart. Asian Pac J Trop Biomed 2019; 9(5): 222-226.

✉Corresponding author: Henrique Douglas Melo Coutinho, Microbiology and Molecular Biology Laboratory, Biological Chemistry Department, Regional University of Cariri – URCA, Crato-CE, Brazil.

Tel: +55(88) 31021212

Fax: +55(88) 31021291

E-mail: hdmcoutinho@gmail.com

Chagas disease is a parasitic infection caused by the flagellate protozoan *Trypanosoma cruzi* (*T. cruzi*) and transmitted through vectors. Chagas disease is a systemic and chronic disease considered to be an endemic tropical disease in 21 countries in Latin America where its main transmission mechanism occurs through triatomine vectors. Secondary transmission mechanisms such as transfusion (through blood transfusion or infected organ transplantation), congenital (mother-child transmission, if the woman has the disease) and oral (through ingestion of foods infected with insect feces) transmission exist. During the *T. cruzi* biological cycle, three evolutionary forms (trypomastigote, epimastigote and amastigote) can be observed[4,5]. According to the WHO, there are approximately 8 million people infected, with 56 thousand new cases per year and 12 thousand annual deaths[6].

Leishmaniasis is characterized as a group of diseases caused by protozoa from more than 20 *Leishmania* species where the parasites have two life cycle stages: promastigote and amastigote. Transmission of these parasites is achieved through a phlebotomine vector[7,8]. There are three main types of leishmaniasis: cutaneous, which causes ulcers on exposed body parts (face, arms and legs); visceral, one of the most severe forms, is characterized by fever, weight loss, spleen and liver enlargement, followed by anemia; and mucocutaneous, where the lesions cause destruction of the nose, mouth, throat and surrounding mucous tissue membranes[9,10]. The WHO noted that in 2014 more than 90% of new cases occurred in six countries: Brazil, Ethiopia, India, Somalia, South Sudan and Sudan[11].

Neglected diseases suffer from a therapeutic disadvantage as they do not attract the interest of pharmaceutical industries[12]. Chagas' disease treatment is achieved by only two drugs, nifurtimox and benznidazole. For leishmaniasis, pentavalent antimonials, amphotericin B and pentamidine are used. In general, these drugs have a high parasite resistance index, often requiring high administration doses, which generate considerable toxicity[13,14]. Thus, the importance of finding new compounds which act as chemotherapeutic agents for the treatment of these diseases is evident.

Moreover, natural products derived from various plant parts can be used as agents for the treatment of infections, mainly due to the bioactive potential of secondary metabolites present in their composition[15]. *Ziziphus joazeiro* (*Z. joazeiro*) Mart. (Rhamnaceae) is a tree species widely used in ethnomedicine with several proven biological activities such as antifungal, antibacterial, antioxidant, antipyretic, anti-inflammatory and astringent activity[16,17].

The objective of this work was to evaluate the *in vitro* antiparasitic activity of the *Z. joazeiro* aqueous extracts derived from its leaves and stem bark against *T. cruzi*, *Leishmania braziliensis* (*L. braziliensis*), and *Leishmania infantum* (*L. infantum*), as well as its cytotoxic potential in mammalian cells, in addition to identifying the chemical composition of the extracts through ultra-efficient liquid chromatography coupled to quadrupole/time of flight system (UPLC-MS-ESI-QTOF).

2. Materials and methods

2.1. Collection area and plant material

The leaves and stem bark were collected from eight *Z. joazeiro* Mart. specimens located in the Sítio Ipueiras, in the rural area of the Brejo Santo municipality, south of Ceará, Brazil, at the foot of the Chapada do Araripe (geographical coordinates, south latitude and west longitude of Greenwich: 1: 442 m, 07°28'54.4"S/39°01'47.2"W; 2: 431 m, 07°28'53.3"S/39°01'46.1"W; 3: 436 m, 07°28'50.5"S/39°01'57.6"W; 4: 440 m, 07°28'42.8"S/39°02'10.2"W; 5: 447 m, 07°28'48.5"S/39°02'12.0"W; 6: 441 m, 07°28'51.4"S/39°02'16.0"W; 7: 439 m, 07°28'54.6"S/39°02'07.6"W; 8: 436 m, 07°28'58.6"S/39°01'48.8"W). The voucher material was deposited in the Herbarium Dárdano de Andrade Lima of the Regional University of Cariri - URCA under n° 13.346 and identified as *Z. joazeiro* Mart. The collection took place in the month of February 2017, from 7:30 to 9:00 in the morning. The plant material was sent to the laboratory, cleaned and weighed.

2.2. Extract preparation

The aqueous extracts from the leaves and stem bark (AEL and AEB) of *Z. joazeiro* were prepared by cold extraction maceration[18]. Fresh leaves were cut to increase their surface area, while the stem barks were dried at room temperature and ground in a mechanical mill. Subsequently, both were added in distilled sterile water and maintained in a container protected from light and air. After 72 h, the extracts were filtered, frozen and taken to a lyophilizer (-60 °C) producing a crude extract of 39.9 g and 111.58 g, respectively.

2.3. Compound identification through UPLC-MS-ESI-QTOF

The analysis was performed on a Waters ACQUITY UPLC system to the Q-TOF Premier mass spectrometer (Waters MS Technologies, Manchester, UK) with electrospray ionization interface (ESI) in negative ionization mode. Chromatographs were run on a Waters Acquity UPLC BEH column (150 mm × 2.1 mm, 1.7 μm), fixed temperature of 40 °C, mobile water phases with 0.1% formic acid (A) and acetonitrile, 1% formic acid (B), gradient ranging from 2% to 95% B (15 min), flow 0.4 mL/min and injection volume 5 μL. The negative ESI mode was acquired in the range of 110-1 180 Da, fixed source temperature at 120 °C, desolvation temperature 350 °C, desolvation gas flow of 500 L/h, extraction cone 0.5 V, voltage capillary of 2.6 kV. Leucine-enkephalin was used as a lock mass. The acquisition mode was MS^E. MS data were collected for *m/z* values in the range of 110-1 180 Da with a scan time of 0.1 over an analysis time of 19 min. The accurate mass and molecular formula assignments were obtained with the MassLynx 4.1 software (Waters MS Technologies).

2.4. Cell lines

The *T. cruzi* assays used CL-B5 parasite strains (clone CL-B5) transfected with the β -galactosidase gene of *Escherichia coli* (LacZ). Epimastigote forms were cultured in infusions of liver tryptase with 10% fetal bovine serum, penicillin and streptomycin at 28 °C, being harvested during the exponential growth phase[19]. For the leishmanicidal activity, promastigotes of *L. braziliensis* and *L. infantum* grown at 26 °C were used in Schneider's medium, supplemented with 10% fetal bovine serum, 2% normal human urine plus penicillin and streptomycin. In the NCTC clone 929 mammalian fibroblast cytotoxicity test, they were cultured in RPMI 1640 medium (Sigma) supplemented with 10% fetal bovine serum, penicillin and streptomycin. Cells in the pre-confluence phase were harvested with trypsin, maintained at 37 °C in a humidified atmosphere of 5% CO₂.

2.5. In vitro trypanocidal and leishmanicidal assays

The trypanocidal assay was performed on 96-well microdilution plates with cultures that did not reach the stationary phase[20]. Epimastigotes were seen (1×10^5 cells/well) on 200 mL of RPMI médium and incubated with the products at 28 °C for 72 h. Subsequently, 50 μ L of chlorophenol red- β -D-galactopyranoside solution was added, incubated at 37 °C for an additional 6 h and then read at 595 nm in a spectrophotometer. Nifurtimox was used as reference drug. Each concentration was tested in triplicate. The percentage of inhibition (%AE) was calculated.

The leishmanicidal assay was based on the method developed by Mikus and Steverding[21] with modifications. Promastigotes (2.5×10^5 parasites/well) were grown in 96-well plastic plates. The extracts were dissolved in dimethyl sulfoxide (DMSO), and different dilutions of up to 200 μ L of final volume were added. After 48 h at 26 °C, 20 μ L of resazurin solution was added and the oxidation-reduction was quantified at 570 to 595 nm. Each concentration was tested in triplicate. In each test, Metronidazole, the reference drug, was used as a control. The antipromastigote percentages (%AP) were calculated.

2.6. Cytotoxicity assays

The method for assessing cell viability was colorimetric, with resazurine[22]. Fibroblasts NCTC 929 were seeded (5×10^4 cells/well) in 96-well microdilution plates with 100 μ L RPMI 1640 medium for 24 h at 37 °C in 5% CO₂ atmosphere. The medium was replaced by different concentrations of the extracts in 200 μ L of medium and then incubated for another 24 h. Growth controls were also included. Subsequently, a 20 μ L volume of 2 mM resazurin solution was added and the plates incubated for 3 h to evaluate cell viability. Each concentration was tested in triplicate. The cytotoxicity of each compound was estimated

by calculating the percentage of cytotoxicity (%C).

2.7. Statistical analysis

The IC₅₀ of the results about cytotoxic effect, anti-leishmania and anti-trypanosoma activities were calculated using the software GraphPad Prism 7.0, applying a sigmoidal regression curve of dose-response.

3. Results

3.1. Compounds identified in the aqueous extracts of the leaves and stem bark of *Z. joazeiro* Mart.

The chromatographic analysis of the *Z. joazeiro* Mart. extracts was determined by UPLC-MS-ESI-QTOF in the negative mode. The results obtained show 12 compounds present in AEB, with the identification of 7 saponin derivatives which belong to the terpenoid classes (Table 1).

Twenty-four compounds were found in AEL, with 12 compounds being identified (Table 1), including four terpenoids, seven flavonoids and one phenolic acid.

3.2. Antipromastigote and antiepipromastigote activities of the extracts of *Z. joazeiro* Mart.

Table 2 presents the antipromastigote and antiepipromastigote results of the *Z. joazeiro* extracts, in addition to the standard drug results. AEB presented more significant results for its antiparasitic activity in comparison to the AEL, especially against the *L. braziliensis* and *L. infantum* promastigote forms.

The AEL presented antiparasitic activity against promastigotes and epimastigotes, with an IC₅₀ > 500 μ g/mL, however, without clinical relevance.

The IC₅₀ values of AEL for antipromastigote activities were 1 241 μ g/mL (*L. braziliensis*) and 33 770 μ g/mL (*L. infantum*) and for antiepipromastigote activity, the IC₅₀ value was 76 640 μ g/mL.

The AEB was the most efficient extract against the promastigote forms with an 85.71% inhibition for *L. braziliensis* and 68.44% for *L. infantum*, at 1 000 μ g/mL concentration. The IC₅₀ values for antipromastigote activities were 327.4 μ g/mL (*L. braziliensis*) and 405.2 μ g/mL (*L. infantum*) and for antiepipromastigote activity, the IC₅₀ value was 1 080 μ g/mL.

3.3. Cytotoxicities of the extracts of *Z. joazeiro* Mart.

The cytotoxicity of AEL was not observed. The AEB showed moderate toxicity, with an IC₅₀ value for cytotoxicity of 333.9 μ g/mL, and inhibition of 61.87% fibroblasts at the concentration of 250 μ g/mL.

Table 1. Identification of compounds by UPLC-QTOF in aqueous extract of stem bark (AEB) and leaves (AEL) of *Z. joazeiro*.

Samples	Quantitative	Classes	Compound
AEB	7	Terpenoids	Saponin derivatives
	5	None	None
AEL	4	Terpenoids	Saponin derivatives
	7	Flavonoids	C-flavone glycosides
			Myricetin- <i>O</i> -glucoside
			Rutin
			Quercetin- <i>O</i> -hexoside
			Quercetin-robnoside
			Ramnazin-3- <i>O</i> -rutinoside
			Ramnazin-hexoside
	1	Phenolic acid	Dihydroxybenzoic acid pentoside
	12	None	None

Table 2. Antipromastigote and antiepipmastigote activities of the extracts of *Z. joazeiro* Mart (mean±SD)(%).

Samples (µg/mL)	%AP (L.b)	%AP (L.i)	%AE
AEL			
1 000	51.11±2.33	3.88±0.86	1.72±0.80
500	23.51±0.43	0.00±0.22	0.00±0.31
250	13.05±0.75	0.00±0.31	0.00±0.61
AEB			
1 000	85.71±0.70	68.44±0.42	43.00±4.74
500	82.16±0.47	67.89±0.20	26.71±0.14
250	42.92±0.78	51.83±0.43	25.78±0.64
Metronidazole			
2.0	100.0	100.0	-
1.0	97.9	97.9	-
Nifurtimox			
1.0	-	-	54.9
0.5	-	-	45.6

AP: antipromastigote percentage; AE: antiepipmastigote; L.b: *L. braziliensis*; L.i: *L. infantum*. AEL: aqueous extracts of leaves of *Z. joazeiro*; AEB: aqueous extracts of stem bark of *Z. joazeiro*.

4. Discussion

The *Ziziphus* genus is known for having a presence of alkaloids and polysaccharides, in addition to a significant number of flavonoids, tannins and saponins in its composition[23,24].

Regarding antiparasitic activity, in a study carried out by Brito *et al*[25] using a *Z. joazeiro* leaf hydroethanolic extract, no relevant antiparasitic activity was observed against *T. cruzi* (IC₅₀: 612.061 µg/mL), *L. braziliensis* (IC₅₀: > 5 000 µg/mL) and *L. infantum* (IC₅₀: 693.67 µg/mL), with the extract exhibiting a low cytotoxic activity.

Gomes *et al*[26] evaluated the antimalarial activity of aqueous extract from *Z. joazeiro* stem bark, observing complete nematode egg inhibition, *Haemonchus* spp., where the IC₅₀ was determined at 1.9 µg/mL of the extract.

In this study, the AEL did not present relevant antiparasitic activity. However, some constituents of the flavonoid class, such as myricetin-*O*-glucoside, quercetin-*O*-hexoside and quercetin-robnoside, which are derived from myricetin and quercetin, have been observed in its composition. When isolated from plant extracts,

myricetin and quercetin were effective in reducing *T. cruzi* strains, in addition to presenting low toxicity to mammalian cells[27]. In this way, it is believed that the constituents present in complexes do not have the effect that their isolated form has.

The cytotoxicity presented by the AEB may be related to the presence of saponins in its chemical composition. Several studies report high cytotoxicity for many saponins[28–30].

Saponins are molecules produced by the plant's secondary metabolism which act primarily as a chemical defense system against herbivores, as well as fungal and bacterial infections[31,32]. Despite reports on its toxicity, these molecules are responsible for a wide variety of biological activities such as molluscicidal, antiparasitic[33], anti-inflammatory, cytotoxic, anti-platelet and anti-diabetic activities[23,26,29,34].

In conclusion, comparative evaluation of the antiparasitic activity of the *Z. joazeiro* aqueous extracts demonstrated the efficacy of the AEB against promastigote forms and an antiparasitic action without clinical relevance of the AEL. The cytotoxicity from the AEB extract observed is of relevant information for the rational use of this plant part since it is widely used by the population. UPLC-QTOF analysis revealed compounds which can be used as a basis for further biological studies against this group of neglected diseases.

Conflict of interest statement

The authors declare that there is no conflict of interest.

References

- [1] World Health Organization. *Working to overcome the global impact of neglected tropical diseases: First WHO report on neglected tropical diseases*. Geneva: World Health Organization; 2011. [Online] Available from: https://www.who.int/neglected_diseases/2010report/en/ [Accessed on 15 Dec 2017].
- [2] Vasconcelos RS, Kovaleski DF, Tesser Junior ZC. Doenças negligenciadas: Revisão da literatura sobre as intervenções propostas. *Saúde Transformação Soc* 2016; **6**(2): 114-131.
- [3] Reis ACMS, Borges DPL, D'Ávila VCGF, Barbosa MS, Ternes YMF, Santiago SB, et al. O cenário de políticas públicas do Brasil diante do quadro das doenças negligenciadas. *Saúde Ciência em Ação* 2016; **2**(2): 99-107.
- [4] Moraes-Braga MFB, Souza TM, Santos KKA, Andrade JC, Guedes GMM, Tintino SR, et al. Citotoxicidade e atividade antiparasitária de *Lygodium venustum* SW. *Acta Toxicol Argent* 2013; **21**(1): 50-56.
- [5] DNDi. *Diseases and projects*. [Online] Available from: <https://www.dndial.org/doencas/doenca-chagas/> [Accessed on 22 Jan 2017].
- [6] World Health Organization. *Información general: Enfermedad de Chagas*. [Online] Available from: http://www.paho.org/hq/index.php?option=com_content&view=article&id=5856&Itemid=41506&lang=es

- [Accessed on 15 Dec 2017].
- [7] Gontijo B, Carvalho MLR. Leishmaniose tegumentar americana. *Rev Soc Bras Med Trop* 2003; **36**(1): 71-80. Doi: 10.1590/S0037-86822003000100011.
- [8] Mennai I, Hanfer M, Esseid C, Benayache S, Ameddah S, Menad A, et al. Chemical composition, *in vitro* antiparasitic, antimicrobial and antioxidant activities of *Frankenia thymifolia* Desf. *Nat Prod Res* 2019; **24**: 1-6. Doi: 10.1080/14786419.2018.1561685.
- [9] Chappuis F, Sundar S, Hailu A, Ghalib H, Rijal S, Peeling RW, et al. Visceral leishmaniasis: What are the needs for diagnosis, treatment and control? *Nat Rev Microbiol* 2007; **5**(11): 873-882. Doi: 10.1038/nrmicro1748.
- [10] World Health Organization. *Clinical forms of the leishmaniases*. [Online] Available from: <http://www.who.int/leishmaniasis/disease/en/> [Accessed on 23 Jan 2018].
- [11] Organização Panamericana da Saúde - OPAS/OMS. *Leishmanioses: Informe Epidemiológico das Américas. Informe Leishmanioses N° 7*. [Online] Available from: http://iris.paho.org/xmlui/bitstream/handle/123456789/34857/LeishReport6_por.pdf?sequence=5 [Accessed on 22 Jan 2018].
- [12] Santos CS, Gomes AMT, Souza FS, Marques SC, Lobo MP, Oliveira DC. Representações sociais de profissionais de saúde sobre doenças negligenciadas. *Esc Anna Nery* 2017; **21**(1): 1-9. Doi: 10.5935/1414-8145.20170016.
- [13] Gontijo CMF, Melo MN. Leishmaniose Visceral no Brasil: quadro atual, desafios e perspectivas. *Rev Bras Epidemiol* 2004; **7**(3): 338-349. Doi: 10.1590/S1415-790X2004000300011.
- [14] Silva EL, Nicoletti MA. Controle e tratamento das doenças negligenciadas: visão da situação atual. *Rev Saúde* 2013; **7**(3): 65-81.
- [15] Freitas AVL, Coelho MFB, Pereira YB, Freitas Neto EC, Azevedo RAB. Diversidade e usos de plantas medicinais nos quintais da comunidade de São João da Várzea em Mossoró, RN. *Rev Bras Pl Med* 2015; **17**(4): 845-856. Doi: 10.1590/1983-084X/14_080.
- [16] Almeida CFCBR, Silva TCL, Amorim ELC, Maia MBS, Albuquerque UP. Life strategy and chemical composition as predictors of the selection of medicinal plants from the Caatinga (Northeast Brazil). *J Arid Environ* 2005; **62**: 127-142.
- [17] Nisar M, Adzu B, Inamullah K, Bashir A, Ihsan A, Gilani AH. Antinociceptive and antipyretic activities of the *Zizyphus oxyphylla* Edgew. Leaves. *Phytother Res* 2007; **21**(7): 693-695. Doi: 10.1002/ptr.2139.
- [18] Matos FJA. *Farmácias vivas*. 4 ed. Fortaleza, CE: Editora UFC; 2002, p. 36-40.
- [19] Le Senne A, Muelas-Serrano S, Fernandez-Portillo C, Escario JA, Gómez-Barrio A. Biological characterization of a beta-galactosidase expressing clone of *Trypanosoma cruzi* CL strain. *Mem Inst Oswaldo Cruz* 2002; **97**: 1101-1105.
- [20] Vega C, Rolón M, Martínez-Fernández AR, Escario JA, Gómez-Barrio A. A new pharmacological screening assay with *Trypanosoma cruzi* epimastigotes expressing beta-galactosidase. *Parasitol Res* 2005; **95**: 296-298.
- [21] Mikus J, Steverding D. A simple colorimetric method to screen drug cytotoxicity against *Leishmania* using dye Alamar Blue®. *Parasitol Int* 2000; **48**: 265-259.
- [22] Rolón M, Seco E, Vega C, Nogal JJ, Escario JA, Gómez-Barrio A, et al. Selective activity of polyene macrolides produced by genetically modified *Streptomyces* on *Trypanosoma cruzi*. *Int J Antimicrob Agents* 2006; **28**: 104-109.
- [23] Kaleem WA, Muhammad N, Khan H, Rauf A. Pharmacological and phytochemical studies of genus *Zizyphus*. *Int J Sci Res* 2014; **21**(8): 1243-1263. Doi: 10.5829/idosi.mejsr.2014.21.08.21099.
- [24] Pu Y, Ding T, Zhang N, Jiang P, Liu D. Identification of bitter compounds from dried fruit of *Zizyphus jujuba* cv. Junzao. *Int J Food Prop* 2017; **20**: 1-34. Doi: 10.1080/10942912.2017.1288133.
- [25] Brito SMO, Coutinho HDM, Talvani A, Coronel C, Barbosa AGR, Veja C, et al. Analysis of bioactivities and chemical composition of *Zizyphus joazeiro* Mart. using HPLC-DAD. *Food Chem* 2015; **1**(186): 185-191. Doi: 10.1016/j.foodchem.2014.10.031.
- [26] Gomes DC, Lima HG, Vaza AV, Santos NS, Santos FO, Dias ÊR, et al. *In vitro* anthelmintic activity of the *Zizyphus joazeiro* bark against gastrointestinal nematodes of goats and its cytotoxicity on Vero cells. *Vet Parasitol* 2016; **15**(226): 10-16. Doi: 10.1016/j.vetpar.2016.06.004.
- [27] Faria RX, Souza ALA, Lima B, Tietbohl LAC, Fernandes CP, Amaral RR, et al. Plants of Brazilian restingas with tripanocide activity against *Trypanosoma cruzi* strains. *J Bioenerg Biomembr* 2017; **49**(6): 473-483. Doi: 10.1007/s10863-017-9733-9.
- [28] Kim AD, Kang KA, Zhang R, Lim CM, Kim HS, Kim DH, et al. Ginseng saponin metabolite induces apoptosis in MCF-7 breast cancer cells through the modulation of AMP-activated protein kinase. *Environ Toxicol Pharmacol* 2010; **30**(2): 134-140. Doi: 10.1016/j.etap.2010.04.008.
- [29] Li N, Wu CF, Xu XY, Liu ZY, Li X, Zhao YQ. Triterpenes possessing an unprecedented skeleton isolated from hydrolyzate of total saponins from *Gynostemma pentaphyllum*. *Eur J Med Chem* 2012; **50**: 173-178. Doi: 10.1016/j.ejmech.2012.01.052.
- [30] Hu Q, Chen YY, Jiao QY, Khan A, Li F, Han DF, et al. Triterpenoid saponins from the pulp of *Sapindus mukorossi* and their antifungal activities Qingwen. *Phytochemistry* 2018; **147**: 1-8. Doi: 10.1016/j.phytochem.2017.12.004.
- [31] Qu L, Wang J, Ruan J, Yao X, Huang P, Wang Y, et al. Spirostane-type saponins obtained from *Yucca schidigera*. *Molecules* 2018; **23**(1): 2-12. Doi: 10.3390/molecules23010167.
- [32] Wina E, Muetzel S, Becker K. The impact of saponins or saponin-containing plant materials on ruminant production-A review. *J Agric Food Chem* 2005; **53**(21): 8093-8105. Doi: 10.1021/jf048053d.
- [33] Moses T, Pollier J, Almagro L, Buyst D, Montagu MV, Pedreño MA, et al. Combinatorial biosynthesis of saponins and saponins in *Saccharomyces cerevisiae* using a C-16 α hydroxylase from *Bupleurum falcatum*. *Proc Natl Acad Sci U S A* 2014; **111**(4): 1634-1639. Doi: 10.1371/journal.pone.0165954.
- [34] Podolak I, Galanty A, Sobolewska D. Saponins as cytotoxic agents: A review. *Phytochem Rev* 2010; **9**(3):425-474. Doi: 10.1007/s11101-010-9183-z.