

## ORIGINAL RESEARCH ARTICLES

## Plant Genetic Resources

# Occurrence of $2n$ microspore production in diploid interspecific hybrids between the wild parental species of peanut (*Arachis hypogaea* L., Leguminosae) and its relevance in the genetic origin of the cultigen

A. V. García<sup>1</sup>  | A. M. Ortiz<sup>1</sup>  | M. C. Silvestri<sup>1,2</sup>  | A. R. Custodio<sup>3</sup> |  
M. C. Moretzsohn<sup>3</sup> | G. I. Lavia<sup>1,2</sup> 

<sup>1</sup> Instituto de Botánica del Nordeste (IBONE, CONICET-UNNE), Sargento Cabral 2131, Corrientes, 3400, Argentina

<sup>2</sup> Facultad de Ciencias Exactas y Naturales y Agrimensura, (UNNE), Corrientes, 3400, Argentina

<sup>3</sup> Empresa Brasileira de Pesquisa Agropecuária, Recursos Genéticos e Biotecnologia, Parque Estação Biológica, Federal District, Brasília, 70770-917, Brazil

## Correspondence

A. V. García, Instituto de Botánica del Nordeste (IBONE, CONICET-UNNE), Sargento Cabral 2131, Corrientes, 3400, Argentina.  
Email: [alevanina.g23@gmail.com](mailto:alevanina.g23@gmail.com)

Assigned to Associate Editor Malleswari Gelli.

Agencia Nacional de Promoción Científica y Tecnológica, Grant/Award Number: PICT-2015-2681; Secretaría General de Ciencia y Técnica de la Universidad Nacional del Nordeste, Corrientes, Argentina

## Abstract

Unreduced ( $2n$ ) gametes are the driving force that leads to polyploidization of plants in nature and an important tool for ploidy breeding. Peanut (*Arachis hypogaea* L., Leguminosae) seems most likely to have originated by sexual polyploidization through interspecific hybridization between the wild diploid relatives—*A. duranensis* Krapov. & W.C. Greg. and *A. ipaënsis* Krapov. & W.C. Greg.—and subsequent binding of the unreduced gametes in the hybrid. Therefore, aiming to contribute to understanding the event of polyploid origin of this important crop, we made diploid hybrids between *A. duranensis* and *A. ipaënsis* and then investigated the production of unreduced gametes in the hybrids by analyzing the size range of pollen, the constitution of the sporads, and the microsporogenesis process. The meiotic abnormalities found, such as persistent bridges, tripolar spindles, and cytomixis processes, would lead to the formation of restitution nuclei in both the first and second meiotic division. These failures in the regularity of meiosis lead to the formation of dyads and triads, and consequently to  $2n$  pollen grains. These findings constitute the first evidence of the formation of unreduced gametes in intergenomic hybrids obtained between the wild progenitors of *A. hypogaea* and enable us to propose a model of origin of the crop through sexual polyploidization.

## 1 | INTRODUCTION

The genus *Arachis* includes 82 species (Krapovickas & Gregory, 1994; Santana & Valls, 2015; Valls & Simpson, 2005, 2017; Valls, Da Costa, & Custodio, 2013), 31 of which

belong to the section *Arachis*. This section includes the peanut cultigen *A. hypogaea* L., which is one of the most cultivated edible legumes worldwide and an excellent source of proteins and oils. Peanut cultivation is concentrated mainly in warm areas of Asia, Africa, and America, with a worldwide production that reached 47 Tg in 2017 (FAOSTAT, <http://fao.org/faostat>). Argentina, with a production of 901,800 t of peanuts in grain (2016–2017, Bolsa

**Abbreviations:** DAPI, 4,6-diamidino-2-phenylindole; PCR, polymerase chain reaction; PMC, pollen mother cell.

de Cereales de Córdoba; <http://www.bccba.com.ar/mani-7165.html>), is one of the main producers and exporters in the world, with 98% of the planted area concentrated in the center and south of the province of Córdoba (Ministerio de Agricultura, Ganadería y Pesca, Argentina; <https://www.argentina.gob.ar/agricultura-ganaderia-y-pesca>).

The cultivated peanut is a segmental allotetraploid species with  $2n = 40$  chromosomes (Bertioli et al., 2019; Gregory & Gregory, 1976; Husted, 1933, 1936). Based on the results obtained through different techniques, different species have been proposed as the likely progenitors of *A. hypogaea* (Lavia, Fernández, & Seijo, 2008). Currently, given the information obtained by classical cytogenetic (Fernández & Krapovickas, 1994), molecular cytogenetic (Robledo & Seijo, 2010; Robledo, Lavia, & Seijo, 2009; Seijo et al., 2004, 2007), molecular analysis (Bertioli et al., 2016, 2019; Grabiele, Chalup, Robledo, & Seijo, 2012; Kochert, Halward, Branch, & Simpson, 1991; Kochert et al., 1996; Moretzsohn et al., 2013), and morphological and geographic analysis (Krapovickas & Gregory, 1994), it is well accepted that the progenitors of *A. hypogaea* (AABB) are *A. duranensis* Krapov. & W.C. Greg. (A genome) and *A. ipaënsis* Krapov. & W.C. Greg. (B genome). This hypothesis has also been supported by the development of a synthetic amphidiploid by crosses between *A. ipaënsis* and *A. duranensis* and subsequent chromosome duplication, as well as by obtaining successful hybrids between the synthetic amphidiploid and *A. hypogaea* (Fávero, Simpson, Valls, & Vello, 2006).

Currently, it is largely accepted that *A. hypogaea* originated from a single allopolyploidization event (Bertioli et al., 2011, 2019; Grabiele et al., 2012; Seijo et al., 2004, 2007). Molecular studies using sequence analysis of non-coding regions of chloroplast DNA and the nontranscribed spacer of the nuclear ribosomal DNA 5S unequivocally identified *A. duranensis* as the female parent of the AABB tetraploid *A. hypogaea* (Grabiele et al., 2012). However, after many attempts, few researchers have obtained successful crosses with *A. ipaënsis* by using *A. duranensis* as the female parent (García, Ortiz, & Lavia, 2016; Mallikarjuna, Senthilvel, & Hoisington, 2011; Simpson, 2017).

Allopolyploidy as a result of interspecific hybridization, accompanied by chromosome duplication, has played a key role in the evolution of many important plants, such as cereal, forage, oilseed, and fiber crops (Jauhar, 2007). Allopolyploids may form either through asexual polyploidization via somatic cell chromosome doubling or through sexual polyploidization. However, the latter represents the main evolutionary route for polyploidization among plant species (De Storme & Geelen, 2013; Jauhar, 2003, 2007; Kreiner, Kron, & Husband, 2017a, 2017b; Ramsey & Schemske, 1998). In nature, sexual polyploids are highly efficient, even more than the polyploids gener-

ated by somatic chromosome duplication. In this context, the phenomenon responsible for sexual polyploidization is the production of unreduced ( $2n$ ) gametes in the parental species of polyploids or their hybrids (Jauhar, 2003). Sexual allopolyploids, depending on the parental contribution, can be formed via two pathways: either bilaterally, after the union of two unreduced gametes ( $2n + 2n$ ), or unilaterally, by an uneven ploidy level combination ( $2n + n$ ) (den Nijs & Peloquin, 1977). Most techniques to detect unreduced gametes focus on pollen, since it is more convenient to isolate and screen than egg cells (Dewitte, Van Laere, & Van Huylenbroeck, 2012). Large pollen has frequently been attributed to  $2n$  pollen, since there is a positive correlation between DNA content and cell volume, which in turn influences pollen diameter (Bretagnolle & Thompson, 1995). Thus,  $2n$  pollen grains are larger than reduced grains, and plants producing  $2n$  pollen grains display bimodal pollen grain size distribution instead of a normal distribution (Dewitte et al., 2012; Ramsey & Schemske, 1998). The analysis of microsporogenesis may also provide a useful method to detect the presence and the frequency of  $2n$  gametes, since the presence of dyads and triads in the sporad stage constitutes strong evidence of unreduced pollen production (Dewitte et al., 2012; Orjeda, Freyre, & Iwanaga, 1990).

All the above allow the assumption that peanut originated by sexual polyploidization, through interspecific hybridization between *A. duranensis* and *A. ipaënsis*, and subsequent binding of the unreduced gametes. Therefore, as a first step to test this hypothesis and contribute to the understanding of the mode of peanut origin, in the present study, we carried out controlled reciprocal crosses between *A. duranensis* and *A. ipaënsis* and identified their hybrids<sup>2x</sup>, both by analyzing the distribution patterns of 4,6-diamidino-2-phenylindole (DAPI) heterochromatin by using fluorescent banding, and by genotyping using microsatellite markers. As a second step, in an attempt to elucidate the contribution of  $2n$  gametes to the origin of peanut, we studied the occurrence of unreduced gametes and estimated the frequency of their production by analyzing the size range of pollen and the constitution of the sporads. Finally, we analyzed the meiotic chromosome behavior of pollen mother cells (PMCs) to detect the underlying cytological mechanisms involved in the origin of unreduced gametes.

The results obtained here (i.e., the presence of unreduced gametes in the artificially synthesized interspecific hybrids between *A. duranensis* and *A. ipaënsis*) constitute the first evidence on the genetic origin of peanut via sexual polyploidization. The results also allow unraveling, in induced diploid hybrids (AB), the evolutionary steps that could have occurred in nature when tetraploid peanut originated.

**TABLE 1** Accessions and species used for crosses and induced F<sub>1</sub> hybrids obtained in this study, as well as chromosome number (2n) and genome type (G)

Species	2n	G	Origin and collection number <sup>a</sup>
<i>A. duranensis</i> Krapov. & W.C. Greg.	20	AA	Argentina, Province Salta, Department Capital. 24°45' S, 65°26' W. VEVNv 14167.
	20	AA	Argentina, Province Salta, Department Gral José de San Martín. 22°15' S; 63°43' W. K 7988.
<i>A. ipaënsis</i> Krapov. & W.C. Greg.	20	BB	Bolivia, Department Tarija, Province Gran Chaco, Ipa. 21°00' S; 63°24' W. KGBPScS 30076
[ <i>A. ipaënsis</i> K30076 × <i>A. duranensis</i> K7988] <sup>2x</sup>	20	BA	IpaDur 1, 2
[ <i>A. ipaënsis</i> K30076 × <i>A. duranensis</i> V14167] <sup>2x</sup>	20	BA	IpaDur 3, 4, 5, 6
[ <i>A. duranensis</i> V14167 × <i>A. ipaënsis</i> K30076] <sup>2x</sup>	20	AB	DurIpa 7, 8, 9, 10, 11, 12

<sup>a</sup>B, D. Banks; Ev, A. Echeverry; G, W.C. Gregory; K, A. Krapovickas; Nv, L. Novara; P, J. Pietrrelli; S, C.E. Simpson; Sc, A. Schinini; V, J.F.M. Valls.

## 2 | MATERIALS AND METHODS

### 2.1 | Plant material

The material used in the crosses is detailed in Table 1.

### 2.2 | Crosses

Crosses were made between November 2013 and April 2014, in the greenhouse of the Instituto de Botánica del Nordeste (IBONE), Corrientes, Argentina. The technique consisted of emasculation of flowers of the female parents in the bud phase (before the floral opening) between 1600 and 1900 h. In the morning of the following day, emasculated flowers were pollinated with pollen grains from the male parent (Fávero et al., 2006). The crosses were reciprocal, taking both species as female and male parents. The fruits were harvested during June–July 2014 and kept in the Germplasm Bank of the IBONE. In early September, all harvested seeds were placed, conveniently separated and identified, on individual germinators with vermiculite.

### 2.3 | Identification of hybrids

The seedlings resulting from the germination of the fruits obtained from the reciprocal crosses between *A. duranensis* and *A. ipaënsis* were identified by fluorescent DAPI banding and microsatellite marker analysis, both of which were used for the separation of hybrids and possible self-pollinated individuals.

The DAPI banding was performed due to the difference in the heterochromatin distribution patterns in the different genomes of the species involved in the crosses. Species with the A genome (*A. duranensis*) have centromeric DAPI+ bands, whereas those with the B genome (*A. ipaënsis*) are devoid of heterochromatic bands (Robledo & Seijo, 2010; Robledo et al., 2009; Seijo et al., 2004). If the individuals resulting from the crosses were hybrids, DAPI+ heterochromatic bands should be found in half of the chromosome complement. Root tips were treated with 2 mM 8-hydroxyquinoline and subsequently fixed in absolute ethanol/glacial acetic acid solution (3:1) for 12 h at 4 °C (Fernández & Krapovickas, 1994, modified). The root apices were digested in 1% (w/v) cellulase (from *Trichoderma viridae*; Onozuka R-10, Serva) plus 10% (v/v) pectinase (from *Aspergillus niger*, Sigma) dissolved in 40% glycerol in 0.01 M citrate buffer (pH 4.8) for 2 h at 37 °C before squashing, and slides were air dried according to Gerber and Schweizer (1988). The DAPI banding was performed according to the protocol described by Schweizer (1976).

Microsatellite markers were chosen due to their codominant nature, ease of use, and speed in providing results. DNA was extracted using the protocol described by Inglis, de Pappas, Resende, and Grattapaglia (2018). The quality and quantity of the DNA were evaluated in 0.8% agarose gel electrophoresis and using a Nanodrop 1000 Spectrophotometer (Thermo Fisher Scientific), respectively. DNA amplification reactions were performed by multiplex polymerase chain reaction (PCR) designed with the Multiplex Manager program (Holleley & Geerts, 2009). The selected parameters included a maximum of three primers each marked with a different fluorescence (6-FAM, HEX, and NED) by reaction and similar annealing temperatures. The

PCR reactions were performed using the QIAGEN multiplex PCR kit (Qiagen), with 2× of master mix, 0.05 μM of primers (forward and reverse), 1 ng of genomic DNA, and adjusting the final volume to 5 μl with ultrapure water. The amplifications were performed in ABI GeneAmp PCR system 9700 thermal cyclers (Applied Biosystems) with the following cycling program: initial denaturation for 15 min at 95 °C, followed by 30 cycles composed of denaturation for 30 s at 94 °C, hybridization for 90 s at 58 °C, and extension for 90 s at 72 °C, followed by a final extension of 30 min at 72 °C. The amplified fragments were resolved by capillary electrophoresis on an ABI 3730 DNA analyzer (Applied Biosystems). For a better resolution, the obtained PCR products were diluted 20× in ultrapure water. Loading samples contained 1 μl of PCR product diluted 20×, 0.2 μl of Rox-labeled size standards, and 8 μl of Hi-di Formamide (Applied Biosystems) and were denatured for 4 min at 95 °C. For the allele sizing, the electropherograms were analyzed with the GeneMapper software (Applied Biosystems).

## 2.4 | Detection and analysis of unreduced gametes

The frequency of unreduced gametes was determined by analyzing the pollen size distribution and the constitution of the sporads from all the hybrids obtained. Pollen stainability with carmine/glycerine (1:1) was used as an estimation of pollen viability (Pittenger & Frolik, 1951). Three mature flower buds for each hybrid individual were analyzed. All pollen grains found were counted, and those regularly shaped and darkly stained were considered to be viable. Pollen viability was expressed as an average percentage of the stained pollen grains/total pollen grains analyzed.

All viable pollen grains per flower bud were measured and counted using ImageJ software (Rasband, 1997). Pollen size was estimated by the Em index (i.e., the distance between the aperture angle and the middle point of the opposite side; Van Campo, 1957). According to their size, viable pollen grains were classified into four categories: small, normal, large  $2n$ , and large  $4n$ . Normal pollen was considered to determine the mean pollen diameter of each individual. Pollen grains showing a size 0.25× smaller than the average size of the individuals were considered as small pollen. Unreduced pollen was detected following Darlington (1937): pollen grains showing a size 1.25× larger than the average size of the individuals were considered as large  $2n$  grains (giant pollen), whereas deeply stained grains showing a size 1.5× larger than the average size of the individuals were considered large  $4n$  grains (jumbo pollen).

To confirm that large pollen grains effectively represent  $2n$  and  $4n$  pollen production, microspores at the sporad stage were screened in all hybrid plants. For this analysis, PMCs from fresh flower buds were stained with carmine/glycerine (1:1). Sporads were classified as tetrads, abnormal tetrads, triads, dyads, monads, and polyads. Tetrads were considered normal when they had four equal-sized cells. Abnormal tetrads include tetrads with four microspores of different sizes, tetrads with collapsed microspores, and tetrads with microcytes.

To evaluate the existence of significant differences for pollen size and the production of  $2n$  gametes among individuals, one-way ANOVA at a significance level of 5% ( $\alpha = .05$ ) after Bartlett's test of homogeneity was performed. Also, Tukey's test at 5% significance was carried out to test differences between each pair of means. All statistical analyses were performed using INFOSTAT software version 10.0 (Di Rienzo et al., 2017).

## 2.5 | Analysis of meiotic behavior

Meiotic analysis was carried out in PMCs from young flower buds, either fresh or fixed in ethanol/acetic acid (3:1). The PMCs were stained and squashed in 2% lacto-propionic orcein (Dyer, 1963). Each anther was processed individually. The meiotic behavior was evaluated from Prophase I to Telophase II, and the frequency of each abnormality was recorded. All PMCs found were analyzed.


## 3 | RESULTS AND DISCUSSION

### 3.1 | Crosses

A total of 220 crosses were made. *Arachis duranensis* was used as the female parent in 153 of the crosses (69.5% of the hybridizations), obtaining pegs in 40.5% of the cases, whereas *A. ipaënsis* was used as the female parent in 67 of the crosses (30.5% of the hybridizations), obtaining pegs in 20.9% of the cases. Unsuccessful results in both species included pollinations that were not effective and pegs that dried due to natural causes.

All the fruits from the different hybrid progenies were collected. A total of 47 seeds were collected from the flower pots of *A. duranensis*, whereas only 10 were found in pots where *A. ipaënsis* was used as the female parent. Also, in the latter, 10 seedlings were found, suggesting the lack of dormancy of those seeds.

Every seed collected was put in individual cups to germinate with mineral substratum. All seeds of *A. ipaënsis* × *A. duranensis* germinated, whereas seeds from the reciprocal



**FIGURE 1** Somatic metaphases of *Arachis* diploids after 4,6-diamidino-2-phenylindole (DAPI) staining: (a) *Arachis duranensis* (AA) with DAPI<sup>+</sup> centromeric bands in all chromosomes, (b) *Arachis ipaënsis* (BB) with chromosomes devoid of bands, (c) the [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrid, and (d) the [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrid, showing half of the chromosomes (those belonging to the A genome) with conspicuous centromeric bands and half of the chromosomes (those of the B genome) without centromeric bands. White arrows indicate “A chromosomes,” and white arrowheads indicate the satellites. Scale bar = 5 μm

crosses did not. The ungerminated seeds were removed from the cups, stored, and resown several months later. This procedure allowed 16 seeds to germinate, suggesting that they presented the physiological characteristics of dormancy.

### 3.2 | Identification of hybrids

The fluorescent DAPI banding technique allowed determination that six of the seedlings germinated in the pot of *A. ipaënsis* as the female parent, and that six of the seedlings germinated in vermiculite from *A. duranensis* × *A. ipaënsis* crosses were hybrids (Table 1), because they presented the characteristic banding pattern (DAPI<sup>+</sup> centromeric bands on chromosomes of the A genome and the absence of DAPI<sup>+</sup> centromeric bands on chromosomes of the B genome, Figure 1). All the hybrids analyzed showed diploid level with  $2n = 20$  chromosomes in the mitotic cells.

A total of 26 polymorphic microsatellite markers were also used for hybrid identification. The 26 loci confirmed that the six seedlings from the *A. duranensis* × *A. ipaënsis* crosses and the six seedlings from the *A. ipaënsis* × *A. duranensis* crosses were true F<sub>1</sub> hybrids. Since 13 of the markers were polymorphic between the accessions K 7988 and V 14167 of *A. duranensis*, it was also possible to identify

that four of these seedlings resulted from crosses between *A. ipaënsis* × *A. duranensis* V 14167 and that two resulted from crosses between *A. ipaënsis* × *A. duranensis* K 7988. As expected, microsatellite markers proved to be efficient in the identification of hybrids. All the individuals of the progenies were undoubtedly identified as the hybrid or self-pollinated condition.

The success rate achieved was calculated as the number of hybrid individuals in relation to the number of pollinations. In the crosses using *A. ipaënsis* as the female parent, we obtained a success rate of 8.95%. Our results are slightly different from Fávero et al. (2006), despite the fact that the same accessions were used as parents in both studies. In the case of *A. duranensis* × *A. ipaënsis* crosses, the success rate could not be calculated because not all the seeds managed to germinate. However, obtaining hybrids with *A. duranensis* as a female parent has not been previously registered. Although several authors have made attempts to achieve this hybrid (Simpson, personal communication, 1984–1986; Stalker, Dhesi, Parry, & Hahn, 1991), every effort was unsuccessful. The crosses obtained in the present work, together with those obtained by Simpson (2017), constitute the first [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrids successfully obtained.

### 3.3 | Estimation of unreduced gametes based on the viability, size, and morphology of pollen grains

The pollen grains of *A. duranensis*, *A. ipaënsis*, and the 12 hybrids derived from the reciprocal interspecific crosses (Table 2) were analyzed to estimate pollen viability and determine variation in their pollen grain size, in order to identify potential unreduced gamete producers.

Both parental species had high pollen viability: *A. duranensis* V 14167 (97.65%), *A. duranensis* K 7988 (98.55%), and *A. ipaënsis* (98.75%). By contrast, the hybrids had very low viability, ranging from 1.45 to 3.96% in [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> and from 1.27 to 3.11% in [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> (Table 2, Supplemental Figure S1). In the genus *Arachis*, pollen viability of interspecific hybrids has traditionally been used to infer the genomic relationships between the species involved in the crosses. Hybrids between species that have the same genome type generally show values of pollen viability between 25 and 70% (Burow, Simpson, Faries, Starr, & Paterson, 2009; Krapovickas & Gregory, 1994; Smartt, Gregory, & Gregory, 1978; Stalker et al., 1991; Tallury et al., 2005), whereas crossings between species with different genome types usually show values below 10% (Burow et al., 2009; Krapovickas & Gregory, 1994; Stalker et al., 1991). As expected, the intergenomic

TABLE 2 Types of pollen grains found and pollen viability evaluated in *A. duranensis*, *A. ipaënsis* and the 12 F<sub>1</sub> hybrids obtained

Individual <sup>a</sup>	n <sup>b</sup>	Unviable pollen grains	Type of viable pollen grains <sup>c</sup>				Viability
			S	No	2n	4n	
<i>A. duranensis</i> V 14167	2,000	2.35	–	–	–	–	97.65
<i>A. duranensis</i> K 7988	2,000	1.45	–	–	–	–	98.55
<i>A. ipaënsis</i> K 30076	2,000	1.25	–	–	–	–	98.75
IpaDur							
1	7,623	97.61	0.09	1.39	0.83	0.08	2.39
2	5,716	96.04	0.12	2.47	1.35	0.02	3.96
3	6,150	97.63	0.21	1.95	0.19	0.02	2.37
4	7,082	98.39	1.41	0.15	0.02	0.03	1.61
5	7,314	98.55	1.24	0.17	0.04	0.00	1.45
6	6,857	97.99	1.28	0.48	0.18	0.07	2.01
DurIpa							
7	7,164	97.89	1.48	0.34	0.14	0.15	2.11
8	6,511	98.11	1.00	0.34	0.40	0.15	1.89
9	6,036	98.47	1.09	0.31	0.13	0.00	1.53
10	8,333	98.73	0.92	0.21	0.10	0.04	1.27
11	6,900	96.89	1.18	0.86	0.82	0.25	3.11
12	6,823	98.40	0.79	0.45	0.23	0.13	1.60

<sup>a</sup>IpaDur, [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup>; DurIpa, [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup>.

<sup>b</sup>n, number of pollen grains analyzed.

<sup>c</sup>S, small size pollen grain; No, reduced normal pollen grain; 2n, giant pollen grain; 4n, jumbo pollen grain.

hybrids obtained between *A. duranensis* (AA) and *A. ipaënsis* (BB) showed values of pollen viability lower than 4% (Table 2), data that support their hybrid condition.


A significant difference in the size of viable pollen grains was detected between both progenitor species ( $F = 128.2$ ,  $p = .0001$ ), with *A. duranensis* having a pollen grain larger than *A. ipaënsis* (Supplemental Figure S1). The average pollen size of *A. duranensis* was  $34.87 \pm 1.90 \mu\text{m}$ , varying from  $31.36$  to  $39.20 \mu\text{m}$ , whereas that of *A. ipaënsis* was  $32.18 \pm 1.69 \mu\text{m}$ , varying from  $27.44$  to  $35.28 \mu\text{m}$ .

In the hybrid individuals, the size of viable pollen grains showed significant variation ( $F = 139.50$ ,  $p \leq .0001$ ), and so they were classified as small, normal, large 2n (giant pollen), or large 4n pollen (jumbo pollen) (Figures 2a–2c). Hybrid individuals also showed a significant difference in the size of normal viable pollen ( $F = 4.53$ ,  $p = .0354$ ), depending on the pollen donor species, with [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrids having a larger pollen size. The average size of normal pollen of [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrids was  $34.95 \pm 2.94 \mu\text{m}$ , varying from  $29.40$  to  $39.20 \mu\text{m}$ , whereas that of [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrids was  $36.10 \pm 2.80 \mu\text{m}$ , varying from  $29.40$  to  $41.16 \mu\text{m}$ . Among the large pollen grains found, those with a size equal to or greater than  $43.68$  and  $45.12 \mu\text{m}$  were classified as 2n pollen or giant pollen


grains in [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> and [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrids, respectively (Figure 2b), whereas those with a size equal to or higher than  $52.45$  and  $54.15 \mu\text{m}$  were classified as 4n pollen or jumbo pollen grains in [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> and [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrids, respectively (Figure 2c). All hybrid individuals analyzed produced giant pollen grains, whereas 80% produced jumbo pollen grains (Table 2). Usually, the presence of large pollen grains results in a bimodal size distribution, as opposed to a normal size distribution (Tondini, Tavoletti, Mariani, & Veronesi, 1993). Concerning that, the analysis of the distribution of pollen grain sizes of the parental species showed an expected normal distribution of the data (Figure 3). In contrast, the frequencies of the pollen grain sizes of the F<sub>1</sub> hybrids were found to have a bimodal distribution, attributed to the high production of normal and large grains in [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> hybrids, and due to the abundant presence of small and large grains in [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrids (Figure 3).

Regarding shape, normal pollen grains were elliptical in both the parental species and hybrids, whereas the large pollen grains detected in hybrids were spherical.

Following the criteria of Kovalsky and Solís Neffa (2012), Dewitte et al. (2012), and Orjeda et al. (1990), three facts



**FIGURE 2** Variation in pollen size and sporad constitution in the hybrids: (a) small and normal viable pollen grains, (b) size difference between normal and  $2n$  pollen grains, and (c) size difference between  $2n$  and  $4n$  pollen grains, (d) normal tetrads, (e) triads with one  $2n$  microspore, (f) dyads with two  $2n$  microspores, and (g) monads with one  $4n$  microspore. S, small; No, normal; Uv, unviable. Scale bar = 20  $\mu\text{m}$




**FIGURE 3** Frequency distribution of pollen grain size in the parental species (*A. duranensis* and *A. ipaënsis*) and the hybrid individuals (IpaDur 1 and DurIpa 10)

suggest that the giant pollen grains detected in the hybrids correspond to  $2n$  gametes. The first is the presence of triads and dyads in the hybrids, indicating that unreduced

gametes certainly occurred and that  $2n$  pollen formation was expected (see below). The second (i.e., the bimodal distribution of pollen size in hybrids) is in accordance with the description for other species with  $2n$  pollen grains. Lastly, besides pollen size, the spherical instead of ellipsoidal pollen shape detected in the giant pollen grains of the hybrids has been associated with the level of ploidy, simplifying the determination of  $2n$  pollen. In this sense, Akutsu, Kitamura, Toda, Miyajima, and Okazaki (2007) observed that by omitting the first or second meiotic division, the restitution nuclei consequently combine tetrads, and the resulting  $2n$  pollen (dyad) is a semisphere, which looks round under the microscope.

The finding of  $2n$  and  $4n$  pollen in the [*A. duranensis*  $\times$  *A. ipaënsis*]<sup>2x</sup> and [*A. ipaënsis*  $\times$  *A. duranensis*]<sup>2x</sup> interspecific hybrids is the first evidence of the production of unreduced gametes in *Arachis* hybrids involving the wild parental species of peanut. In this sense, the importance of this finding lies in the fact that all artificially developed hybrids produced  $2n$  pollen grains (Table 2, Figures 2B and 2c). The ANOVA test showed that the production frequency of  $2n$  gametes did not differ statistically between the F<sub>1</sub> hybrids ( $F = 1.55$ ,  $p = .17$ ), but three of them (IpaDur1, IpaDur2, and DurIpa11) showed the highest frequency (between 0.82 and 1.35%). Another relevant result is that, although all F<sub>1</sub> hybrids showed low pollen viability (<4%), in the three abovementioned individuals,  $2n$  pollen represented 34.73,



**FIGURE 4** Percentages of  $2n$  pollen grains in relation to viable grains in the 12  $F_1$  hybrid individuals analyzed. Individuals 1–6 are [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup> and 7–12 are [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup>

34.1, and 26.37% of the viable grains (Figure 4), respectively. By contrast, in the rest of the  $F_1$  hybrids,  $2n$  pollen represented between 1.24 and 21.16% of the viable grains. Therefore, although pollen viability of the hybrids was low, in some of them,  $2n$  gametes represented a significant proportion of viable grains, thus with a high probability of participating in the fertilization. These findings suggest that the three  $F_1$  hybrids with the highest production of  $2n$  gametes could be used as diploid parents in the genetic improvement of peanut and for the development of new ones, as well as to provide a bridge to transfer desirable genes from wild diploid parents to the gene pool of peanut.

### 3.4 | Estimation of unreduced gametes derived from sporads

The sporad constitution of the 12 hybrids derived from reciprocal interspecific crosses was analyzed to ascertain if the giant pollen grains found in the hybrids corresponded to unreduced gametes, according to the abovementioned proposed criteria (Dewitte et al., 2012; Kovalsky & Solís Neffa, 2012; Orjeda et al., 1990). This analysis revealed, in addition to the expected normal tetrads, the presence of triads (up to 15.98%), dyads (up to 26.52%), and monads (up to 2.50%) (Table 3, Figures 2d–2g). The presence of triads and dyads indicates the production of one and two  $2n$  microspores, respectively, and that the production of  $2n$  pollen is expected. These data are in agreement with previous studies with hybrids, such as *Begonia* (Dewitte, Van Laere, Van Huylbroeck, & Van Bockstaele, 2010) and *Brassica* (Mason, Nelson, Yan, & Cowling, 2011). In addition, the presence of monads indicates that  $4n$  microspores also occurred, based on the assumption that each monad results in one jumbo pollen grain (Kovalsky & Solís Neffa, 2012). Among sporads with unreduced microspores, only the frequency of production of dyads differed statistically

( $F = 23.95$ ,  $p < .0001$ ) between  $F_1$  hybrids (Table 3), with the individuals DurIpa 9 (37.29%) and IpaDur 2 (20.83%) being the highest producers.

Additionally, abnormal tetrads (tetrads with four microspores of different sizes, tetrads with collapsed microspores, and tetrads with microcytes) and polyads were found (Table 3, Supplemental Figure S2), a fact that probably leads to the production of abnormal gametes (see below).

### 3.5 | Cytological mechanisms involved in the formation of $2n$ gametes

To identify possible mechanisms for the production of  $2n$  gametes, a detailed analysis of the meiosis of the hybrids was performed. This analysis revealed a large number of meiotic abnormalities at different stages (Table 4), which can be grouped into two major categories: those that would produce aneuploid gametes (Supplemental Figure S2), and those that would originate unreduced gametes (Figure 5).

The origin of aneuploid gametes has been related to out-of-plate and nucleus chromosomes, laggard chromosomes, chromosomes with precocious migration, multipolar spindles and cytomixis (Guan, Wang, Cheng, Liu, & Li, 2012; Mursalimov & Deineko, 2018; Ortiz, Seijo, Fernández, & Lavia, 2011; Riso-Pascotto, Pagliarini, & Do Valle, 2005; Wang, Kang, & Zhu, 2010). In our analysis, one tripolar spindle in anaphase I (6.95%), and two tripolar spindles in anaphase II (up to 11.11%) were found. Although these types of spindle alterations are rarely observed in plants (De Storme & Geelen, 2013), we assume that the presence of these alterations would also contribute to the formation of aneuploid gametes. Multipolar spindles have been reported in several plant species (Reis, Sousa, & Viccini, 2016; Riso-Pascotto et al., 2005; Wang et al., 2010) and are correlated with abnormal cytokinesis, which leads to the formation of polyads with unbalanced microspores (Riso-Pascotto et al., 2005; Tilquin, de Brouwer, & Horvat, 1984), such as telophase II with five or six nuclei and the hexads detected in the hybrids. On the other hand, out-of-plate chromosomes were detected in metaphase I (up to 51.02%), anaphase I (up to 18.18%), metaphase II (up to 25%), and anaphase II (up to 7.80%). Meiotic abnormalities related to irregular chromosome segregation, such as precocious chromosome migration to the poles in metaphase I (up to 38.09%) and metaphase II (up to 3.85%), and laggard chromosomes at anaphase I and II (up to 24.65%), were also detected. All the abovementioned meiotic abnormalities could alter the regular chromosome segregation, and the chromosomes that ultimately cannot be incorporated into the telophase



TABLE 3 Analysis of the constitution of sporads in the 12 hybrid individuals studied

Hybrid individual <sup>a</sup>	<i>n</i> <sup>b</sup>	Type of sporad <sup>c</sup>					
		Te	Te ≠	Pol	Tr	Dy	Mo
%							
IpaDur							
1	280	31.43	56.78	0.00	6.43	2.86ab <sup>d</sup>	2.50
2	48	41.67	27.08	0.00	10.42	20.83c	0.00
3	313	71.56	25.88	0.64	1.92	0.00a	0.00
4	650	51.38	40.00	5.85	2.42	0.31ab	0.00
5	625	51.68	41.44	1.76	4.16	0.96ab	0.00
6	554	54.69	34.12	8.12	2.53	0.54ab	0.00
DurIpa							
7	409	51.83	33.26	2.69	11.25	0.97ab	0.00
8	363	30.85	36.33	2.47	17.37	12.13bc	0.55
9	724	33.70	16.85	4.28	7.05	37.29d	0.83
10	713	38.85	38.15	16.27	6.31	0.42ab	0.00
11	1,111	58.86	28.44	5.31	2.79	4.51ab	0.09
12	1,045	61.81	31.68	2.96	3.16	0.39ab	0.00
<i>F</i>					1.62	23.95	0.49
<i>P</i>					.1704	<.0001	.8896

<sup>a</sup>IpaDur, [*A. ipaënsis* × *A. duranensis*]<sup>2x</sup>; DurIpa, [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup>.

<sup>b</sup>*n*, number of pollen grains analyzed.

<sup>c</sup>Te, tetrad; Te ≠, abnormal tetrads; Pol, polyad; Tr, triad; Dy, dyad; Mo, monad.

<sup>d</sup>Means followed by a common letter are not significantly different by the Tukey's test at the 5% ( $\alpha = .05$ ) level of significance.

I and II nuclei, leading to the origin of micronuclei (Ortiz et al., 2011; Pagliarini, 2000; Seijo & Solis Neffa, 2006; Wang et al., 2015). In our analysis, one micronucleus was observed at anaphase I, telophase I, and metaphase II (up to 12.5%), and one to two micronuclei were observed at anaphase II and telophase II (up to 8.33%). The recorded frequency of micronuclei was lower than that expected from the number of PMCs that presented segregation and spindle abnormalities (Table 4). This suggests that a large proportion of the chromosomes irregularly segregated should have been included in at least one of the main nuclei. However, those inclusions did not guarantee the restitution of the euploid chromosome number of the main nuclei. In this sense, the counting at anaphase I, prophase II, metaphase II, anaphase II, and telophase II revealed PMCs with plates and nuclei with an unbalanced number of chromosomes (Table 4). Finally, cytomixis was observed in all the divisional stages of PMCs, a fact that would contribute to the abnormal meiotic behavior, since the transfer of different amounts of chromatin results in unbalanced complements of the PMCs involved (Guan et al., 2012; Qiu, Liao, Liu, Mao, & Liu, 2017). Therefore, all the abovementioned meiotic abnormalities would lead to the formation of microcytes, as well as of sporads with more than four microspores (pentads and hexads) or sporads with four microspores with different sizes (Table 3).

The ultimate consequence of these irregularities is the high number of unviable pollen grains detected in the hybrids (Table 2). This would explain the great diversity of pollen grain sizes found and would also be the main cause of the low pollen viability of the hybrids analyzed.

Concerning the origin of unreduced gametes, detailed cytological studies in several plant species have revealed that these phenomena may be the result of a plethora of cellular failures. According to the review of De Storme and Geelen (2013), these are generally subdivided into three main classes: (a) alterations in meiotic spindle dynamics, (b) defects in meiotic cell plate formation, and (c) omission of meiosis I or II. The formation of unreduced gametes has also been associated with both chromosomal bridges (Bajer, 1964) and the phenomenon of cytomixis, although this has not yet been confirmed experimentally (Dewitte et al., 2012; Mursalimov & Deineko, 2015, 2018).

The meiotic irregularities linked to the formation of unreduced gametes found in the hybrids analyzed here are individually described below.

### 3.5.1 | Chromosome bridges

The formation of bridges was observed in up to 64.10% of the PMCs in anaphase I (Figure 5a) and could be preserved

TABLE 4 Meiotic abnormalities recorded in five hybrid individuals studied

Phase <sup>a</sup>	Abnormality	Hybrid individual													
		IpaDur		1		2		3		6		DurIpa			
		n	%	n	%	n	%	n	%	n	%	n	%		
PI	Cytomixis	(188)	19.68	(2)	100	–	–	(215)	46.05	–	–	–	–		
		(72)	–	(49)	–	(4)	25.00	(63)	–	–	–	–			
		–	–	–	–	–	–	–	38.09	–	–	–	–		
MI	Cell with only one univalent 1–7 Chromosomes with precocious migration	–	–	–	–	–	–	–	–	–	–	–			
		–	–	–	–	–	–	–	–	–	–	–			
		–	–	–	–	–	–	–	–	–	–	–			
AI	1–7 Out-of-plate chromosomes Cytomixis	25.00	–	–	51.02	25.00	–	–	–	–	–	–			
		8.34	–	(72)	4.09	(39)	20.52	(70)	12.86	–	–	–			
		–	–	–	6.94	–	–	–	–	–	–	–			
TI	1–5 Out-of-plate chromosomes Unbalanced plates	18.18	–	–	11.11	–	–	–	–	–	–	–			
		68.18	–	–	8.33	2.56	–	–	–	–	–				
		–	–	–	2.78	–	–	–	–	–	–				
PII	1 Micronucleus 1 Bivalent with precocious migration	–	–	–	6.94	–	–	–	–	–	–	–			
		–	–	–	–	–	–	–	–	–	–				
		–	–	–	–	–	–	–	–	–	–				
MII	1 Tripolar spindle 1–5 Chromosome bridges Cytomixis	–	–	–	–	2.56	–	–	–	–	–	–			
		4.55	–	–	23.61	64.10	–	–	–	–	–				
		–	–	–	22.22	–	–	–	–	–	–				
MIII	1 Micronucleus 1–4 Out-of-nucleus chromosomes 1 Chromosome bridge	–	–	(40)	12.50	9.68	–	–	–	–	–	–			
		–	–	–	5.00	–	–	–	–	–	–				
		–	–	–	12.50	19.35	–	–	–	–	–				
MIV	1 Nucleus of restitution (4n) Cytomixis	–	–	–	35.00	3.23	–	–	–	–	–	–			
		–	–	–	–	32.26	–	–	–	–	–				
		–	–	–	–	–	–	–	–	–	–				
MVI	Unbalanced nuclei 1 Residual bridge 1 Nucleus of restitution (4n)	(3)	–	–	–	0.92	(109)	60.00	(7)	–	–	–			
		–	–	–	–	2.75	–	–	–	–	–				
		–	–	–	–	4.59	–	–	–	–	–				
MVII	Cytomixis Unbalanced plates 1–5 Out-of-plate chromosomes Chromosomes with precocious migration	66.67	–	–	–	7.34	–	–	–	–	–	–			
		–	–	(8)	50.00	25.42	(26)	23.08	–	–	–				
		–	–	–	25.00	16.95	–	–	–	–	–				
MVIII	1 Micronucleus 1–4 Residual bridges 1 Nucleus of restitution (4n) Cytomixis	–	–	–	–	–	–	–	–	–	–	–			
		–	–	–	–	–	–	–	–	–	–				
		–	–	–	–	–	–	–	–	–	–				
MIX	1 Micronucleus 1–4 Residual bridges 1 Nucleus of restitution (4n) Cytomixis	–	–	–	12.50	33.90	–	–	–	–	–	–			
		–	–	–	–	3.39	–	–	–	–	–				
		–	–	–	–	–	–	–	–	–	–				
MXX	1 Micronucleus 1–4 Residual bridges 1 Nucleus of restitution (4n) Cytomixis	–	–	–	–	–	–	–	–	–	–	–			
		–	–	–	–	–	–	–	–	–	–				
		–	–	–	–	–	–	–	–	–	–				


(Continues)

TABLE 4 (Continued)

Phase <sup>a</sup>	Abnormality	Hybrid individual											
		IpaDur						DurIpa					
		1 <i>n</i> <sup>b</sup>	%	2 <i>n</i>	%	3 <i>n</i>	%	6 <i>n</i>	%	7 <i>n</i>	%		
AII	1–6 Laggard chromosomes	–	–	(4)	–	(77)	24.65	–	(44)	6.82	–	–	
	1–4 Out-of-plate chromosomes	–	–	–	–	–	7.80	–	–	–	–	–	
	Unbalanced plates	–	–	–	–	–	6.49	–	–	–	–	30.00	
	1–2 Micronuclei	–	–	–	–	–	3.89	–	–	–	–	–	
	1–4 Out-of-plate chromosomes + 1 chromosome bridge + 1 residual bridge	–	–	–	–	–	1.29	–	–	–	–	–	
	2 Tripolar spindles	–	–	–	–	–	1.29	–	–	6.82	–	10.00	
	1–2 Chromosome bridges	–	–	–	75.00	–	9.08	–	–	6.82	–	–	
	1 Tripolar spindle	–	–	–	25.00	–	3.89	–	–	6.82	–	–	
	1–2 Residual bridges	–	–	–	–	–	14.27	–	–	–	–	–	
	1 Nucleus of restitution (2 <i>n</i> ) + 2 nuclei ( <i>n</i> )	–	–	–	–	–	1.29	–	–	–	–	20.00	
	1 Tripolar spindle + 1 residual bridge	–	–	–	–	–	1.29	–	–	–	–	–	
	1–2 Chromosome bridges +1 residual bridge	–	–	–	–	–	–	–	–	15.91	–	–	
	2 Nucleus of restitution (2 <i>n</i> )	–	–	–	–	–	–	–	–	–	–	1.00	
TII	Cytomixis	–	–	–	–	–	–	–	–	11.36	–	–	
	1–2 Micronuclei	(33)	–	–	–	(12)	8.33	–	(182)	2.75	–	(76)	
	5–6 nuclei	–	–	–	–	–	–	–	–	0.55	–	11.84	
	Unbalanced nuclei	–	–	–	–	–	–	–	–	24.18	–	9.22	
	1 Nucleus of restitution (4 <i>n</i> )	–	48.49	–	–	–	25.00	–	–	3.85	–	9.22	
	2 nuclei of restitution (2 <i>n</i> )	–	27.27	–	–	–	16.67	–	–	4.95	–	48.68	
	1 Nucleus of restitution (2 <i>n</i> ) and 2 nuclei ( <i>n</i> )	–	12.12	–	–	–	–	–	–	9.89	–	14.47	
	Cytomixis	–	–	–	–	–	–	–	–	20.88	–	1.31	
Total		318	35.22	176	71.02	331	55.89	646	65.17	150	89.33		

<sup>a</sup>P, prophase; M, metaphase; A, anaphase; T, telophase; I, first meiotic division; II, second meiotic division.

<sup>b</sup>*n*, absolute values of pollen mother cells analyzed.



**FIGURE 5** Meiotic abnormalities involved in the origin of unreduced gametes: (a) anaphase I and (b) telophase I with chromosome bridges; (c) metaphase II and (d) anaphase II with residual bridges; (e) anaphase II with chromosome bridges; (F) anaphase II with chromosome and residual bridges, (g) Anaphase II with tripolar spindle; (h) cytomixis in metaphase I showing chromosome migration; (i) telophase I, (j) prophase II, and (k) metaphase II with one restitution nucleus; (l) anaphase II and (m) teophase II with two  $2n$  nuclei; (n) telophase II with a  $4n$  restitution nucleus; (o) anaphase II and (p) telophase II with one restitution nucleus and two  $n$  nuclei. Scale bar = 10  $\mu$ m

during telophase I (Figure 5b), like those detected in up to 33.34% of the PMCs. During the meiotic division process, the chromatin continuity of the bridges may not be interrupted, which is why the presence of residual bridges is noticed during meiosis II. This was detected in up to 2.75% of the PMCs in prophase II, 33.90% in metaphase II (Figure 5c), and 14.27% in anaphase II (Figure 5d). Likewise, bridges can be generated during meiosis II, like those

found in up to 75% of the PMCs in anaphase II (Figure 5e). Finally, bridge formation may occur in the second meiotic division of cells showing residual bridges of the first division, as observed in up to 15.91% of the PMCs in anaphase II (Figure 5f).

It has been determined that the persistence of some bridges in telophase I and II can bring the poles closer together and thus establish a chromatin continuity

between them, promoting the formation of the nuclear membrane around the nuclei involved (Seijo, personal communication, 2002). Zhang, Cao, Zhou, and Jia (2017) proposed that chromosome stickiness in thick chromosomal bridges restricts the separation during telophase I. This phenomenon would explain the formation of the restitution nuclei detected in the individuals analyzed in telophase I and in later meiotic phases. This nuclear restitution mechanism has already been cited in different genera such as *Begonia* (Dewitte, Eeckhaut, Van Huylenbroeck, & Van Bockstaele, 2010) and *Arachis* (Ortiz et al., 2011), and in both cases, the results showed that this mechanism is involved in the origin of unreduced gametes.

### 3.5.2 | Tripolar spindles

In several plant species, it has been proposed that the aberrations in the formation of the meiotic spindle constitute one of the main causes of the nuclear restitution in the microsporogenesis process (Bretagnolle & Thompson, 1995; Zhang & Kang, 2010). One of the spindle defects leading to meiotic nonreduction are tripolar spindles (Rim & Beuselinck, 1996). In our analysis, one tripolar spindle was detected in up to 25% of the PMCs in anaphase II (Figure 5g). In addition, in meiosis II, this phenomenon was found to be associated with a residual bridge in up to 1.29% of PMCs.

According to Zhang and Kang (2010), the presence of tripolar and fused spindles may explain the formation of triads and dyads. Likewise, some PMCs can exhibit absence of cytokinesis, producing high frequencies of this type of sporads (Zhang et al., 2017). In tripolar spindles, unilateral defects in metaphase II spindle orientation lead to a rejoining of chromatids at one pole and normal separation at the other pole, producing triads with one  $2n$  spore and two  $n$  spores. Due to the abovementioned information, these irregularities are currently the main mechanisms involved in the formation of unreduced gametes (De Storme & Geelen, 2013). In this work, the tripolar spindles detected in anaphase II of Individuals 2, 3 and 6 would explain the presence of telophase II with a nucleus of restitution ( $2n$ ) and two  $n$  nuclei. Consequently, this is consistent with the finding of triads that will produce  $2n$  and  $n$  pollen grains.

### 3.5.3 | Cytomixis

In this work, the migration of chromatin from one of the PMCs to another was observed in all meiotic phases (Table 4, Figure 5h), showing the highest percentages dur-

ing meiotic prophase, up to 100% of the PMCs in prophase I and 66.67% in prophase II. Cytomixis has been proposed as one of the mechanisms of unreduced gamete formation (Dewitte et al., 2012; Mursalimov & Deineko, 2015), as observed in *Dactylis* (Falistocco, Tosti, & Falcinelli, 1995), *Sorghum bicolor* (L.) Moench (Ghaffari, 2006), *Arachis pintoi* Krapov. & W.C. Greg. (Lavia, Ortiz, Robledo, Fernández, & Seijo, 2011), *Triticum* and *Secale* (Sidorchuk, Novikovskaya, & Deineko, 2016), *Lippia alba* (Mill.) N.E. Br. ex Britton & P. Wilson (Reis et al., 2016), and *Lilium* (Zhang et al., 2017).

It has been proposed that the mechanism of formation of unreduced gametes by cytomixis is the development of binucleated meiocytes, formed when an entire nucleus migrates to the recipient cell (Sidorchuk et al., 2016; Mursalimov & Deineko, 2018). These nuclei show no signs of damage in the zygo-pachytene phase and progressively continue meiotic division (Mursalimov & Deineko, 2015), being observable until the nuclear membrane disappears in metaphase I, after which they are undetectable. Consequently, it is likely that a single meiotic spindle will form for both nuclei, leading to the formation of unreduced pollen. Although this has not been experimentally confirmed, it has been demonstrated that in plants with a high cytomixis rate, the pollen produced is more than twice the size of normal pollen (Mursalimov & Deineko, 2018). This allows us to assume that cytomixis could also be a possible cause of the generation of unreduced gametes in the hybrids analyzed.

### 3.5.4 | Restitution nuclei

Our analysis suggests that the high frequency of irregularities such as bridge persistence, tripolar spindle formation, and cytomixis is the meiotic mechanism that would originate restitution nuclei. All hybrid individuals presented evidence of nuclear restitution in both meiotic divisions. The analysis of PMCs showed that nuclear restitution began in the first meiotic division (see above), as restitution nuclei were found in up to 35% of the PMCs in telophase I (Figure 5i), and then observed as a single nucleus in prophase II (Figure 5j) and as a single plate in metaphase II (Figure 5k) in up to 57.14 and 56.14% of the PMCs, respectively. At the end of meiotic division, during anaphase II (Figure 5l) and telophase II (Figure 5m), these unreduced nuclei may either undergo meiotic division and give rise to two  $2n$  nuclei (up to 48.68% of the PMCs) or suffer a second nuclear restitution process, giving rise to  $4n$  restitution nuclei, as those found in the present study (up to 48.49% of the PMCs, Figure 5n). In this sense, Dewitte, Eeckhaut, et al. (2010) showed that *Begonia* ‘Orococo’ mainly produces first division restitution dyads and  $2n$  pollen grains,

as a consequence of an omitted meiosis I. In addition, it has been proposed that, if a complete loss of meiotic cytokinesis occurs, tetranuclear spores are formed, developing  $4n$  gametes (De Storme & Geelen, 2013). Consequently, these mechanisms would explain the finding of both dyads and monads, as well as the formation of  $2n$  and  $4n$  pollen grains.

We also found that the process of nuclear restitution could originate in the second meiotic division, because one nucleus of restitution and two haploid nuclei were found in up to 22.22% of the PMCs in anaphase II (Figure 5o) and 14.47% of the PMCs in telophase II (Figure 5p). This can occur if the chromosomes located on one of the metaphase II plates segregate towards the poles, while the chromosomes on the other plate undergo nuclear restitution (see above), resulting in the formation of triads, and therefore in  $2n$  and  $n$  pollen grains. Several authors proposed that second division restitution (i.e., incomplete separation of nuclei during meiosis II) gives rise to  $2n$  pollen grains (Bretagnolle & Thompson, 1995; Dewitte, Eeckhaut, et al., 2010), as observed in this work.

An important fact to point out is the high production of  $2n$  restitution nuclei found in the [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrid. This phenomenon occurred in up to 57.14% of the PMCs in prophase II, 56.14% of the PMCs in metaphase II, 22.22% of the PMCs in anaphase II, and 48.68% of the PMCs in telophase II. Although the rate of production of dyads and triads and the formation of  $2n$  pollen grains of this hybrid was not high, it should be noted that we were able to analyze the meiosis of a single individual, which, taking into account the records of sporads and pollen of other hybrids of the same cross, could indicate that they may have high rates of production of restitution nuclei. These results suggest that hybrids in which *A. duranensis* was used as the female parent produced a greater number of restitution nuclei.

In summary, the meiotic abnormalities found in the meiosis of intergenomic hybrids between *A. duranensis* and *A. ipaënsis* (persistent bridges, tripolar spindles, and cytomixis) are direct evidence of the formation of restitution nuclei in both the first and second meiotic division. These failures in the regularity of meiosis lead to the formation of monads, dyads, and triads, and consequently to the formation of  $2n$  and  $4n$  gametes.

### 3.6 | Origin of *A. hypogaea*

Several authors have studied the origin of *A. hypogaea* for many years. Based on the morphological characteristics of this species, Krapovickas and Gregory (1994) identified six botanical varieties: two belonging to the

subspecies *hypogaea* (*hypogaea* and *hirsuta*), and four belonging to the subspecies *fastigiata* (*fastigiata*, *vulgaris*, *aequatoriana*, and *peruviana*). The objective of elucidating the origin of peanut is to find out which are its wild progenitors, what type of polyploid it is, whether it was a single event of polyploid origin, where and when the event occurred, and how this event happened. Currently, cytogenetic, phylogenetic, and molecular evidence indicate that the wild species involved in the origin of peanut are *A. duranensis* and *A. ipaënsis* (Bertioli et al., 2016; Grabiele et al., 2012; Kochert et al., 1996; Moretzsohn et al., 2013; Robledo et al., 2009; Seijo et al., 2007). Recent research has also established that *A. hypogaea* is a segmental allotetraploid, and that the six botanical varieties of the crop, as well as the immediate wild ancestor *A. monticola*, have suffered a single event of polyploid origin, involving *A. duranensis* as the female parent and *A. ipaënsis* as the male parent (Bertioli et al., 2019; Grabiele et al., 2012; Seijo et al., 2007, 2018). It has been hypothesized that the probable center of origin of *A. hypogaea* was the south of Bolivia and northwest of Argentina, which is the current distribution area of the diploid species involved in its origin (Dillehay, Rossen, Andres, & Williams, 2007; Grabiele et al., 2012; Krapovickas & Gregory, 1994; Simpson, 2001). Furthermore, analysis of nucleotide divergence has determined that, ~9,400 years ago, *A. ipaënsis* was carried by humans into the area of *A. duranensis*, enabling their hybridization and formation of *A. hypogaea* (Bertioli et al., 2016).

Despite the great efforts made in elucidating all the abovementioned questions, there are no accurate data on how the event of polyploidization of the crop occurred. The model of origin proposed suggests that it was an event of hybridization between wild species followed by a chromosomal duplication or fusion of unreduced gametes, giving rise to a wild tetraploid with two complements of each genome, A and B. After the origin of this wild allotetraploid, *A. hypogaea* arose by domestication. In this context, the ability of [*A. duranensis* × *A. ipaënsis*]<sup>2x</sup> hybrids to produce  $2n$  pollen constitutes relevant evidence to support the fact that the most probable mechanism for the origin of peanut was sexual polyploidization. In this sense, this crop could have originated as a result of the production of a diploid AB hybrid through natural hybridization between *A. duranensis* (AA) and *A. ipaënsis* (BB). This amphiploid would have the ability to produce polyploid individuals either by unilateral sexual polyploidization through the union of a normal gamete ( $n$ ) with an unreduced one ( $2n$ ), triploid bridge, or by bilateral sexual polyploidization through the formation and union of two  $2n$  gametes.

### ACKNOWLEDGMENTS

This research was carried out with funding from Secretaría General de Ciencia y Técnica de la Universidad Nacional

del Nordeste, Corrientes, Argentina (PI 16P003), and the Agencia Nacional de Promoción Científica y Tecnológica, Argentina (PICT-2015-2681). C. Silvestri and A. García received fellowships from CONICET. G.I. Lavia and A.M. Ortiz are research staff members of CONICET.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## ORCID

A. V. García  <https://orcid.org/0000-0002-9441-0478>

A. M. Ortiz  <https://orcid.org/0000-0001-5670-7097>

M. C. Silvestri  <https://orcid.org/0000-0001-8161-2583>

G. I. Lavia  <https://orcid.org/0000-0003-0765-7809>

## REFERENCES

- Akutsu, M., Kitamura, S., Toda, R., Miyajima, I., & Okazaki, K. (2007). Production of 2n pollen of Asiatic hybrid lilies by nitrous oxide treatment. *Euphytica*, 155, 143–152. <https://doi.org/10.1007/s10681-006-9317-y>
- Bajer, A. (1964). Cine-micrographic studies on dicentric chromosomes. *Chromosoma*, 15, 630–651. <https://doi.org/10.1007/BF00319996>
- Bertioli, D. J., Seijo, G., Freitas, F. O., Valls, J. F. M., Leal-Bertioli, S. C. M., & Moretzsohn, M. C. (2011). An overview of peanut and its wild relatives. *Plant Genetic Resources-Characterization and Utilization*, 9, 134–149. <https://doi.org/10.1017/S1479262110000444>
- Bertioli, D. J., Jenkins, J., Clevenger, J., Dudchenko, O., Gao, D., Seijo, G., ... Schmutz, J. (2019). The genome sequence of segmental allotetraploid peanut *Arachis hypogaea*. *Nature Genetics*, 51, 877–884. <https://doi.org/10.1038/s41588-019-0405-z>
- Bertioli, D. J., Cannon, S. B., Froenicke, L., Huang, G., Farmer, A. D., Cannon, E. K. S., ... Ozias-Akins, Peggy (2016). The genome sequences of *Arachis duranensis* and *Arachis ipaënsis*, the diploid ancestors of cultivated peanut. *Nature Genetics*, 48, 438–446. <https://doi.org/10.1038/ng.3517>
- Bretagnolle, F., & Thompson, J. D. (1995). Gametes with the somatic chromosome number: Mechanisms of their formation and role in the evolution of autopolyploid plants. *New Phytologist*, 129, 1–22. <https://doi.org/10.1111/j.1469-8137.1995.tb03005.x>
- Burow, M. D., Simpson, C. E., Faries, M. W., Starr, J. L., & Paterson, A. H. (2009). Molecular biogeographic study of recently described B- and A-genome *Arachis* species, also providing new insights into the origins of cultivated peanut. *Genome*, 52, 107–119. <https://doi.org/10.1139/G08-094>
- Darlington, C. D. (1937). *Recent advances in cytology* (2nd ed.). London: J. and A. Churchill.
- De Storme, N., & Geelen, D. (2013). Sexual polyploidization in plants: Cytological mechanisms and molecular regulation. *New Phytologist*, 198, 670–684. <https://doi.org/10.1111/nph.12184>
- den Nijs, T. P. M., & Peloquin, S. J. (1977). Polyploid evolution via 2n gametes. *American Potato Journal*, 54, 377–386. <https://doi.org/10.1007/BF02852758>
- Dewitte, A., Eeckhaut, T., Van Huylenbroeck, J., & Van Bockstaele, E. (2010). Meiotic aberrations during 2n pollen formation in *Begonia*. *Heredity*, 104, 215–223. <https://doi.org/10.1038/hdy.2009.111>
- Dewitte, A., Van Laere, K., & Van Huylenbroeck, J. (2012). Use of 2n gametes in plant breeding. In I. Y. Abdurakhmonov (Ed.), *Plant breeding* (pp. 59–86). Rijeka, Croatia: InTech Open Access Publisher. <https://doi.org/10.5772/29827>
- Dewitte, A., Van Laere, K., Van Huylenbroeck, J., & Van Bockstaele, E. (2010). Inheritance of 2n pollen formation in an F<sub>1</sub> and F<sub>2</sub> population of begonia hybrids. *Acta Horticulturae*, 855, 95–100. <https://doi.org/10.17660/ActaHortic.2010.855.11>
- Di Rienzo, J. A., Casanoves, F., Balzarini, M. G., Gonzalez, L., Tablada, M., & Robledo, C. W. (2017). *InfoStat*. Córdoba, Argentina: Universidad Nacional de Córdoba.
- Dillehay, T. D., Rossen, J., Andres, T. C., & Williams, D. E. (2007). Pre-ceramic adoption of peanut, squash, and cotton in northern Peru. *Science*, 316, 1890–1893. <https://doi.org/10.1126/science.1141395>
- Dyer, A. F. (1963). The use of lacto-propionic orcein in rapid squash methods for chromosome preparations. *Stain Technology*, 38, 85–90. <https://doi.org/10.3109/10520296309061169>
- Falisticco, E., Tosti, N., & Falcinelli, M. (1995). Cytomixis in pollen mother cells of diploid *Dactylis*, one of the origins of 2n gametes. *Journal of Heredity*, 86, 448–453. <https://doi.org/10.1093/oxfordjournals.jhered.a111619>
- Fávero, A. P., Simpson, C. E., Valls, J. F. M., & Vello, N. A. (2006). Study of the evolution of cultivated peanut through crossability studies among *Arachis ipaënsis*, *A. duranensis*, and *A. hypogaea*. *Crop Science*, 46, 1546–1552. <https://doi.org/10.2135/cropsci2005.09-0331>
- Fernández, A., & Krapovickas, A. (1994). Cromosomas y evolución en *Arachis* (Leguminosae). *Bonplandia*, 8, 187–220.
- García, A. V., Ortiz, A. M., & Lavia, G. I. (2016). Origen genético de *Arachis hypogaea* L.: Evidencias citogenéticas de poliploidización sexual. *Journal of Basic and Applied Genetics*, 27, 241. Retrieved from [https://sag.org.ar/jbag/wp-content/uploads/2020/01/V.XXXVIII\\_2016\\_SupplI\\_19092016.pdf](https://sag.org.ar/jbag/wp-content/uploads/2020/01/V.XXXVIII_2016_SupplI_19092016.pdf)
- Gerber, G., & Schweizer, D. (1988). Cytochemical heterochromatin differentiation in *Sinapis alba* (Cruciferae) using a simple air-drying technique for producing chromosome spreads. *Plant Systematics and Evolution*, 158, 97–106. <https://doi.org/10.1007/BF00936336>
- Ghaffari, S. M. (2006). Occurrence of diploid and polyploid microspores in *Sorghum bicolor* (Poaceae) is the result of cytomixis. *African Journal of Biotechnology*, 5, 1450–1453. <https://doi.org/10.4314/ajb.v5i16.43137>
- Grabiele, M., Chalup, L., Robledo, G., & Seijo, G. (2012). Genetic and geographic origin of domesticated peanut as evidenced by 5S rDNA and chloroplast DNA sequences. *Plant Systematics and Evolution*, 298, 1151–1165. <https://doi.org/10.1007/s00606-012-0627-3>
- Gregory, W. C., & Gregory, M. P. (1976). Groundnut. In N. W. Simmonds (Ed.), *Evolution of crop plants* (pp. 151–154). Harlow, UK: Longman Group.
- Guan, J. Z., Wang, J. J., Cheng, Z. H., Liu, Y., & Li, Z. Y. (2012). Cytomixis and meiotic abnormalities during microsporogenesis are responsible for male sterility and chromosome variations in *Houttuynia cordata*. *Genetics and Molecular Research*, 11, 121–130. <https://doi.org/10.4238/2012.January.17.2>
- Holley, C. E., & Geerts, P. G. (2009). Multiplex Manager 1.0: A cross-platform computer program that plans and optimizes

- multiplex PCR. *Biotechniques*, 46, 511–517. <https://doi.org/10.2144/000113156>
- Husted, L. (1933). Cytological studies on the peanut, *Arachis*. I. Chromosome number and morphology. *Cytologia*, 5, 109–117. <https://doi.org/10.1508/cytologia.5.109>
- Husted, L. (1936). Cytological studies on the peanut, *Arachis*. II. Chromosome number and morphology and behavior, and their application to the problem of the origin of the cultivated forms. *Cytologia*, 7, 396–423. <https://doi.org/10.1508/cytologia.7.396>
- Inglis, P. W., de, M., Pappas, C. R., Resende, L. V., & Grattapaglia, D. (2018). Fast and inexpensive protocols for consistent extraction of high quality DNA and RNA from challenging plant and fungal samples for high-throughput SNP genotyping and sequencing applications. *PLOS ONE*, 13(10). <https://doi.org/10.1371/journal.pone.0206085>
- Jauhar, P. P. (2003). Formation of 2n gametes in durum wheat haploids: Sexual polyploidization. *Euphytica*, 133, 81–94. <https://doi.org/10.1023/A:1025692422665>
- Jauhar, P. P. (2007). Meiotic restitution in wheat polyhaploids (amphihaploids): A potent evolutionary force. *Journal of Heredity*, 98, 188–193. <https://doi.org/10.1093/jhered/esm011>
- Kochert, G., Halward, T., Branch, W. D., & Simpson, C. E. (1991). RFLP variability in peanut (*Arachis hypogaea* L.) cultivars and wild species. *Theoretical and Applied Genetics*, 81, 565–570. <https://doi.org/10.1007/BF00226719>
- Kochert, G., Stalker, H. T., Gimenes, M., Galgano, L., Lopes, C. R., & Moore, K. (1996). RFLP and cytogenetic evidence on the origin and evolution of allotetraploid domesticated peanut, *Arachis hypogaea* (Leguminosae). *American Journal of Botany*, 83, 1282–1291. <https://doi.org/10.1002/j.1537-2197.1996.tb13912.x>
- Kovalsky, I. E., & Solís Neffa, V. G. (2012). Evidence of 2n microspore production in a natural diploid population of *Turnera sidoides* subsp. *carnea* and its relevance in the evolution of the *T. sidoides* (Turneraceae) autopolyploid complex. *Journal of Plant Research*, 125, 725–734. <https://doi.org/10.1007/s10265-012-0493-7>
- Krapovickas, A., & Gregory, W. C. (1994). Taxonomía del género *Arachis* (Leguminosae). *Bonplandia*, 8, 1–186.
- Kreiner, J. M., Kron, P., & Husband, B. C. (2017a). Evolutionary dynamics of unreduced gametes. *Trends in Genetics*, 33, 583–593. <https://doi.org/10.1016/j.tig.2017.06.009>
- Kreiner, J. M., Kron, P., & Husband, B. C. (2017b). Frequency and maintenance of unreduced gametes in natural plant populations: Associations with reproductive mode, life history and genome size. *New Phytologist*, 214, 879–889. <https://doi.org/10.1111/nph.14423>
- Lavia, G. I., Fernández, A., & Seijo, J. G. (2008). Cytogenetic and molecular evidences on the evolutionary relationships among *Arachis* species. In A. K. Sharma & A. Sharma (Eds.), *Plant genome: Biodiversity and evolution* (pp. 101–134). Enfield, NH: Science Publishers.
- Lavia, G. I., Ortiz, A. M., Robledo, G., Fernández, A., & Seijo, G. (2011). Origin of triploid *Arachis pintoii* (Leguminosae) by autopolyploidy evidenced by FISH and meiotic behaviour. *Annals of Botany*, 108, 103–111. <https://doi.org/10.1093/aob/mcr108>
- Mallikarjuna, N., Senthilvel, S., & Hoisington, D. (2011). Development of new sources of tetraploid *Arachis* to broaden the genetic base of cultivated groundnut (*Arachis hypogaea* L.). *Genetic Resources and Crop Evolution*, 58, 889–907. <https://doi.org/10.1007/s10722-010-9627-8>
- Mason, A. S., Nelson, M. N., Yan, G., & Cowling, W. A. (2011). Production of viable male unreduced gametes in *Brassica* interspecific hybrids is genotype specific and stimulated by cold temperatures. *BMC Plant Biology*, 11. <https://doi.org/10.1186/1471-2229-11-103>
- Moretzsohn, M. C., Gouvea, E. G., Inglis, P. W., Leal-Bertioli, S. C. M., Valls, J. F. M., & Bertioli, D. J. (2013). A study of the relationships of cultivated peanut (*Arachis hypogaea*) and its most closely related wild species using intron sequences and microsatellite markers. *Annals of Botany*, 111, 113–126. <https://doi.org/10.1093/aob/mcs237>
- Mursalimov, S. R., & Deineko, E. V. (2015). How cytomixis can form unreduced gametes in tobacco. *Plant Systematics and Evolution*, 301, 1293–1297. <https://doi.org/10.1007/s00606-014-1150-5>
- Mursalimov, S., & Deineko, E. V. (2018). Cytomixis in plants: Facts and doubts. *Protoplasma*, 255, 719–731. <https://doi.org/10.1007/s00709-017-1188-7>
- Orjeda, G., Freyre, R., & Iwanaga, M. (1990). Production of 2n pollen in diploid *Ipomoea trifida*, a putative wild ancestor of sweet potato. *Journal of Heredity*, 81, 462–467. <https://doi.org/10.1093/oxfordjournals.jhered.a111026>
- Ortiz, A. M., Seijo, J. G., Fernández, A., & Lavia, G. I. (2011). Meiotic behavior and pollen viability of tetraploid *Arachis glabrata* and *A. nitida* species (Section Rhizomatosae, Leguminosae): Implications concerning their polyploid nature and seed set production. *Plant Systematics and Evolution*, 292, 73–83. <https://doi.org/10.1007/s00606-010-0397-8>
- Pagliarini, M. S. (2000). Meiotic behavior of economically important plant species: The relationship between fertility and male sterility. *Genetics and Molecular Biology*, 23, 997–1002. <https://doi.org/10.1590/S1415-47572000000400045>
- Pittenger, T. H., & Frolik, E. F. (1951). Temporary mounts for pollen abortion determinations. *Stain Technology*, 26, 181–184. <https://doi.org/10.3109/10520295109113205>
- Qiu, Y., Liao, L., Liu, S., Mao, D., & Liu, R. (2017). Differences on the microsporogenesis and tapetal development of male fertile and cytoplasmic male sterile pepper (*Capsicum annuum* L.). *Grana*, 56, 215–227. <https://doi.org/10.1080/00173134.2016.1248860>
- Ramsey, J., & Schemske, D. W. (1998). Pathways, mechanisms, and rates of polyploid formation in flowering plants. *Annual Review of Ecology, Evolution, and Systematics*, 29, 467–501. <https://doi.org/10.1146/annurev.ecolsys.29.1.467>
- Rasband, W. S. (1997). *ImageJ*. Bethesda, MD: National Institutes of Health. Retrieved from <http://imagej.nih.gov/ij>
- Reis, A. C., Sousa, S. M., & Viccini, L. F. (2016). High frequency of cytomixis observed at zygotene in tetraploid *Lippia alba*. *Plant Systematics and Evolution*, 302, 121–127. <https://doi.org/10.1007/s00606-015-1249-3>
- Rim, Y. W., & Beuselinck, P. R. (1996). Cytology of 2 N pollen formation and pollen morphology in diploid *Lotus tenuis* (Fabaceae). *American Journal of Botany*, 83, 1057–1062. <https://doi.org/10.1002/j.1537-2197.1996.tb12803.x>
- Risso-Pascotto, C., Pagliarini, M. S., & Do Valle, C. B. (2005). Multiple spindles and cellularization during microsporogenesis in an artificially induced tetraploid accession of *Brachiaria ruziziensis* (Gramineae). *Plant Cell Reports*, 23, 522–527. <https://doi.org/10.1007/s00299-004-0867-y>
- Robledo, G., Lavia, G. I., & Seijo, G. (2009). Species relations among wild *Arachis* species with the A genome as revealed by FISH mapping of rDNA loci and heterochromatin detection. *Theoretical and*



- Applied Genetics*, 118, 1295–1307. <https://doi.org/10.1007/s00122-009-0981-x>
- Robledo, G., & Seijo, G. (2010). Species relationships among the wild B genome of *Arachis* species (section *Arachis*) based on FISH mapping of rDNA loci and heterochromatin detection: A new proposal for genome arrangement. *Theoretical and Applied Genetics*, 121, 1033–1046. <https://doi.org/10.1007/s00122-010-1369-7>
- Santana, S. H., & Valls, J. F. M. (2015). *Arachis veigae* (Fabaceae), the most dispersed wild species of the genus, and yet taxonomically overlooked. *Bonplandia*, 24, 139–150.
- Schweizer, D. (1976). Reverse fluorescent chromosome banding with chromomycin and DAPI. *Chromosoma*, 58, 307–324. <https://doi.org/10.1007/BF00292840>
- Seijo, J. G., Kovalsky, E. I., Chalup, L. M. I., Samoluk, S. S., Fávero, A., & Robledo, G. A. (2018). Karyotype stability and genome-specific nucleolar dominance in peanut, its wild 4× ancestor, and a synthetic AABB polyploid. *Crop Science*, 58, 1671–1683. <https://doi.org/10.2135/cropsci2018.02.0088>
- Seijo, J. G., Lavia, G. I., Fernández, A., Krapovickas, A., Ducasse, D. A., Bertoli, D. J., & Moscone, E. A. (2007). Genomic relationships between the cultivated peanut (*Arachis hypogaea*, Leguminosae) and its close relatives revealed by double GISH. *American Journal of Botany*, 94, 1963–1971. <https://doi.org/10.3732/ajb.94.12.1963>
- Seijo, J. G., Lavia, G. I., Fernández, A., Krapovickas, A., Ducasse, D., & Moscone, E. A. (2004). Physical mapping of the 5S and 18S-25S rRNA genes by FISH as evidence that *Arachis duranensis* and *A. ipaënsis* are the wild diploid progenitors of *A. hypogaea* (Leguminosae). *American Journal of Botany*, 91, 1294–1303. <https://doi.org/10.3732/ajb.91.9.1294>
- Seijo, J. G., & Solís Neffa, V. G. (2006). Cytogenetic studies in the rare South American *Lathyrus hasslerianus* Burk. (Leguminosae). *Cytologia*, 71, 11–19. <https://doi.org/10.1508/cytologia.71.11>
- Sidorchuk, Y. V., Novikovskaya, A. A., & Deineko, E. V. (2016). Cytomixis in the cereal (Gramineae) microsporogenesis. *Protoplasma*, 253, 291–298. <https://doi.org/10.1007/s00709-015-0807-4>
- Simpson, C. E. (2001). Use of wild *Arachis* species/introgression of genes into *A. hypogaea* L. *Peanut Science*, 28, 114–116. <https://doi.org/10.3146/10095-3679-28-2-12>
- Simpson, C. E. (2017). Finally, the cross that made *Arachis monticola* Krapov. & Rigoni and/or *Arachis hypogaea* L. In T. Baughman & K. Cutchins (Eds.), *49th Proceedings of the American Peanut Research and Education Society, Inc.* (p. 66). Tifton, GA: American Peanut Research and Education Society.
- Smartt, J., Gregory, W. C., & Gregory, M. P. (1978). The genomes of *Arachis hypogaea*. 1. Cytogenetic studies of putative genome donors. *Euphytica*, 27, 665–675. <https://doi.org/10.1007/BF00023701>
- Stalker, H. T., Dhese, J. S., Parry, D. C., & Hahn, J. H. (1991). Cytological and interfertility relationships of *Arachis* section *Arachis*. *American Journal of Botany*, 78, 238–246. <https://doi.org/10.1002/j.1537-2197.1991.tb15751.x>
- Tallury, S., Hilu, K., Milla, S., Friend, S., Alsaghir, M., Stalker, H. T., & Quandt, D. (2005). Genomic affinities in *Arachis* section *Arachis* (Fabaceae): Molecular and cytogenetic evidence. *Theoretical and Applied Genetics*, 111, 1229–1237. <https://doi.org/10.1007/s00122-005-0017-0>
- Tilquin, J. P., de Brouwer, K., & Horvat, F. (1984). Unusual cytological patterns in microsporogenesis in a cultivar of *Fuchsia*. *Theoretical and Applied Genetics*, 67, 413–417. <https://doi.org/10.1007/BF00263403>
- Tondini, F., Tavoletti, S., Mariani, A., & Veronesi, F. (1993). A statistical approach to estimate the frequency of n, 2n and 4n pollen grains in diploid alfalfa. *Euphytica*, 69, 109–114. <https://doi.org/10.1007/BF00021733>
- Valls, J. F. M., & Simpson, C. E. (2005). New species of *Arachis* (Leguminosae) from Brasil, Paraguay, and Bolivia. *Bonplandia*, 14, 35–64. <http://doi.org/10.30972/bon.141-21387>
- Valls, J. F. M., & Simpson, C. E. (2017). A new species of *Arachis* (Fabaceae) from Mato Grosso, Brasil, related to *Arachis matiensis*. *Bonplandia*, 26, 143–149. <https://doi.org/10.30972/bon.2622575>
- Valls, J. F. M., Da Costa, L. C., & Custodio, A. R. (2013). A novel trifoliolate species of *Arachis* (Fabaceae) and further comments on taxonomic section *Trierectoides*. *Bonplandia*, 22, 91–97.
- Van Campo, M. (1957). Palynologie africaine I. *Bulletin de l'Institut Francais d'Afrique Noire*, 19, 659–678.
- Wang, J., Kang, X., & Zhu, Q. (2010). Variation in pollen formation and its cytological mechanism in an allotriploid white poplar. *Tree Genetics & Genomes*, 6, 281–290. <https://doi.org/10.1007/s11295-009-0248-3>
- Wang, J., You, H., Tian, J., Wang, Y., Liu, M., & Duan, W. (2015). Abnormal meiotic chromosome behavior and gametic variation induced by intersectional hybridization in *Populus* L. *Tree Genetics & Genomes*, 11, 1–11. <https://doi.org/10.1007/s11295-015-0880-z>
- Zhang, X., Cao, Q., Zhou, P., & Jia, G. (2017). Meiotic chromosome behavior of the male-fertile allotriploid lily cultivar ‘Cocossa’. *Plant Cell Reports*, 36, 1641–1653. <https://doi.org/10.1007/s00299-017-2180-6>
- Zhang, Z., & Kang, X. (2010). Cytological characteristics of numerically unreduced pollen production in *Populus tomentosa* Carr. *Euphytica*, 173, 151–159. <https://doi.org/10.1007/s10681-009-0051-0>

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** García AV, Ortiz AM, Silvestri MC, Custodio AR, Moretzsohn MC, Lavia GI. Occurrence of 2n microspore production in diploid interspecific hybrids between the wild parental species of peanut (*Arachis hypogaea* L., Leguminosae) and its relevance in the genetic origin of the cultigen. *Crop Science*. 2020;60:2420–2436. <https://doi.org/10.1002/csc2.20233>