

## Variants in *GH*, *IGF1*, and *LEP* genes associated with body traits in Santa Inês sheep

Alessandro Lima Machado<sup>1</sup>, Ariana Nascimento Meira<sup>1</sup>, Adriana de Farias Jucá<sup>1</sup>, Hymerson Costa Azevedo<sup>2</sup>, Evandro Neves Muniz<sup>2</sup>, Luiz Lehmann Coutinho<sup>3</sup>, Gerson Barreto Mourão<sup>3</sup>, Victor Breno Pedrosa<sup>4</sup>, Luís Fernando Batista Pinto<sup>1\*</sup>

<sup>1</sup>Universidade Federal da Bahia – Depto. de Zootecnia, Av. Adhemar de Barros, 500 – 40170-110 – Salvador, BA – Brasil.

<sup>2</sup>Embrapa Tabuleiros Costeiros, Av. Beira Mar, 3250 – 49025-040 – Aracaju, SE – Brasil.

<sup>3</sup>Universidade de São Paulo/ESALQ – Depto. de Zootecnia, Av. Pádua Dias, 11 – 13418-900 – Piracicaba – SP – Brasil.

<sup>4</sup>Universidade Estadual de Ponta Grossa – Depto. de Zootecnia, Av. General Carlos Cavalcanti, 4748 – 84030-900 – Ponta Grossa, PR – Brasil.

\*Corresponding author <luisfbp@ufba.br>

Edited by: Paulo Cesar Sentelhas

Received August 10, 2019

Accepted November 11, 2019

**ABSTRACT:** Growth hormone (*GH*), insulin-like growth factor 1 (*IGF1*), and leptin (*LEP*) can be candidate genes for association studies because they play vital roles in the metabolism process. Thus, this study aimed to identify variants in these genes associated with body traits in Santa Inês sheep. The following were recorded: body weight at 100 (BW100) and 240 days (BW240), average daily gain (ADG), withers (WH) and croup (CH) heights, body length (BL), thoracic (TG) and leg (LG) girths, thoracic (TW) and croup (CW) widths, body depth (BD), rib eye area (REA), fat thickness (FT), and carcass finishing score (CFS). Single-locus association analysis was performed with 11 variants in *IGF1*, 18 in *LEP*, and 16 in *GH*. Moreover, two haplotypes in *IGF1* and one haplotype in *LEP* were evaluated in haplotype association analysis. The single-locus analysis revealed 23 suggestive additive effects ( $p < 0.05$ ), but no additive effect was found at the Bonferroni threshold. Haplotype association analysis revealed 19 additive effects, of which ten were at the Bonferroni threshold ( $p < 0.0074$ ). In *IGF1* gene, haplotype replacements were associated with ADG 20.51(7.37), CH 4.09(1.21), WH 3.52(1.20), BL 3.94(1.19), TG 3.88(1.30), TW 1.13(0.36), and LG 3.40(1.08); while in the *LEP* gene the haplotype replacement was associated with BW100 1.83(0.51), BD -2.51(0.56), and CFS -0.24(0.06). Therefore, there are haplotypes in *IGF1* and *LEP* genes associated with body traits in Santa Inês sheep, which can be useful in marker-assisted selection.

**Keywords:** hormone, carcass, growth, haplotype, ovine

## Introduction

The Callipyge, Carwell, Double Muscling (Cockett et al., 2005), and Myostatin (Hickford et al., 2010) are major genes that affect growth and carcass traits in sheep. However, the variants in these genes do not explain total genetic variation related to either growth or carcass traits in sheep. Additionally, certain polymorphisms in these genes are typical of a number of specific sheep breeds (Cockett et al., 2005). Therefore, it is essential to study other candidate genes. In this context, the growth hormone (*GH*), insulin-like growth factor 1 (*IGF1*), and leptin (*LEP*) genes can be candidates for association studies with either growth or carcass traits in sheep. The transcripts of these genes play vital roles in the metabolism process (Tuersunjiang et al., 2016). Consequently, these genes can explain the genetic variance of several phenotypes in livestock.

The *GH* gene encodes the growth hormone which, through its specific receptor (Sahu et al., 2017), is the major regulator of IGF-I synthesis in the liver (Teran et al., 2016). IGF-1 is the main mediator of the effects of the growth hormone on muscle and bone tissues. Therefore, the GH/IGF is the major axis controlling the growth of animals (Yakar et al., 2018). A number of polymorphisms in *GH* were associated with growth (Abdelmoneim et al., 2017) and carcass traits (Gorlov et al., 2017) in sheep. In addition, associations between variants in the *IGF1* gene with sheep growth traits were reported (Trukhachev et al., 2016).

Leptin has multiple functions, such as regulating appetite, body weight, maturation of the reproductive

axis, neuroendocrine adaptations to fasting, and regulating glucose homeostasis (Caron et al., 2018). The vital role of leptin in controlling appetite can cause an impact on the variables directly associated with food intake, such as body weight and fat deposition. Thus, variants in the *LEP* gene were associated with growth (Hajihosseini et al., 2012), morphometric (Bakhtiar et al., 2017), and carcass traits (Barzehkar et al., 2009) in different sheep breeds.

Certain variants in the *IGF1* gene were associated with internal carcass length, rib yield, and neck weight in Santa Inês sheep; while a variant in the *GH* gene was associated with weights and yields of primal cuts such as rib, loin, leg, and neck (Meira et al., 2019). Additive effects of polymorphisms in the *LEP* gene were also reported for weights and yields of these primal cuts, as well as weights and yields of the carcass (Meira et al., 2018). However, growth, morphometric, and in vivo ultrasound carcass traits were not evaluated in previous association studies on *IGF1*, *GH*, and *LEP* genes in Santa Inês sheep. Thus, this study aimed to identify polymorphisms in the *GH*, *IGF*, and *LEP* genes associated with body weight, morphometric traits, and carcass in vivo traits in Santa Inês sheep.

## Materials and Methods

### Animals and phenotypes

This study was carried out with the approval of the Ethical Committee for Animal Use from Veterinary Medicine and Animal Science School of the Federal University of Bahia (protocol number 02/2010).

Deoxyribonucleic Acid (DNA) was extracted from 192 Santa Inês lambs, all males, of which 74 were born in 2010, 15 in 2011, and 17 in 2012, in Frei Paulo, SE, Brazil (10°32'56" S, 37°32'02" W, altitude of 277 m). The other 86 lambs were raised in 2014 in São Gonçalo dos Campos, BA, Brazil (12°25'58" S, 38°58'01" W, altitude of 233 m). All animals received diets formulated according to the National Research Council (NRC, 2007) to meet the nutritional requirements for lambs with estimated weight gains of 170 g d<sup>-1</sup>.

The morphometric traits were measured in lambs at 240 days of age using a tape and a measuring stick. Withers (WH) (distance from the highest point of the thoracic vertebra to the ground) and the croup (CH) (distance from the coxal tuberosity to the ground) heights; body length (BL) (distance from supraglenoid tuberosity to sciatic tuberosity); thoracic (TW) (distance between the supraglenoid tuberosities) and croup (CW) (distance between the coxal tuberosities) widths; thoracic (TG) (contour of the thoracic cavity, adjacent to the shoulder blades) and leg (LG) (mid-thigh contour) girths; and the body depth (BD) (distance from the thoracic vertebrae to the sternum) were the morphometric traits evaluated.

The carcass finishing score (CFS) was accessed *in vivo* with values from 1 to 5, by a single recorder as follows: score 1 (very lean carcass), 2 (lean), 3 (medium), 4 (fat), and 5 (very fat). Additionally, ultrasonography was performed to obtain the rib eye area (REA) and fat thickness (FT) at 240 days of age. The REA and FT ultrasound images, between the 12<sup>th</sup> and 13<sup>th</sup> ribs on the left side of the lambs, were recorded. The length (A) and maximum depth (B) of the muscles were measured for estimating the REA using the equation  $\frac{A}{2} * \frac{B}{2} * \pi$ . Furthermore, the average daily gain (g d<sup>-1</sup>) was calculated using the difference between the body weights obtained at 100 (weaning) and 240 days of age, divided by the number of days between these measurements. A descriptive statistical summary of the traits can be found in Table 1.

### Genotyping

For the extraction of DNA a sample of 5 mL of blood was collected in vacutainer tubes containing Ethylenediaminetetraacetic acid (EDTA). The DNA extraction was performed using a salt precipitation method and proteinase K solutions following the Embrapa protocol (Oliveira et al., 2007). The primer design for amplification of the genes was determined by observing the available sequence in the National Center for Biotechnology Information (NCBI) database for *LEP* (GeneID:443534), *IGF1* (GeneID:443318), and *GH* (GeneID:443329) genes of the sheep genome (version Oar\_v4.0). The oligonucleotides were designed using the Primer 3 software package (<http://bioinfo.ut.ee/primer3-0.4.0/>). Net Primer was used to test the quality of the sequences. After selecting the forward and reverse primers, the Basic Local Alignment Search Tool (BLAST) was used for sequence alignment in NCBI (<http://blast.ncbi.nlm.nih.gov/Blast.cgi>) and to

**Table 1** – Sample size (N), mean and standard deviation (SD) of the growth and carcass traits in Santa Inês sheep.

Traits	N	Mean	SD
Withers height (cm)	184	66.26	5.69
Croup height (cm)	184	66.93	5.64
Body length (cm)	184	56.27	8.94
Thoracic width (cm)	184	17.77	2.11
Croup width (cm)	180	15.77	3.36
Leg girth (cm)	184	40.47	8.23
Thoracic girth (cm)	184	73.21	4.76
Body depth (cm)	180	25.23	2.13
Body weight at 100 days (kg)	172	20.56	4.15
Body weight at 240 days (kg)	184	34.03	6.27
Average daily gain (g d <sup>-1</sup> )	171	136.99	62.11
Rib eye area (cm <sup>2</sup> )	97	7.165	1.64
Fat thickness (cm)	99	0.198	0.04
Carcass finish score	181	2.32	0.37

confirm the similarity with the *Ovis aries* sequence. The forward (*GTGCTGCTTTTGTGATTCTTG*) and reverse (*GATAGAAGAGATGCGAGGAGGA*) primers were used for the amplification of 4,550 bp of *IGF1* between the positions 171037883 and 171113228. For the *GH* gene, the forward (*GCTGCTGACACCTTCAAAGA*) and reverse (*TGACCCCTCAGGTACGTCTCC*) primers were used for the amplification of 1,194 bp between the positions 47485651 and 47487536. For the *LEP* gene, the forward (*GGACCCCTGTACCGATTCCCT*) and reverse (*CAAACCTCAGGAGGGTGA*) primers were used to amplify 2,045 bp located between the positions 92501195 and 92503239.

Touchdown-PCR were performed on the three genes, using 15 µL of the reaction mixture, containing 0.3 mM of each of the primer, Taq polymerase, and 100 ng of the template DNA. The PCR conditions for *LEP* gene were as follows: initial denaturation of 98 °C for 5 min, followed by 20 cycles with denaturation at 98 °C for 10 s, annealing at 67 °C down to 57 °C, varying at -0.5 °C at each cycle for 30 s, and extension at 72 °C for 3 min. Immediately after the initial cycles, another 20 cycles were followed with denaturation at 98 °C for 10 s, followed by annealing at 57 °C for 30 s, extension at 72 °C for 3 min, and a final extension of 72 °C for 5 min. For the *IGF1*, an initial denaturation 98 °C/5 min, followed by ten denaturation cycles at 98 °C/10 s, annealing at 61 °C to 56 °C, reducing -0.5 °C at each cycle, for 30 s, and extension at 72 °C/4 min. Following another 30 cycles with denaturation temperature at 98 °C/10 s, annealing at 56 °C/30 s, and extension at 72 °C/4 min, ending with an extension at 72 °C/5 min. For *GH* amplification, the initial denaturation at 98 °C/5 min, followed by 20 denaturation cycles at 98 °C/10 s, annealing at 63 °C to 53 °C, reducing -0.5 °C at each cycle, for 30 s, and extension at 72 °C/1 min. A further 20 cycles with denaturation temperature at 98 °C/10 s were followed by annealing at 53 °C/30 s, and extension at 72 °C/1 min, ending with an extension at 72 °C/5 min.

After PCR, the amplicons were purified with magnetic beads, and the recommended volume of AgencourtAMPure XP (Beckman Coulter, Brea, USA) was used to homogenize the beads to bind to the amplified products. Immediately after this step, the samples were purified with 70 % ethanol to remove the contaminants. The samples were quantified with Qubit® fluorometer (Life Technologies, Carlsbad, USA) and diluted to 0.2 ng  $\mu\text{L}^{-1}$  for library preparation. The Nextera® XT DNA sample preparation and the Nextera® XT index (Illumina, San Diego, USA) were used to prepare the library. All steps performed followed the Nextera XT protocol. Sequencing was performed on the MiSeq platform (Illumina, San Diego, USA) using the MiSeq Reagent Kit v2 (500 cycles).

The qualities of the reads were verified by the FastQC software program ([https://dnacore.missouri.edu/PDF/FastQC\\_Manual.pdf](https://dnacore.missouri.edu/PDF/FastQC_Manual.pdf)). For the first data filtering, the SeqyClean software tool version 1.3.12 (Zhbannikov et al., 2013) was used, adopting a quality parameter of 24 (Phred score) for each base and a minimum length of 50 bp. Subsequently, the reads were aligned against the reference sheep genome deposited in the NCBI (version Oar\_v4.0) by using the Bowtie2 program (Langmead and Salzberg, 2012). Finally, the functional annotation was performed using the variant effect predictor (VEP) for the online annotation of Ensembl in order to identify the locations of the mutations across different regions of the genome and the possible functional effects of the variants. For the nomenclature of the variants, we followed the recommendations of the Human Genome Variation Society (HGVs).

### Haplotype and Hardy-Weinberg Equilibrium

The Hardy-Weinberg Equilibrium (HWE) was tested by comparing the predicted and observed heterozygosities. The predicted heterozygosity (PH) was obtained using the equation:  $PH = 2 * (1 - MAF) * MAF$ , where MAF was the minor allelic frequency. The Haploview software (version 4.2) was used to test the HWE and to search for linkage disequilibrium (LD) blocks. The haplotype analysis revealed two LD blocks in *IGF1* and one LD block in *LEP*, but no LD blocks were found in the *GH* gene.

### Association analysis

Before the association analysis, all traits had been analyzed using the model:  $y_{ijkl} = \mu + F_i + Y_j + M_k + \alpha_{ijkl}(A) + \varepsilon_{ijkl}$  where  $y_{ijkl}$  is the phenotypic value,  $\mu$  the general average,  $F_i$  the farm effect,  $Y_j$  the year of birth effect,  $M_k$  the month of birth effect,  $\alpha_{ijkl}(A)$  the covariate animal's age, and  $\varepsilon_{ij}$  the error term. This analysis aims to identify possible record errors and test the assumptions of the analysis of variance (ANOVA). The MIXED procedure in SAS (Statistical Analysis System, version 9.3) was used in ANOVA. All the fixed effects tested were significant ( $p < 0.01$ ). Therefore,

it is important to include them in the association analysis models. Also, a principal component analysis to evaluate structuration was performed as reported by Price et al. (2006). The result (Figure 1) did not indicate structuration.

The single-locus association analysis was carried out using the software Qxpack 5 tool (Pérez-Enciso and Misztal, 2011), which performs a likelihood ratio test. The general model can be described as  $y = \beta X + \sum_{k=1}^n Z\delta_k + \varepsilon$ , where  $y$  is the vector containing the records of the traits,  $\beta$  the vector of solutions for the fixed effects,  $\delta$  the vector of solutions for the genetic effects for any of the  $n$  QTL (quantitative trait loci) that affect the trait, and  $\varepsilon$  the vector of the residues.  $X$  and  $Z$  are the incidence matrices that correlate observations in  $y$  to the solutions in the vectors in  $\beta$  and  $\delta_k$ , respectively. The fixed effects included in the model were: (a) farm (2 levels), (b) year (4 levels), (c) month of birth (12 levels), and (d) the covariate age of the animal. The additive and dominance effects of the QTLs were also tested. The additive effect was calculated as the contrast between the genotypes ( $SS - RR$ ), where the allelic variant ( $R$ ) was found in the reference gene sequence, while ( $S$ ) is the allelic variant found in Santa Inês sheep. The dominance effect was calculated as the contrast

$$\left[ SR - \left( \frac{SS + RR}{2} \right) \right].$$

Only variants with MAF  $\geq 2$  % and HWE ( $p \geq 0.05$ ) were used in this analysis. For the single-locus analysis, forty-five variants (18 in *LEP*, 11 in *IGF1*, and 16 in *GH*) that showed HWE ( $p > 0.05$ ) and MAF  $\geq 1$  % were employed (Table 2). Thus, the Bonferroni threshold for single-locus analysis was 0.0005.

The haplotype association analysis was performed as reported by Lake et al. (2003), and the haplo.glm subroutine of the haplo.stat package (version 1.7.7) was utilized in this analysis. Only haplotypes with a frequency greater than 4 % were tested. The haplotype analysis revealed two LD blocks in *IGF1*, one in *LEP*, and no LD blocks were found in the *GH* gene. In the haplotype association analysis, three LD blocks were used (Table 3). Therefore, the Bonferroni threshold was 0.0074.




Figure 1 – The first (PC1) and second (PC2) principal components.

**Table 2** – Variants identified in the *GH*, *IGF1* and *LEP* genes in Santa Inês sheep.

Gene	NCBI Number	Site	Heterozygosity		HWE <sup>1</sup> (p-value)	MAF <sup>2</sup>
			Observed	Expected		
<i>GH</i>	rs1135847308	Exon 5	0.016	0.026	5.23E-02	0.013
<i>GH</i>	rs397514078	Intron 4	0.037	0.036	1.00E+00	0.018
<i>GH</i>	rs397514077	Intron 4	0.052	0.051	1.00E+00	0.026
<i>GH</i>	rs397514076	Intron 4	0.225	0.200	1.39E-01	0.113
<i>GH</i>	rs397514102	Exon 4	0.105	0.099	1.00E+00	0.052
<i>GH</i>	rs1092944696	Intron 3	0.173	0.175	1.00E+00	0.097
<i>GH</i>	rs1092437056	Intron 3	0.173	0.158	4.41E-01	0.086
<i>GH</i>	rs1135847309	Intron 3	0.099	0.095	1.00E+00	0.050
<i>GH</i>	rs397514070	Exon 3	0.120	0.113	9.89E-01	0.060
<i>GH</i>	rs397514069	Intron 2	0.094	0.090	1.00E+00	0.047
<i>GH</i>	rs589527314	Intron 2	0.298	0.289	9.08E-01	0.175
<i>GH</i>	rs1087440770	Intron 2	0.110	0.104	1.00E+00	0.055
<i>GH</i>	rs1135847310	Intron 2	0.021	0.021	1.00E+00	0.010
<i>GH</i>	rs397514066	Intron 2	0.052	0.051	1.00E+00	0.026
<i>GH</i>	rs397514065	Intron 2	0.157	0.145	5.81E-01	0.079
<i>GH</i>	rs397514064	Intron 2	0.157	0.145	5.81E-01	0.079
<i>IGF1</i>	rs430457475	Intron 1	0.388	0.389	1.00E+00	0.265
<i>IGF1</i>	rs1135847304	Intron 1	0.224	0.199	1.95E-01	0.112
<i>IGF1</i>	rs595347398	Intron 1	0.024	0.023	1.00E+00	0.012
<i>IGF1</i>	rs412470350	Intron 1	0.494	0.499	9.82E-01	0.482
<i>IGF1</i>	rs402300271	Intron 1	0.118	0.111	1.00E+00	0.059
<i>IGF1</i>	rs418030625	Intron 1	0.259	0.242	6.30E-01	0.141
<i>IGF1</i>	rs425204511	Intron 1	0.265	0.247	5.70E-01	0.144
<i>IGF1</i>	rs403521045	Intron 1	0.194	0.194	1.00E+00	0.109
<i>IGF1</i>	rs430449367	Intron 1	0.400	0.395	1.00E+00	0.271
<i>IGF1</i>	rs421570650	Intron 1	0.065	0.074	4.24E-01	0.038
<i>IGF1</i>	rs600588782	Intron 1	0.035	0.035	1.00E+00	0.018
<i>LEP</i>	rs408463464	Intron 2	0.524	0.491	4.71E-01	0.435
<i>LEP</i>	rs398357543	Intron 2	0.377	0.371	1.00E+00	0.246
<i>LEP</i>	rs409675427	Intron 2	0.068	0.066	1.00E+00	0.034
<i>LEP</i>	rs421064645	Intron 2	0.482	0.455	5.50E-01	0.351
<i>LEP</i>	rs410864710	Intron 2	0.503	0.493	9.26E-01	0.440
<i>LEP</i>	rs422219521	Intron 2	0.351	0.352	1.00E+00	0.228
<i>LEP</i>	rs400734857	Intron 2	0.382	0.368	7.87E-01	0.243
<i>LEP</i>	rs423196216	Intron 2	0.084	0.099	1.62E-01	0.052
<i>LEP</i>	rs406003615	Intron 2	0.387	0.371	7.15E-01	0.246
<i>LEP</i>	rs413205084	Intron 2	0.267	0.309	9.86E-02	0.191
<i>LEP</i>	rs424642048	Intron 2	0.251	0.299	5.13E-02	0.183
<i>LEP</i>	rs593178720	Intron 2	0.073	0.071	1.00E+00	0.037
<i>LEP</i>	rs403103423	Intron 2	0.366	0.349	6.62E-01	0.225
<i>LEP</i>	rs418548121	Intron 2	0.393	0.374	6.46E-01	0.249
<i>LEP</i>	rs1135847360	Intron 2	0.366	0.360	1.00E+00	0.236
<i>LEP</i>	rs596008192	Intron 2	0.372	0.363	9.36E-01	0.238
<i>LEP</i>	rs429879457	Intron 2	0.492	0.495	1.00E+00	0.450
<i>LEP</i>	rs404287904	Intron 2	0.366	0.366	1.00E+00	0.241

<sup>1</sup>HWE = Hardy-Weinberg equilibrium; <sup>2</sup>MAF = minor allele frequency.

## Results

### Associations analysis - *GH* gene

Haplotype association analysis was not carried out for *GH* because no LD blocks were detected in this gene. Additionally, after Bonferroni correction, no association

was found with the single-locus analysis approach in *GH*. However, the variant *rs589527314*, a substitution C/A found in the intron-2 of the *GH* gene, had a suggestive additive effect ( $p < 0.05$ ) on BW100 (Table 4). The A allele was associated with a higher value of these traits, and the difference between the CC and AA genotypes was 3.2 kg.

**Table 3** – Haplotypes in the *IGF1* and *LEP* genes with frequencies  $\geq 1$  %.

<i>IGF1</i>				<i>LEP</i>	
Block-1		Block-2		Block-1	
Haplotype	Frequency	Haplotype	Frequency	Haplotype	Frequency
TCC	0.479	GCG	0.856	ACGGATCGATGAGCAG	0.406
TCT	0.257	ATT	0.109	GCAAATCGATTAGCGG	0.199
GCT	0.149	ATG	0.032	GTGAGCCCGCATGA	0.134
GAT	0.112			GCAAATCGATGAGCGG	0.045
				GTGAGCCACGCATGA	0.045
				GCAAATGATGAGCGG	0.044
				GTGAGCCCGTGCATGA	0.029
				ACAGATCGATGAGCAG	0.016
				GCAGATCGATTAGCGG	0.010
				GCAAATCGATTAGCAG	0.010
				GTGAGCCCATGCATGA	0.010
				GTGAACCCGTGCATGA	0.010

**Table 4** – Additive effect (a) and standard error (SE) of polymorphisms in *GH*, *IGF1*, and *LEP* genes associated with growth, carcass and morphometric traits in Santa Inês sheep.

Trait	variants	NCBI Number	Intron	a	SE	LRT	p-value
<i>GH</i>							
BW100	g.47486819C > A	rs589527314	2	1.600	0.7076	5.02	0.0251
<i>IGF1</i>							
WH	g.171110428C > T	rs412470350	1	-0.941	0.4481	4.35	0.0370
CH	g.171110428C > T	rs412470350	1	-1.429	0.4970	8.06	0.0045
TW	g.171110428C > T	rs412470350	1	-0.576	0.2088	7.45	0.0064
LG	g.171110428C > T	rs412470350	1	-1.330	0.6091	4.70	0.0302
ADG	g.171110428C > T	rs412470350	1	-14.739	5.1931	7.79	0.0053
<i>LEP</i>							
BW100	g.92501543A > G	rs421064645	2	1.186	0.5561	4.47	0.0344
CFS	g.92501407C > T	rs398357543	2	0.117	0.0425	7.37	0.0066
CFS	g.92502245A > G	rs422219521	2	0.122	0.0435	7.60	0.0058
CFS	g.92502283T > C	rs400734857	2	0.116	0.0425	7.19	0.0073
CFS	g.92502623G > C	rs406003615	2	0.109	0.0424	6.42	0.0113
CFS	g.92502947A > C	rs418548121	2	0.106	0.0421	6.14	0.0132
CFS	g.92503024G > A	rs1135847360	2	0.124	0.0432	7.92	0.0049
CFS	g.92503025C > T	rs596008192	2	0.128	0.0429	8.65	0.0033
CFS	g.92503086G > A	rs404287904	2	0.128	0.0429	8.65	0.0033
CW	g.92501407C > T	rs398357543	2	0.706	0.2726	6.58	0.0103
CW	g.92502245A > G	rs422219521	2	0.787	0.2815	7.64	0.0057
CW	g.92502283T > C	rs400734857	2	0.714	0.2752	6.62	0.0101
CW	g.92502623G > C	rs406003615	2	0.725	0.2749	6.82	0.0090
CW	g.92502947A > C	rs418548121	2	0.685	0.2740	6.14	0.0132
CW	g.92503024G > A	rs1135847360	2	0.744	0.2785	7.00	0.0082
CW	g.92503025C > T	rs596008192	2	0.737	0.2774	6.93	0.0085
CW	g.92503086G > A	rs404287904	2	0.677	0.2746	5.98	0.0145
FT	g.92501407C > T	rs398357543	2	0.015	0.006	5.93	0.0149
FT	g.92502245A > G	rs422219521	2	0.017	0.006	7.59	0.0059
FT	g.92502283T > C	rs400734857	2	0.013	0.006	4.72	0.0298
FT	g.92503024G > A	rs1135847360	2	0.016	0.006	6.61	0.0101
FT	g.92503025C > T	rs596008192	2	0.016	0.006	6.87	0.0088
FT	g.92503086G > A	rs404287904	2	0.016	0.006	6.87	0.0088

LRT = likelihood ratio test; BW100 = Body weight at 100 days of age; WH and CH = withers and croup heights; TW and CW = thoracic and croup widths; LG = leg girth; ADG = average daily gain; CFS = carcass finishing score; FT = fat thickness.

#### Associations analysis - *IGF1* gene

Single-locus analysis in *IGF1* did not find an additive effect at the Bonferroni threshold. However, suggestive additive effects ( $p < 0.05$ ) of the variant

*rs412470350* on WH, CH, TW, LG, and ADG were found (Table 4). This variant is a C > T substitution in the intron-1 of *IGF1* gene. For all traits, the C allele was associated with a higher mean value. The differences



between the *CC* and *TT* genotypes were: 1.88 cm (WH), 2.86 cm (CH), 1.15 cm (TW), 2.66 cm (LG), and 29.48 g d<sup>-1</sup> (ADG). In addition, twelve haplotype replacements in LD block-1 of *IGF1* were found (Table 5). The replacement of *TCC* by *TCT* was associated ( $p < 0.0075$ ) with ADG, CH, WH, BL, TG, TW, and LG, where regression coefficients ( $\beta$ ) and standard errors (SE) were 20.5079  $\pm$  7.3741, 4.0859  $\pm$  1.2121, 3.5227  $\pm$  1.2002, 3.9393  $\pm$  1.1920, 3.8814  $\pm$  1.3036, 1.1302  $\pm$  0.3577, and 3.3994  $\pm$  1.0808, respectively. Moreover, another five suggestive ( $p < 0.05$ ) haplotype replacements were found.

#### Associations analysis - *LEP* gene

In the *LEP* gene, no single-locus effect was found at the Bonferroni threshold. However, 23 suggestive additive effects ( $p < 0.05$ ) were found (Table 4). The variants *rs398357543*, *rs422219521*, *rs400734857*, *rs1135847360*, *rs596008192*, and *rs404287904*, all in intron-2, had suggestive additive effects ( $p < 0.05$ ) on CFS, CW, and FT (Table 4). Moreover, the variants *rs406003615* and *rs418548121*, also located in intron-2, were associated with both CFS and CW. The difference between homozygous ranged from 0.212 (*rs418548121*) to 0.256 (*rs596008192* and *rs404287904*) scores of CFS, from 0.026 (*rs400734857*) to 0.034 cm (*rs422219521*) of

FT, and from 1.35 (*rs404287904*) to 1.57 cm (*rs422219521*) of CW. In addition, the *rs421064645*, a substitution A/G in intron-2, had a suggestive additive effect ( $p < 0.05$ ) on BW100, where the difference between *AA* and *GG* was 2.37 kg. Finally, associations ( $p < 0.0075$ ) with CFS (-0.2441  $\pm$  0.0611), BW100 (1.8270  $\pm$  0.5143) and BD (-2.5103  $\pm$  0.5621) were found when the most frequent haplotype (*ACGGATCGATGAGCAG*) in the *LEP* gene was replaced by other haplotypes (Table 5).

## Discussion

#### *GH* gene

The variant *rs589527314* in intron-2 of the *GH* gene was associated with BW100 in Santa Inês sheep, which is supported by previous association studies. Variants in intron-2 were associated with body weight in Black Tibetan sheep (Han et al., 2016), and birth weight, body weight at 120 days, and ADG from birth to 120 days in Harri sheep (Abdelmoneim et al., 2017). Other regions of *GH* were also associated with body weight in sheep. The variant in intron-1 was associated with weaning weight and ADG in Nilagiri sheep (Cauveri et al., 2016). On exon 4, a PCR-SSCP associated with body weight in Makooei sheep was reported (Moradian et al., 2013), and one single nucleotide polymorphism was associated with body weight in Tibetan and Poll Dorset breeds (Jia et al., 2014). Moreover, a PCR-SSCP in exon 5 was associated with body weight in Baluchi sheep at six months of age (Tahmoorespur et al., 2011), while a PCR-RFLP was associated with body weight and ADG in Donggala and East Java breeds (Malewa et al., 2014) and Jambi Province sheep (Depison et al., 2017). These many associations between variants in *GH* and growth traits are supported by biological processes such as insulin secretion (Zhang et al., 2004), response to food (Berryman et al., 2004), and positive regulation of multicellular organism growth (Harvey, 2010), which depends on growth hormone activity.

#### *IGF1* gene

The variant *rs412470350* and haplotype replacements were associated with different body traits in Santa Inês sheep (Tables 4 and 5). The molecular function and biological process associated with the *IGF1* gene supports these results. Yakar et al. (2002), demonstrated that circulating levels of IGF-1 directly regulate bone growth and density in mice, while Sun et al. (2014) found an association between *IGF1* gene expression and body weight in Hu sheep. In addition, the *IGF1* gene has important molecular functions for animal growth, such as cell proliferation, maintenance of skeletal muscle satellite cells for regeneration, and an essential role in muscle hypertrophy (Philippou et al., 2007). Previous studies carried out with other sheep breeds also supported our results. A variant (*rs430457475*) in intron-1 was associated with WH and CH in Russian Merino breed (Trukhachev et al.,

**Table 5** – Regression coefficients ( $\beta$ ) and standard errors (SE) estimated by haplotype association analysis in the *IGF1* and *LEP* genes in Santa Inês sheep.

Trait	Haplotype replacement	$\beta$	SE	p-value
<i>IGF1</i>				
ADG	<i>TCC</i> > <i>TCT</i>	20.5079	7.3741	0.006*
CH	<i>TCC</i> > <i>GAT</i>	5.1563	1.9695	0.010
CH	<i>TCC</i> > <i>GCT</i>	3.1585	1.5026	0.037
CH	<i>TCC</i> > <i>TCT</i>	4.0859	1.2121	0.001*
WH	<i>TCC</i> > <i>GAT</i>	4.0960	1.9502	0.037
WH	<i>TCC</i> > <i>GCT</i>	2.9829	1.4880	0.047
WH	<i>TCC</i> > <i>TCT</i>	3.5227	1.2002	0.004*
BL	<i>TCC</i> > <i>TCT</i>	3.9393	1.1920	0.001*
TG	<i>TCC</i> > <i>TCT</i>	3.8814	1.3036	0.003*
TW	<i>TCC</i> > <i>GCT</i>	1.1406	0.4437	0.011
TW	<i>TCC</i> > <i>TCT</i>	1.1302	0.3577	0.002*
LG	<i>TCC</i> > <i>TCT</i>	3.3994	1.0808	0.002*
<i>LEP</i>				
BW100	H1 > GCAAATTGATGAGCGG	1.8270	0.5143	< 0.0001*
BW240	H1 > GCAAATCGATTAGCGG	-2.0353	0.9165	0.028
REA	H1 > GCAAATCGATTAGCGG	-2.4400	0.2049	0.016
CFS	H1 > GTGAGCCCGCATGA	-0.2441	0.0611	< 0.0001*
TG	H1 > GCAAATCGATGAGCGG	-5.2915	1.9733	0.008
BD	H1 > GCAAATTGATGAGCGG	-2.5103	0.5621	< 0.0001*
BL	H1 > GCAAATTGATGAGCGG	2.4830	0.9563	0.01

\*Significant effect at 5 % of Bonferroni correction; H1 = haplotype ACGGATCGATGAGCAG; ADG = average daily gain; CH and WH = withers and croup heights; BL = body length; TG = Thoracic girth; TW = thoracic widths; LG = leg girth; BW100 and BW240 = Body weight at 100 and 240 days of age, respectively; REA = rib eye area; CFS = carcass finishing score; and BD = body depth.

2016). Furthermore, association between a PCR-RFLP in the regulatory region of *IGF1* with BL, TW, and body weight in Balami sheep, and WH in Yankasa sheep were also reported (Raji et al., 2017). A PCR-SSCP in exon-1 was associated with ADG and BL in Makui sheep (Hajihosseini et al., 2013), and ADG in both Baluchi (Tahmoorespur et al., 2009) and Makoei (Negahdary et al., 2013) sheep breeds.

### LEP gene

Variants in the *LEP* gene of Santa Inês sheep were associated with several body traits in the current study. These results are supported by molecular functions such as hormonal activities (Wauters et al., 2000) that regulate vital metabolic processes; and the activity of growth factors that stimulate cells to grow or proliferate. Furthermore, the *LEP* gene is a component of several biological processes, especially those related to fat metabolism, such as the lipid metabolic process and beta-oxidation of fatty acids (Havel, 2004), which may also explain the associations observed here for FT and CFS. The leptin also plays a vital role in biological processes that are related to the negative regulation of appetite (Blundell et al., 2001), adult eating behavior and feeding behavior (Williamson et al., 2005), intestinal absorption (Yarandi et al., 2011), and bone growth (Upadhyay et al., 2015), which can explain the effects on growth and morphometric traits in Santa Inês sheep.

Previous studies also support our results with the *LEP* gene. In intron-2, associations are reported with REA in the Suffolk breed (Boucher et al., 2006) and with cold carcass weight, fat-tail, and total body fat in the Shal and Zelsheep breeds (Barzehkar et al., 2009). Additionally, the variants *g.92501372G>A*, *rs398357543*, *rs421064645*, and *rs1135847360* in intron-2 of *LEP* gene were associated with several carcass traits recorded post-mortem in Santa Inês sheep, such as cold and hot carcass weights and yields, internal carcass length, leg and neck yields, neck weight, and CFS (Meira et al., 2018). Polymorphisms in other regions of the *LEP* gene, especially in exon 3, were also associated with economic traits. A PCR-SSCP in exon 3 of *LEP* gene was associated with breeding values for TG and rump length in Makoei sheep (Sadeghi et al., 2014), body weight and ADG in Makoei sheep (Hajihosseini et al., 2012), body weight at 90 days in Baluchi sheep (Tahmoorespur et al., 2010), and body weight in Kermani sheep with 3, 6, 9, and 12 months of age (Shojaei et al., 2010). Some variants in the exon 3 were associated with BL in Sanjabu sheep (Bakhtiar et al., 2017) feed conversion rate, residual feed intake and feed intake in crossbreed Awassi-Merino sheep (Jonas et al., 2016), and cold carcass weight in Santa Inês sheep (Quirino et al., 2016).

The present study found evidence of intronic variants in the *IGF1*, *LEP*, and *GH* genes associated with body traits in Santa Inês sheep. No previous

studies identified small RNA transcripts encoded by the intronic regions associated with body traits in the current study. Moreover, the variants, here associated with body traits, are not in splice sites. Therefore, linkage disequilibrium between the polymorphism found here, and the causal variant is the main hypothesis that explains the additive effects found.

## Acknowledgments

To the Bahia State Foundation for Research Support (FAPESB) for the financial support to projects APP0116/2009 and PNE0006/2014; to the Brazilian National Council for Scientific and Technological Development (CNPq) for financial support to projects 562551/2010-7 and 474494/2010-1; to Embrapa Coastal Tablelands for the infrastructure of the experimental farm; and to Dr. Luiz Lehmann Coutinho for the infrastructure of Functional Genomics Core facility of University of São Paulo/ESALQ (USP/ESALQ). Luiz Coutinho, Gerson Mourão, Victor Pedrosa, and Luis Pinto are recipient of a CNPq Productivity Scholarship. This study was also financed in part by the Coordination for the Improvement of Higher Level Personnel (CAPES) - Finance Code 001.

## Authors' Contributions

**Conceptualization:** Pinto, L.F.B. **Data acquisition:** Coutinho, L.L.; Azevedo, H.C.; Muniz, E.N.; Meira, A.N.; Jucá, A.F. **Data analysis:** Machado, A.L.; Meira, A.N. **Design of methodology:** Pinto, L.F.B.; Machado, A.L. **Writing and editing:** Machado, A.L.; Mourão, G.B.; Pedrosa, V.B.; Coutinho, L.L.; Pinto, L.F.B.

## References

- Abdelmoneim, T.S.; Brooks, P.H.; Afifi, M.; Swelum, A.A.A. 2017. Sequencing of growth hormone gene for detection of polymorphisms and their relationship with body weight in Harri sheep. *Indian Journal of Animal Research* 51: 205-211.
- Bakhtiar, R.; Abdolmohammadi, A.; Hajarian, H.; Nikousefat, Z.; Kalantar-Neyestanaki, D. 2017. Identification of 332G > A polymorphism in exon 3 of the leptin gene and partially effects on body size and tail dimension in Sanjabu sheep. *International Journal of Bioengineering and Life Sciences* 11: 506-509.
- Barzehkar, R.; Salehi, A.; Mahjoubi, F. 2009. Polymorphisms of the ovine leptin gene and its association with growth and carcass traits in three Iranian sheep breeds. *Iranian Journal of Biotechnology* 7: 241-246.
- Berryman, D.E.; List, E.O.; Coshcigano, K.T.; Behar, K.; Kim, J.K.; Kopchick, J.J. 2004. Comparing adiposity profiles in three mouse models with altered GH signaling. *Growth Hormone & IGF Research* 14: 309-318.
- Blundell, J.E.; Goodson, S.; Halford, J.C.G. 2001. Regulation of appetite: role of leptin in signaling systems for drive and satiety. *International Journal of Obesity* 25: S29-S34.

- Boucher, D.; Palin, M.F.; Castongua, F.; Gariépy, C.; Pothier, F. 2006. Detection of polymorphisms in the ovine leptin (*LEP*) gene: association of single nucleotide polymorphism with muscle growth and meat quality traits. *Canadian Journal of Animal Science* 86: 31-35.
- Caron, A.; Lee, S.; Elmquist, J.K.; Gautron, L. 2018. Leptin and brain-adipose crosstalks. *Nature Reviews Neuroscience* 19: 153-165.
- Cauveri, D.; Sivaselvam, S.N.; Karthickeyan, S.M.K.; Tirumurugan, K.G.; Kumanan, K.; Venkataramanan, R. 2016. Single nucleotide polymorphisms in *GH* (growth hormone) gene associated with growth traits in Nilagiri sheep of Tamil Nadu. *International Journal of Science, Environment and Technology* 5: 4097-4103.
- Cockett, N.E.; Smit, M.A.; Bidwell, C.A.; Segers, K.; Hadfield, T.L.; Snowden, G.D.; Georges, M.; Charlier, C. 2005. The callipyge mutation and other genes that affect muscle hypertrophy in sheep. *Genetics Selection Evolution* 37: S65-S81.
- Depison; Anwar, S.; Jamsari; Arnim; Yurnalis. 2017. Association of growth hormone gene polymorphism with quantitative characteristics of thin-tailed sheep using PCR-RFLP in Jambi province. *African Journal of Biotechnology* 16: 1159-1167.
- Gorlov, I.F.; Kolosov, Y.A.; Shirokova, N.V.; Getmantseva, L.V.; Slozhenkina, M.I.; Mosolova, N.I.; Bakoev, N.F.; Leonova, M.A.; Kolosov, A.Y.; Zlobina, E.Y. 2017. Association of the growth hormone gene polymorphism with growth traits in Salsk sheep breed. *Small Ruminant Research* 150: 11-14.
- Hajihosseainlo, A.; Hashemi, A.; Razavi-Sheshdeh, S.A.; Pirany, N. 2013. Association of the polymorphism in the 5' flanking region of the ovine IGF-I gene with growth and development traits in Makui sheep of Iran. *European Journal of Zoological Research* 2: 19-24.
- Hajihosseainlo, A.; Hashemi, A.; Sadeghi, S. 2012. Association between polymorphism in exon 3 of leptin gene and growth traits in the Makooei sheep of Iran. *Livestock Research for Rural Development* 24: 543-546.
- Han, Y.C.; Sun, Y.G.; Li, Q. 2016. Growth hormone polymorphisms and growth traits in Chinese Tibetan sheep *Ovis aries*. *Genetics and Molecular Research* 15: 15038397.
- Harvey, S. 2010. Extrapituitary growth hormone. *Endocrine* 38: 335-359.
- Havel, P.J. 2004. Update on adipocyte hormones: regulation of energy balance and carbohydrate/lipid metabolism. *Diabetes* 53: S143-S151.
- Hickford, J.G.H.; Forrest, R.H.; Zhou, H.; Fang, Q.; Han, J.; Frampton, C.M.; Horrell, A.L. 2010. Polymorphisms in the ovine myostatin gene (*MSTN*) and their association with growth and carcass traits in New Zealand Romney sheep. *Animal Genetics* 41: 64-72.
- Jia, J.L.; Zhang, L.P.; Wu, J.P.; Ha, Z.J.; Li, W.W. 2014. Study of the correlation between *GH* gene polymorphism and growth traits in sheep. *Genetics and Molecular Research* 13: 7190-7200.
- Jonas, E.; Martin, G.B.; Celi, P.; Li, L.; Soattin, M.; Thomsson, P.C.; Raadsma, H.W. 2016. Association of polymorphisms in leptin and leptin receptor genes with circulating leptin concentrations, production and efficiency traits in sheep. *Small Ruminant Research* 136: 78-86.
- Lake, S.L.; Lyon, H.; Tantisira, K.; Silverman, E.K.; Weiss, S.T.; Laird, N.M.; Schaid, D.J. 2003. Estimation and tests of haplotype-environment interaction when linkage phase is ambiguous. *Human Heredity* 55: 56-65.
- Langmead, B.; Salzberg, S.L. 2012. Fast gapped-read alignment with Bowtie 2. *Nature Methods* 4: 357-359.
- Malewa, A.D.; Hakim, L.; Maylinda, S.; Husain, M.H. 2014. Growth hormone gene polymorphisms of Indonesia fat tailed sheep using PCRRFLP and their relationship with growth traits. *Livestock Research for Rural Development* 26: 6.
- Meira, A.N.; Montenegro, H.; Coutinho, L.L.; Mourão, G.B.; Azevedo, H.C.; Muniz, E.N.; Machado, A.L.; Sousa-JR, L.P.; Pedrosa, V.B.; Pinto, L.F.B. 2019. Single nucleotide polymorphisms in the *GH* and *IGF1* genes associated with carcass traits in Santa Inês sheep. *Animal* 13: 460-468.
- Meira, A.N.; Moreira, G.C.M.; Coutinho, L.L.; Mourão, G.B.; Azevedo, H.C.; Muniz, E.N.; Machado, A.L.; Sousa-Jr, L.P.; Pedrosa, V.B.; Pinto, L.F.B. 2018. Carcass and commercial cut yield of Santa Inês sheep affected by polymorphisms of the *LEP* gene. *Small Ruminant Research* 166: 121-128.
- Moradian, C.; Mohamadi, N.; Razavi-Sheshdeh, S.A.; Hajihosseainlo, A.; Ashrafi, F. 2013. Effects of genetic polymorphism at the growth hormone gene on growth traits in Makooei sheep. *European Journal of Experimental Biology* 3: 101-105.
- Negahdary, M.; Hajihosseainlo, A.; Ajdary, M. 2013. PCR-SSCP variation of *IGF1* and *PIT1* genes and their association with estimated breeding values of growth traits in Makooei sheep. *Genetics Research International* 2013: 272346.
- National Research Council [NRC]. 2007. Nutrient Requirements of Small Ruminants: Sheep, Goats, Cervids and New World Camelids. Natl. Acad. Press, Washington, DC, USA.
- Oliveira, M.D.S.; Regitano, L.D.A.; Roese, A.D.; Anthonisen, D.G.; Patrocínio, E.D.; Parma, M.M.; Scagliusi, S.M.M.; Timoteo, W.H.B.; Belicuas, S. 2007. Theoretical and practical background and protocols for DNA extraction and amplification by polymerase chain reaction technique = Fundamentos teórico-práticos e protocolos de extração e de amplificação de DNA por meio da técnica de reação em cadeia de polimerase. *Embrapa Pecuária Sudeste, São Carlos, SP, Brazil* (in Portuguese).
- Pérez-Enciso, M.; Miszta, I. 2011. Qxpk.5: old mixed model solutions for new genomics problems. *BMC Bioinformatics* 12: 202.
- Philippou, A.; Maridaki, M.; Halapas, A.; Koutsilieris, M. 2007. The role of the insulin-like growth factor 1 (IGF-1) in skeletal muscle physiology. *In vivo* 21: 45-54.
- Price, A.L.; Patterson, N.J.; Plenge, R.M.; Weinblatt, M.E.; Shadick, N.A.; Reich, D. 2006. Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics* 8: 904-909.
- Quirino, C.R.; Costa, R.L.D.; Pacheco, A.; Madella-Oliveira, A.F.; Beltrame, R.T.; Azevedo, A.S.; Bartholazzi-Junior, A.; Vega, W.H.O. 2016. Identification of polymorphisms in the myostatin and leptin genes of Santa Inês breed and crossbreed sheep and association with carcass traits. *Bioscience Journal* 32: 699-704.
- Raji, A.O.; Mohammed, A.; Igwebuike, J.U.; Alphonsus, C. 2017. Association of IGF 1 gene polymorphisms with some morphometric traits of Nigerian indigenous sheep breeds. *Nigerian Journal of Biotechnology* 34: 97-104.



- Sadeghi, S.; Hajhosseinlo, A.; Bohlouli, M. 2014. Haplotype association of ovine leptin gene on breeding value of body measurements in Makooei sheep breed. *Biotechnology in Animal Husbandry* 30: 233-242.
- Sahu, A.R.; Jeichitra, V.; Rajendran, R.; Raja, A. 2017. Polymorphism of growth hormone receptor (GHR) gene in Nilagiri sheep. *Tropical Animal Health and Production* 49: 281-285.
- Shojaei, M.; Abadi, M.M.; Fozi, M.A.; Dayani, O.; Khezri, A.; Akhondi, M. 2010. Association of growth trait and leptin gene polymorphism in Kermani sheep. *Journal of Cell and Molecular Research* 2: 67-73.
- Sun, W.; Su, R.; Li, D.; Musa, H.H.; Kong, Y.; Ding, J.T.; Ma, Y.H.; Chen, L.; Zhang, Y.F.; Wu, W.Z. 2014. Developmental changes in IGF-I and MyoG gene expression and their association with meat traits in sheep. *Genetics and Molecular Research* 13: 2772-2783.
- Tahmoorespur, M.; Taheri, A.; Gholami, H.; Ansary, M. 2011. PCR-SSCP Variation of *GH* and *STAT5A* genes and their association with estimated breeding values of growth traits in Baluchi sheep. *Animal Biotechnology* 22: 37-43.
- Tahmoorespur, M.; Taheri, A.; Valeh, M.V.; Saghi, D.A.; Ansary, M. 2010. Assessment relationship between leptin and ghrelin genes polymorphisms and estimated breeding values EBVs of growth traits in Baluchi sheep. *Journal of Animal and Veterinary Advances* 9: 2460-2465.
- Tahmoorespur, M.; Valeh, M.V.; Nassiry, M.R.; Moussavi, A.H.; Ansary, M. 2009. Association of the polymorphism in the 5' flanking region of the ovine IGF-I gene with growth traits in the Baluchi sheep. *South African Journal of Animal Science* 39: 97-101.
- Teran, E.; Chesner, J.; Rapaport, R. 2016. Growth and growth hormone: an overview. *Growth Hormone & IGF Research* 28: 3-5.
- Trukhachev, V.; Skripkin, V.; Kvochko, A.; Kulichenko, A.; Kovalev, D.; Pisarenko, S.; Volynkina, A.; Selionova, M.; Aybazov, M.; Shumaenko, S.; Omarov, A.; Mamontova, T.; Yatsyk, O.; Krivoruchko, A. 2016. Polymorphisms of the *IGF1* gene in Russian sheep breeds and their influence on some meat production parameters. *Slovenian Veterinary Research* 53: 2.
- Tuersunjiang, N.; Odhiambo, J.F.; Shasa, D.R.; Smith, A.M.; Nathaniels, P.W.; Ford, S.P. 2016. Maternal obesity programs reduced leptin signaling in the pituitary and altered *GH/IGF1* axis function leading to increased adiposity in adult sheep offspring. *Plos One* 12: e0181795.
- Upadhyay, J.; Farr, O.M.; Mantzoros, C.S. 2015. The role of leptin in regulating bone metabolism. *Metabolism* 64: 105-113.
- Yakar, S.; Rosen, C.J.; Beamer, W.G.; Ackert-Bicknell, C.L.; Wu, Y.; Liu, J.L.; Ooi, G.T.; Setser, J.; Frystyk, J.; Boisclair, Y.R.; Leroith, D. 2002. Circulating levels of IGF-1 directly regulate bone growth and density. *Journal of Clinical Investigation* 110: 771-781.
- Yakar, S.; Werner, H.; Rosen, C.J. 2018. 40 Years of IGF1 Insulin-like growth factors: actions on the skeleton. *Journal of Molecular Endocrinology* 61: T115-T137.
- Yarandi, S.S.; Hebbar, G.; Sauer, C.G.; Cole, C.R.; Ziegler, T.R. 2011. Diverse roles of leptin in the gastrointestinal tract: modulation of motility, absorption, growth, and inflammation. *Nutrition* 27: 269-275.
- Wauters, M.; Considine, R.V.; Van-Gaal, L.F. 2000. Human leptin: from an adipocyte hormone to an endocrine mediator. *European Journal of Endocrinology* 143: 293-311.
- Williamson, D.A.; Ravussin, E.; Wong, M.L.; Wagner, A.; Dipaoli, A.; Caglayan, S.; Ozata, M.; Martin, C.; Walden, H.; Arnett, C.; Licinio, J. 2005. Microanalysis of eating behavior of three leptin deficient adults treated with leptin therapy. *Appetite* 45: 75-80.
- Zhang, Q.; Kohler, M.; Yang, S.N.; Zhang, F.; Larsson, O.; Berggren, P.O. 2004. Growth hormone promotes Ca<sup>2+</sup>-induced Ca<sup>2+</sup> release in insulin-secreting cells by ryanodine receptor tyrosine phosphorylation. *Molecular Endocrinology* 18: 1658-1669.
- Zhbannikov, I.Y.; Hunter, S.S.; Settles, M.L. 2013. SEQYCLEAN User Manual. Available at: <http://sourceforge.net/projects/seqclean/files/> [Accessed Dec 13, 2016] Woessner Junior, J.F. 1961. The determination of hydroxyproline in tissue and protein samples containing small proportions of this amino acid. *Archives of Biochemistry and Biophysics* 93: 440-447.