

Ricardo de Oliveira Dart, *Brazil*  
Ricardo Simão Diniz Dalmolin, *Brazil*  
Roza Orozakunova, *Kyrgyzstan*  
Sergio Radic, *Chile*  
Shishkov Toma, *Bulgaria*  
Skye Angela Wills, *United States of America*  
Stephen Roecker, *United States of America*  
Susana Valle, *Chile*  
Suzann Kienast-Brown, *United States of America*  
Svitlana Nakisko, *Ukraine*  
Thomas W. Kuyper, *Netherlands*  
Vadym Solovei, *Ukraine*  
Vasenev Ivan, *Russian Federation*  
Vasyl Cherlinka, *Ukraine*  
Veronica Reynoso De La Mora, *Mexico*  
Vitalii Lebed, *Ukraine*  
William Andrés Cardona, *Colombia*  
Xiaoyuan Geng, *Canada*  
Yan Li, *China*  
Ying Zhang, *China*  
Yiyi Sulaeman, *Indonesia*  
Yurii Zalavskiy, *Ukraine*  
Yusuf Yigini, *FAO-Global Soil Partnership*  
Yusuke Takata, *Japan*  
Zheng Sun, *China*

### **Chapter 3.** **Status and challenges of black soils**

Lead Authors:

**Pavel Krasilnikov**, *Russian Federation*

Contributing Authors:

**Ademir Fontana**, *Brazil*  
**Ahmad Landi**, *Islamic Republic of Iran*  
**Ahmet R. Mermut**, *Canada*  
**Beata Labaz**, *Poland*  
**Bożena Smreczak**, *Poland*  
**Carlos Roberto Pinheiro**, *Brazil*  
**Cornelius Wilhelm (Cornie) Van Huyssteen**, *South Africa*  
**Curtis Monger**, *United States of America*  
**Flávio Pereira de Oliveira**, *Brazil*  
**Héctor J. M. Morras**, *Argentina*  
**Hussam HAG Husein**, *Syrian Arab Republic*  
**Jorge Ivelic-Sáez**, *Chile*  
**Kathia Peralta**, *Mexico*  
**Lei Wang**, *China*  
**Lúcia Helena Cunha dos Anjos**, *Brazil*  
**Luís Antônio Coutrim dos Santos**, *Brazil*  
**Marco Pfeiffer**, *Chile*  
**Marcos Gervasio Pereira**, *Brazil*

**Martha Marina Bolanos-Benavides**, *Colombia*  
**Miguel Angel Taboada**, *Argentina*  
**Milton César Costa Campos**, *Brazil*  
**Ricardo Simão Diniz Dalmolin**, *Brazil*  
**Roza Orozakunova**, *Kyrgyzstan*  
**Sergejus Ustinov**, *FAO-Global Soil Partnership*  
**Sergio Radic**, *Chile*  
**Susana Valle**, *Chile*  
**Thomas W. Kuyper**, *Netherlands*  
**Vasyl Cherlinka**, *Ukraine*  
**Wilian Demetrio**, *Brazil*  
**William Andrés Cardona**, *Colombia*  
**Ying Zhang**, *China*  
**Yuriy Dmytruk**, *Ukraine*  
**Yusuke Takata**, *Japan*

### **Chapter 4.** **Sustainable management of black soils: from practices to policies**

Lead Authors:

**Luca Montanarella**, *Joint Research Centre, European Commission*

**William Bill May**, *Canada*

**Yuxin Tong**, *FAO-Global Soil Partnership*

Contributing Authors:

**Ademir Fontana**, *Brazil*  
**Anatoly Klimanov**, *Russian Federation*  
**Anna Kontoboytseva**, *Russian Federation*  
**Arcangelo Loss**, *Brazil*  
**Bayarsukh Noov**, *Mongolia*  
**Beata Labaz**, *Poland*  
**Bożena Smreczak**, *Poland*  
**Cai Hongguang**, *China*  
**Carlos Clerici**, *Uruguay*  
**Carolina Olivera Sanchez**, *FAO-Global Soil Partnership*  
**Deliang Peng**, *China*  
**Elena Timofeeva**, *Russian Federation*  
**Élvio Giasson**, *Brazil*  
**Enkhtuya Bazarradnaa**, *Mongolia*  
**Fan Wei**, *China*  
**Fernando Fontes**, *Uruguay*  
**Gonzalo Pereira**, *Uruguay*  
**Hakkı Emrah Erdogan**, *Türkiye*  
**Hussam Hag Husein**, *Syrian Arab Republic*  
**Ievgen Skrylnyk**, *Ukraine*  
**Jaroslava Sobocká**, *Slovakia*  
**Jianhua Qu**, *China*  
**Jihong Liu Clarke**, *Norway*  
**Jingkuan Wang**, *China*

# 3. Status and challenges of black soils

## 3.1 A global overview of black soils

Although black soils account for only 8.2 percent of the Global Land Area (FAO, 2022a), they are of paramount importance for food security, as highlighted by the UN Sustainable Development Goal 2 (i.e. to end hunger, achieve food security and improve nutrition, and promote sustainable agriculture by 2030). The general category of black soils includes three main soil groups Chernozem, Kastanozem and Phaeozem, according to the World Reference Base for Soil Resources (WRB). Black soils are characterized by their thick, dark-coloured, and humus-rich topsoil. In general terms, black soils have granular and subangular blocky structure, optimal bulk density, and high amounts of plant nutrients. However all these favorable properties are only present in soils within virgin or quasi-pristine ecosystems, which are now rare (Montanarella *et al.*, 2021). There are other soil types also considered as black soils, as for example, swelling soils (Vertisols), volcanic soils (Andisols), anthropogenic soils, among others. Not all of them strictly comply with some of the conditions indicated for the Category 1 of black soil definition (such as having been formed under grassland vegetation) (FAO, 2019), but they all have some characteristics in their profiles that allow them to be classified as black soils, such as having a thick, dark-coloured, and humus-rich topsoil.

Apart from being highly productive lands, black soils are responsible for multiple ecosystem services such as water retention, maintenance of soil biodiversity from microorganism to megafauna, and soil fertility, and prevention of soil compaction and waterlogging. One of the most valuable services is accumulation of

great amounts of SOC in a relatively stable form. Black soils are one of the most important pools of carbon accounting for 8.27 percent (56 PgC) of the total global SOC stock in the top 30 centimetres of the soil (FAO, 2022d).

These carbon stores are, however, endangered by the processes of organic carbon loss due to the accelerated humus oxidation under cultivation. In many places the loss of humus and nutrient mining are the most important threats to black soils because these soils are considered as highly fertile “by nature” and thus have not needed application of organic and mineral fertilizers. Black soils are under further threat from various physical, chemical, and biological degradation processes (FAO and ITPS, 2015).

Some of these processes are easily reversible through sustainable soil management practices, such as nutrient imbalance, compaction, and structural degradation. However, other processes are difficult to reverse. First, soil loss due to erosion (wind, water and meltwater), is the most widespread threat in all world’s soils. Wind erosion is a problem that tremendously affected the Midwestern of the United States of America (the infamous Dust Bowl in the 1930s) and west Siberia and north Kazakhstan in the ex-Union of Soviet Socialist Republics (USSR) during the development of virgin lands in 1950s. Currently, soil salinization is becoming a growing problem, especially in irrigated areas in the most arid parts of the distribution of black soils. Second, land use change for food production exacerbates unsustainable management practices such as aggressive tillage and overgrazing furthering losses due to erosion. Diffuse pollution processes affect black soils devoted to fibre agriculture. This happens for various reasons, including the use of inappropriate fertilization technologies with high doses of nitrogen and phosphorus fertilizers and manures, or the excessive or inadequate use of herbicides and pesticides whose decomposition products are potential contaminants in soils, streams and groundwater. Finally, many black soils are threatened by soil sealing, due to the advance of urban areas and infrastructure in overpopulated regions or countries. This advance makes thousands of hectares of previously black soils destined for food production disappear.



© Xingzhu Ma

**Photo 3.1** Black soil in Nenjiang county of Heilongjiang province, China



© Xingzhu Ma



## 3.2 Multiple benefits of black soils

### 3.2.1 Ecosystem services (ES)

Soils are involved in most of the ecosystem services (ES) that enable life on Earth, such as the provision of food, fibre, bioenergy, and water; the regulation of climate, gas, floods, droughts, land degradation, water quality, and pests and diseases; the support of nutrient cycles, and habitat for organisms; and the cultural non-material benefits such as recreational, spiritual, and religious values (See Figure 3.2.1a). Black soils have distinctive soil properties that are key for providing essential ES, for example, high soil organic matter contents and cation exchange capacities, better soil physical properties (soil structure, porosity, hydraulic conductivity, and infiltration) and habitats for soil


organisms, which ensure the provision of food, fuel and fibre and freshwater, the regulation of climate, erosion control and water purification, and the support of nutrient cycling (Adhikari and Hartemink, 2016).

Although all soils are responsible for and intervene in the provision of ES, black soils have a preponderant role in the provision of healthy food, nutrient and water reserves, habitat for organisms, among other functions. That is why the loss of organic matter due to unsustainable management practices is likely causes a greater impact than in other less fertile soils.

The linkages between soil and ecosystem services were represented by Adhikari and Hartemink (2016) through a diagram (See Figure 3.2.1a) that conceptualizes soils as a complex system which provides multiple benefits for the environment and society, and the need to study it in a holistic approach to understand the multiple interactions between soil functions, ecosystem services and human well-being.








**Figure 3.2.1a** A conceptual diagram linking key soil properties to ecosystem services through soil functions for the well being of humans

Source: Adhikari, K. & Hartemink, A.E. 2016. *Linking soils to ecosystem services—A global review*. Geoderma, 262: 101–111. <https://doi.org/10.1016/j.geoderma.2015.08.009>

Anthropic soil erosion driven by conversion of natural ecosystems to agroecosystems and mechanical tillage, has numerous adverse impacts on ecosystem services. There are: 1) on-site impacts like the degradation of soil quality, reduction of agronomic productivity, and

the decrease in use-efficiency of inputs; and 2) off-site impacts like accelerated soil erosion that promotes eutrophication and contamination, and sedimentation of reservoirs and waterways, and emissions of CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O (Lal, 2014) (Figure 3.2.1b).



**Figure 3.2.1b** Adverse effects of accelerated erosion on ecosystem functions and services

Source: Lal, R. 2014. *Soil conservation and ecosystem services*. International Soil and Water Conservation Research, 2(3): 36-47. [https://doi.org/10.1016/S2095-6339\(15\)30021-6](https://doi.org/10.1016/S2095-6339(15)30021-6)



### 3.2.2 Climate change mitigation and adaptation

Black soils can contribute to both the mitigation of and adaptation to climate change. On the one hand, black soils have a high potential to mitigate climate change due to their high SOC sequestration potential (See Figure 3.2.2). According to FAO's Global Soil Organic Carbon map (GSOCmap), in the top 30 centimetres SOC stock of black soils is on average 56 PgC (or 77.24 tonnes C/ha), which is higher than the average of SOC stock in all mineral soils (FAO, 2022b). On the other hand, black soils have been cultivated since many centuries in Europe and Asia, and in the last 150 to 200 years in America and Oceania. After extensive and intensive cultivation (for cereals, pastures, ranges, and forage systems), black soils have significant losses of SOC (See Figure 3.2.2). According to various estimates, SOC loss of up to 50 percent of initial SOC occurred after conversion from a natural system to intensive farming, as happened in intensively cropped soils of the United States of America (Gollany *et al.*, 2011). This SOC loss results from of inappropriate land use and unsustainable management practices, leading to declining soil quality. The decline in soil quality is generally characterized by poor topsoil structure, increased soil erosion, resulting in emissions of carbon into the atmosphere exacerbating climate change (Lal, 2019).


Much of the carbon loss was caused using aggressive tillage systems, but also by the replacement of perennial vegetation (grasslands, forests) by annual crops, which in general generate lower carbon returns to the soil and hydrological imbalances (Fan *et al.*, 2017). When grasslands are converted to croplands, they lose on average 36 percent of their SOC stocks after 20 years (Poehlau *et al.*, 2011).

Soil organic carbon sequestration represents 25 percent of the total potential of climate change mitigation

solutions (23.8 Gt of CO<sub>2</sub>e per year) (Bossio *et al.*, 2020). Forty percent of potential solutions for climate change mitigation through soil carbon is to maintain the existing SOC stocks, and the remaining 60 percent is rebuilding the depleted SOC stocks. The historical loss of substantial amounts of SOC confers black soils a low enough baseline to achieve significant SOC gains on the path to recovery. Nature-based solutions based on SOC-centered sustainable management practices have multiples benefits, and no tradeoffs have been identified (Smith *et al.*, 2020). The major potential for SOC sequestration is in black soils devoted to annual crops. This potential is mainly due to the large yield gaps and/or large historic SOC losses (Amelung *et al.*, 2020). After an appropriate land use and soil management, these black soils can increase their SOC and improve their quality. As a result, the rise of atmospheric CO<sub>2</sub> can be mitigated in black soil regions (Liu *et al.*, 2012). In conclusion, sustainable use and management of black soils toward maintaining or increasing their SOC stocks could be key for climate change mitigation and adaptation.

Black soils will contribute to mitigation through increases in carbon by sequestration in their profiles resulting from the adoption of nature-based practices such as those reviewed by Smith *et al.*, (2020). Many of these practices, such as improved cropland management, improved forest management, and increased SOC content, are based on more intensification and do not create demand for more land conversion. This land productivity increase for food production can avoid emissions that would occur through expansion of the agricultural land area (Mueller *et al.*, 2012), or by reducing the greenhouse gas intensity of products (Bennetzen, Smith and Porter, 2016). Improved cropland, grazing, and livestock management have moderate carbon mitigation potential, although their impact can be high because of the high number of hectares that they occupy.

# Blacksoils, a **sink** or a **source** of carbon?



**Figure 3.2.2** Duality of black soils as carbon sinks or emitters as a function of management practices  
 Source: Author's elaboration



### 3.2.3 Human well-being

Black soils contribute to human well-being by providing food security, filtering water, protection against chemicals and pathogens, and cultural ecosystem services (Brevik and Sauer, 2015) (See Figure 3.2.3). After hundreds of years of farming, black soils continue to be a symbol of healthy and nutritional food in many local cultures (Liu *et al.*, 2012). Black soils contain sufficient nutrients and provide nutritious food to people living there and in other regions, thus avoiding negative effects on human health (Steffan *et al.*, 2018).

There is evidence of the contribution of black soils made by ancient civilizations, as found in the Amazon region where pre-Columbian indigenous communities cultivated lowlands hundred years ago, left a legacy of charcoal, fish bones, and organic matter. The highly fertile soils, now called Amazonian Dark Earths evolved through these materials (Kern *et al.*, 2019; Anne, 2015; Schmidt *et al.*, 2014). Cultural values associated with black soils are observed in northeast China, where people associate them with healthy and positive characters to enhance the value of their personality, products, and culture (Cui *et al.*, 2017).



© Zhiqiang Zhou

# Multiple benefits of black soils




Figure 3.2.3 Multiple benefits of black soils

Source: Author's elaboration



### 3.2.4 Food production and food security

Global analysis shows that out of the total land dedicated to growing crops, 17 percent of the farmland is currently occupied by black soils (Chernozems, Kastanozems and Phaeozems) (IUSS Working Group WRB, 2006), and out of the total area covered by black soils, one third of the black soil area is used as croplands (FAO, 2022a), due, in part, to its inherent fertility. This high fertility often leads to underestimation of the risks of degradation, although these soils are strongly affected by irreversible degradation processes such as erosion, nutrient imbalance, compaction and structural degradation (FAO and ITPS, 2015).


In cold regions of east Europe and Eurasia, there are black soils with high inherent fertility (Chernozems). If the annual weather such as precipitation and temperature is supportive, these soils can ensure food security for the countries (Avetov *et al.*, 2011; Kogan, Adamenko and Kulbida, 2011; Kobza and Pálka, 2017). Food security is a national priority in China, a country where black soils are considered the food basket since the 1950s. They have been responsible for the production of 15.9 percent of rice, 33.6 percent of

maize, and 33.9 percent of soybeans in 2014 (Bureau of Statistics of China, 2015). In the United States of America, black soils cover 31.2 million hectares, and 42 percent of them are used for crop production (Soil Survey Staff, 2014; FAO, 2022a). In the southern cone of South America, most of the black soils sustain the production of grain and oilseed crops, orchards, forage, and crops for fibre production. They are also used for cattle raising and dairy farming, feeding the cattle with grains, forage crops or natural pastures (Durán, 2010; Durán *et al.*, 2011; Rubio, Pereyra and Taboada, 2019).

A set of international initiatives, such as the International Network on black soils, the 4 x 1 000 Initiative (Soussana *et al.*, 2019), and the framework of the Global Soil Partnership (Rojas *et al.*, 2016) have highlighted the need to maintain healthy soils and address threats to more fertile soils in order to cope with a 60 percent increase in food demand by 2050. Black soils are very fertile, and they are considered “the world crop basket” or “giant pandas in cultivated farmland” (Zhang and Liu, 2020) (See Figure 3.2.4). They are expected to receive increased use pressure in future decades that require better management practices and governance.



# Black soils for food security



**Figure 3.2.4** Black soils as a key player for global food security

Source: Author's elaboration




### 3.3 Main threats to black soils

The Status of the World's Soil Resources report (FAO and ITPS, 2015) highlighted the most significant threats to soil functions at the global scale, specifically soil erosion, loss of SOC and nutrient imbalance, and the current outlook is that this situation will worsen unless concerted actions are taken by all, the private sector, governments, international organizations and academia.

Black soils are not the exception and are affected by all global threats. As already mentioned, most of the cultivated black soils have already lost at least half of their carbon stocks and suffer from moderate to severe erosion processes, among other degradation processes. Other ongoing soil threats are soil nutrient unbalances, soil sealing and soil biodiversity loss (See Figure 3.3a).

#### World's soils are **under threat**



**Figure 3.3a** Global assessment of the four main threats to soil by FAO regions

**Source:** Montanarella, L., Pennock, D.J., McKenzie, N., Badraoui, M., Chude, V., Baptista, I., Mamo, T., Yemefack, M., Aulakh, m.s., Yagi, K., Hong, Suk Young., Vijjarnsorn, P., Zhang, G., Arrouays, D., Black, H., Krasilnikov, P., JSobocká, A., Alegre, J., Henriquez, C.R., Mendonça-Santos, M.L., Taboada, M., Espinosa-Victoria, D., AlShankiti, A., AlaviPanah, S.K., Elsheikh, E.A.E.M., Hempel, J., Arbestain, M.C., Nachtergaele, F. & Ronald V. 2016. World's soils are under threat. SOIL, 2(1): 79-82. <https://doi.org/10.5194/soil-2-79-2016>

### 3.3.1 Soil organic carbon loss

Land use changes and unsustainable management practices lead to generalized significant soil organic carbon (SOC) losses in black soils. SOC changes appear as the second main threat in South America due to deforestation, intensive cultivation of grasslands and monocultures, in northeast China due to land use change and degradation of grassland, and in Europe due to the replacement of the natural vegetation; all of these are regions where black soils are predominant or at least conspicuous (See Figure 3.3a).

The black soils of Ukraine provide a well-documented example of SOC loss. Since 1970 there have been significant changes in the reserves of organic matter in Ukraine. Average losses of SOC due to irrational land use over 140 years since the time of V. V. Dokuchae have reached 22 percent about 19 percent in the Steppe, and more than 20 percent in Polissya (Baliuk and Kucher, 2019).

According to Yatsuk (2015, 2018), the largest losses of humus occurred from the 1960s to 1980s, due to the intensification of agricultural production by increasing the area of row crops, especially sugar beets and corn. During this period, the annual losses of humus reached 0.55 to 0.60 tonnes/ha. These processes of soil dehumidification on agricultural lands continue. According to the results of agrochemical certification of agricultural lands during the last five rounds (1986 to 2010) the humus content in the soils of Ukraine decreased by 0.22 percent in absolute terms and is 3.14. In terms of soil and climatic zones, the largest decrease in humus content occurred in the soils of the steppe zone, dropping from 3.72 to 3.40 percent, (by 0.32 percent in absolute terms). In the Forest-Steppe these changes are slightly smaller but given the loss of humus are significant 0.19 percent. However, the dynamics of losses in the period up to 2015 is somewhat slowing down due to the introduction of new management practices (Yatsuk, 2018).

Studies from Canada also document the complex balance between organic matter additions and losses in black soils. In studies by Landi *et al.*, (2003 a, b, 2004), the net primary production of vegetation (NPP) of seeded forage grasses (based on dry matter) for the black Chernozems annual averaged about 490 g/m<sup>2</sup> for above ground and 206 g/m<sup>2</sup> below ground. The amount of organic C to a 1.2 m depth is nearly 150 MgC/ha. Annual average rates in the three soils studied 1.18 gC/M<sup>2</sup>. Many researchers have suggested that prairie soils have lost about 30 percent of their organic matter under cultivation. The loss is estimated to be about 1.5 to 2 kg/m<sup>2</sup> by Mann (1986). Considering these losses

over 80 years of agriculture practices, the annual rate of loss is about 19 to 25 gC/m<sup>2</sup>. This is ten to thirty times greater than the accumulation rate in the Black Chernozems. This rate is likely to be higher at the early stage of C losses and before levelling off and reaching. Therefore, it may take only a few hundred years to lose the majority of the SOC. Organic C in subsoils is older than the SOC in A horizon and, therefore, can represent a vegetation composition different than that of today (Mermut and Acton, 1984).


Losses of SOC can also occur due to grazing. In soils of the Anaime moorland (Tolima, Colombia) (between 0 and 30 cm deep) pastures in use stored SOC of 34.4 tonnes/ha, while pastures without use for 20 years stored 22 tonnes/ha. One explanation is the possible increase of biomass in the fine roots that when decomposed provide greater carbon content to the soil (Maia *et al.*, 2010), while lack of pasture renewal probably decreases the contribution of senescent roots (Andrade, Espinosa and Moreno, 2014). However, this aspect requires further study (Castañeda-Martín and Montes-Pulido, 2017). Avellaneda-Torres, Leon-Sícard and Torres-Rojas, (2018) and Otero *et al.*, (2011) found the behaviour of organic C showed a trend of moorlands > potato farms > cattle farms. Similar results were found in Chingaza natural national park (NNP) and Nevados NNP, where the C was lower in the soil profiles of conserved highland ecosystems than in non-conserved ecosystems. The decrease in easily oxidised organic C in soils under potato and cattle farming might have been caused by the loss of native vegetation cover due to cattle farming relative to the Paramo, which exposed the soil to environmental factors such as water, air, and solar radiation and likely increased erosion (Otero *et al.*, 2011).

Restoring SOC stocks in black soils via reasonable management such as conservation tillage, manure and compost fertilization, and biomass management is crucial to sustainable development and is important for environmental stability (Xu *et al.*, 2020). Among the causes of the decline in SOC are land use change, aggressive tillage, inadequate cropping system management (such as monocultures), and limited replacement of nutrients (FAO and ITPS, 2015) (See Figure 3.3b). Studies in the Russian Federation and Ukraine show that the loss of soil vegetation cover favours erosion processes, and soil organic matter content (SOM) can decline by 15, 25, and 40 percent in weak, medium, and severely eroded black soils (Iutynskaya and Patyka, 2010).

Using a quantitative global SOM-crop yield potential model, Oldfield, Bradford and Wood (2019) found

that wheat and maize yields are greater with higher concentrations of SOC and level off at approximately 2 percent SOC. Potential yield increases through higher SOC concentrations amount to 32 percent of the projected yield gap for maize and 60 percent of that for wheat.

The Status of the World's Soil Resources report (FAO and ITPS, 2015) concluded that a priority action should be to stabilize or increase the global SOM stocks (SOC and soil organisms). Locally appropriate SOC-improving management practices should be identified by each country and facilitate their implementation towards a national-level goal of achieving a stable or positive net SOC balance. Black soils should be prioritized to maintain and increase SOC stocks.



**Figure 3.3b** Mayor drivers of SOC loss  
 Source: Author's elaboration



# 15 | Recarbonization of global soils (RECSOIL)



RECSOIL is a FAO innovative initiative with the aim to boost soil health through the maintenance and enhancement of SOC stocks (FAO and ITPS, 2021). It unlocks the potential of SOC to provide multiple benefits through key ecosystem services. Healthy soils directly contribute to enhance food security and farm income, reducing poverty and malnutrition, providing essential ecosystem services, contribute to the achievement of the SDGs, fight climate change, and build soils' resilience to extreme climatic events and to pandemics. Black soils are the most productive carbon-rich soils and contain 8.2 percent of the world's SOC stocks. Their SOC sequestration potential is 10 percent of the global annual potential (FAO, 2022). However, this is not evenly distributed throughout the world. For example, in Europe and Eurasia, black soils account for 66 percent of the potential SOC sequestration, while only reaching 10 percent in Latin America and the Caribbean. Therefore, it is critical to prioritize those areas to restore and maintain SOC stock and avoid losses. That can be done through the implementation of initiatives such as RECSOIL at country level to unlock the potential of these precious soils for climate changes adaptation and mitigation, and halt greenhouse gases emissions.

### 3.3.2 Soil erosion

Globally soil erosion was identified as the most severe threat, leading to poorer water quality in developed regions and to lower crop yields in many developing regions (Montanarella *et al.*, 2016). Figure 3.3a shows that soil erosion is the first main threat in regions where black soils are predominant or co-dominant, for example, South and North America, eastern Europe and northeast China.

Erosion induced by rainfall and wind degrades the quality of all soils, including black soils (See Figure

3.3.2). Due to the degree of severity that has occurred (such as deep gullies, total soil loss), many studies have been carried out during the last decade in the black soil region. The dominant soil erosion processes are due to water, wind, and snow meltwater, with water erosion on hillside farmland being the major contributor to soil erosion (Xu *et al.*, 2010). Ouyang *et al.*, (2018) observed that from 1979 to 2014, cropping system conversion from forestry to dry lands increased erosion losses from 204 to 421 tonnes per km<sup>2</sup> per year (Ouyang *et al.*, 2018). These losses can be controlled by basin, contour, rat tunnel, and conservation tillage,



in combination with terraces and strip cultivation. Crop productivity can be increased by fertilizer or manure application (Liu *et al.*, 2011).

In Japan, Taniyama (1990) reported that there was about 40 to 50 cm of topsoil loss by water erosion during the 16 years after land use change from forest to vegetable plantation in the Andosols of the northern Kanto region, Japan. Counter measures for soil erosion adopted in Japan include contour farming, cover cropping to prevent bare land, greenbelt farming, terracing. The Ministry of Agriculture, Forestry and Fishery (MAFF) in Japan has announced basic principles of soil water erosion and sediment control, namely enhancement of rain water percolation to reduce surface flow water, minimization of surface flow velocity, construction of channel networks to drain rain water rain water safely and decreasing soil erodibility.


More than 90 percent of the agricultural area of the Pampas region in Argentina is currently cultivated under no-till, more recently combined with the cultivation of “cover crops” during the fallow period. This tends to reduce the loss of organic carbon and soil erosion due to the generalization of no-till practices.

Wind erosion is a phenomenon that mainly affects soils in semi-arid and arid areas, which often have low levels

of plant cover and organic matter (Skidmore, 2017). In any case, climatic cycles with drought can generate predisposing causes of wind erosion, even in black soils, as happened with the dust bowl during the 1930s in the United States of America (Lee and Gill, 2015).

The combined effect of water and wind in Ukraine has been severe. The average annual soil loss from water and wind erosion is 15 tonnes/ha. This means that the country’s soil cover loses about 740 million tonnes of the top, fertile soil layer every year (Baliuk *et al.*, 2010). The amount of land in Ukraine damaged by water erosion is up to 32 percent of the total area (13.3 million hectares). Of these hectares, 4.5 million hectares with medium and heavily washed soils, as well as 68 000 hectares that have completely lost the humus horizon. More than 6 million hectares are systematically affected by wind erosion, and up to 20 million hectares in years with dust storms. A particularly potentially dangerous zone in Ukraine is the southern Steppe (the main zone with Kastanozems and Calcic Glosic Chernozems). Thus, the number of days per year with dust storms in the southern steppe zone is 159, northern and central is 88, Forest-Steppe is about 33 days (Baliuk *et al.*, 2010).

## Land use change



**Figure 3.3.2** Main soil erosion processes of black soils

Source: Author's elaboration





© Andrei Borisovich Ryzanov

**Photo 3.3.2a** Wind erosion in Liaoning province, China



© Andrei Borisovich Ryzanov

**Photo 3.3.2b** Water erosion in Laetoli Gorge, the United Republic of Tanzania




© Irina Kalimova

**Photo 3.3.2c** Wind erosion in Jiusan farm, China caused by land conversion and aggressive tillage



© Irina Kalimova

**Photo 3.3.2d** Water erosion in the Russian Federation





## 16 | The infamous Dust Bowl!



Dust storm approaching Stratford, Texas. Dust Bowl surveying in Texas.

The Dust Bowl was one of the largest sandstorms in the history of the United States of America that happened in the 1930s (History, 2020). This phenomenon began in the southern Great Plains and was caused by intensive farming, poor agricultural practices, and was associated with a period of severe droughts (History, 2020). Soil erosion and desertification on these lands caused massive dust storms affecting the states of Oklahoma, Kansas, Texas, New Mexico, and Colorado (Texas and Kansas have black soils) and reaching cities such as Washington, DC, and New York (Findmypast, 2015; SSSA, 2015). About 1.2 billion tonnes of soil were lost between 1934 and 1935 in the southern Great Plains (Britannica, 2022). These dust storms caused several adverse effects such as respiratory diseases causing the death of people and animals, with farmlands becoming unusable, and hunger and poverty spreading across several states (SSSA, 2015). Many people migrated to other places like California to escape the drought and the dust, and to find work (Findmypast, 2015). Following this huge catastrophe, the Soil Conservation Service (later the USDA Natural Resources Conservation Service) was founded to encourage farmers to adopt erosion mitigation strategies implementing sustainable management practices (reduced tillage, leaving crop residues in fields, strip cropping and crop rotation) to conserve soil and minimize erosion (SSSA, 2015). Desertification increasingly threatens significant land areas worldwide, affecting more than 100 countries, including the United States of America (SSSA, 2015).

Source: History. 2020. *Dust Bowl*. In: HISTORY. Cited 6 June 2022. <https://www.history.com/topics/great-depression/dust-bowl>

### 3.3.3 Soil nutrient imbalance

Nutrient imbalances include both deficits and excesses of soil nutrients. Imbalances were judged by Montanarella *et al.*, (2016) as the second main threat in North America, and the third greatest threat in most of Africa, in regions where black soils are conspicuous (See Figure 3.3.3).


Nitrogen (N) and phosphorus (P) fertilizer use need to be increased in infertile tropical and semi-tropical soils where the most food insecurity is found. Soil nutrient levels have decreased in the Argentine Pampas, because of the lack of resupply of nutrients, causing soil fertility to be exhausted in many places. Due to economic reasons, fertilizer use and soil testing was not historically widespread in Argentina and the level of nutrients such as N, P, calcium (Ca), magnesium (Mg) and zinc (Zn) has decreased (Rubio *et al.*, 2019; Lavado and Taboada, 2009). Nutrient stocks have also noticeably decreased in black soils of the Russian Federation (Grekov *et al.*, 2011; Medvedev, 2012), Ukraine (Balyuk and Medvedev, 2012), and Brazil (Rezapour and Alipour, 2017).

On the other hand, in other parts of the world with black soils, excessive N fertilization and decreasing N recovery rates by crops have caused dramatic increases in non-point source pollution from agriculture (Ju *et al.*, 2004). Nutrient excesses often originate in the use of high doses of synthetic fertilizers and organic

manures containing N and P, with the consequent risks of pollution and therefore the eutrophication of groundwater and surface water (See Figure 3.3.3).

Soil phosphorus is strongly affected by interaction with soil minerals and organic matter. Due to this, P added as fertilizer (14.2 Tg P/year) and manure (9.6 Tg P/year) collectively exceeded P removal by harvested crops (12.3 Tg of P/year) at the global scale (Zang *et al.*, 2017). However, almost 30 percent of the global cropland area, particularly in Europe and South America, is deficient in soil P, either total or extractable by crops. Soil P deficits are common in areas producing forage crops used as livestock feed (MacDonald *et al.*, 2011). On the other hand, high P fertilizer application relative to crop P use resulted in a greater proportion of intense P surpluses (>13 kg of P/ha/year) in many areas with black soils. Together with N excesses, P surpluses represent a risk of eutrophication of freshwater and marine ecosystems (Dodds and Smith, 2016; Ngatia *et al.*, 2019). In Japan the overuse of chemical fertilizer causes nutrient imbalance in Andosols. Potassium is excessively accumulated in upland soils, and the ratio of magnesium to potassium is low (Japanese Soil Conservation Research Project Nationwide Council, 2021). The level of available phosphate in upland soils tends to be higher than the governmental recommendation (Japanese Soil Conservation Research Project Nationwide Council, 2021).





**Figure 3.3.3** Effects of intensive use as a trigger on nutrient imbalance

Source: Author's elaboration





### 3.3.4 Soil compaction

Excessive soil compaction is a direct consequence of intense agricultural traffic of heavy machinery in the fields (See Figure 3.3.4), and a higher soil susceptibility because of organic matter content decreases and lower aggregate stability (Gupta and Allmaras, 1987; Montanarella *et al.*, 2016). Soil compaction is indicated by increases in soil bulk density and soil penetrometer resistance and decreases in soil macroporosity and water infiltration rates, among changes in other soil properties (Gupta and Allmaras, 1987; Liu *et al.*, 2010), with important consequences on crop yields (Liu *et al.*, 2010; Peralta, Alvarez and Taboada, 2021). Evidence of soil compaction and physical deterioration of black soils is widely available. After 75 years of cultivation, water-stable aggregation declined by 27 percent and clay content by 27 percent in black soils of the Russian Federation (Balashov and Buchkina, 2011). This soil physical decline reached 40 percent in Ukraine, where many soils have a compacted layer (Balyuk and Medvedev, 2012). Likewise, intensive cultivation and summer fallowing have degraded the Canadian prairie soils, resulting in poor surface structure (Agriculture and Agri-Food Canada, 2003). Without significant variation, 14 to 20 percent higher bulk density and 10 to 22 percent lower porosity values were observed in cultivated black soils compared to forestlands in Brazil (Rezapour and Alipour, 2017).

Soil physical degradation has covered almost the entire area of distribution of black soils in Ukraine. This is due to a number of factors, including excessive ploughing of agricultural land (78 percent) (Medvedev, 2012; Yatsuk, 2015; Yatsuk, 2018; National report on the state of the environment in Ukraine in 2018, 2020), due to their suboptimal structure, intensive mechanical tillage led to widespread physical degradation. Physical degradation is manifested in the destructing of the upper layer, blocky (cloddy) after ploughing, swimming

and crusting, the presence of a plough pan, and subsoil compaction. Physically degraded soils are prone to erosion, poor water retention properties, thus limiting the development of plant root systems (Baliuk *et al.*, 2010; Medvedev, 2012). In addition, with the current trends of climate change (aridization and warming) in Ukraine, there is already a de facto shift of natural-climatic zones from 100 to 150 km to the north, which brings new threats to desertification (Zatula and Zatula, 2020). These processes were already clearly traced 20 years ago (Pylypenko *et al.*, 2002).


Soil compaction is not only a consequence of tillage, as even under continuous conservation tillage farming it was repeatedly observed (Peralta, Alvarez and Taboada, 2021). In this case, the process affects the first layer of the soil, promotes planar aggregates and associated voids in crop rotations with long fallow periods (Alvarez *et al.*, 2014; Peralta, Alvarez and Taboada, 2021).

Additionally, burning, intensive grazing, tilling, and replacement of the natural grassland with more nutritive grass species in Colombia significantly affected water balance of the Colombian moorlands areas (Sarmiento and Frolich, 2002). Phenomena typically accompanying pasture farming and tillage, such as soil compaction and soil crusting, additionally alter the infiltration rates, water storage, and regulation capacity of moorlands.

The soils of the Pampas region in Argentine provide an example of the consequences of physical soil degradation. Continuous cultivation has generated soil physical degradation as sealing and compaction, increasing the processes of water erosion. Since the soils of the Pampas region, particularly in the Undulating Pampa, have been put into cultivation, they have lost an average of 50 percent of their original organic matter, while the total phosphorus of the surface horizon would have decreased by 80 percent (Sainz-Rozas, Echeverria and Angelini, 2011; Lavado, 2016).



# Good soil structure vs. compacted soil



**Figure 3.3.4** Effects of soil compaction

Source: Author's elaboration



**Photo 3.3.4a** Soil compaction caused by heavy machinery in Jilin province, China



**Photo 3.3.4b** Soil compaction caused by heavy machinery in Jilin province, China




### 3.3.5 Salinization

Salinization is a related processes that result from both natural (primary) and human-induced (secondary) processes (See Figure 3.3.5). In black soils the cases are due to: a) hydrological imbalances caused by changes in land use (such as the replacement of perennial vegetation such as forests, grasslands and pastures by annual crops), or associated with climate change, which causes the rise of saline groundwater to the surface (Taboada *et al.*, 2021); or b) the use of irrigation water with moderate to high salt content (Choudhary and Kharche, 2018; Bilanchyn *et al.*,

2021). In both cases, the increases in pH, electrical conductivity, and percentage of exchangeable sodium, decrease the quality of black soils. However, human-induced salinization owing to inappropriate soil and fertilizer management are the main challenges in regions of black soils. Secondary salinization of irrigated soils, accompanied by a reduction of the humus-rich layer depth was reported in the Russian Federation (Grekov *et al.*, 2011; Medvedev, 2012). In other cases, soil salinity in black soils is associated with swelling-shrinking processes and is not shown by saltpans on surface (Choudhary and Kharche, 2018).

## Soils affected by human-induced salinity



**Figure 3.3.5** Soils affected by human-induced salinity


Source: Author's elaboration



### 3.3.6 Acidification

The acidification of black soils is most commonly due to an excessive extraction of exchangeable bases (Ca, Mg, K) by crops without adequate replenishment or a consequence of N fertilization (See Figure 3.3.6). Acidification only appears as the first main threat in Oceania (See Figure 3.3a). In the regions of Cherkassy and Sumy (Ukraine), soil pH dropped 0.3 to 0.5 units after 40 to 50 years cultivation (Grekov *et al.*, 2011; Medvedev, 2012). In black soils of northeast China, from 2005 to 2014, a trend of acidification due to overuse of N fertilizers was detected in intensive cropping systems (Tong, 2018). It is interesting to note that in Chinese croplands, N-induced acidification was also associated with an accrual of soil organic matter (Zhang and Liu, 2020), and a great decrease in soil inorganic carbon (Raza *et al.*, 2020). Andosols in Japan can experience nutrient issues associated with acidification. In general, allophanic

Silandic Andosols are originally weakly acidic, and aluminum toxicity does not occur frequently in plants on these soils. However, Silandic Andosols can become strongly acidic following the heavy application of chemical fertilizer (Fujii, Mori and Matsumoto, 2021). Strongly acidic Andosols with an accumulation of acidic materials dissolves a part of the active Al fraction in the soils, which causes Al toxicity and thereby leads to the shallow rooting of Al-susceptible crops (Fujii, Mori and Matsumoto, 2021). Additionally, the soil productivity of strongly acidic Andosols is lower than that of the original weakly acidic soil; for example, the number of bacteria decreases (e.g, from  $160 \times 10^6$  cfu /g for weakly acidic soil to  $10 \times 10^6$  cfu /g for strongly acidic soil) as do the levels of readily mineralizable soil nitrogen (Matsuyama *et al.*, 2005). To improve these nutritional imbalances, the appropriate use of soil amendments based on soil diagnosis is desired (Japanese Soil Conservation Research Project Nationwide Council, 2021).



# Soil acidification


## Causes

- Anthropic
  - Excessive inputs of nitrogen (as  $\text{NH}_4^+$ )
  - Legume monoculture
  - Acid rain
  - Removal of all harvested materials and no reposition of exchangeable bases
- Natural
  - High rainfall and weathering oxidation

## Effects

- pH drop (< 5.5)
- Leaching of nitrates ( $\text{NO}_3^-$ )
- Soil pollution
- Toxicity:  $\text{Al}^{3+}$ ,  $\text{H}^+$ , Fe and Mn sesquioxides
- Deficiency: P,  $\text{Ca}^{2+}$ , Mo,  $\text{Mg}^{2+}$

## Soil pH chart



**Figure 3.3.6** Causes and effects of soil acidification


Source: Author's elaboration



### 3.3.7 Soil biodiversity loss

Unlike other aspects of soil science, soil biodiversity remains poorly understood, in terms of anthropogenic impacts on the diversity of microbes and soil fauna that live unseen in soils. Only a small fraction of the immense morphological diversity of soil organisms is

known, and this is especially true for microorganisms. Examples of this biological richness include bacteria (a), microscopic (b) and fruiting bodies of fungi (c), viruses (d), algae (e), protists (f), nematodes (g), mites (h), springtails (i), enchytraeids (j), earthworms (k), mealybugs (l), termites (m), ants (n), and mammals (o), among many others (Figure 3. 3.7a).



**Figure 3.3.7a** Overview of the most common soil biodiversity groups


Examples shown are bacteria (A), microscopic (B) and fruiting bodies of fungi (C), viruses (D), algae (E), protists (F), nematodes (G), mites (H), springtails (I), enchytraeids (J), earthworms (k), mealybugs (L), termites (M), ants (N), and mammals (O), among many others

Source: Author's elaboration

Most of black soils evolved supporting grassland vegetation with an active rhizosphere around the fibrous root system of the dominant grasses (Tisdall and Oades, 1982; Oades, 1993). These grasslands were characterized by floristic richness in plant communities, and an enormous soil biodiversity (from microbes to megafauna) which plays a key role in the provision of essential ecosystem services as carbon sequestration, nutrient cycling (carbon, nitrogen, phosphorus, and sulphur), water retention, provision of nutritious food, among others.

One of the consequences of the transition to crops from grasslands on black soils is the loss of much of the original biodiversity, at levels that are not well known because these changes occurred a long time ago. It is difficult to think that these soils will recover their pristine or near pristine state, so one of the future challenges is how to recover at least part of this enormous lost biodiversity.

The effects of reducing soil biodiversity on soil functioning was studied in an experiment by Wagg *et al.*, (2014). A broad soil biodiversity gradient was reproduced in grassland microcosms (Figure 3.3.7b). Some groups of soil organisms (nematodes and mycorrhizal fungi) were eliminated within the gradient, while fungal and bacterial communities reduced in abundance and richness. Plant species diversity decreased strongly with the reduction of soil biodiversity and the simplification of soil communities. This supports previous findings that plant community composition is driven by the diversity and species composition of various groups of soil organisms. As is expected (and repeatedly seen in real examples), carbon sequestration also decreased along the gradient. Changes in soil biodiversity and soil community composition also influenced processes related to nutrient cycling.



**Figure 3.3.7b** Change in soil community characteristics (abundance and richness) of various guilds of soil organisms in grassland communities

Means  $\pm$  SEM are expressed as a ratio of the most complete soil treatment (dashed line), such that 0 represents no detection. The coloured lines highlight the general trend of changes in soil community characteristics along the gradient.

**Source:** Wagg, C., Bender, S.F., Widmer, F. & Van Der Heijden, M.G. 2014. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences*, 111(14): 5266–5270. <https://doi.org/10.1073/pnas.1320054111>

This simplification of soil communities due to cultivation has also been observed in the field. A valuable example of changes in arthropod communities (spiders, ants and carabids, among others) is provided by a study on black soils in northern China (Gao *et al.*, 2021). Due to intensive agricultural practices, a more simplified species richness and biodiversity was achieved at the local scale.

A well-documented effect of soil organisms is on the aggregation of the soil. The structure of the topsoil of black soils presents a hierarchical organization of aggregate sizes that depends on this enormous

biodiversity. In fact, soil macroaggregates and soil clumps (units > 250 microns) depend for their stability on the binding and bonding mechanisms from cementing agents excreted in the soil rhizosphere and entanglement by fine roots and mycorrhizal hyphae. Soil mesopores (0.2 to 30 microns in diameter) and macropores (larger than 30 microns) are the habitat of many of the microorganisms and microfauna that make up soil biodiversity (Degens, 1997; Kay, 1990; Chantigny *et al.*, 1997). The protection of SOC within aggregates is a key element of increasing SOC levels to achieve carbon sequestration.





## 17 | What is the relationship between wars and soil pollution?

The use of artillery possesses a severe risk to soil degradation and pollution especially with the use of mechanization and modern weapon technologies. The most common sources of pollution affected by armed conflicts are the following contaminants:

Sources of pollution	Soil contaminants
Conventional explosives	TNT, RDX, HMX
Fragmentation shells, bullets, cartridge cases and shotgun pellets	Copper, iron, lead and zinc
Armour-piercing projectiles	Depleted uranium
Incendiary weapons	White phosphorus and Napalm

Soil pollution can come from the use of nitro aromatic explosive compounds (FAO and UNEP, 2021). These compounds have a high persistence and once entered the soil, they tend to remain, harming the local biota and reducing the soil health and fertility. The negative effects from the use of incendiary weapons containing white phosphorus comes from their co-contaminants and residues of combustion. Such weapons may result in soil polluted with trace elements, hydrocarbons, organic solvents, surfactants, synthetic phenols, cyanide, dioxins, and radionuclides reducing soil fertility, crop yield and possessing risk to human health and the environment. The use of depleted uranium, one of the least studied forms of uranium, can penetrate the soil as deep as 50 cm. The dust that is emitted from the depleted uranium disperses and contaminates and polluting the soil over large area. Claims have been made that the depleted uranium dust can travel up to 40 km. After an attack with depleted uranium ammunition, this dust will be deposited on the ground and other surfaces as partially oxidized depleted uranium fragments of different sizes, and as uranium oxide dust.

The sources of soil pollution are very varied and range from the primary sector to the final stages of the life cycle of everyday products. For this reason, in order to prevent and reduce soil pollution by the armed conflicts, greater efforts must be made to reduce the use and production of toxic chemicals in the ammunition, to regulate and control industries and verify that their emissions do not introduce contaminants into the environment and that production and consumption systems move towards more sustainable schemes in which waste production is reduced.

Source: FAO & UNEP, 2021. *Global assessment of soil pollution – Summary for policy makers*. Rome, FAO. <https://doi.org/10.4060/cb4827en>

## 3.4 Challenges

The challenges faced by black soils arise from their preponderant role in food and fibre production under agricultural and livestock use. As mentioned before, the main threats that operate on them are erosion by water, tillage and wind, loss of SOC and organic matter, and nutrient imbalance. In addition, physical-structural deterioration should not be ruled out, and in some regions soil salinization, pollution with excess fertilizers and agrochemicals, and soil sealing due to urban advancement also occurs (FAO and ITPS

2015; Montanarella *et al.*, 2016). Main drivers of soil degradation are land use changes and unsuitable land management, unsustainable management practices and the lack of policies.

According to the FAO and ITPS, (2015) report, soil degradation should be minimized and degraded soils restored in those regions where people are most vulnerable, or where food production is critical, Global SOC and SOM stocks should be stabilized or increased, and that we should act to stabilize or reduce global N and P fertilizer use while simultaneously increasing fertilizer use in regions of nutrient deficiency.



**Photo 3.3.7** Black soils in Zhaoguang farm of Heilongjiang province, China

### 3.4.1 Land use change and land management

Due to projected increases in food demand crop production will need to increase by 70 to 110 percent by 2050 (Royal Society of London, 2009; Tilman *et al.*, 2011). As very fertile soils, black soils will be among the main actors in these increases in production and closing of gaps. To meet this objective, sustainable intensification of the current productive systems rather than agricultural expansion into forests and pastures is critical (Fischer and Connor, 2018; Guilpart *et al.*, 2017). Many black soils are also considered soils of high environmental value, where the protection of large carbon stocks and the restoration of these stocks should be included in overall soil resilience programmes to monitor, restore and maintain soil fertility and soil functions, and to enhance the key ecosystem services provided by these soils (Smith *et al.*, 2016).

As a successful example, in recent years, the Ministries of Agriculture, Science and Technology, Land and Resources, and the four provinces of black soil areas in northeast China have actively implemented a wide range of measures to protect and enhance black soils. These include high standard farmland construction, soil and water conservation, soil testing and formulated fertilization, soil organic matter increases, conservation tillage (reduced tillage and no-till), subsoiling for soil compaction alleviation and soil preparation, straw returning, and increasing the use of organic fertilizers (Li *et al.*, 2021). The comprehensive goal of black soil protection and utilization is to control the loss and degradation of black soils and keep water and fertility (Han *et al.*, 2018).

The adoption of restorative land use and recommended management practices are key to strengthen numerous ecosystem services provided by black soils, such as improving water quality and renewability, increasing below and above-ground biodiversity, enhancing soil resilience to climate change and extreme events, and mitigating climate change by sequestering carbon in soil and reducing CO<sub>2</sub>, CH<sub>4</sub> and N<sub>2</sub>O emissions (Lal, 2014).

### 3.4.2 Unsustainable management practices

The most cited unsustainable management practices for black soils are those that cause the main threats, that is, erosion, loss of SOC and SOM, nutrient imbalance, and salinization and sodification. Under agricultural management, these practices are usually aggressive

tillage methods based on plowshares and disc plows, harrows, and so on, monocultures, non-replenishment of nutrients with fertilizers, and the disappearance of pastures. Overgrazing and non-use of rational grazing systems is the most frequent cause on land destined for grazing. Soil pollution by pesticide residues is a major cause of degradation (Smith *et al.*, 2016).

### 3.4.3 Climate change and black soils

Climate change is very much related to the condition of black soils. From one side, climate change negatively impacts black soils. For instance, interactions between the increasing temperature and decreasing precipitation in black soil region led to reduced accumulation of soil organic matter, which results in poor soil fertility (Gong *et al.*, 2013). On the other side, the unsustainable management of black soils causes the loss of soil organic carbon and emits greenhouse gases to the atmosphere, exacerbating climate change. Evidence in black soils showed that organic matter amendments and tillage management can mitigate negative and exploit positive effects of climate change on crop production by enhancing soil quality (Song *et al.*, 2015; Menšík *et al.*, 2019; Farkas, *et al.*, 2018). Unfortunately, those practices are not often adopted by local governments and farmers due to multiple management and economic obstacles. The restoration of degraded black soils should be highlighted and inputted in the global climate change agenda in offsetting anthropogenic emissions and SOC sequestration (Lal, 2021).

### 3.4.4 Lack of policies

Many of the black soils appear in countries with low levels of soil governance, or where different laws and regulations relating to soil and water may exist, but with poor enforcement. Undoubtedly, the lack of effective policies in these sites is a major challenge to preserve the quality and health of black soils for agricultural use, and thus food security.

The implementation of practices for recovery or restoration of black soils mostly depends on good governance (in North America) and the availability of financial resources. This factor limits implementation of sustainable practices in all the affected countries, but mostly in developed countries. The problem lies in the fact that most of the food insecurity problems are not in the more developed countries but in the less developed parts of the world. This is the main challenge for sustainable management, not only for black soils, but also for all the productive soils in the world. Sustainable soil management should increase the supply of healthy food.



# References

- Abrar, M.M., Xu, M., Shah, S.A.A., Aslam, M.W., Aziz, T., Mustafa, A., Ashraf, M.N., Zhou, B. & Ma, X. 2020. Variations in the profile distribution and protection mechanisms of organic carbon under long-term fertilization in a Chinese Mollisol. *Science of the Total Environment*, 723: 138181.
- Allen, V. G., Batello, C., Berretta, E. J., Hodgson, J., Kothmann, M., Li, X., Melvor, J., Milne, J., Morris, C., Peeters, A. & Sanderson, M. 2011. An international terminology for grazing lands and grazing animals. *Grass and Forage Science*, 66(1): 2–28.
- Adhikari, K. & Hartemink, A.E. 2016. Linking soils to ecosystem services—A global review. *Geoderma*, 262: 101–111. <https://doi.org/10.1016/j.geoderma.2015.08.009>
- Agriculture & Agri-Food Canada. 2003. Prairie soils: the case for conservation. Cited 15 September 2020. <http://www.rural-gc.agr.ca/pfra/soil/prairiesoils.htm>
- Agriculture & Agri-Food Canada. 2011. *Soil erosion indicators in Canada*. Government of Canada, Ottawa.
- Almeida, J.A. 2017. Solos das pradarias mistas do sul do Brasil (Pampa Gaúcho). In: N. Curi, J.C. Ker, R.F. Novais, P. Vidal-Torrado & C.E.G.R. Schaefer, eds. *Pedologia: Solos dos Biomas Brasileiros*, pp. 407–466. 1ª Edição. Viçosa, MG: Sociedade Brasileira de Ciência do Solo.
- Alvarez, C. R., Taboada, M. A., Gutierrez Boem, F. H., Bono, A., Fernandez, P. L. & Prystupa, P. 2009. Topsoil properties as affected by tillage systems in the Rolling Pampa region of Argentina. *Soil Science Society of America Journal*, 73(4): 1242–1250.
- Alvarez, C.R., Taboada, M.A., Perelman, S. & Morrás, H.J.M. 2014. Topsoil structure in no-tilled soils in the Rolling Pampa, Argentina. *Soil Research*, 52(6): 533–542. <https://doi.org/10.1071/SR13281>
- Amelung, W., Bossio, D., de Vries, W., Kögel-Knabner, I., Lehmann, J., Amundson, R., Bol, R. et al., 2020. Towards a global-scale soil climate mitigation strategy. *Nature Communications*, 11(1): 5427. <https://doi.org/10.1038/s41467-020-18887-7>
- Amiro, B., Tenuta, M., Hanis-Gervais, K., Gao, X., Flaten, D., Rawluk, C. & Lupwayi, N. 2017. Agronomists' views on the potential to adopt beneficial greenhouse gas nitrogen management practices through fertilizer management. *Canadian Journal of Soil Science*, 97(4): 801–804.
- Andrade, B.O., Koch, C., Boldrini, I.I., Vélez-Martin, E., Hasenack, H., Hermann, J.M., Kollmann, J., Pillar, V.D. & Overbeck, G.E. 2015. Grassland degradation and restoration: a conceptual framework of stages and thresholds illustrated by southern Brazilian grasslands. *Natureza & Conservação*, 13: 95–104.
- Andrade, H., Espinosa, E. & H. Moreno. 2014. Impact of grazing in soil organic storage carbon in high lands of Anaime, Tolima, Colombia. *Zootecnia Tropical (Venezuela)*. 32(1):7–21.
- Anne, S.B. 2015. The secret of black soil. DW, 20 January 2015. In: *DW.COM*. Cited 30 May 2022. <https://www.dw.com/en/the-secret-of-black-soil/a-18199797>
- Antonenko, D.A., Nikiforenko, Y.Y., Melnik, O.A., Yurin, D.A. & Danilova, A.A. 2022. Organomineral compost and its effects for the content of heavy metals in the top layer leached chernozem. *IOP Conference Series: Earth and Environmental Science*, pp. 012028. IOP Publishing.
- Aquino, R.E., Campos, M.C.C., Oliveira, I.A., Marques Júnior, J. & Silva, D.M.P. 2014. Variabilidade espacial de atributos físicos de solos antropogênico e não antropogênico na região de Manicoré, AM. *Bioscience Journal*, 30(5): 988–997.
- Arshad, M.A., Soon, Y.K. & Azooz, R.H. 2002. Modified no-till and crop sequence effects on spring wheat production in northern Alberta, Canada. *Soil and Tillage Research*, 65: 29–36.
- Assefa, B.A., Schoenau J.J. & Grevers M.C.J. 2004. Effects of four annual applications of manure on Black Chernozemic soils. *Canadian Biosystems Engineering*, 46(6): 39–46.
- Avellaneda-Torres, L.M., León-Sicard, T.E. & Torres-Rojas, E. 2018. Impact of potato cultivation and cattle farming on physicochemical parameters and enzymatic activities of Neotropical high Andean Páramo ecosystem soils. *Science of The Total Environment*, Volume 631–632: 1600–1610. <https://doi.org/10.1016/j.scitotenv.2018.03.137>.
- Avetov, N.A., Alexandrovskii, A.L., Alyabina, I.O., Dobrovolskii, G.V. & Shoba, S.A. 2011. *National Atlas of Russian Federation's soils*. Moscow, Astrel.
- Avila, L.A., Martini, L.F.D., Mezzomo, R.F., Refatti, J.P., Campos, R.L., Cezimbra, D.M., Machado, S.L.O., Massey, J., Carlesso, R.L. & Marchesan, E. 2015. Rice water use efficiency and yield under continuous and intermittent irrigation. *Agronomy Journal*, 107: 442–458.

- Azooz, R.H. & Arshad, M.A.** 1998. Effect of tillage and residue management on barley and canola growth and water use efficiency. *Canadian Journal of Soil Science*, 78: 649–656.
- Baethgen, W. & Morón, A.** 2000. Carbon sequestration in agricultural production systems of Uruguay: observed data and CENTURY model simulation runs. *Anales de la V Reunión de la Red Latinoamericana de Agricultura Conservacionista*. Florianópolis, Brasil.
- Bailey, A.W., McCartney, D. & Schellenberg, M.P.** 2010. Management of Canadian Prairie Rangeland. 13 October 2020. (also available at [https://www.beefresearch.ca/files/pdf/factsheets/991\\_2010\\_02\\_TB\\_RangeMgmt\\_EWEB\\_2\\_.pdf](https://www.beefresearch.ca/files/pdf/factsheets/991_2010_02_TB_RangeMgmt_EWEB_2_.pdf))
- Balashov, E. & Buchkina, N.** 2011. Impact of short- and long-term agricultural use of chernozem on its quality indicators. *International Agrophysics*, 25(1).
- Baliuk, S. A. & Kucher, A. V.** 2019. Spatial features of soil cover as a basis for sustainable soil management (In Ukrainian). *Ukrainian Geographical Journal*, 3 (107): 3–14. <https://doi.org/https://doi.org/10.15407/ugz2019.03.003>
- Baliuk, S. A., Miroshnychenko, M. M. & Medvedev, V. V.** 2018. Scientific bases of stable management of soil resources of Ukraine (In Ukrainian). *Bulletin of Agricultural Science*, 11: 5–12. <https://doi.org/https://doi.org/10.31073/agrovisnyk201811-01>.
- Baliuk, S., Nosonenko, A., Zakharova, M., Drozd, E., Vorotyntseva, L. & Afanasyev, Y.** 2017. Criteria and parameters for forecasting the direction of irrigated soil evolution. *Soil science working for a living*, pp. 149–158. Springer.
- Baliuk, S.A. & Miroshnychenko, M.M.** 2016. *Fertilizer systems of crops in agriculture at the beginning of XXI Century*. Kyiv, Ukraine, Alpha-stevia express.
- Balyuk S.A. & Medvedev, V.V.** 2012. *Strategy of balanced use, reproduction and management of soil resources of Ukraine (In Ukrainian)*. Kiev, Agrarian science.
- Balyuk, S.A. & Medvedev, V.V.** 2015. *The concept of organization and functioning of soil monitoring in Ukraine taking into account the European experience (scientific publication)* (In Ukrainian). NSC Sokolovsky Institute of Soil Science and Agrochemistry. Kharkiv, TOV “Smuhasta typohrafiya”.
- Balyuk, S.A., Medvedev, V.V. & Miroshnychenko, M.M.** 2018. *The concept of achieving a neutral level of degradation of lands (soils) of Ukraine* (In Ukrainian). NSC IGA. Kharkiv, Brovin O.V.
- Balyuk, S.A., Medvedev, V.V., Miroshnychenko, M.M., Skrylnyk, E.V., Tymchenko, D.O., Fateev, A.I., Khristenko, A.O. & Tsapko, Yu. L.** 2012. Ecological condition of soils of Ukraine (In Ukrainian). *Ukrainian Geographical Journal*, 2: 38–42.
- Balyuk, S.A., Medvedev, V.V., Tarariko, O.G., Grekov, V.O. & Balaev, A.D.** 2010. *National report on the state of soil fertility of Ukraine* (In Ukrainian). MAPU, State Center for Fertility, NAAS, NSC IGA, NULES.
- Banik, C., Koziel, J.A., De, M., Bonds, D., Chen, B., Singh, A. & Licht, M.A.** 2021. Biochar-Swine Manure Impact on Soil Nutrients and Carbon Under Controlled Leaching Experiment Using a Midwestern Mollisols. *Front. Environ. Sci*, 9(10.3389).
- Baron, V.S., Mapfumo, E., Dick, A.C., Naeth, M.A., Okine, E.K. & Chanasyk, D.S.** 2002. Grazing intensity impacts on pasture carbon and nitrogen flow. *Journal of Range Management*, 55: 535–541.
- Bedendo, D.**, 2019. Soils of Entre Ríos. In G. Rubio, R. Lavado, & F. Pereyra, eds. *The Soils of Argentina*, Chapter 4, pp. 165–173. Springer, Switzerland.
- Behling, H.** 2002. South and southeast Brazilian grasslands during Late Quaternary times: a synthesis. *Palaeogeogr.Palaeoclimatol. Palaeoecol.* 177: 19–27
- Belyuchenko, I.S. & Antonenko, D.A.** 2015. The influence of complex compost on the aggregate composition and water and air properties of an ordinary chernozem. *Eurasian soil science*, 48(7): 748–753.
- Bender, M.** 1971. Variation in the  $^{13}\text{C}/^{12}\text{C}$  ratios of plants in relation to the pathway of photosynthetic carbon dioxide fixation. *Phytochemistry*, 10: 1239–124.
- Bennetzen, E.H., Smith, P. & Porter, J.R.** 2016. Decoupling of greenhouse gas emissions from global agricultural production: 1970–2050. *Global Change Biology*, 22(2): 763–781. <https://doi.org/10.1111/gcb.13120>
- Bieganowski, A., Witkowska-Walczak, B., Glinski, J., Sokolowska, Z., Slawinski, C., Brzezinska, M. & Włodarczyk, T.** 2013. Database of Polish arable mineral soils: a review. *International Agrophysics*, 27(3).

- Bilanchyn, Y., Tsurkan, O., Tertyk, M., Medinets, V., Buyanovskiy, A., Soltys, I. & Medinets, S.** 2021. *Post-irrigation state of black soils in south-western Ukraine*. In: D. Dent & B. Boincean, eds. *Regenerative Agriculture*, pp.303–309. Cham, Springer International Publishing.
- Blackshaw, R.E., Molnar, L.J. & Moyer, J.R.** 2010. Suitability of legume cover crop-winter wheat intercrops on the semi-arid Canadian prairies. *Canadian Journal of Plant Science*, 90(4): 479–488.
- Bockheim, J. G. & Hartemink, A. E.** 2017. Soil-forming processes. In *The Soils of Wisconsin*, pp. 55–65.
- Boroday, I. I.** 2019. The main factors of soil degradation in Ukraine (In Ukrainian). Proceedings of the International Scientific and Practical Conference Youth and Technological Progress in Agriculture. *Innovative developments in the agricultural sphere*, 2: 228–229.
- Borodina, O., Kyryzyuk, S., Yarovy, V., Ermoliev, Y. & Ermolieva, T.** 2016. Modeling local land uses under the global change (In Ukrainian). *Economics and Forecasting*, 1: 117–128. <https://doi.org/https://doi.org/10.15407/eip2016.01.117>.
- Bossio, D.A., Cook-Patton, S.C., Ellis, P.W., Fargione, J., Sanderman, J., Smith, P., Wood, S., Zomer, R. J., von Unger, M., Emmer I. M. & Griscom, B.W.** 2020. The role of soil carbon in natural climate solutions. *Nature Sustainability*, 3(5): 391–398. <https://doi.org/10.1038/s41893-020-0491-z>
- Bradshaw, B., Dolan, H. & Smit, B.** 2004. Farm-level adaptation to climatic variability and change: crop diversification in the Canadian prairies. *Climatic change*, 67(1): 119–141.
- Breiman, L.** 2001. Random forests. *Machine learning*, 45(1): 5–32.
- Brevik, E.C. & Sauer, T.J.** 2015. The past, present, and future of soils and human health studies. *SOIL*, 1(1): 35–46. <https://doi.org/10.5194/soil-1-35-2015>
- Britannica.** 2022. Dust Bowl. In: *Encyclopedia Britannica*. Cited 6 June 2022. <https://www.britannica.com/place/Dust-Bowl>. Accessed 12 October 2022.
- Brooks, J. R., Flanagan, L. B. Buchmann, N. & Ehleringer, J. R.** 1997. Carbon Isotope composition of boreal plants: functional grouping of life forms. *Oecologia*, 110: 301–311.
- Bruulsema, T.W., Peterson, H.M. & Prochnow, L.I.** 2019. The science of 4R nutrient stewardship for phosphorus management across latitudes. *Journal of Environmental Quality*, 48(5): 1295–1299.
- Bui, E.N. & Moran, C.J.** 2001. Disaggregation of polygons of surficial geology and soil maps using spatial modelling and legacy data. *Geoderma*, 103(1–2): 79–94.
- Buytaert, W., Iñiguez, V., Celleri, R., De Bièvre, B., Wyseure, G. & Deckers, J.** 2006a. Analysis of the water balance of small paramo catchments in south Ecuador. In *Environmental Role of Wetlands in Headwaters*; Springer: Dordrecht The Netherlands, 271–281.
- Buytaert, W., Célleri, R., De Bièvre, B., Cisneros, F., Wyseure, G., Deckers, J. & Hofstede, R.** 2006b. Human impact on the hydrology of the Andean páramos. *Earth-Science Reviews*, 79: 53–72.
- Cabrera, A.L. & Willink, A.** 1980. *Biogeografía da America Latina*. Second ed. OEA, Washington.
- Cai, H.G., Mi, G.H. & Zhang, X.Z.** 2012. Effect of different fertilizing methods on nitrogen balance in the black soil for continuous maize production in northeast China. *Journal of Maize Sciences*. 18(01): 89–97. (In Chinese)
- Campbell, C.A., Biederbeck, V.O., Selles, F., Schnitzer, M. & Stewart, J.W.B.** 1986. Effect of manure and P fertilizer on properties of a Black Chernozem in southern Saskatchewan. *Canadian journal of soil science*, 66(4): 601–614.
- Campbell, C.A., Biederbeck, V.O., Zentner, R.P. & Lafond, G.P.** 1991. Effect of crop rotations and cultural practices on soil organic matter, microbial biomass and respiration in a thin black Chernozem. *Canadian Journal of Soil Science*, 71: 363–376.
- Campbell, C.A., Selles, F., Lafond, G.P., Biederbeck, V.O. & Zentner, R.P.** 2001. Tillage – fertilizer changes: Effect on some soil quality attributes under long-term crop rotations in a thin Black Chernozem. *Canadian Journal of Soil Science*, 81(2):157–165.
- Campos, M.C.C., Alho, L.C., Silva, D.A.P., Silva, M.D.R., Cunha, J.M. & Silva, D.M.P.** 2016. Distribuição espacial do efluxo de CO<sub>2</sub> em área de terra preta arqueológica sob cultivo de cacau e café no município de Apuí, AM, Brasil. *Revista Ambiente & Água*, 11(4): 788–798.



- Campos, M.C.C., Ribeiro, M.R., Souza Júnior, V.S., Ribeiro Filho, M.R., Souza, R.V.C.C. & Almeida, M.C.** 2011. Caracterização e classificação de terras pretas arqueológicas na Região do Médio Rio Madeira. *Bragantia*, 70(3): 598–609.
- Cárdenas, C. de los A.** 2013. El fuego y el pastoreo en el páramo húmedo de Chingaza (Colombia): efectos de la perturbación y respuestas de la vegetación. Universitat Autònoma de Barcelona. PhD dissertation.
- Castañeda-Martín, A.E. & Montes-Pulido, C.R.** 2017. Carbono almacenado en páramo andino. *Entramado*, 13 (1): 210–221. <http://dx.doi.org/10.18041/entramado.2017v13n1.25112>.
- Cattani, D.J.** 2019. Potential of perennial cereal rye for perennial grain production in Manitoba. *Canadian Journal of Plant Science*, 99(6): 958–960.
- Chaney, N.W., Wood, E.F., McBratney, A.B., Hempel, J.W., Nauman, T.W., Brungard, C.W. & Odgers, N.P.** 2016. POLARIS: A 30-meter probabilistic soil series map of the contiguous United States. *Geoderma*, 274: 54–67.
- Chang, Q., Wang, L., Ding, S., Xu, T., Li, Z., Song, X., Zhao, X., Wang, D. & Pan, D.** 2018. Grazer effects on soil carbon storage vary by herbivore assemblage in a semi-arid grassland. *Journal of Applied Ecology*, 55(5): 2517–2526.
- Chantigny, M.H., Angers, D.A., Prévost, D., Vézina, L.-P. & Chalifour, F.-P.** 1997. Soil aggregation and fungal and bacterial biomass under annual and perennial cropping systems. *Soil Science Society of America Journal*, 61(1): 262–267. <https://doi.org/10.2136/sssaj1997.03615995006100010037x>
- Chaturika, J.S., Kumaragamage, D., Zvomuya, F., Akinremi, O.O., Flaten, D.N., Indraratne, S.P. & Dandeniyi, W.S.** 2016. Woodchip biochar with or without synthetic fertilizers affects soil properties and available phosphorus in two alkaline, chernozemic soils. *Canadian journal of soil science*, 96(4): 472–484.
- Chen, F. H.** 2012. Foundations on expansive soils (Vol. 12). Elsevier.
- Chen, Y., Zhang, X., He, H., Xie, H., Yan, Y., Zhu, P., Ren, J. & Wang, L.** 2010. Carbon and nitrogen pools in different aggregates of a Chinese Mollisol as influenced by long-term fertilization. *Journal of Soils and Sediments*, 10(6): 1018–1026.
- Choudhary, O.P. & Kharche, V.K.** 2018. Soil salinity and sodicity. *Soil science: an introduction*, 12: 353–384.
- Cicek, H., Entz, M.H., Martens, J.R.T. & Bullock, P.R.** 2014. Productivity and nitrogen benefits of late-season legume cover crops in organic wheat production. *Canadian Journal of Plant Science*, 94(4): 771–783.
- CIESIN.** 2018. Gridded Population of the World, Version 4 (GPWv4): Population Density, Revision 11. Center for International Earth Science Information Network. Cited 31 March 2022. <https://doi.org/10.7927/H49C6VHW>. Accessed 31st March 2022.
- Ciolacu, T.** 2017. Current state of humus in arable chernozems of Moldova. *Scientific Papers-Series A, Agronomy*, 60: 57–60.
- Clément, C.C., Cambouris, A.N., Ziadi, N., Zebarth, B.J. & Karam, A.** 2020. Nitrogen source and rate effects on residual soil nitrate and overwinter no<sub>3</sub>-n losses for irrigated potatoes on sandy soils. *Canadian Journal of Soil Science*, 100(1): 44–57.
- Cohen, J.C.P., Beltrão, J.C., Gandu, A.W. & Silva, R.R.** 2007. Influência do desmatamento sobre o ciclo hidrológico na Amazônia. *Ciência e Cultura*, 59(3): 36–39.
- Collantes, M.B. & Faggi, A.M.** 1999. Los humedales del sur de Sudamérica. In: A.I. Malvárez, ed. *Tópicos sobre humedales subtropicales y templados de Sudamérica*, pp. 15–25. Montevideo, Uruguay, UNESCO.
- Conceição, P.C., Bayer, C., Castilhos, Z.M.S., Mielniczuk, J. & Guterres, D.B.** 2007. Estoques de carbono orgânico num Chernossolo Argilúvico manejado sob diferentes ofertas de forragem no Bioma Pampa Sul-Riograndense. In *Anais do 3Ind Congresso Brasileiro de Ciência do Solo*. Gramado, Rio Grande do Sul.
- Cordeiro, F.R.** 2020. Funções de Pedotransferência para Padronização de Base de Dados, Critérios de Classificação Taxonômica e Susceptibilidade Magnética em Terra Preta de Índio. Department of Soil. Universidade Federal Rural do Rio de Janeiro. Master dissertation.
- Corporación Nacional Forestal (CONAF).** 2006. Catastro de uso del suelo y vegetación, región de Magallanes y Antártica Chilena. *Monitoreo y actualización 2006*. Santiago de Chile.
- Cuervo-Barahona, E.L., Cely-Reyes, G.E. & Moreno-Pérez, D.F.** 2016. Determinación de las fracciones de carbono orgánico en el suelo del páramo La Cortadera, Boyacá. *Ingenio Magno*, 7(2): 139–149.

- Cui, W.L., Wang, J.J., Zhu, J. & Kong, F.Z. 2017. "Lishu black land culture" continues to heat up. *Jilin Daily*. [http://jjuban.moa.gov.cn/fwllm/qgxxlb/qg/201709/t20170914\\_5815758.htm](http://jjuban.moa.gov.cn/fwllm/qgxxlb/qg/201709/t20170914_5815758.htm)
- Cumba, A., Imbellone, P. & Ligier, A. 2005. Propiedades morfológicas, físicas, químicas y mineralógicas de suelos del sur de Corrientes. *Revista de la Asociación Geológica Argentina*, 60 (3): 579–590.
- Cunha, J.M., Campos, M.C.C., Gaio, D.C., Souza, Z.M., Soares, M.D.R., Silva, D.M.P. & Simões, E.L. 2018. Spatial variability of soil respiration in Archaeological Dark Earth areas in the Amazon. *Catena*, 162(5): 148–156.
- Cunha, J.M., Gaio, D.C., Campos, M.C.C. Soares, M.D.R., Silva, D.M.P. & Lima, A.F.L. 2017. Atributos físicos e estoque de carbono do solo em áreas de Terra Preta Arqueológica da Amazônia. *Revista Ambiente & Água*, 12 (3): 263–281.
- Cunha, L., Brown, G.G., Stanton, D.W.G., Da Silva, E., Hansel, F.A., Jorge, G., McKey, D., Vidal-Torrado, P., Macedo, R., Velasquez, E., James, S., Samuel, W. & Lavelle, P.K. 2016. Soil animals and pedogenesis: the role of earthworms in anthropogenic soils. *Soil Science*, 181(3–4): 110–125. <https://doi.org/10.1097/SS.000000000000144>
- Degens, B.P. 1997. Macro-aggregation of soils by biological bonding and binding mechanisms and the factors affecting these: a review. *Australian Journal of Soil Research*, 35: 431–459. <https://doi.org/10.1071/S96016>
- Demattê, J.L.I., Vidal-Torrado, P. & Sparovek, G. 1992. Influência da drenagem na morfogênese de solos desenvolvidos de rochas básicas no município de Piracicaba (SP). *Rev. Bras. Ci Solo*, 16: 241–247.
- Demetrio, W.C., Conrado, A.C., Acioli, A.N.S., Ferreira, A.C., Bartz, M.L.C., James, S.W., da Silva, E., Maia, Lilliane S., Martins, Gilvan C., Macedo, Rodrigo S., Stanton, David W. G., Lavelle, P., Velasquez, E., Zangerlé, A., Barbosa, R., Tapia-Coral, S.C., Muniz, A.W., Santos, A., Ferreira, T., Segalla, R., Decaëns, T., Nadolny, H.S., Peña-Venegas, C.P., Maia, C.M.B.F., Pasini, A., Mota, A.F., Taube Júnior, P.S., Silva, T.A.C., Rebellato, L., de Oliveira Júnior, R.C., Neves, E.G., Lima, H.P., Feitosa, R.M., Torrado, P.V., McKey, D., Clement, C.R., Shock, M.P., Teixeira, W.G., Motta, A.C.V., Melo, V.F., Dieckow, J., Garrastazu, M.C., Chubatsu, L.S., Kille, P., TPI Network, Brown, G.G. & Cunha, L. 2021. A "Dirty" Footprint: Macroinvertebrate diversity in Amazonian anthropic soils. *Global Change Biology*, 27(19): 4575–4591. <https://doi.org/10.1111/gcb.15752>
- Deng, F., Wang, H., Xie, H., Bao, X., He, H., Zhang, X. & Liang, C. 2021. Low-disturbance Farming Regenerates Healthy Deep Soil towards Sustainable Agriculture. *bioRxiv*: 828673.
- Derpsch, R. 2003. Conservation tillage, no-tillage and related technologies. In *Conservation agriculture*, pp. 181–190. Springer, Dordrecht.
- Derpsch, R., Friedrich, T., Kassam, A. & Hongwen, L. 2010. Current status of adoption of no-till farming in the world and some of its main benefits. *International Journal of Agricultural and Biological Engineering*, 3: 1–25.
- Díaz Barradas, M.C., García Novo, F., Collantes, M.B & Zunzunegui, M. 2001. Vertical structure of a wet grassland under and non-grazed conditions in Tierra del Fuego. *J Veg Sci*, 12: 385–390.
- Díaz-Zorita, M., Duarte, G. A. & Grove, J. H. 2002. A review of no-till systems and soil management for sustainable crop production in the subhumid and semiarid Pampas of Argentina. *Soil and Tillage Research*, 65(1), 1–18.
- Dick, C., Cattani, D. & Entz, M.H. 2018. Kernza intermediate wheatgrass (*Thinopyrum intermedium*) grain production as influenced by legume intercropping and residue management. *Canadian Journal of Plant Science*, 98(6):1376–1379.
- Dick, W. A. & Gregorich, E. G. 2004. Developing and maintaining soil organic matter levels. *Managing soil quality: Challenges in modern agriculture*, 103: 120.
- Ding, J., Jiang, X., Ma, M., Zhou, B., Guan, D., Zhao, B., Zhou, J., Cao, F., Li, L. & Li, J. 2016. Effect of 35 years inorganic fertilizer and manure amendment on structure of bacterial and archaeal communities in black soil of northeast China. *Applied soil ecology*, 105: 187–195.
- Ding, X., Han, X., Liang, Y., Qiao, Y., Li, L. & Li, N. 2012. Changes in soil organic carbon pools after 10 years of continuous manuring combined with chemical fertilizer in a Mollisol in China. *Soil and Tillage Research*, 122: 36–41.
- Ding, X., Zhang, B., Zhang, X., Yang, X. & Zhang, X. 2011. Effects of tillage and crop rotation on soil microbial residues in a rainfed agroecosystem of northeast China. *Soil and Tillage Research*, 114(1): 43–49.
- Dmytruk, Y. 2021. Report on multiple cross-sectoral LDN monitoring benefits developed (In Ukrainian). GCP/UKR/004/GEF (unpublished).

- Dodds, W. K., Blair, J. M., Hnebry, G. M., Koelliker, J. K., Ramundo, R. & Tate, C. M. 1996. Nitrogen transport from tallgrass Prairie Watersheds. *Environmental Quality J.*, 25: 973- 981. On line. 1537-2537
- Dodds, W.K. & Smith, V.H. 2016. Nitrogen, phosphorus, and eutrophication in streams. *Inland Waters*, 6(2): 155-164. <https://doi.org/10.5268/IW-6.2.909>
- Domzał, H., Gliński, J. & Lipiec, J. 1991. Soil compaction research in Poland. *Soil and Tillage Research*, 19(2-3): 99-109.
- Dörner, J., Dec, D., Thiers, O., Paulino, L., Zúñiga, F., Valle, S., Martínez, O. & Horn, R. 2016. Spatial and temporal variability of physical properties of Aquands under different land uses in southern Chile. *Soil Use and Management* 32, 411-421. <https://doi.org/10.1111/sum.12286>.
- Dumont, B., Carrère, P., Ginane, C., Farruggia, A., Lanore, L., Tardif, A., Decuq, F., Darsonville, O. & Louault, F. 2011. Plant-herbivore interactions affect the initial direction of community changes in an ecosystem manipulation experiment. *Basic and Applied Ecology*, 12(3):187-194.
- Durán, A, Morrás, H., Studdert, G. & Liu, X. 2011. Distribution, properties, land use and management of Mollisols in South America. *Chinese Geographical Science*, 21 (5): 511-530.
- Duran, A. 2010. *An overview of South American Mollisols: Soil formation, classification, suitability and environmental challenges*. In: *Proceedings of the International Symposium on Soil Quality and Management of World Mollisols*. northeast Forestry University Press, Harbin.
- Durand, R. & Dutil, P. 1971. Soil evolution in a calcic and magnesian clay material in the Der country, Haute-Marne. *Sci Sol*, 1: 65-78.
- Dusén, P. 1903. Die Pflanzenvereine der Magellansländern nebst einem Beitrage zur Ökologie der Magellanischen Vegetation. *Svenska Exped Magellansländerna*, 3: 351-521.
- Duulatov, E., Pham, Q. B., Alamanov, S., Orozbaev, R., Issanova, G. & Asankulov, T. 2021. Assessing the potential of soil erosion in Kyrgyzstan based on RUSLE, integrated with remote sensing. *Environmental Earth Sciences*, 80(18): 1-13.
- Dybdal, S.E. 2019. Sinograin II project: Tomorrow's development collaboration. In: Nibio. Cited 3 June 2022. <https://www.nibio.no/nyheter/sinograin-ii-project-tomorrows-development-collaboration>
- Dybdal, S.E. 2020. Black soil – China's giant panda in cultivated land – Nibio. In: Nibio. Cited 3 June 2022. <https://www.nibio.no/en/news/black-soil--chinas-giant-panda-in-cultivated-land>
- Eckmeier, E., Gerlach, R. Gehrt, E. & Schmidt, M. W. I 2007. Pedogenesis of Chernozems in Central Europe a review. *Geoderma*, 288-299.
- Eldridge, D.J., Delgado-Baquerizo, M., Travers, S.K., Val, J. & Oliver, I. 2017. Do grazing intensity and herbivore type affect soil health? Insights from a semi-arid productivity gradient. *Journal of Applied Ecology*, 54: 976-985.
- Engel, R. E., Romero, C. M., Carr, P. & Torrión, J. A. 2019. Performance of nitrate compared with urea fertilizer in a semiarid climate of the northern great plains. *Canadian Journal of Soil Science*, 99(3): 345-355.
- Entz, M.H., Baron, V.S., Carr, P.M., Meyer, D.W., Smith Jr, S.R. & McCaughey, W.P. 2002. Potential of forages to diversify cropping systems in the northern Great Plains. *Agronomy Journal*, 94(2): 240-250.
- Erickson, C.L. 2008. Amazonia: the historical ecology of a domesticated landscape. In: H. Silverman & W. Isbell, eds. *Handbook of South American Archaeology*, pp. 157-183. Springer.
- Eswaran, H., Almaraz, R., van den Berg, E. & Reich, P. 1997. An assessment of the soil resources of Africa in relation to productivity. *Geoderma*, 77: 1-18.
- Evans, P. & Halliwell, B. 2001. Micronutrients: oxidant/antioxidant status. *British journal of nutrition*, 85(S2): S67-S74.
- Fan, R., Liang, A., Yang, X., Zhang, X., Shen, Y. & Shi, X. 2010. Effects of tillage on soil aggregates in black soils in northeast China. *Scientia Agricultura Sinica*, 43(18): 3767-3775.
- Fan, R., Zhang, X., Liang, A., Shi, X., Chen, X., Bao, K., Yang, X. & Jia, S. 2012. Tillage and rotation effects on crop yield and profitability on a Black soil in northeast China. *Canadian Journal of Soil Science*, 92(3): 463-470.
- FAO. 2020. Environment Statistics. Mineral and Chemical Fertilizers: 1961-2018 [online]. [Cited 12 March 2021]. <http://www.fao.org/economic/ess/environment/data/mineral-and-chemical-fertilizers/en/>
- Fan, Y., Miguez-Macho, G., Jobbágy, E.G., Jackson, R.B. & Otero-Casal, C. 2017. Hydrologic regulation of plant rooting depth. *Proceedings of the National Academy of Sciences*, 114(40): 10572-10577.



- FAO & ITPS. 2015. Status of the World's Soil Resources (SWSR) – Main report. *Food and Agriculture Organization of the United Nations and Intergovernmental Technical Panel on Soils*, Rome, Italy. <https://www.fao.org/documents/card/en/c/6814873-efc3-41db-b7d3-2081a10ede50/>
- FAO & ITPS. 2021. *Recarbonizing global soils – A technical manual of recommended management practices*. Rome, FAO. <https://doi.org/10.4060/cb6386en>
- FAO & UNEP. 2021. *Global assessment of soil pollution – Summary for policy makers*. Rome, FAO. <https://doi.org/10.4060/cb4827en>
- FAO, ISRIC & JRC. 2012. *Harmonized world soil database*. Harmonised World Soil Database (version 1.2).
- FAO. 2002. *Captura de carbono en los suelos para un mejor manejo de la tierra*. Informes sobre recursos mundiales de suelos. Roma.
- FAO. 2015. *Healthy soils are the basis for healthy food production*. Rome, Italy, FAO. <https://www.fao.org/documents/card/en/c/645883cd-ba28-4b16-a7b8-34babbb3c505/>
- FAO. 2017. Global Soil Organic Carbon Map. In: *FAO Land and Water Division*. Rome. Cited 5 December 2017. <https://www.fao.org/world-soil-day/about-wsd/wsd-2017/global-soil-organic-carbon-map/en/>
- FAO. 2019. *Black Soils definition*. Cited 20 October 2020. <http://www.fao.org/global-soil-partnership/intergovernmental-technical-panel-soils/gsoc17-implementation/internationalnetworkblacksoils/more-on-black-soils/definition-what-is-a-black-soil/en/>
- FAO. 2020. *Soil testing methods manual – Soil Doctors Global Programme – A farmer-to-farmer training programme*. Rome. <https://doi.org/10.4060/ca2796en>
- FAO. 2022a. *Global Map of Black Soils*. Rome, Italy, FAO. <https://www.fao.org/documents/card/en/c/cc0236en>
- FAO. 2022b. Global Soil Laboratory Network (GLOSOLAN). In: *Food and Agriculture Organization of the United Nations*. Cited 12 October 2022. <https://www.fao.org/global-soil-partnership/glosolan/en/>
- FAO. 2022c. *Global Soil Organic Carbon Sequestration Potential Map – SOCseq v.1.1*. Technical report. Rome. <https://doi.org/10.4060/cb9002en>
- FAO. 2022d. *Global Soil Organic Carbon Map – GSOCmap v.1.6: Technical report*. Rome, FAO. <https://books.google.com.mx/books?id=ML1qEAAAQBAJ>
- FAO-UNESCO. 1981. *Soil map of the world 1:5 000 000*. FAO, Rome.
- Farkas, C., Hagyló, A., Horváth, E. & Várallyay, G. 2008. A Chernozem soil water regime response to predicted climate change scenarios. *Soil and Water Research*, 3 (Special Issue 1).
- Farsang, A., Babcsányi, I., Ladányi, Z., Perei, K., Bodor, A., Csányi, K.T. & Barta, K. 2020. Evaluating the effects of sewage sludge compost applications on the microbial activity, the nutrient and heavy metal content of a Chernozem soil in a field survey. *Arabian Journal of Geosciences*, 13(19): 1–9.
- Fey, M.V. 2010. *Soils of South Africa*. Cambridge, Cambridge University Press.
- Fileccia, T., Guadagni, M., Hovhera, V. & Bernoux, M. 2014. *Ukraine: Soil Fertility to Strengthen Climate Resilience*. Washington, DC, World Bank and FAO.
- Filipová L. 2011. Soil and vegetation of meadow wetlands (Vegas) in the South of the Chilean Patagonia. Faculty of Science Department of Botany, University Olomouc. PhD dissertation.
- Filipová L., Hédl R. & Covacevich N. 2010. Variability of the soil types in meadow wetlands in the south of the Chilean Patagonia. *Chilean Journal of Agricultural Research*, 70(2): 266–277.
- Findmypast. 2015. 1939: The year the dust settled. In: *Findmypast – Genealogy, Ancestry, History blog from Findmypast*. Cited 6 June 2022. <https://www.findmypast.com/blog/history/1939-the-year-the-dust-bowl-settled>
- Fischer, R.A. & Connor, D.J. 2018. Issues for cropping and agricultural science in the next 20 years. *Field Crops Research*, 222: 121–142. <https://doi.org/10.1016/j.fcr.2018.03.008>
- Focht, T. & Medeiros R.B. 2012. Prevention of natural grassland invasion by *Eragrostis plana* Nees using ecological management practices. *Revista Brasileira de Zootecnia*, 41: 1816–1823.
- Follett, R. F. 2001. Soil management concepts and carbon sequestration in cropland soils. *Soil and tillage research*, 61(1–2): 77–92.
- Foster, P. 2001. The potential negative impacts of global climate change on tropical montane cloud forests. *Earth-Science Reviews*, 55: 73–106.

- Freitas, P.L. de & Landers, J.N.** 2014. The transformation of agriculture in Brazil through development and adoption of Zero Tillage Conservation Agriculture. *International Soil and Water Conservation Research*, 2: 35–46.
- Fujii H., Mori S., & Matsumoto Y.** 2021. Tohoku region. In: R. Hatano, H. Shinjo & Y. Takata, eds. *The Soil of Japan*, pp. 69–134, Springer.
- Fujino A. & Matsumoto E.** 1992. *Topsoil erosion on the cropland in the Sugadaira Basin, Central Japan* (In Japanese). Bulletin of Environmental Research Center, the University of Tsukuba, 16: 69–77.
- Fujita T., Okuda T. & Fujie K.** 2007. Influence of colian dust brought from northern Asia continent on the parent materials in a fine-textured soil developed on the bedrock of the tertiary rock near cape Saruyama, in Noto peninsula, central Japan (In Japanese with English summary). *Pedologist*, 51: 97–103.
- Galán, S.** 2003. Manejo y Enriquecimiento del Bosque a Partir del Uso de las Chagras y Rastrojos de un Núcleo Familiar Indígena en Araracuara, Medio Río Caquetá (Amazonia colombiana). *Departament of Ecology. Pontificia Universidad Javeriana, Bogotá*:37.
- Gao, M., Guo, Y., Liu, J., Liu, J., Adl, S., Wu, D. & Lu, T.** 2021. Contrasting beta diversity of spiders, carabids, and ants at local and regional scales in a black soil region, northeast China. *Soil Ecology Letters*, 3(2): 103–114. <https://doi.org/10.1007/s42832-020-0071-1>
- Gao, X., Asgedom, H., Tenuta, M. & Flaten, D. N.** 2015. Enhanced efficiency urea sources and placement effects on nitrous oxide emissions. *Agronomy Journal*, 107(1): 265–277.
- Gao, X., Shaw, W. S., Tenuta, M. & Gibson, D.** 2018. Yield and Nitrogen Use of Irrigated Processing Potato in Response to Placement, Timing and Source of Nitrogen Fertilizer in Manitoba. *American Journal of Potato Research*, 95(5): 513–525.
- Garcia-Franco, N., Hobbey, E., Hübner, R. & Wiesmeier, M.** 2018. Climate-smart soil management in semiarid regions. *Soil management and climate change*, pp. 349–368. Elsevier.
- Geng, X., VandenBygaart, A.J. & He, J.** 2021. Soil organic carbon sequestration potential assessment using Roth-C model from the agriculture land of Canada. FAO.
- German, L.A.** 2003. Historical contingencies in the coevolution of environment and livelihood: contributions to the debate on Amazonian Black Earth. *Geoderma*, 111(3): 307–331.
- Giani, L., Makowsky, L. & Mueller, K.** 2014. Plaggic Anthrosol: Soil of the Year 2013 in Germany: An overview on its formation, distribution, classification, soil function and threats. *Journal of Plant Nutrition and Soil Science*, 177(3): 320–329.
- Glaser, B. & Birk, J.J.** 2012. State of the scientific knowledge on properties and genesis of Anthropogenic Dark Earths in Central Amazonia (terra preta de Índio). *Geochimica et Cosmochimica acta*, 82: 39–51.
- Glaser, B., Haumaier, L., Guggenberger, G. & Zech, W.** 2001. The ‘Terra Preta’ Phenomenon: A Model for Sustainable Agriculture in the Humid Tropics. *Naturwissenschaften*, 88: 37–41.
- Glauber, J., Laborde, D. & Mamun, A.** 2022. From bad to worse: How Russia-Ukraine war-related export restrictions exacerbate global food insecurity. In: *International Food Policy Research Institute (IFPRI)*. Cited 1 June 2022. <https://www.ifpri.org/blog/bad-worse-how-export-restrictions-exacerbate-global-food-security>
- Gollany, H.T., Rickman, R.W., Liang, Y., Albrecht, S.L., Machado, S. & Kang, S.** 2011. Predicting agricultural management influence on long-term soil organic carbon dynamics: Implications for biofuel production. *Agronomy Journal*, 103(1): 234–246. <https://doi.org/10.2134/agronj2010.0203s>
- Gong, H., Meng, D., Li, X. & Zhu, F.** 2013. Soil degradation and food security coupled with global climate change in northeastern China. *Chinese Geographical Science*, 23(5): 562–573.
- Gregg, J.S. & Izaurralde, R.C.** 2010. Effect of crop residue harvest on long-term crop yield, soil erosion and nutrient balance: trade-offs for a sustainable bioenergy feedstock. *Biofuels*, 1(1): 69–83.
- Gregorich, E. G. & Anderson, D. W.** 1985. Effects of cultivation and erosion on soils of four toposequences in the Canadian prairies. *Geoderma*, 36: 343–354.
- Grekov, Datsko, L.V., Zhilkin, V.A., Maistrenko, M.I. & Datsko, M.O.** 2011. *Methodical instructions for soil protection* (In Ukrainian). Kyiv, The State Center of Soil Fertility Protection. 108 pp.
- Guilpart, N., Grassini, P., Sadras, V.O., Timsina, J. & Cassman, K.G.** 2017. Estimating yield gaps at the cropping system level. *Field Crops Research*, 206: 21–32. <https://doi.org/10.1016/j.fcr.2017.02.008>
- Guo, Y., Amundson, R., Gong, P. & Yu, Q.** 2006. Quantity and Spatial Variability of Soil Carbon in the Conterminous United States. *Soil Sci. Soc. Am. J.* 70: 590–600.

- Guo, Y., Luo, L., Chen, G., Kou, Y. & Xu, H.** 2013. Mitigating nitrous oxide emissions from a maize-cropping black soil in northeast China by a combination of reducing chemical N fertilizer application and applying manure in autumn. *Soil Science and Plant Nutrition*, 59(3): 392–402.
- Gupta, S.C. & Allmaras, R.R.** 1987. Models to assess the susceptibility of soils to excessive compaction. In: B.A. Stewart, ed. *Advances in Soil Science*. pp. 65–100. New York, NY, Springer New York.
- Halde, C., Bamford, K.C. & Entz, M.H.** 2015. Crop agronomic performance under a six-year continuous organic no-till system and other tilled and conventionally-managed systems in the northern Great Plains of Canada. *Agriculture, Ecosystems & Environment*, 213: 121–130.
- Han, J., Mao, K., Xu, T., Guo, J., Zuo, Z. & Gao, C.** 2018. A Soil Moisture Estimation Framework Based on the CART Algorithm and Its Application in China. *Journal of Hydrology*, 563. <https://doi.org/10.1016/j.jhydrol.2018.05.051>
- Han, X., Wang, S., Veneman, P.L. & Xing, B.** 2006. Change of organic carbon content and its fractions in black soil under long-term application of chemical fertilizers and recycled organic manure. *Communications in Soil Science and Plant Analysis*, 37(7–8): 1127–1137.
- Han, Y., Chen, X., Wang, E. & Xia, X.** 2019. Optimum biochar preparations enhance phosphorus availability in amended Mollisols of northeast China. *Chilean journal of agricultural research*, 79(1): 153–164.
- Han, Z.M., Deng, M.W., Yuan, A.Q., Wang, J.H., Li, H. & Ma, J.C.** 2018. Vertical variation of a black soil's properties in response to freeze-thaw cycles and its links to shift of microbial community structure. *Sci. Total Environ*, 625: 106–113.
- Hansen, M. C., P. V. Potapov, R. Moore, M. Hancher, S. A. Turubanova, A. Tyukavina, D. Thau, S. V. Stehman, S. J. Goetz, T. R. Loveland, A. Kommareddy, A. Egorov, L. Chini, C. O. Justice, & Townshend, J. R. G.** 2013. "High-Resolution Global Maps of 21st-Century Forest Cover Change." *Science* 342 (15 November): 850–53. 10.1126/science.1244693. <https://glad.earthengine.app/view/global-forest-change>.
- Hao, X., Han, X., Wang, S. & Li, L.** 2022. Dynamics and composition of soil organic carbon in response to 15 years of straw return in a Mollisol. *Soil and Tillage Research*, 215: 105221.
- Hartemink, A.E., Krasilnikov, P. & Bockheim, J.G.** 2013. Soil maps of the world. *Geoderma*, 207: 256–267.
- Hayes, W.A.** 1985. Conservation Tillage Systems and Equipment Requirements. In F. D'Itri, ed. *A Systems Approach to Conservation Tillage*. Boca Raton, the USA, CRC Press.
- Haynes, R. J. & Naidu, R.** 1998. Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions: a review. *Nutrient cycling in agroecosystems*, 51(2): 123–137.
- Herrero-Jáuregui, C. & Oesterheld, M.** 2018. Effects of grazing intensity on plant richness and diversity: A meta-analysis. *Oikos*, 127(6), 757–766.
- Hincapié, J.C.A., Castillo, C.B., Argüello, S.C., Aguilera, D.P.R., Holguín, F.S., Triana, J.V. & Lopera, A.** 2002. Transformación y cambio en el uso del suelo en los páramos de Colombia en las últimas décadas. In: C, Castaño, ed. *Páramos y ecosistemas alto andinos de Colombia en condición hotspot y global climatic tensor*, pp. 211–333. Bogotá, IDEAM.
- History.** 2020. Dust Bowl. In: *HISTORY*. Cited 6 June 2022. <https://www.history.com/topics/great-depression/dust-bowl>
- Hofstede, R.G. & Rossenaar, A.J.** 1995. Biomass of grazed, burned, and undisturbed Paramo Grasslands, Colombia. II. Root mass and aboveground: Belowground ratio. *Arct. Alp. Res*, 27: 13–18.
- Hofstede, R.G.** 1995. The effects of grazing and burning on soil and plant nutrient concentrations in Colombian paramo grasslands. *Plant Soil*, 173: 111–132.
- Hofstede, R.G.** 2001. El Impacto de las actividades humanas sobre el Páramo. In: *Los Páramos del Ecuador, particularidades, problemas y perspectivas*, pp. 161–182. Quito, Ecuador, Editorial Abya-Yala.
- Holland, J.M.** 2004. The environmental consequences of adopting conservation tillage in Europe: reviewing the evidence. *Agriculture, ecosystems & environment*, 103(1): 1–25.
- Holmes, K.W., Griffin, E.A. & Odgers, N.P.** 2015. Large-area spatial disaggregation of a mosaic of conventional soil maps: evaluation over Western Australia. *Soil Research*, 53(8): 865–880.
- Horn, S.P. & Kappelle, M.** 2009. Fire in the paramo ecosystems of Central and South America. In *Tropical Fire Ecology*, pp. 505–539. Heidelberg, Berlin, Germany, Springer.



- Hospodarenko, H., Trus, O. & Prokopchuk, I.** 2012. Humus Conservation Conditions in a Field Crop Rotation. *Biological Syst.*, 4: 31–34.
- Hothorn, T.** 2022. CRAN Task View: Machine Learning & Statistical Learning. Cited 7 March 2022. <https://CRAN.R-project.org/view=MachineLearning>
- Hou, D.** 2022. China: protect black soil for biodiversity. *Nature*, 604(7904): 40–40. <https://doi.org/10.1038/d41586-022-00942-6>
- Haiwi, M.** 2001. Soils of the Syrian Arab Republic. Soil resources of Southern and Eastern Mediterranean countries. *CIHEAM, Bari*, 227–242.
- Imbellone, P. & Mormeneo, L.** 2011. Vertisoles hidromórficos de la planicie costera del Río de la Plata, Argentina. *Ciencia del Suelo*, 29: 107–127.
- Instituto Nacional de Estadística (INE).** 2014. *Estadística Pecuaria, período 2008–2013 y primer semestre 2014*. Santiago, Chile.
- Instituto Nacional De Estadística (INE).** 2007. *XII Censo Agropecuario y Forestal*. Santiago, Chile.
- IPCC.** 2019. Summary for Policymakers. In: Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems [P.R. Shukla, J. Skea, E. Calvo Buendia, V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)]. In press.
- Isbell, R. F.** 1991. Australian vertisols. *Characterization, classification and utilization of cold Aridisols and Vertisols. Proc. VI ISCOM, USDA-SCS*. National Soil Survey Center, Lincoln NB:73–80.
- Iturraspe, R. & Uriuolo.** 2000. Caracterización de las cuencas hídricas de Tierra del Fuego. Actas del XVIII Congreso Nacional del Agua. Junio de 2000, Termas de Río Hondo, Santiago del Estero.
- IUSS Working Group WRB.** 2006. World Reference Base. World reference base for soil resources. Available at: <https://www.fao.org/soils-portal/data-hub/soil-classification/world-reference-base/en/>
- IUSS Working Group WRB.** 2015. World Reference Base for Soil Resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps. World Soil Resources Reports No. 106. FAO, Rome.
- Iutynskaya, G. A. & Patyka V. F.** 2010. Soil biology: problems and perspectives (In Ukrainian). Agricultural chemistry and soil science. *Proceedings of Soil Science Council*. Vol. 1, Zhitomir, Ruta, 2008 pp.
- Ivelic-Sáez, J., Dörner, J., Arumí, J.L., Cisternas, L., Valenzuela, J., Muñoz, E., Clasing, R., Valle, S., Radic, S., Alonso, H., López, R., Uribe, H., Muñoz, R., Ordoñez, I. & Carrasco, J.** 2021. Balance hídrico de humedales de uso agropecuario: El primer paso para el mejoramiento en la gestión hídrica a nivel predial en Magallanes”. Una investigación multidisciplinaria. *Centro Regional de Investigación Kampenaike. Boletín INIA N°435*, pp. 162. Punta Arenas, Chile.
- Japanese Soil Conservation Research Project Nationwide Council.** 2012. National Farmland Soil Guidebook (In Japanese). Japan Soil Association, Tokyo, 121 p.
- Jat, M. L., Cathala, M. K., Ladha, J. K., Saharawat, Y. S., Jat, A. S., Kumar, V., Sharma, S. K., Kumar, V. & Gupta, R.** 2009. Evaluation of precision land leveling and double zero-till systems in the rice-wheat rotation: Water use, productivity, profitability and soil physical properties. *Soil and Tillage Research*, 105(1): 112–121.
- Jian, J., Du, X., Reiter, M.S. & Stewart, R.D.** 2020. A meta-analysis of global cropland soil carbon changes due to cover cropping. *Soil Biology and Biochemistry*, 143: 107735. <https://doi.org/10.1016/j.soilbio.2020.107735>
- Jin, L., Wei, D., Yin, D., Zhou, B., Ding, J., Wang, W., Zhang, J., Qiu S., Zhang C., Li, Y., An, Z., Gu, J. & Wang, L.** 2020. Investigations of the effect of the amount of biochar on soil porosity and aggregation and crop yields on fertilized black soil in northern China. *Plos one*, 15(11): e0238883.
- Johnson, W. G., Davis, V. M., Kruger, G. R. & Weller, S. C.** 2009. Influence of glyphosate-resistant cropping systems on weed species shifts and glyphosate-resistant weed populations. *European Journal of Agronomy*, 31(3): 162–172.
- Jones, A., Breuning-Madsen, H., Brossard, M., Dampha, A., Deckers, J., Dewitte, O., Gallali, T., Hallett, S., Jones, R., Kilasara, M., Le Roux, P., Micheli, E., Montanarella, L., Spaargaren, O., Thiombiano, L., Van Ranst, E., Yemefack, M. & Zougmoré, R.** 2013. Soil Atlas of Africa. *European Commission*, pp. 176. Publications Office of the European Union, Luxembourg.

- Ju, X., Liu, X., Zhang, F. & Roelcke, M.** 2004. Nitrogen fertilization, soil nitrate accumulation, and policy recommendations in several agricultural regions of China. *Ambio*, 33(6): 300–305. <https://doi.org/10.1579/0044-7447-33.6.300>
- Kahimba, F.C., Ranjan, R.S., Froese, J., Entz, M. & Nason, R.** 2008. Cover crop effects on infiltration, soil temperature, and soil moisture distribution in the Canadian Prairies. *Applied engineering in agriculture*, 24(3): 321–333.
- Kämpf, N., Woods, W., Sombroek, W., Kern, D. & T. Cunha, T.** 2003. Classification of Amazonian Dark Earths and other ancient anthropic soils. In J. Lehmann, D.K.B. Glaser D.K.B. & W. Woods, eds. *Amazonian Dark Earths: Origin, Properties, Management*, pp. 77–102. The Netherlands, Kluwer Academic Publishers.
- Kay, B.D.** 1990. Rates of change of soil structure under different cropping systems. In: Stewart, B.A, eds. *Advances in Soil Science*, vol 12. New York, NY, Springer. [https://doi.org/10.1007/978-1-4612-3316-9\\_1](https://doi.org/10.1007/978-1-4612-3316-9_1)
- Kazakova, I.** 2016. The impact of global changes at soil resources and agricultural production (In Ukrainian). *Agricultural and Resource Economics: International Scientific E-Journal*, 2(1): 21–44.
- Kern, D.C. & Kämpf, N.** 1989. Old Indian settlements on the formation of soils with archaeological black earth at Oriximina region (In Portuguese). *Para, Brazil. Revista Brasileira de Ciencia do Solo*, 13: 219–225.
- Kern, J., Giani, L., Teixeira, W., Lanza, G. & Glaser, B.** 2019. What can we learn from ancient fertile anthropic soil (Amazonian Dark Earths, shell mounds, Plaggen soil) for soil carbon sequestration? *Catena*, 172: 104–112. <https://doi.org/10.1016/j.catena.2018.08.008>
- Kobza, J. & Pálka, B.** 2017. Contribution to black soils in Slovakia according to INBS criteria. [In Slovak: Príspevok k tmavým pôdam na Slovensku podľa kritérií INBS]. *Proceedings of Soil Science and Conservation Research Institute*, 29: 34–42.
- Kogan, F., Adamenko, T. & Kulbida, M.** 2011. Satellite-based crop production monitoring in Ukraine and regional food security. In F. Kogan, A. Powell & O. Fedorov, eds. *Use of satellite and in-situ data to improve sustainability*. pp. 99–104. NATO Science for Peace and Security Series C: Environmental Security. Paper presented at, 2011, Dordrecht. [https://doi.org/10.1007/978-90-481-9618-0\\_11](https://doi.org/10.1007/978-90-481-9618-0_11)
- Kostić, M.M., Tagarakis, A.C., Ljubičić, N., Blagojević, D., Radulović, M., Ivošević, B. & Rakić, D.** 2021. The Effect of N Fertilizer Application Timing on Wheat Yield on Chernozem Soil. *Agronomy*, 11(7): 1413.
- Krasilnikov, P., Martí, J.-J. I., Arnold, R. & Shoba, S.** 2009. *A handbook of soil terminology, correlation and classification*. London, Sterling, UK, Earthscan. <https://doi.org/10.4324/9781849774352>
- Krasilnikov, P., Sorokin, A., Golozubov, O. & Bezuglova, O.** 2018. Managing chernozems for advancing sdgs. In R. Lal, R. Horn & T. Kosaki, eds. *Soil and Sustainable Development Goals*, pp. 175–188. GeoEcology Essays, Catena-Schweizerbart Stuttgart.
- Krupenikov, I.A.** 1992. *The soil layer of Moldova: past, present, management, forecast* [In Slovak: Moldova: Proshloe, nastoyashchec, upravlenie, prognoz].
- Kucher, A.** 2017. Adaptation of the agricultural land use to climate change (In Ukrainian). *Agricultural and Resource Economics: International Scientific E-Journal*, 3(1): 119–138.
- Lafond, G.P., Brandt S.A., Clayton G.W., Irvine R.B. & May W.E.** 2011a. Rainfed Farming Systems on the Canadian Prairies. In: Tow P., Cooper I., Partridge I., Birch C. (eds) *Rainfed Farming Systems*. Dordrecht, the UK, Springer.
- Lafond, G.P., Walley, F., May, W.E. & Holzapfel, C.B.** 2011b. Long term impact of no-till on soil properties and crop productivity on the Canadian prairies. *Soil and Tillage Research*, 117: 110–123.
- Lal, R.** 2014. Soil conservation and ecosystem services. *International Soil and Water Conservation Research*, 2(3): 36–47. [https://doi.org/10.1016/S2095-6339\(15\)30021-6](https://doi.org/10.1016/S2095-6339(15)30021-6)
- Lal, R.** 2019. Accelerated soil erosion as a source of atmospheric CO<sub>2</sub>. *Soil and Tillage Research*, 188: 35–40. <https://doi.org/10.1016/j.still.2018.02.001>
- Lal, R.** 2021. Managing Chernozem for Reducing Global Warming. In: D. Dent & B. Boincean, eds. *Regenerative Agriculture*. Cham, Springer International Publishing, 2021. [https://doi.org/10.1007/978-3-030-72224-1\\_7](https://doi.org/10.1007/978-3-030-72224-1_7)
- Lal, R., Monger, C., Nave, L. & Smith, P.** 2021. The role of soil in regulation of climate. *Phil. Trans. R. Soc. B*, 376: 20210084.
- Landi, A., Mermut, A. R., & Anderson, D. W.** 2003a. Origin and Rate of Pedogenic Carbonate Accumulation in Saskatchewan Soils, Canada. *Geoderma*. 117:143–156.

- Landi, A., Anderson D. W. & Mermut A. R.** 2003b. Organic carbon storage and stable isotope composition of soils along a grassland to forest environmental gradient in Saskatchewan. *Can. J. Soil Sci*, 83: 405–414.
- Landi, A., Mermut A. R. & Anderson, D. W.** 2004. Carbon Dynamics in a Hummocky Landscape from Saskatchewan. *SSSAJ*, 68: 175–184.
- Laos, F., Satti, P., Walter, I., Mazzarino, M.J. & Moyano, S.** 2000. Nutrient availability of composted and noncomposted residues in a Patagonian Xeric Mollisol. *Biology and Fertility of Soils*, 31(6): 462–469.
- Laufer, D., Loibl, B., Märlander, B. & Koch, H.-J.** 2016. Soil erosion and surface runoff under strip tillage for sugar beet (*Beta vulgaris* L.) in Central Europe. *Soil and Tillage Research*, 162: 1–7.
- Lavado, R.** 2016. Degradación de suelos argentinos. In F. Pereyra & M. Torres Duggan, eds. *Suelos y Geología Argentina. Una visión integradora desde diferentes campos disciplinarios*. AACs-AGA, UNDAV Ediciones, pp. 313–328.
- Lavado, R.S. & Taboada, M. A.** 2009. The Argentinean Pampas: A key region with a negative nutrient balance and soil degradation needs better nutrient management and conservation programs to sustain its future viability as a world agroresource. *Journal of Soil and Water Conservation*, 64(5), 150A–153A. <https://doi.org/10.2489/jswc.64.5.150A>
- Lawinfochina.** 2022. Black Soil Protection Law of the People’s Republic of China, Cited 24 June 2022. <https://www.lawinfochina.com/display.aspx?id=38784&lib=law>
- Leah, T. & Cerbari, V.** 2015. Cover crops–Key to storing organic matter and remediation of degraded properties of soils in Moldova. *Scientific Papers-Series A, Agronomy*, 58: 73–76.
- Lee, J. & Gill, T.** 2015. Multiple causes of wind erosion in the Dust Bowl. *Aeolian Research*, 19: 15–36. <https://doi.org/10.1016/j.aeolia.2015.09.002>
- Lehmann, J. & Joseph, S.** 2015. *Biochar for environmental management: science, technology and implementation*. Routledge.
- Lehmann, J.** 2009. Terra Preta Nova: where to from here?. In W. Woods *et al.*, eds. *Amazonian Dark Earths: Wim Sombroek’s vision*, pp. 473–486. Springer.
- Li, H., Yao, Y., Zhang, X., Zhu, H. & Wei, X.** 2021. Changes in soil physical and hydraulic properties following the conversion of forest to cropland in the black soil region of northeast China. *Catena*, 198: 104986.
- Li, H., Zhu, H., Qiu, L., Wei, X., Liu, B. & Shao, M.** 2020. Response of soil OC, N and P to land-use change and erosion in the black soil region of the northeast China. *Agriculture, Ecosystems & Environment*, 302: 107081.
- Li, N., Lei, W., Sheng, M., Long, J. & Han, Z.** 2022. Straw amendment and soil tillage alter soil organic carbon chemical composition and are associated with microbial community structure. *European Journal of Soil Biology*, 110: 103406.
- Li, P., Kong, D., Zhang, H., Xu, L., Li, C., Wu, M., Jiao, J., Li, D., Xu, L., Li, H. & Hu, F.** 2021. Different regulation of soil structure and resource chemistry under animal- and plant-derived organic fertilizers changed soil bacterial communities. *Applied Soil Ecology*, 165: 104020. <https://doi.org/10.1016/j.apsoil.2021.104020>
- Li, S., Liu, X. & Ding, W.** 2016. Estimation of organic nutrient sources and availability for land application. *Better Crops*, 100: 4–6.
- Li, S., Lobb, D.A. & Lindstrom, M.J.** 2007. Tillage translocation and tillage erosion in cereal-based production in Manitoba, Canada. *Soil and Tillage Research*, 94(1): 164–182.
- Li, S., Liu, X. & He, P.** 2017. Analyses on nutrient requirements in current agriculture production in China. *Journal of Plant Nutrition and Fertilizers*, 23: 1416–1432.
- Licht, M.A. & Al-Kaisi, M.** 2005. Strip-tillage effect on seedbed soil temperature and other soil physical properties. *Soil and Tillage Research*, 80: 233–249.
- Lima, H.N., Schaefer, C.E.R., Mello, J.W.V., Gilkes, R.J. & Ker, J.C.** 2002. Pedogenesis and Pre-Colombian Land Use of “Terra Preta Anthrosols” (“Indian black earth”) of Western Amazonia. *Geoderma*, 110: 1–17.
- Lins, J., Lima, H.P., Baccaro, F.B., Kinupp, V. F., Shepard Jr, G. H. & Clement, C.R.** 2015. Pre-Columbian floristic legacies in modern homegardens of Central Amazonia. *Plos one*, 10(6): 1–10.
- Liu, H., Wang, D., Wang, S., Meng, K., Han, X., Zhang, L. & Shen, S.** 2001. Changes of crop yields and soil fertility under long-term application of fertilizer and recycled nutrients in manure on a black soil. *Ying Yong Sheng tai xue bao= The Journal of Applied Ecology*, 12(1): 43–46.



- Liu, J., Yu, Z., Yao, Q., Hu, X., Zhang, W., Mi, G., Chen, X. & Wang, G. 2017. Distinct soil bacterial communities in response to the cropping system in a Mollisol of northeast China. *Applied soil ecology*, 119: 407–416.
- Liu, S., Fan, R., Yang, X., Zhang, Z., Zhang, X. & Liang, A. 2019. Decomposition of maize stover varies with maize type and stover management strategies: A microcosm study on a black soil (Mollisol) in northeast China. *J. Environ. Manage.* 234: 226–236.
- Liu, X., Burras, C., Kravchenko, Y., Durán, A., Huffman, T., Morras, H., Studdert, G., Zhang, X., Cruse, R. & Yuan, X. 2012. Overview of Mollisols in the world: distribution, land use and management. *Can. J. Soil. Sci.* 92: 383–402.
- Liu, X., Herbert, S.J., Jin, J., Zhang, Q. & Wang, G. 2004. Responses of photosynthetic rates and yield/quality of main crops to irrigation and manure application in the black soil area of northeast China. *Plant and Soil*, 261(1): 55–60.
- Liu, X., Lee Burras, C., Kravchenko, Y.S., Duran, A., Huffman, T., Morras, H., Studdert, G., Xhang, X., Cruse, R.M. & Yuan, X.H. 2012. Overview of Mollisols in the world: Distribution, land use and management. *Canadian Journal of Soil Science*, 92(3): 383–402. <https://doi.org/10.4141/cjss2010-058>
- Liu, X., Zhang, S., Zhang, X., Ding, G., & Cruse, R. M. 2011. Soil erosion control practices in northeast China: A mini-review. *Soil and Tillage Research*, 117, 44–48. <https://doi.org/10.1016/j.still.2011.08.005>
- Liu, X., Zhang, X., Wang, Y., Sui, Y., Zhang, S., Herbert, S.J. & Ding, G. 2010. Soil degradation: a problem threatening the sustainable development of agriculture in northeast China. *Plant, Soil and Environment*, 56(2): 87–97.
- Lupwayi, N.Z., May, W.E., Kanashiro, D.A. & Petri, R.M. 2018. Soil bacterial community responses to black medic cover crop and fertilizer N under no-till. *Applied Soil Ecology*, 124: 95–103.
- MacDonald, G.K., Bennett, E.M., Potter, P.A. & Ramankutty, N. 2011. Agronomic phosphorus imbalances across the world's croplands. *Proceedings of the National Academy of Sciences*, 108(7): 3086–3091. <https://doi.org/10.1073/pnas.1010808108>
- Maia, S.M., Ogle, S.M., Cerri, C.C. & Cerri, C.E. 2010. Changes in soil organic carbon storage under different agricultural management systems in the Southwest Amazon Region of Brazil. *Soil and Tillage Research*, 106 (2): 177–184.
- Malhi, S. S. & Lemke, R. 2007. Tillage, crop residue and N fertilizer effects on crop yield, nutrient uptake, soil quality and nitrous oxide gas emissions in a second 4-yr rotation cycle. *Soil and Tillage Research*, 96: 269–283.
- Malhi, S. S., Grant, C. A., Johnston, A. M. & Gill, K. S. 2001. Nitrogen fertilization management for no-till cereal production in the Canadian Great Plains: a review. *Soil & Tillage Research*, 60(3–4): 101–122.
- Malhi, S. S., Nyborg, M., Goddard, T. & Puurveen, D. 2011a. Long-term tillage, straw management and N fertilization effects on quantity and quality of organic C and N in a Black Chernozem soil. *Nutrient Cycling in Agroecosystems*, 90(2): 227–241.
- Malhi, S. S., Nyborg, M., Solberg, E. D., Dyck, M. F. & Puurveen, D. 2011b. Improving crop yield and N uptake with long-term straw retention in two contrasting soil types. *Field Crop Research*, 124(3): 378–391.
- Malhi, S.S., Brandt, S.A., Lemke, R., Moulin, A.P. & Zentner, R.P. 2009. Effects of input level and crop diversity on soil nitrate-N, extractable P, aggregation, organic C and N, and nutrient balance in the Canadian Prairie. *Nutrient Cycling in Agroecosystems* 84: 1–22.
- Mamytov A.M. & Bobrov V.P. 1977. Black Earths of Central Asia (In Russian). Frunze, USSR.
- Mamytov, A.M. & Mamytova, G.A. 1988. Soils of the Issyk-Kul Basin and the adjacent territory (In Russian). Frunze, USSR.
- Mamytov, A.M. 1973. Features of Soil Formation in Mountainous Conditions (In Russian). Kirghiz Institute of Soil Science, vol. IV. Frunze, USSR.
- Mann, L. K. 1986. Changes in soil carbon storage after cultivation. *Soil Sci.* 142: 279–288.
- Manojlović, M., Aćin, V. & Šeremešić, S. 2008. Long-term effects of agronomic practices on the soil organic carbon sequestration in Chernozem. *Archives of Agronomy and Soil Science*, 54(4): 353–367.
- Mapfumo, E. Chanasyk, D.S., Naeth, M.A. & Baron, V.S. 1999 Soil compaction under grazing of annual and perennial forages. *Canadian Journal of Soil Science*, 79: 191–199.
- MARA (Ministry of Agriculture and Rural Affairs). 2020. northeast Black Soil Conservation Tillage Action Plan, adopted by the Ministry of Agriculture and Rural Affairs. Cited 28 June 2022. [http://www.moa.gov.cn/nybgb/2020/202004/202005/t20200507\\_6343266.htm](http://www.moa.gov.cn/nybgb/2020/202004/202005/t20200507_6343266.htm)

- MARA (Ministry of Agriculture and Rural Affairs).** 2021. National Implementation Plan on Black Soil Protection (2021–2025), adopted by the Ministry of Agriculture and Rural Affairs of People’s Republic of China. Cited 28 June 2022. <http://www.moa.gov.cn/ztlz/gdzlbhys/htdbhl/202108/P020210804604124115741.pdf>
- Maranhão, D.D., Pereira, M.G., Collier, L.S., Anjos, L.H. dos, Azevedo, A.C. & Cavassani, R. de S.** 2020. Pedogenesis in a karst environment in the Cerrado biome, northern Brazil. *Geoderma*, 365: 114169.
- Martens, J. T., Entz, M. & Wonneck, M.** 2013. *Ecological farming systems on the Canadian prairies. A path to profitability, sustainability and resilience*. Manitoba: University of Manitoba.
- Matsui K., Takata Y., Matsuura S. & Wagai R.** 2021a. Soil organic carbon was more strongly linked with soil phosphate fixing capacity than with clay content across 20 000 agricultural soils in Japan: a potential role of reactive aluminum revealed by soil database approach. *Soil Sci. Plant Nutr.*, 67: 233–242.
- Matsui K., Takata Y., Maejima Y., Kubotera H., Obara H. & Shirato Y.** 2021b. Soil carbon and nitrogen stock of the Japanese agricultural land estimated by the national soil monitoring database (2015–2018). *Soil Sci. Plant Nutr.* (In press)
- Matsumoto Y.** 1992. Soil conservation conducted by actual furrowing practice on steep farmland of Kuroboku soil (In Japanese). *J. Jap. Soc. Soil Phys.*, 66: 55–63.
- Matsuyama N., Saigusa M. et al.,** 2005. Acidification and soil productivity of allophanic andosols affected by application of fertilizers. *Soil Sci Plant Nutr.*, 51: 117–123.
- McConkey, B.G., Liang, B.C., Campbell, C.A., Curtin, D., Moulin, A., Brandt, S.A. & Lafond, G.P.** 2003. Crop rotation and tillage impact on carbon sequestration in Canadian prairie soils. *Soil and Tillage Research*, 74: 8190.
- McMichael, C. H., Palace, M. W., Bush, M. B., Braswell, B., Hagen, S., Neves, E. G., Silman M. R., Tamanaha E. K. & Czarnecki, C.** 2014. Predicting pre-Columbian anthropogenic soils in Amazonia. *Proceedings of the Royal Society B: Biological Sciences*, 281(1777): 20132475.
- Medvedev, V.V.** 2012. Soil monitoring of the Ukraine. *The Concept. Results. Tasks. (2nd rev. and adv. edition)*. Kharkiv: CE “City printing house.
- Melo, A.F.D., Souza, C.M.M., Rego, L.G.S., Lima, N.S. & Moura, I.N.B.M.** 2017. Pedogênese de chernossolos derivados de diferentes materiais de origem no oeste potiguar. *Revista Agropecuária Científica no Semiárido*. 13: 229–235.
- Meng, Q., Zhao, S., Geng, R., Zhao, Y., Wang, Y., Yu, F., Zhang, J. & Ma, J.** 2021. Does biochar application enhance soil salinization risk in black soil of northeast China (a laboratory incubation experiment)? *Archives of Agronomy and Soil Science*, 67(11): 1566–1577.
- Menšík, L., Hlisenikovsky, L. & Kunzová, E.** 2019. The state of the soil organic matter and nutrients in the long-term field experiments with application of organic and mineral fertilizers in different soil-climate conditions in the view of expecting climate change. In *Organic fertilizers-history, production and applications*. IntechOpen.
- Merante, P., Dibari, C., Ferrise, R., Sánchez, B., Iglesias, A., Lesschen, J. P. Peter, K., Jagadeesh Y., Pete S. & Bindi, M.** 2017. Adopting soil organic carbon management practices in soils of varying quality: Implications and perspectives in Europe. *Soil and Tillage Research*, 165: 95–106.
- Mermut A. R. & Acton, D. F.** 1984. The Age of Some Holocene Soils on the Ear Lake Terraces in Saskatchewan. *Canadian J. Soil Science*. 64, 163–172.
- Milić, S., Ninkov, J., Zeremski, T., Latković, D., Šeremešić, S., Radovanović, V. & Žarković, B.** 2019. Soil fertility and phosphorus fractions in a calcareous chernozem after a long-term field experiment. *Geoderma*, 339: 9–19.
- Ministry of Natural Resources and Environment of the Russian Federation.** 2022. Central Black Earth State Reserve named after Professor V.V. Alekhine [In Russian]. In: *Ministry of Natural Resources and Environment of the Russian Federation*. Russia. Cited 7 June 2022. <http://zapoved-kursk.ru/>
- Miroshnychenko, M. & Khodakivska, O.** 2018. Black soils in Ukraine. International Symposium on Black Soils (ISBS18): Protect Black Soils, Invest in the Future. Charbin
- Misra, R.V., Roy, R.N. & Hiraoka, H.** 2003. *On-farm composting methods*. Rome, Italy: UN-FAO. (also available at: <http://www.fao.org/docrep/007/y5104e/y5104e00.htm#Contents>)
- Modernel, P., Rossing, W.A.H., Corbeels, M., Dogliotti, S., Picasso V. & Tittonell, P.** 2016. Land use change and ecosystem service provision in Pampas and Campos grasslands of southern South America. *Environmental Research Letters*, 11–113002.

- Mokhtari, M. & Dehghani, M.** 2012. Swell-shrink behavior of expansive soils, damage and control. *Electronic Journal of Geotechnical Engineering*, 17: 2673–2682.
- Monger, H.C., Kraimer, R.A., Khresat, S., Cole, D.R., Wang, X.J. & Wang, J.P.** 2015a. Sequestration of inorganic carbon in soil and groundwater. *Geology*, 43:375–378. doi:10.1130/G36449.1.
- Monger, H.C., Sala, O.E., Duniway, M., Goldfus, H., Meir, I.A., Poch, R.M. & Vivoni, E.R.** 2015b. Legacy effects in linked ecological–soil–geomorphic systems of drylands. *Frontiers in Ecology and the Environment*, 13(1): 13–19.
- Montanarella, L., Panagos, P. & Scarpa, S.** 2021. The Relevance of Black Soils for Sustainable Development. In D. Dent & B. Boincean, eds. *Regenerative Agriculture*, pp. 69–79. Cham, Springer.
- Montanarella, L., Pennock, D.J., McKenzie, N., Badraoui, M., Chude, V., Baptista, I., Mamo, T., Yemefack, M., Aulakh, m.s., Yagi, K., Hong, Suk Young., Vijarnsorn, P., Zhang, G., Arrouays, D., Black, H., Krasilnikov, P., JSobocká, A., Alegre, J., Henriquez, C.R., Mendonça-Santos, M.L., Taboada, M., Espinosa-Victoria, D., AlShankiti, A., AlaviPanah, S.K., Elsheikh, E.A.E.M., Hempel, J., Arbestain, M.C., Nachtergaele, F. & Ronald V.** 2016. World's soils are under threat. *SOIL*, 2(1): 79–82. <https://doi.org/10.5194/soil-2-79-2016>
- Moon, D.** 2020. Soil Science I. In: The American Steppes: The Unexpected Russian Roots of Great Plains Agriculture, 1870s–1930s. pp. 188–225. Studies in Environment and History. Cambridge, Cambridge University Press. <https://doi.org/10.1017/9781316217320.006>
- Mora, S.** 2003. Archaeobotanical methods for the study of Amazonian Dark Earths. In J. Lehmann, D. Kern, B. Glaser, & W. Woods, eds., *Amazonian Dark Earths: Origin, Properties, Management*, pp. 205–225. Netherlands, Kluwer Academic Publishers.
- Morales, M., Otero, J., Van der Hammen, T., Torres, A., Cadena, C., Pedraza, C., Rodríguez, N., Franco, C., Betancourth, J.C., Olaya, E., Posada, E. & L. Cárdenas.** 2007. Atlas de páramos de Colombia. Instituto de Investigación de Recursos Biológicos Alexander von Humboldt, pp. 208. Bogotá, D. C.
- Morcote-Ríos, G. & Sicard, T.L.** 2012. *Las Terras Pretas del Igarapé Takana. Un Sistema de Cultivo Precolombino en Leticia-Amazonas*. Universidad Nacional de Colombia, Bogotá, Colombia.
- Moretti, L., Morrás, H., Pereyra, F. & Schulz, G.** 2019. Soils of the Chaco Region. In G. Rubio, R. Lavado, & F. Pereyra, eds. *The soils of Argentina*, Chapter 10, pp. 149–160. World Soils Book Series, Switzerland, Springer.
- Morrás, H. & Moretti, L.** 2016. A new soil-landscape approach to the genesis and distribution of Typic and Vertic Argiudolls in the Rolling Pampa of Argentina. In A. Zinck, G. Metternich, G. Bocco, & H. del Valle eds. *Geopedology – An Integration of Geomorphology and Pedology for Soil and Landscape Studies*, pp. 193–209.
- Morrás, H.** 2017. Propiedades químicas y físicas de suelos hidromórficos de la fracción norte de los Bajos Submeridionales. In E. Taleisnik & R. Lavado, eds. *Ambientes salinos y alcalinos de la Argentina*, pp. 29–54. Recursos y aprovechamiento productivo. Buenos Aires, Orientación Gráfica Editora.
- Morrás, H.** 2020. Modelos composicionales y áreas de distribución de los aportes volcánicos en los suelos de la Pampa Norte (Argentina) en base a la mineralogía de arenas. In P. Imbellone & O. Barbosa, eds. *Suelos y Vulcanismo*, pp. 127–167. Buenos Aires, Asociación Argentina de la Ciencia del Suelo.
- Morris, N.L., Miller, P.C.H., Orson, J.H. & Froud-Williams, R. J.** 2010. The adoption of non-inversion tillage systems in the United Kingdom and the agronomic impact on soil, crops and the environment—A review. *Soil and Tillage Research*, 108(1–2): 1–15.
- Mueller, N.D., Gerber, J.S., Johnston, M., Ray, D.K., Ramankutty, N. & Foley, J.A.** 2012. Closing yield gaps through nutrient and water management. *Nature*, 490(7419): 254–257. <https://doi.org/10.1038/nature11420>
- Nanzyo, M., Dahlgren R. & Shoji S.** 1993. “Chemical characteristics of volcanic ash soils.” In S. Shoji, M. Nanzyo and R. Dahlgren, eds. *Volcanic ash soils – Genesis, Properties and Utilization*, pp. 145–187. The Netherlands, Elsevier.
- National Bureau of statistics of China.** 2015. China Statistical Yearbook. <http://www.stats.gov.cn/tjsj/ndsj/2015/indexeh.htm>
- National report on the state of the environment in Ukraine in 2018.** 2020. Kiev. Ministry of Ecology and Natural Resources of Ukraine (In Ukrainian). Cited 1 May 2021. <https://mepr.gov.ua/news/35937.html>
- Nauman, T.W. & Thompson, J.A.** 2014. Semi-automated disaggregation of conventional soil maps using knowledge driven data mining and classification trees. *Geoderma*, 213: 385–399.



- Neall, V. E. 2009. Volcanic soils. *Land use, land cover and soil sciences*, 7: 23–45.
- Neves, E.G., Petersen, J.B., Bartone, R.N. & Heckenberger, M.J. 2004. *The timing of Terra preta formation in the Central Amazon: archaeological data from three sites*. In B. Glaser, & W. Woods, eds. *Explorations in Amazonian Dark Earths*, pp. 125–134.
- Ngatia, L., Grace III, J. M., Moriasi, D., & Taylor, R. 2019. Nitrogen and phosphorus eutrophication in marine ecosystems. *Monitoring of marine pollution*, 1–17.
- Nowatzki, J., Endres, G. & DeJong-Hughes, J. 2017. *Strip Till for Field Crop Production* Pages 1–10. Fargo, US, North Dakota State University express.
- Nunes, M.R., Van Es, H.M., Schindelbeck, R., Ristow, A.J. & Ryan, M. 2018. No-till and cropping system diversification improve soil health and crop yield. *Geoderma*, 328: 30–43.
- Oades, J.M. 1993. The role of biology in the formation, stabilization and degradation of soil structure. In: L. Brussaard & M.J. Kooistra, eds. *Soil Structure/Soil Biota Interrelationships*. pp. 377–400. Amsterdam, Elsevier. <https://doi.org/10.1016/B978-0-444-81490-6.50033-9>
- O'Donnell, J. A., Aiken, G. R., Butler, K. D., Guillemette, F., Podgorski, D. C. & Spencer, R. G. 2016. DOM composition and transformation in boreal forest soils: The effects of temperature and organic-horizon decomposition state. *Journal of Geophysical Research: Biogeosciences*, 121(10): 2727–2744.
- Okuda, T., Fujita, T., Fujie, K., Kitagawa, Y., Saito, M. & Naruse, T. 2007. Influence of eolian dust brought from the Precambrian area in northern Asia on the parent materials in a fine-textured soil developed on the tertiary rock in Mt. Horyu, Noto peninsula, central Japan (In Japanese with English summary). *Pedologist*, 51:104–110.
- Oldfield, E.E., Bradford, M.A. & Wood, S.A. 2019. Global meta-analysis of the relationship between soil organic matter and crop yields. *SOIL*, 5(1): 15–32. <https://doi.org/10.5194/soil-5-15-2019>
- Oliveira, I.A., Campos, M.C.C., Freitas, L. & Soares, M.D.R. 2015a. Caracterização de solos sob diferentes usos na região sul do Amazonas. *Acta Amazonica*, 45(3): 1–12.
- Oliveira, I.A., Campos, M.C.C., Marques Junior, J., Aquino, R.E., Teixeira, D.B. & Silva, D.M.P. 2015b. Use of scaled semivariograms in the planning sample of soil chemical properties in southern Amazonas, Brazil. *Rev. Bras. Ci Solo*, 39(5): 31–39.
- OpenLandMap/global-layers. 2022. In: *GüLab*. Cited 4 April 2022. <https://gitlab.com/openlandmap/global-layers>
- Otero, J.D., Figueroa, A., Muñoz, F.A. & Peña, M.R. 2011. Loss of soil and nutrients by surface runoff in two agro-ecosystems within an Andean paramo area. *Ecol. Eng.*, 37 (12): 2035–2043, 10.1016/j.ecoleng.2011.08.001.
- Ouyang, W., Wu, Y., Hao, Z., Zhang, Q., Bu, Q. & Gao, X. 2018. Combined impacts of land use and soil property changes on soil erosion in a mollisol area under long-term agricultural development. *The Science of the Total Environment*, 613–614: 798–809. <https://doi.org/10.1016/j.scitotenv.2017.09.173>
- Overbeck, G.E., Müller, S.C., Fidelis, A., Pfadenhauer, J., Pillar, V.D., Blanco, C.C., Boldrini, I.I., Both, R. & Foerneck, E.D. 2007. Brazil's neglected biome: The South Brazilian Campos. *Perspectives in Plant Ecology, Evolution and Systematics*. 9:101–116.
- Overbeck, G.E., Müller, S.C., Pillar, V.D. & Pfadenhauer, J. 2005. Fine-scale post-fire dynamics in southern Brazilian subtropical grassland. *Journal of Vegetation Science*, 16: 655–664.
- Overbeck, G.E., Müller, S.C., Pillar, V.D. & Pfadenhauer, J. 2006. Floristic composition, environmental variation and species distribution patterns in burned grassland in southern Brazil. *Braz. J. Biol.*, 66: 1073–1090.
- Pape, J.C. 1970. Plaggen soils in the Netherlands. *Geoderma*, 4: 229–255.
- Peña-Venegas, C.P. & Vanegas-Cardona, G.I. 2010. *Dinámica de los suelos amazónicos: Procesos de degradación y alternativas para su recuperación*. Instituto Sinchi. Bogotá, Colombia.
- Peña-Venegas, C.P., Stomph, T.J., Verschoor, G., Echeverri, J.A. & Struik, P.C. 2016. Classification and Use of Natural and Anthropogenic Soils by Indigenous Communities of the Upper Amazon Region of Colombia. *Hum Ecol*, 44: 1–15. <https://doi.org/10.1007/s10745-015-9793-6>.
- Pepo, P., Vad, A. & Berényi, S. 2006. Effect of some agrotechnical elements on the yield of maize on chernozem soil. *Cereal Research Communications*, 34(1): 621–624.

- Peralta, G., Alvarez, C.R. & Taboada, M.Á.** 2021. Soil compaction alleviation by deep non-inversion tillage and crop yield responses in no tilled soils of the Pampas region of Argentina. A meta-analysis. *Soil and Tillage Research*, 211: 105022. <https://doi.org/10.1016/j.still.2021.105022>
- Pereira, M.G., Schiavo J.A., Fontana A., Dias Neto, A.H. & Miranda, L.P.M.** 2013. Caracterização e classificação de solos em uma toposequência sobre calcário na serra da Bodoquena, MS. *Rev. Bras. Ci.Solo*, 37: 25–36.
- Pereyra, F. & Bouza, P.** 2019. Soils from the Patagonian Region. In G. Rubio, R. Lavado, & F. Pereyra, eds. *The soils of Argentina Chapter 7*, pp. 101–121. Switzerland, Springer, World Soils Book Series.
- Pikul, J.L., Chilom, G., Rice, J., Eynard, A., Schumacher, T.E., Nichols, K., Johnson, J.M.F., Wright, S., Caesar, T. & Ellsbury, M.** 2009. Organic matter and water stability of field aggregates affected by tillage in South Dakota. *Soil Science Society of America Journal*, 73: 197–206.
- Pillar, V.D., Tornquist, C.G. & Bayer, C.** 2012. The southern Brazilian grassland biome: soil carbon stocks, fluxes of greenhouse gases and some options for mitigation. *Brazilian Journal of Biology*, 72:673–681.
- Pinto, L.F.S. & Kämpf, N.** 1996. Solos derivados de rochas ultrabásicas no ambiente subtropical do Rio Grande do Sul. *Revista Brasileira de Ciência do Solo*, 20: 447–458.
- Plisko, I.V., Bigun, O.M., Lebed, V.V., Nakisko, S.G. & Zalavsky, Y.V.** 2018. Creation of a national map of organic carbon reserves in the soils of Ukraine. *Agrochemistry and soil science*, 87: 57–62.
- Podolsky, K., Blackshaw, R.E. & Entz, M.H.,** 2016. A comparison of reduced tillage implements for organic wheat production in Western Canada. *Agronomy Journal*, 108(5): 2003–2014.
- Poeplau, C. & Don, A.** 2015. Carbon sequestration in agricultural soils via cultivation of cover crops—A meta-analysis. *Agriculture, Ecosystems & Environment*, 200: 33–41.
- Poeplau, C., Don, A., Vesterdal, L., Leifeld, J., Van Wesemael, B., Schumacher, J. & Gensior, A.** 2011. Temporal dynamics of soil organic carbon after land-use change in the temperate zone – carbon response functions as a model approach. *Global Change Biology*, 17(7): 2415–2427. <https://doi.org/10.1111/j.1365-2486.2011.02408.x>
- Polidoro, J. C., Freitas, P.L. de, Hernani, L.C., Anjos, L. H. C., Rodrigues, R. de A. R., Cesário, F.V., Andrade, A. G. & Ribeiro, J. L.** 2021. Potential impact of plans and policies based on the principles of Conservation Agriculture on the control of soil erosion in Brazil. *Land Degradation & Development*. 32: 1–12.
- Polupan, M.I., Velichko, V.A. & Solovey, V.B.** 2015. Development of Ukrainian agronomic soil science: genetic and production bases (In Ukrainian). Kyiv, Ahrarna nauka.
- Polupan, N.I.** 1988. *Soils of Ukraine and increase of their fertility: Vol. 1. Ecology, regimes and processes, classification and genetic and production aspects* (In Russian). Kiev, Urogaj.
- Poulenard, J., Podwojewski, P., Janeau, J.L. & Collinet, J.** 2001. Runoff and soil erosion under rainfall simulation of andisols from the ecuadorian páramo: effect of tillage and burning. *Catena*, 45(3): 185–207.
- Pretty, J.** 2008. Agricultural sustainability: concepts, principles and evidence. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 363(1491): 447–465.
- Pugliese, J.Y., Culman, S.W. & Sprunger, C.D.** 2019. Harvesting forage of the perennial grain crop kernza (*Thinopyrum intermedium*) increases root biomass and soil nitrogen cycling. *Plant and Soil*, 437(1–2): 241–254.
- Pylypenko, H.P., Varlamova, N.Y., Borshch, O.V. & Borshch, A.V.** 2002. Aridization and desertification of the steppes of southern Ukraine (In Ukrainian). *Bulletin of Odessa National University*, 7 (4): 45–51.
- Qiao, Y., Miao, S., Zhong, X., Zhao, H. & Pan, S.** 2020. The greatest potential benefit of biochar return on bacterial community structure among three maize-straw products after eight-year field experiment in Mollisols. *Applied Soil Ecology*, 147: 103432.
- Qin, Z., Yang, X., Song, Z., Peng, B., Zwieten, L.V., Yue, C., Wu, S., Mohammad, M.Z. & Wang, H.** 2021. Vertical distributions of organic carbon fractions under paddy and forest soils derived from black shales: Implications for potential of long-term carbon storage. *Catena*, 198: 105056.
- Raza, S., Miao, N., Wang, P., Ju, X., Chen, Z., Zhou, J. & Kuzakov, Y.** 2020. Dramatic loss of inorganic carbon by nitrogen-induced soil acidification in Chinese croplands. *Global change biology*, 26(6): 3738–3751. <https://doi.org/10.1111/gcb.15101>

- Reifenberg, A.** 1947. *The Soils of Palestine*. Rev. 2nd ed. Thomas Murbery and Co., London.
- Research Conference material.** 2015. Strengthening Mongolia's Pastureland Rehabilitation Capacity. Ulaanbaatar, Mongolia.
- Rezapour, S. & Alipour, O.** 2017. Degradation of Mollisols quality after deforestation and cultivation on a transect with Mediterranean condition. *Environmental Earth Sciences*, 76(22): 755. <https://doi.org/10.1007/s12665-017-7099-2>
- Richter, D.D. & Babbar, L.I.** 1991. Soil Diversity in the Tropics. *Advances in Ecological Research*, 21: 315–389.
- Ritter, J.** 2012. *Soil erosion – Causes and effects*, OMAFRA Factsheet, Queens Printer for Ontario, Toronto. Nov 11, 2020. (also available at <http://www.omafra.gov.on.ca/english/engineer/facts/12-053.htm>)
- Roecker, S., Ferguson, C. & Wills, S.** Chapter 2 "Portrait of Black Soils." USA. The Global Status of Black Soils. A FAO Report (this publication)
- Roesch, L.F.W., Vieira, F.C.B., Pereira, V.A., Schünemann, A.L., Teixeira, I.F., Senna, A.J.T. & Stefeno, V.M.** 2009. The Brazilian Pampa: A Fragile Biome. *Diversity*, 1: 82–198.
- Rogovska, N., Laird, D.A., Rathke, S.J. & Karlen, D.L.** 2014. Biochar impact on Midwestern Mollisols and maize nutrient availability. *Geoderma*, 230: 340–347.
- Rojas, R.V., Achouri, M., Maroulis, J. & Caon, L.** 2016. Healthy soils: a prerequisite for sustainable food security. *Environmental Earth Sciences*, 75(3): 180. <https://doi.org/10.1007/s12665-015-5099-7>
- Romero, C.M., Hao, X., Li, C., Owens, J., Schwinghamer, T., McAllister, T.A. & Okine, E.** 2021. Nutrient retention, availability and greenhouse gas emissions from biochar-fertilized Chernozems. *Catena*, 198: 105046.
- Royal Society of London.** 2009. Reaping the benefits: Science and the sustainable intensification of global agriculture. The Royal Society. London SW1Y 5AG
- Rubio, G., Lavado, R. S. & Pereyra, F. X.** 2019. *The soils of Argentina*. Springer International Publishing.
- Rubio, G., Lavado, R., Pereyra, F., Taboada, M., Moretti, L., Rodríguez, D., Echeverría, H. & Panigatti, J.** 2019. Future Issues. In Rubio, G., Lavado, R. & Pereyra, F. eds. *The soils of Argentina*. Springer, World Soils Book Series, Switzerland, Chapter 19, pp. 261–263
- Rubio, G., Pereyra, F. & Taboada, M.** 2019. Soils of the Pampean Region. In G. Rubio, R. Lavado, & F. Pereyra, eds. *The soils of Argentina*, pp. 81–100. Switzerland, Springer, World Soils Book Series.
- Russell, A.E., Laird, D.A., Parkin, T.B. & Mallarino, A.P.** 2005. Impact of nitrogen fertilization and cropping system on carbon sequestration in Midwestern Mollisols. *Soil Science Society of America Journal*, 69(2): 413–422.
- Rusu A.** 2017. The influence of the straw applied as a fertilizer on the humus in the ordinary chernozem. In: Proceedings of the International Scientific Conference, dedicated to the 120th anniversary of the birth of Academician Ion Dicusar, 6–7 September 2017, Chisinau, Republic of Moldova. [in Romanian]
- Ryan, J., Pala, M., Masri, S., Singh, M. & Harris, H.** 2008. Rainfed wheat-based rotations under Mediterranean conditions: Crop sequences, nitrogen fertilization, and stubble grazing in relation to grain and straw quality. *European Journal of Agronomy*, 28(2): 112–118.
- Ryan, M.R., Crews, T.E., Culman, S.W., DeHaan, L.R., Hayes, R.C., Jungers, J.M. & Bakker, M.G.** 2018. Managing for Multifunctionality in Perennial Grain Crops. *BioScience*, 68: 294–304.
- Saigusa M., Matsuyama N. & Abe A.** 1992. Distribution of Allophanic Andosols and Nonallophanic Andosols in Japan based on the data of soil survey reports on reclaimed land (In Japanese with English summary). *Jpn J Soil Sci Plant Nutr* 63: 646–651.
- Sainz-Rozas, H., Echeverría, H. & Angelini, H.** 2011. Niveles de materia orgánica y pH en suelos agrícolas de la Región Pampeana y extra-Pampeana argentina. *Ciencia del Suelo*, 29 (1):29–37.
- Santos, H.G., Jacomine, P.K.T., Anjos, L.H.C., Oliveira, V.A., Lumberras, J.F., Coelho, M.R., Almeida, J.A, Araújo Filho, J.C., Oliveira, J.B. & Cunha, T.J.F.** 2018a. *Brazilian Soil Classification System. 5th ed. rev. and exp. E-book*. Brasília: Embrapa.
- Santos, L.A.C., Araújo, J.K.S., Souza Júnior, V.S., Campos, M.C.C., Corrêa, M.M. & Souza, R.A.S.** 2018b. Pedogenesis in an Archaeological Dark Earth – Mulatto Earth Catena over Volcanic Rocks in Western Amazonia, Brazil. *Rev. Bras. Ci Solo*, 42(3): 1–18.
- Santos, L.A.C., Campos, M.C.C., Aquino, R.E., Bergamin, A.C., Silva, D.M.P., Marques Junior, J. & Franca, A.B.C.** 2013. Caracterização e gênese de terras pretas arqueológicas no sul do Estado do Amazonas. *Rev. Bras. Ci Solo*, 37(5): 825–836.



- Sarmiento, F.O. & Frolich, L.M.** 2002. Andean cloud forest tree lines: Naturalness, agriculture and the human dimension. *Mt. Res. Dev.* 22: 278–287.
- Sato, M., Tállai, M., Kovács, A.B., Vágó, I., Kátai, J., Matsushima, M.Y., Sudo, S. & Inubushi K.** 2022. Effects of a new compost-chemical fertilizer mixture on CO<sub>2</sub> and N<sub>2</sub>O production and plant growth in a Chernozem and an Andosol. *Soil Science and Plant Nutrition*, 68(1): 175–182.
- Sayegh, A. H. & Salib, A. J.** 1969. Some physical and chemical properties of soils in the Beqa'a plain, Lebanon. *Journal of Soil Science*, 20: 167–175.
- Schmidt, M.J., Rapp Py-Daniel, A., de Paula Moraes, C., Valle, R.B.M., Caromano, C.F., Texeira, W.G., Barbosa, C.A., Barbosa, C.A., Fonseca, J.A., Magalhães, M.P., Santos, D.S.C., Silva, R.S., Guapindaia, V.L., Moraes, B., Lima, H.P., Neves, E.G. & Heckenberger, M.J.** 2014. Dark earths and the human built landscape in Amazonia: a widespread pattern of anthrosol formation. *Journal of Archaeological Science*, 42: 152–165. <https://doi.org/10.1016/j.jas.2013.11.002>
- Schönbach, P., Wan, H., Gierus, M., Bai, Y., Müller, K., Lin, L., Susenbeth, A. & Taube, F.** 2011. Grassland responses to grazing: effects of grazing intensity and management system in an Inner Mongolian steppe ecosystem. *Plant and Soil*, 340(1): 103–115.
- Sedov, S., Solleiro-Rebolledo, E., Morales-Puente, P., Arias-Herrera, A., Vallejo-Gómez, E. & Jasso-Castañeda, C.** 2003. Mineral and organic components of the buried paleosols of the Nevado de Toluca, Central Mexico as indicators of paleoenvironments and soil evolution. *Quaternary International*, 106: 169–184.
- Servicio Agrícola y Ganadero (SAG).** 2003. El pastizal de Tierra del Fuego. Guía de uso, condición actual y propuesta de seguimiento para determinación de tendencia. *Punta Arenas, XII Región de Magallanes y Antártica Chilena*. Chile, Gobierno de Chile.
- Servicio Agrícola y Ganadero (SAG).** 2004a. El pastizal de Magallanes. Guía de uso, condición actual y propuesta de seguimiento para determinación de tendencia. *XII Región de Magallanes y Antártica Chilena, Punta Arenas*. Chile, Gobierno de Chile.
- Servicio Agrícola y Ganadero (SAG).** 2004b. El pastizal de Última Esperanza y Navarino. Guía de uso, condición actual y propuesta de seguimiento para determinación de tendencia. *XII Región de Magallanes y Antártica Chilena*. Punta Arenas, Chile, Gobierno de Chile.
- Sherwood, S. & Uphoff, N.** 2000. Soil health: research, practice and policy for a more regenerative agriculture. *Applied Soil Ecology*, 15(1): 85–97.
- Shiono, T.** 2015. Soil Erosion and Sediment Control Measures for Farmlands in Japan, MARCO International Symposium 2015: Next Challenges of Agro-Environmental Research in Monsoon Asia, 26–28 August 2015. Tsukuba, Japan.
- Shiono, T., Okushima S., Takagi A. & Fukumoto M.** 2004. Influence of Cabbage Cultivation on Soil Erosion in a Ridged Field with Kuroboku Soil (In Japanese). *Journal of Irrigation Engineering and Rural Planning*, 230: 1–9.
- Shoji, S.** 1984. Andosols, Exploring its Today's Issues, Kagakutoseibutsu (In Japanese). *Chemistry and Biology*. 22, 242–250.
- Shoji, S., Nanzyo M. & Dahlgren R.A.** 1993. Volcanic ash soils-genesis, properties and utilization. *Developments in Soil Science 21*, Elsevier, Amsterdam.
- Shpedt, A. A. & Aksenova, Y. V.** 2021. Modern assessment of soil resources of Kyrgyzstan. In *IOP Conference Series: Earth and Environmental Science Vol. 624, No. 1*, p. 012233). IOP Publishing.
- Silveira, V.C.P., Gonzalez, J.A. & Fonseca, E.L.** 2017. Land use changes after the period commodities rising price in the Rio Grande do Sul State. *Brazil. Ciência Rural*, 47(4): e20160647.
- Skidmore, E. L.** 2017. Wind erosion. In *Soil erosion research methods*, pp. 265–294. Routledge.
- Smith, A.** 2022. A Russia-Ukraine war could ripple across Africa and Asia. In: *Foreign Policy*. Cited 1 June 2022. <https://foreignpolicy.com/2022/01/22/russia-ukraine-war-grain-exports-africa-asia/>
- Smith, K.L.** 1999. *Epidemiology of Anthrax in the Kruger National Park, South Africa: Genetic Diversity and Environment*. LSU Historical Dissertations and Theses. 6962.
- Smith, P., Calvin, K., Nkem, J., Campbell, D., Cherubini, F., Grassi, G., Korotkov, V., Hoang, A.L., Lwasa, S., McElwee, P., Nkonya, E., Saigusa, N., Soussana, J., Taboada, M.A., Manning, F.C., Nampanzira, D., Arias-Navarro, C., Vizzarri, M., House, J., Roe, S., Cowie, A., Rounsevell, M. & Arneeth, A.** 2020. Which practices co-deliver food security, climate change mitigation and adaptation, and combat land degradation and desertification? *Global Change Biology*, 26(3): 1532–1575. <https://doi.org/10.1111/gcb.14878>

- Smith, P., House, J.I., Bustamante, M., Sobocká, J., Harper, R., Pan, G., West, P.C., Clark, J.M., Adhya, T., Rumpel, C., Paustian, K., Kuikman, P., Cotrufo, M.F., Elliott, J.A., McDowell, R., Griffiths, R.I., Asakawa, S., Bondeau, A., Jain, A.K., Meersmans, J. & Pugh, T.A.M. 2016. Global change pressures on soils from land use and management. *Global Change Biology*, 22(3): 1008–1028. <https://doi.org/10.1111/gcb.13068>
- Snyder, R.L. & Melo-Abreu, J. de. 2005. Frost protection: fundamentals, practice and economics. Volume 1. FAO.
- Soil Science Society of America (SSSA). 2008. Glossary of soil science terms. [online]. [Cited 19 February 2020]. <https://www.soils.org/publications/soils-glossary>
- Soil Science Society of America (SSSA). 2020. Glossary of terms. [Online]. [Cited 27 November 2020]. <https://www.soils.org/publications/soils-glossary#>
- Soil Survey Staff. 1999. *Soil taxonomy: A basic system of soil classification for making and interpreting soil surveys*. Second edition. U.S. Department of Agriculture Handbook No. 436. Natural Resources Conservation Service.
- Soil Survey Staff. 2014. Keys to Soil Taxonomy, 12th ed. USDA-Natural Resources Conservation Service, Washington, DC.
- Sojka, R.E. & Bjorneberg, D.L. 2017. *Encyclopedia of Soil Science*. Boca Raton, USA, CRC Press.
- Sollenberger, L. E., Kohman, M. M., Dubeux, Jr. J. C. B. & Silveira, M. L. 2019. Grassland Management Affects Delivery of Regulating and Supporting Ecosystem Services. *Crop Science*, 59: 441–459.
- Sollenberger, L.E., Agouridis, C.T., Vanzant, E.S., Franzluebbers, A.J. & Owens, L.B. 2012. *Conservation outcomes from pastureland and hayland practices: Assessment, Recommendations, and Knowledge Gaps*. Lawrence, KS, Allen Press.
- Sombroek, W. I. M., Ruivo, M. D. L., Fearnside, P. M., Glaser, B. & Lehmann, J. 2003. Amazonian dark earths as carbon stores and sinks. *In Amazonian dark earths*, pp: 125–139. Dordrecht, Springer.
- Song, Z., Gao, H., Zhu, P., Peng, C., Deng, A., Zheng, C., AbdulMannaf, M., Nurullislam, M. & Zhang, W. 2015. Organic amendments increase corn yield by enhancing soil resilience to climate change. *The Crop Journal*, 3(2): 110–117.
- Sorokin, A., Owens, P., Láng, V., Jiang, Z.-D., Michéli, E. & Krasilnikov, P. 2021. “Black soils” in the Russian Soil Classification system, the US Soil Taxonomy and the WRB: Quantitative correlation and implications for pedodiversity assessment. *Catena*, 196: 104824.
- Soussana, J.F., Lutfalla, S., Ehrhardt, F., Rosenstock, T., Lamanna, C., Havlík, P., Richards, M., Wollenberg, Eva (Lini), Chotte, Jean-Luc., Torquebiau, Emmanuel., Ciaís, Philippe., Smith, Pete. & Lal, Rattan. 2019. Matching policy and science: Rationale for the ‘4 per 1000 – soils for food security and climate’ initiative. *Soil and Tillage Research*, 188: 3–15. <https://doi.org/10.1016/j.still.2017.12.002>
- SSSA (Soil Science Society of America). 2015. Desertification and the American Dust Bowl. Cited 6 June 2022. <https://www.soils.org/files/sssaiys/dust-bowl-activity.pdf>
- St. Luce, M., Lemke, R., Gan, Y., McConkey, B., May, W.E., Campbell, C., Zentner, R., Wang, H., Kroebel, R., Fernandez, M. & Brandt, K. 2020. Diversifying cropping systems enhances productivity, stability, and nitrogen use efficiency. *Agronomy Journal*, 112(3): 1517–1536.
- Stainsby, A., May, W.E., Lafond, G.P. & Entz, M.H. 2020. Soil aggregate stability increased with a self-regenerating legume cover crop in low-nitrogen, no-till agroecosystems of Saskatchewan, Canada. *Canadian Journal of Soil Science*, 100: 314–318.
- Steffan, J.J., Brevik, E.C., Burgess, L.C. & Cerdà, A. 2018. The effect of soil on human health: an overview. *European Journal of Soil Science*, 69(1): 159–171. <https://doi.org/10.1111/ejss.12451>
- Stewart, W. M., Dibb, D. W., Johnston, A. E. & Smyth, T. J. 2005. The contribution of commercial fertilizer nutrients to food production. *Agronomy journal*, 97(1): 1–6.
- Strauch, B. & Lira, R. 2012. Bases para la producción ovina en Magallanes. *Instituto de Investigaciones Agropecuarias, Boletín INIA N° 244*, pp. 154. Punta Arenas, Chile, Centro Regional de Investigación Kampenaike.
- Stupakov, A.G., Orekhovskaya, A.A., Kulikova, M.A., Manokhina, L.A., Panin, S.I. & Geltukhina, V.I. 2019. Ecological and agrochemical bases of the nitrogen regime of typical chernozem depending on agrotechnical methods. *Conference Series: Earth and Environmental Science*, pp. 052027. IOP Publishing.

- Sun, B., Jia, S., Zhang, S., McLaughlin, N.B., Zhang, X., Liang, A., Chen, X., Wei, S. & Liu, S. 2016. Tillage, seasonal and depths effects on soil microbial properties in black soil of northeast China. *Soil and Tillage Research*, 155: 421–428.
- Taboada, M.Á., Costantini, A.O., Busto, M., Bonatti, M. & Sieber, S. 2021. Climate change adaptation and the agricultural sector in South American countries: Risk, vulnerabilities and opportunities. *Rev. Bras. Cienc. Solo*, 45. <https://doi.org/10.36783/18069657rbcs20210072>
- Tahat, M.M., Alananbeh, K.M., Othman, Y.A. & Leskovar, D.I. 2020. Soil health and sustainable agriculture. *Sustainability*, 12: 48–59.
- Takata, Y., Kawahigashi, M., Kida, K., Tani, M., Kinoshita, R., Ito, T., Shibata, M., Takahashi, T., Fujii, K., Imaya, A., Obara, H., Maejima, Y., Kohyama, K. & Kato, T. 2021. Major Soil Types, In R. Hatano, H. Shinjo & Y. Takata eds. *The Soil of Japan*, pp. 69–134, Springer.
- Taniyama, S. 1990. The Future Direction of Agricultural Engineering in Japan With relation to the 7th ICID Afro-Asian regional conference in Tokyo. *Journal of Irrigation Engineering and Rural Planning*, 1990 (19): 1–6.
- Tarzi, J. G. & Paeth, R. C. 1975. Genesis of a Mediterranean red and a white rendzina soil from Lebanon. *Soil Science*, 120: 272–277.
- Teixeira, W.G. & Martins, G.C. 2003. Soil physical characterization. In Lehmann *et al.*, eds. *Amazonia Dark Earths: Origin, properties, management*. pp. 271–286. Printed in Netherlands, Kluwer Academic Publishers.
- Tenuta, M., Gao, X., Flaten, D. N. & Amiro, B. D. 2016. Lower nitrous oxide emissions from anhydrous ammonia application prior to soil freezing in late fall than spring pre-plant application. *Journal of Environmental Quality*, 45(4): 1133–1143.
- Thiessen Martens, J.R. & Entz, M.H. 2001. Availability of late-season heat and water resources for relay and double cropping with winter wheat in prairie Canada. *Canadian Journal of Plant Science*, 81(2): 273–276.
- Thiessen Martens, J.R., Entz, M.H. & Hoepfner, J.W. 2005. Legume cover crops with winter cereals in southern Manitoba: Fertilizer replacement values for oat. *Canadian Journal of Plant Science*, 85: 645–648.
- Thiessen Martens, J.R., Entz, M.H. & Wonneck, M.D. 2015. Redesigning Canadian prairie cropping systems for profitability, sustainability, and resilience. *Canadian Journal of Plant Science*, 95(6): 1049–1072.
- Thiessen Martens, J.R., Hoepfner, J.W. & Entz, M.H. 2001. Legume cover crops with winter cereals in southern Manitoba: Establishment, productivity, and microclimate effects. *Agronomy Journal*, 93(5): 1086–1096.
- Thorup-Kristensen, K., Dresbøll, D. B. & Kristensen, H. L. 2012. Crop yield, root growth, and nutrient dynamics in a conventional and three organic cropping systems with different levels of external inputs and N re-cycling through fertility building crops. *European Journal of Agronomy*, 37(1): 66–82.
- Tilman, D., Balzer, C., Hill, J. & Befort, B.L. 2011. Global food demand and the sustainable intensification of agriculture. *Proceedings of the National Academy of Sciences*, 108(50): 20260–20264. <https://doi.org/10.1073/pnas.1116437108>
- Tisdall, J.M. & Oades, J.M. 1982. Organic matter and water-stable aggregates in soils. *Journal of Soil Science*, 33(2): 141–163. <https://doi.org/10.1111/j.1365-2389.1982.tb01755.x>
- Tong, Y. 2018. Influence of crop conversion on SOC, soil pH and soil erosion in mollisols region of Songnen Plain. China Agricultural University. PhD dissertation.
- Tong, Y., Liu, J., Li, X., Sun, J., Herzberger, A., Wei, D., Zhang, W., Dou, Z. & Zhang, F. 2017. Cropping system conversion led to organic carbon change in China's Mollisols regions. *Scientific Reports*, 7(1): 1–9.
- Twerdoff, D.A., Chanasyk, D.S., Mapfumo, E., Naeth, M.A. & Baron, V.S. 1999a. Impacts of forage grazing and cultivation on near surface relative compaction. *Canadian Journal of Soil Science*, 79: 465–471.
- Twerdoff, D.A., Chanasyk, D.S., Naeth, M.A. & Baron, V.S. 1999b. Soil water regimes under rotational grazing of annual and perennial forages. *Canadian Journal of Soil Science*, 79: 627–637.
- Unified Land Fund Classification Report. 2019. Ulaanbaatar. Department of Land Affairs and Geodesy under the Ministry of Construction and Urban Development. University of Saskatchewan. 2020. Soils of Saskatchewan. University of Saskatchewan. November 24, 2020. (also available at <https://soilsofsask.ca/soil-classification/chemozemic-soils.php>)



- USDA. 2014. *Keys to Soil Taxonomy*. Soil Survey Staff. Twelfth Edition.
- Vaisman, I., Entz, M.H., Flaten, D.N. & Gulden, R.H. 2011. Blade roller–green manure interactions on nitrogen dynamics, weeds, and organic wheat. *Agronomy Journal*, 103(3): 879–889
- Valle, S., Radic, S. & Casanova, M. 2015. Soils associated to three important grazing vegetal communities in South Patagonia. *Agrosur*, 43(2): 89–99.
- Van der Hammen, T., Pabón-Caicedo, J.D., Gutiérrez, H. & J.C. Alarcón. 2002. El cambio global y los ecosistemas de alta montaña de Colombia. In: C. Castaño Uribe, ed. *Páramos y ecosistemas altoandinos de Colombia en condición hotspot y global climatic tensor*, pp. 163–209. Bogotá, D.C., Colombia, IDEAM.
- Van der Merwe, G.M.E., Laker, M.C. & Bühmann, C. 2002a. Factors that govern the formation of melanic soils in South Africa. *Geoderma*, 107: 165–176.
- Van der Merwe, G.M., Laker, M.C. & Bühmann, C. 2002b. Clay mineral associations in melanic soils of South Africa. *Soil Research*, 40: 115–126.
- Van Hofwegen, G., Kuyper, T.W., Hoffland, E., Van den Broek, J.A. & Becx, G.A. 2009. Opening the black box: Deciphering carbon and nutrient flows in Terra Preta. In *Amazonian Dark Earths: Wim Sombroek's Vision*, pp. 393–409. Springer, Dordrecht.
- Van Poollen, H.W. & Lacey, J.R. 1979. Herbage Response to Grazing Systems and Stocking Intensities. *Journal of Range Management*, 32(4): 250–253.
- Venterea, R.T., Maharjan, B. & Dolan, M.S. 2011. Fertilizer source and tillage effects on yield-scaled nitrous oxide emissions in a corn cropping system. *Journal of Environmental Quality*, 40(5): 1521–1531.
- Veum, K.S., Kremer, R.J., Sudduth, K.A., Kitchen, N.R., Lerch, R.N., Baffaut, C., Stott, D.E., Karlen, D.L. & Sadler, E.J. 2015. Conservation effects on soil quality indicators in the Missouri Salt River Basin. *Journal of Soil and Water Conservation*, 70(4):232–246.
- Viglizzo, E., Frank, F., Carreño, L., Jobbagy, E., Pereyra, H., Clattz, J., Pincén, D. & Ricard, F. 2010. Ecological and environmental footprint of 50 years of agricultural expansion in Argentina. *Global Change Biology*, doi: 10.1111/j.1365–2486.2010.02293.x
- Villarreal, R., Lozano, L.A., Polich, N., Salazar, M.P., Barraco, M. & Soracco, C.G. 2022. Cover crops effects on soil hydraulic properties in two contrasting Mollisols of the Argentinean Pampas region. *Soil Science Society of America Journal*.
- Vinton, M.A. & Burke, I.C. 1995. Interactions between Individual Plant Species and Soil Nutrient Status in Shortgrass Steppe. *Ecology*, 76(4): 1116–1133.
- Voronov, S.I. & Mamytova, B.A. 1987. *Humus state and calculation of humus balance in soils of Chui valley of Kyrgyz SSR* (In Russian). In: *Scientific-applied questions of preservation and increase of fertility of soils of Kyrgyzstan*. Frunze, Kyrgyz SSR, USSR.
- Vyatkin, K.V., Zalavsky, Y.V., Bigun, O.N., Lebed, V.V., Sherstyuk, A.I., Plisko, I.V. & Nakisko, S.G. 2018. Creation of a national map of organic carbon reserves in the soils of Ukraine using digital methods of soil mapping (In Russian). *Soil Science and Agrochemistry*, 2: 5–17.
- Vyn, T.J. & Raimbault, B.A. 1993. Long-term effect of five tillage systems on corn response and soil structure. *Agronomy Journal*, 85: 1074–1079.
- Wagg, C., Bender, S.F., Widmer, F. & Van Der Heijden, M.G. 2014. Soil biodiversity and soil community composition determine ecosystem multifunctionality. *Proceedings of the National Academy of Sciences*, 111(14): 5266–5270. <https://doi.org/10.1073/pnas.1320054111>
- Waller, S. S. & Lewis, J. K. 1979. Occurrence of C3 and C4 photosynthetic pathways in North America grasses. *J. Range Manage.* 32: 12–28.
- Wang, J., Monger, C., Wang, X., Serena, M. & Leinauer, B. 2016. Carbon Sequestration in Response to Grassland–Shrubland–Turfgrass Conversions and a Test for Carbonate Biomineralization in Desert Soils, New Mexico, USA. *Soil Sci. Soc. Amer. J.* 80: 1591–1603.
- Wang, Z., Mao, D., Li, L., Jia, M., Dong, Z., Miao, Z., Ren, C. & Song, C. 2015. Quantifying changes in multiple ecosystem services during 1992–2012 in the Sanjiang Plain of China. *Science of the Total Environment*, 514: 119–130.
- Weesies, G.A., Schertz, D.L. & Kuenstler, W.F. 2017. Erosion: Agronomic practices. In R.B. Lal, ed. *Encyclopedia of Soil Science*. Boca Raton, USA, CRC Press.

- Wei, D., Qian, Y., Zhang, J., Wang, S., Chen, X., Zhang, X. & Li, W. 2008. Bacterial community structure and diversity in a black soil as affected by long-term fertilization. *Pedosphere*, 18(5): 582–592.
- Wen, Y., Kasielke, T., Li, H., Zhang, B. & Zepp, H. 2021. May agricultural terraces induce gully erosion? A case study from the black soil Region of northeast China. *Sci. Total Environ.* 750: 141715.
- Winckell, A., Zebrowski, C. & Delaune, M. 1991. Evolution du modèle Quaternaire et des formations superficielles dans les Andes de l'Équateur. *Géodynamique* 6: 97–117.
- Woods, W.I. & Mann, C.C. 2000. Earthmovers of the Amazon. *Science* 287: 786–789.
- World Bank. 2022. "War in the Region" Europe and Central Asia Economic Update (Spring). Washington, DC, World Bank. <https://www.worldbank.org/en/region/eca/publication/europe-and-central-asia-economic-update>
- Xie, H., Li, J., Zhu, P., Peng, C., Wang, J., He, H. & Zhang, X. 2014. Long-term manure amendments enhance neutral sugar accumulation in bulk soil and particulate organic matter in a Mollisol. *Soil Biology and Biochemistry*, 78: 45–53.
- Xinhua News Agency. 2022. Black soil protection law comes into act on August 1. Beijing, China. Cited 24 June 2022. [http://www.news.cn/legal/2022-06/24/c\\_1128773849.htm](http://www.news.cn/legal/2022-06/24/c_1128773849.htm)
- Xu, X., Pei, J., Xu, Y. & Wang, J. 2020. Soil organic carbon depletion in global Mollisols regions and restoration by management practices: a review. *Journal of Soils and Sediments*, 20(3): 1173–1181. <https://doi.org/10.1007/s11368-019-02557-3>
- Xu, X., Xu, Y., Chen, S., Xu, S. & Zhang, H. 2010. Soil loss and conservation in the black soil region of northeast China: a retrospective study. *Environmental science & policy*, 13(8): 793–800. <https://doi.org/10.1016/j.envsci.2010.07.004>
- Yagasaki, Y. & Shirato, Y. 2014. Assessment on the rates and potentials of soil organic carbon sequestration in agricultural lands in Japan using a process-based model and spatially explicit land-use change inventories – Part 1: Historical trend and validation based on nationwide soil monitoring. *Biogeosciences*, 11: 4429–4442.
- Yang, W., Guo, Y., Wang, X., Chen, C., Hu, Y., Cheng, L. & Gu, S. 2017. Temporal variations of soil microbial community under compost addition in black soil of northeast China. *Applied Soil Ecology*, 121: 214–222.
- Yang, Z., Guan, Y., Bello, A., Wu, Y., Ding, J., Wang, L. & Yang, W. 2020. Dynamics of ammonia oxidizers and denitrifiers in response to compost addition in black soil, northeast China. *PeerJ*, 8: e8844.
- Yao, Q., Liu, J., Yu, Z., Li, Y., Jin, J., Liu, X. & Wang, G. 2017. Three years of biochar amendment alters soil physiochemical properties and fungal community composition in a black soil of northeast China. *Soil Biology and Biochemistry*, 110: 56–67.
- Yatsuk, I.P. 2015. *Periodic report on the state of soils on agricultural lands of Ukraine according to the results of the 9th round (2006–2010) of agrochemical survey of lands* (In Ukrainian). Kyiv, DU «Derzhgruntokhorona».
- Yatsuk, I.P. 2018. Scientific bases of restoration of natural potential of agroecosystems of Ukraine (In Ukrainian). Institute of Agroecology and Nature Management of the National Academy of Agrarian Sciences of Ukraine, Kyiv. PhD dissertation.
- Yusufbekov, Y. 1968. *Improvement of Pastures and Hayfields of Pamir and Alay Valley* (In Russian). Dushanbe, Donish, Tajik SSR, USSR.
- Zanaga, D., Van De Kerchove, R., De Keersmaecker, W., Souverijns, N., Brockmann, C., Quast, R., Wevers J., Grosu, A., Vergnaud, S., Cartus, O., Santoro, M., Fritz, S., Georgieva, I., Lesiv, M., Carter, S., Herold, M., Li, L., Tsendbazar, N., Ramoino, F. & Arino, O. 2021. ESA WorldCover 10 m 2020 v100. *Zenodo*. Geneve, Switzerland.
- Zárate, M. 2003. Loess of southern South America. *Quaternary Science Review*, 22: 1987–2006.
- Zatula, V.I. & Zatula, N.I. 2020. Aridization of Ukraine's climate and its impact on agriculture (In Ukrainian). The Impact of Climate Change on Spatial Development of Earth's Territories: Implications and Solutions. 3rd International Scientific and Practical Conference, 121–124.
- Zentner, R. P., Lafond, G. P., Derksen, D. A., Nagy, C. N., Wall, D. D. & May, W. E. 2004. Effects of tillage method and crop rotation on non-renewable energy use efficiency for a thin Black Chernozem in the Canadian Prairies. *Soil and Tillage Research*, 77(2): 125–136.
- Zentner, R. P., Stephenson, J., Campbell, C., Bowren, K., Moulin, A. & Townley-Smith, L. 1990. Effects of rotation and fertilization on economics of crop production in the Black soil zone of north-central Saskatchewan. *Canadian Journal of Plant Science*, 70(3): 837–851.

- Zhang, J., Beusen, A.H.W., Van Apeldoorn, D.F., Mogollón, J.M., Yu, C. & Bouwman, A.F. 2017. Spatiotemporal dynamics of soil phosphorus and crop uptake in global cropland during the 20th century. *Biogeosciences*, 14(8): 2055–2068. <https://doi.org/10.5194/bg-14-2055-2017>
- Zhang, J., An, T. & Chi, F. 2019. Evolution over years of structural characteristics of humic acids in black soil as a function of various fertilization treatments. *J. Soils Sediments*, 19: 1959–1969.
- Zhang, S., Li, Q., Lü, Y., Zhang, X. & Liang, W. 2013. Contributions of soil biota to C sequestration varied with aggregate fractions under different tillage systems. *Soil Biology and Biochemistry*, 62: 147–156.
- Zhang, S., Wang, Y. & Shen, Q. 2018. Influence of straw amendment on soil physicochemical properties and crop yield on a consecutive mollisol slope in northeastern China. *Water*, 10(5): 559.
- Zhang, W., Gregory, A., Whalley, W.R., Ren, T. & Gao, W. 2021. Characteristics of soil organic matter within an erosional landscape under agriculture in northeast China: stock, source, and thermal stability. *Soil Tillage Res*, 209: 104927.
- Zhang, X.Y. & Liu X. B. 2020. Key Issues of Mollisols Research and Soil Erosion Control Strategies in China. *Bulletin of Soil and Water Conservation*, 40(4): 340–344.
- Zhang, Y., Hartemink, A.E., Huang, J. & Minasny, B. 2021b. Digital Soil Morphometrics. In: *Reference Module in Earth Systems and Environmental Sciences*. Elsevier. <https://doi.org/10.1016/B978-0-12-822974-3.00008-2>
- Zhang, Z., Zhang, X., Jhao, J., Zhang, X. & Liang, W. 2015. Tillage and rotation effects on community composition and metabolic footprints of soil nematodes in a black soil. *European Journal of Soil Biology*, 66: 40–48.
- Zhou, G., Zhou, X., He, Y., Shao, J., Hu, Z., Liu, R., Zhou, H. & Hosseinibai, S. 2017. Grazing intensity significantly affects belowground carbon and nitrogen cycling in grassland ecosystems: a meta-analysis. *Global Change Biology*, 23(3): 1167–1179.
- Zhou, J., Jiang, X., Zhou, B., Zhao, B., Ma, M., Guan, D. & Qin, J. 2016. Thirty four years of nitrogen fertilization decreases fungal diversity and alters fungal community composition in black soil in northeast China. *Soil Biology and Biochemistry*, 95: 135–143.
- Zimmermann, M., Meir, P., Silman, M.R., Fedders, A., Gibbon, A., Malhi, Y. & Zamora, F. 2010. No differences in soil carbon stocks across the tree line in the Peruvian Andes. *Ecosystems*, 13(1): 62–74.
- Zúñiga-Escobar, O., Peña-Salamanca, E.J., Torres-González, A.M., Cuero-Guepando, R. & Peña-Ospina, J.A. 2013. Assessment of the impact of anthropic activities on carbon storage in soils of high montane ecosystems in Colombia. *Agronomía Colombiana*, 31(1): 112–119.
- Zvomuya, F., Rosen C.J., Russelle M.P. & Gupta S.C. 2003. Nitrate leaching and nitrogen recovery following application of polyolefin-coated urea to potato. *Journal of Environmental Quality*, 32: 480–489

