

III International Symposium on Tropical and Subtropical Fruits

# 3<sup>rd</sup> INTERNATIONAL SYMPOSIUM ON TROPICAL AND SUBTROPICAL FRUITS

*FRUITS FOR A HEALTHY WORLD*  
*12 – 17 September 2004*  
*Fortaleza, CE, Brazil*

## PROGRAM AND ABSTRACTS

**Editors**

Fernando Antonio Souza de Aragão  
Ricardo Elesbão Alves

Program and abstracts...

2004

PC - 2004.00486



29680 - 2

**Fortaleza**  
**2004**

**P101**

**Effects of gibberellic acid, crop-set and girdling on the quality of bunches of table grape cv. 'Marroo Seedless' in the São Francisco river valley**

*Patrícia Coelho de Souza Leão, Davi José Silva, Emanuel Élder G. da Silva*

Embrapa Semi-Árido, C.P 23, CEP 56300-970, Petrolina-PE, Brazil, patricia@cpatsa.embrapa.br, davi@cpatsa.embrapa.br

Financial support: CNPq, Banco do Nordeste, Improcrop

The purpose of this research was to evaluate the effects of the use of gibberellic acid, Crop-Set bio-stimulant and girdling to improve yield and quality of the marketable bunches of the seedless grape cv. Marroo Seedless in the São Francisco River Valley, Northeast of Brazil. The trial was carried out throughout two cycles (2002-2003) in the Bebedouro Experimental Station, Embrapa Semi-Árido, Petrolina, PE. The trial was laid out in a randomized complete block design with three replicates, each replicate consisting of a four-tree plot. The treatments were: gibberellic acid in one dosage with two time applications (5 + 40 mg/L), Crop-Set in two doses 0.1 and 0.2 % and trunk girdling, isolated or combined to each other. The cv. Marroo Seedless didn't show any response to the treatments with gibberellic acid, Crop-Set and girdling for none of the variables evaluated in 2002, except for total titratable acidity. In 2003, the berry weight and length were bigger in the treatments where gibberellic acid was present. The berry weight in the gibberellic acid and girdling + gibberellic acid + Crop-Set 0.2 % treatments differed significantly of the untreated control. The best result for berry length (23.09 mm) was showed in the girdling + gibberellic acid + Crop-Set 0.2 % treatment. Total titratable acidity and total soluble solids/ total soluble solids ratio presented significant differences among the treatments.

**P102**

**Effects of gibberellic acid, crop-set and girdling on the quality of bunches of table grape cv. 'Perlette' in the São Francisco river valley**

*Patrícia Coelho de Souza Leão, Davi José Silva, Emanuel Élder G. da Silva*

Embrapa Semi-Árido, C.P 23, CEP 56300-970, Petrolina-PE, Brazil, patricia@cpatsa.embrapa.br, davi@cpatsa.embrapa.br

Financial support: CNPq, Banco do Nordeste, Improcrop

The aim of this study was to evaluate the effects of the gibberellic acid, bio-stimulant Crop-Set and girdling applied during bloom and post-bloom stage to improve yield and quality of the marketable bunches of the seedless grape cv. Perlette in the São Francisco River Valley, Northeast of Brazil. The trial was carried out during two growing seasons (2001-2002) in the Bebedouro Experimental Station, Embrapa Semi-Árido, Petrolina, PE. The trial was laid out in a randomized complete block design with three replicates, each replicate consisting of a four-tree plot. The treatments were: gibberellic acid in one dosis with three time applications (5 + 20 + 40 mg/L), Crop-Set in two dosis 0.1 and 0.2% and trunk girdling, isolated or combined to each other. It were evaluated the bunch and berry weight, berry length and diameter, rachis and pedicels weight, yield per plant, number of bunches per plant and chemical composition of the fruits. In the cycle of 2001, the largest berry weight and length were obtained in the treatment gibberellic acid + Crop Set 0.2%. In 2002, just the berry length and diameter answered to the application of the treatments, and the best results were showed with girdling + gibberellic acid. The rachis and pedicels of bunches treated with gibberellic acid became stronger, resulting in the largest rachis weight. There were no effects on other variables.