



Global Science Books

Breathing New Life into Science



Dynamic Soil, Dynamic Plant

Print ISSN: 1749-6500

Submit Manuscripts to:
dynsoildynplant@yahoo.com

Instructions for Authors
Description, Scope and Editorial Board
Online Journal
Current & Back Numbers (incl. Free Manuscript Samples)
Order form (Journal/PDFs)

[Journals Top Page](#) >

Copyright © Global Science Books 2005-2011. All Rights Reserved

Organic Matter of the Anthropogenic Dark Earths of Amazônia

Beáta Emöke Madari^{1*} • Tony Jarbas Ferreira Cunha² • Ricardo Soares³

¹ Embrapa Rice and Beans, Rod. GO 462, km 12, 75375-000 Santo Antônio de Goiás-GO, Brazil

² Embrapa Semi Arid, Rod. BR-42, km 152, 56302-970 Petrolina-PE, Brazil

³ Institute of Chemistry (GPEAA-GGQ-IQ), Federal Fluminense University (UFF), Outeiro de São João Batista, 24020-150 Niterói-RJ, Brazil

Corresponding author: * madari@cnpaf.embrapa.br

ABSTRACT

The anthropogenic dark earths (ADE, terra preta de índio or terra preta arqueológica, as are often referred to in Portuguese) feature unique properties that are interesting not only from a pedological point of view but have important agronomic and economic relevance. Among other properties, the organic matter (SOM) of these soils is considerably different from that of other mineral soils. Besides the quantity of SOM, its origin, composition and, consequently, function in the soil is different. Carbonized biomass (pyrogenic carbon) is a principal constituent of ADE that was, to our best knowledge, gradually incorporated into these soils, 4 to 10 thousand years ago, by native dwellers of the Amazon region. The carbon of pyrogenic origin was not only accumulated in ADE, but it also played important role in the transformation of organic matter and the humification process. As a result of this, the SOM of ADE is relatively stable, compared to the SOM of adjacent ferralsols, and at the same time it has chemical reactivity that is the major responsible for the higher cation exchange capacity and good buffer properties of these soils. These properties have very important environmental and agronomic relevance in the tropics due to organic carbon conservation in the soil and enhanced soil fertility. The ADE soils are excellent model for the development of modern soil fertility management practices. There are inconsistencies in the names of the earths, please check.

Keywords: physical and chemical characteristics of soil organic matter, pyrogenic carbon, soil color, soil fertility, soil organic matter fractions, soil structure, terra preta de índio

Abbreviations: ADE, anthropogenic dark earth soil; CEC, cation exchange capacity; DP/MAS NMR, direct polarization magic angle spinning nuclear magnetic resonance; EPR, electronic paramagnetic resonance; FOR, free organic radicals; HA, humic acid; SOM, soil organic matter; TM, terra mulata; TPI, terra preta de índio; VACP/MAS NMR, variable angle cross polarization magic angle spinning nuclear magnetic resonance

CONTENTS

INTRODUCTION	1
Characterization of the <i>archaeo-anthropedogenic</i> horizon of Terra Preta	2
The organic matter of the Terra Preta.....	3
The distribution of C among humic substances fractions in TPI.....	4
Composition and physical-chemical properties of the HA fraction of Terra Preta	4
The origin of the molecular structural characteristics of the HAs in Terra Preta.....	5
FINAL CONSIDERATIONS.....	6
REFERENCES.....	7

INTRODUCTION

The organic matter in mineral soils (SOM) rarely exceeds 8%, except some soils in specific climatic or terrain conditions, like Organosols that contain more than 20% organic matter, by definition (IUSS Working Group WRB 2006) to a minimum depth of 40 cm (Embrapa 2006). The distribution of organic matter in the soil is variable, in depth (vertically in the soil profile) and horizontally. This distribution of the C in the soil is dependent on various factors, like soil type, terrain morphology, soil cover or use, climatic conditions, predominant natural vegetation, soil management practices, among others. The quality of organic matter is also of great importance, it is source of nutrients and energy for microorganisms and conditions soil functions. Among these generally great emphasis is given to water retention, soil structure and its stability, porosity, retention and availability of macro- and micronutrients due to the fact that SOM levels are regulating cation exchange capacity (CEC), principally in soils under tropical climate whose mineral

matrix is dominated by low activity clay minerals (kaolinite, iron and aluminum oxides like gothite and hematite). In the soils of Brazil SOM can contribute to CEC even in 80%, considering the whole soil matrix. Hence, variations in the quantity and quality of SOM are able to have a great effect on soil properties and processes. SOM also has an important role in nutrient cycling whose dynamics is still little known. For this reason, SOM management with focus on its conservation and improving its quality is fundamental for the maintenance and sustainability of tropical agroecosystems.

A great proportion of the soils that are suitable for agricultural production at the Amazon Region are acid with low CEC and, consequently, low fertility and production potential, that is a limiting factor for environmental and economic sustainability.

In this same region can be found the so-called anthropogenic dark earths (Terra Preta de Índio - TPI or simply Terra Preta) that feature an anthropogenic A horizon, also called *archaeo-anthropedogenic* horizon, denomination proposed

by Kämpf *et al.* (2003). Different hypothesis were proposed as possible processes of formation of TPI but the actually most accepted is that these soils were formed by pre-historic populations (Woods and McCann 2001), however, there is still no consolidated opinion whether their activity was intentional (Neves *et al.* 2003). The areas where TPI occur now were places of pre-historic villages where great amounts of residues of vegetal and animal origin together with residues of cooking (fire) places (ash and charcoal). More recently, Birk *et al.* (2011), using biomarkers (5 β -cholestan-3 β -ol) in TPI compared to reference soils, showed that faeces were deposited on TPI and contributed to their genesis. This great quantity of organic matter contributed, most probably, to the formation of highly fertile soils with high levels of stable SOM of pyrogenic origin and also high levels of available phosphorus calcium, magnesium and zinc (Kern 1988; Kern and Kämpf 1989; Kern 1996; Costa and Kern 1999). Apparently, in TPI there are proper microecosystems that retain and make macro- and microelements available for a longer period of time than adjacent soils with no *archaeo-anthropedogenic* horizon, even under tropical climate and agricultural use.

Sombroek (1966) mentioned the so called Terra Mulata (TM) that are soils featuring high levels of P and carbon (C) but do not contain artifacts, however their formation was apparently associated to anthropogenic activity. It is believed that areas of TM had been utilized for agriculture and TPI, where high quantity of artifacts can be found, had been the actual day to day living habitat, the village. According to McCann *et al.* (2001) TMs have similar levels of organic C compared to TPI but lower levels of available P and Ca. Examples of TM around the Negro and Urubu Rivers had similar productivities than TPIs of the same regions (Lehmann *et al.* 2003). Because TM was also formed by human activity and has similar characteristics to TPI, Kämpf *et al.* (2003) suggested the use of the term *archaeo-anthrosol* for all Amazonian anthropogenic soils.

The TPI is not a homogeneous group of soils regarding fertility and production potential. There might be great differences even between TPIs of a region and there is a great variability within one TPI site (Lehmann *et al.* 2003). Despite these differences some important soil fertility properties are shared by most TPI: high organic C levels, SOM with particular physical-chemical properties and high levels of available P, Ca and micronutrients, compared to adjacent soils like Oxisols among others.

One of the responsible factors for the different behaviours of these anthropogenic soils is the higher organic C content and principally the different quality of this organic C that has pyrogenic origins. The objective of this paper is to describe the organic matrix of TPI in comparison to that in adjacent soils that have no *archaeo-anthropedogenic* horizon, and to investigate the effect of the differences in SOM on the fertility properties of these soils.

Characterization of the *archaeo-anthropedogenic* horizon of Terra Preta

The characterization of soil horizons that had undergone anthropogenic influence comparing with similar horizons of similar but not anthropogenic soils is important to observe possible functions of SOM in TPI. Hence, characteristics that are different between TPI and adjacent soils are described below.

Soil color. The color of TPI in the *archaeo-anthropedogenic* horizon is, in general, as the name of these soils indicate, is black to very dark brownish grey (5YR 2.5/1; 7.5YR 2/0 to 3/1; 10YR 2/0 to 3/2) (Kämpf and Kern 2005), different from the color of adjacent non-anthropogenic soils whose color is generally determined by the color of the clay minerals of the soil matrix. This very characteristic property of TPI is due to the quantity (Kern and Kämpf 1989) and quality of the SOM present in these soils. Data from 56 soil profiles showed that while the mean organic C concentra-

tion in the surface 20 cm of TPIs was 30 g kg⁻¹ (CV=66%), in non-anthropogenic soils of the Amazon it was 17 g kg⁻¹ (N=47, CV=58%) at the same depth (Madari *et al.* 2003). In a general manner, the correlation between soil color and SOM concentration is low ($r^2 = 0.31$; Schulze *et al.* 1993), however, there is a tendency for having darker color in SOM rich soils. In TPI the distribution between the chemical SOM fractions is different from that in adjacent soils. In TPI the more recalcitrant fractions, that are also more stable and persistent in the soil (humins and HAs (HAs)), predominate in relation to the more soluble fractions (fulvic acids). The stability of organic matter is related to some molecular structures that are also responsible for the stability and persistence of SOM in TPI for a longer period of time as these are less sensitive to microbial decomposition. These molecular structures are, for example, hetero-polycondensed, amorphous structures with high concentration of double conjugated links (Kumada 1965) and poly-aromatic structures. The more stable fractions of SOM, which predominate in TPI, have darker colors than the less stable ones (fulvic acids and low molecular weight organic acids). As mentioned before, the anthropogenic dark earths also have elevated concentrations of charred materials (Glaser *et al.* 2000). The presence of these accentuate dark coloration as well (Schmidt and Noack 2000).

Soil structure. Chemical aspects of sustainable soil fertility management are very important but soil management for adequate physical status is equally relevant. The physical management of soils is highly relevant to soil fertility, agricultural productivity and the quality of the environment (Lal 2000). Soils under tropical climate feature, in general, good structure under natural vegetation. Examples are the kaolinitic clayey Ferralsols. However, under inadequate management, like frequent plowing, the originally well developed structure rapidly deteriorates, leading to crusting, compaction and subsequent erosion. The anthropogenic dark earths also feature excellent physical properties. The bulk density of TPI in the superficial soil layers of the *archaeo-anthropedogenic* horizon is generally low (1.08 ± 0.23 g cm⁻³, N = 10), the porosity ($58.83 \pm 9.07\%$, N = 10) and water retention capacity is high (Teixeira and Martins 2003).

An interesting question is the effect of the pyrogenic carbon on soil physical properties. On one hand char, due to its porosity and, consequently, to its high surface area, can significantly improve water retention, especially in sandy soils. On the other hand, the aromatic structure of char that has hydrophobic characteristics may reduce water penetration into the pore space of soil aggregates, that way increasing aggregate stability (Glaser *et al.* 2002). Probably, due to the mixture of these characteristics the effect of charcoal on the physical properties of medium textured soils in most of the cases is not detectable and in clayey textured soils is negative (Tryon 1948). Almendros *et al.* (1992) described the evolution pathway of humic substances as affected by fire that could significantly contribute to the formation of soils with hydrophobic characteristics (repellent to water). Hence, the high concentration of pyrogenic C in association with its physico-chemical properties, such as high stability and reactivity at the same time (Zech *et al.* 1990; Golchin *et al.* 1997; Schmidt *et al.* 1999; Poirier *et al.* 2000), in fact may have contributed to the improvement of TPI properties, not only the chemical but physical ones as well.

Soil fertility. Compared to non-anthropogenic soils TPI, in general, feature clear tendency to have higher concentrations of organic C, Ca and Mg, higher pH and base saturation and smaller concentrations or non of exchangeable Al. These properties of TPI are responsible for the better fertility and production potential of TPI. Hence, however there is some overlapping between the fertility properties of TPI and adjacent soils, TPI generally feature higher fertility (Madari *et al.* 2003). The capacity of the soil to adsorb nutrients (Ca, Mg, base saturation), diminish toxicity of Al

and the stabilization of pH are processes that are fundamentally controlled by organic matter. Pabst (1992), investigating the principal differences between TPI and regular Ferralsols, also identified that those attributes that are associated with SOM (pH, organic C concentration, "humus structure" – stability to microbial decomposition, N, base saturation, CEC and available P) are the ones that make the most difference. While SOM concentration in TPI is a relevant factor, its composition and characteristics also contribute to the higher fertility of TPI.

The organic matter of the Terra Preta

Processes of stabilization and mobilization affect the accumulation and decomposition of SOM and the equilibrium between these processes are of extreme importance for the formation and maintenance of the fertility of soils under tropical climatic conditions. These processes regulate recalcitrance and reactivity, or in other words, the capacity of SOM to interact with other materials and to be protected against chemical or biological decomposition (Sollins *et al.* 1996). The higher accumulation of organic C in TPI, compared to adjacent soils, indicate that at least a part of its organic matter has characteristics that give stability or little availability to the C associated to it. On the other hand, fertility data on TPI indicate that the SOM in these soils also has elevated chemical reactivity compared to the soils that have no *archaeo-anthropogenic* horizon. In the foregoing paragraphs the characteristics of the different SOM fractions will be described as they are in TPI.

Chars. Chars are generally reported as chemically inert and hydrophobic materials and, due to their high porosity, have large surface area (200-400 m² g⁻¹, Kishimoto and Sugiura 1985). However, the reactivity, hydrophobic characteristics and structure of chars depend greatly on the characteristics of the material of which they are originated, on the conditions of their formation like temperature and duration of carbonization, humidity of the material, availability of oxygen, among others. However chars can be considered stable in the soil, compared to other forms of organic C, they also undergo biodegradation and transformation. Bird *et al.* (1999) suggested that pyrogenic C can be degraded significantly within the timescale of decades or centuries in soil under tropical climate. The half-life of carbonized particles greater than 2 mm was estimated to be shorter than 50 years and of particles larger than 2 mm shorter than 100 years. The degradation of char makes it possible that this material also takes part in the global C cycle. If this is not the case, that would implicate it in the accumulation of C in this form on the surface of the Earth in less than a thousand centuries (Kuhlbusch and Crutzen 1995 in Glaser *et al.* 2002). The biodegradation of chars in soil is a relatively slow process that results in the mobilization of C and in the alteration of the surface properties of chars, hence increasing the concentration of chemically reactive sites.

The quantity of char that is produced during carbonization depends on the originating material and duration and temperature of carbonization. Fearnside *et al.* (1999) reported that char formation during natural fire of a secondary Forest in the Amazon was between 1.6 and 1.8% of the original biomass. Fire in primary Forest resulted in 3.5-4.7% char (Fearnside *et al.* 1993; Graça 1999). In controlled conditions of carbonization, between 300 and 500°C in laboratory or in different carbonizing equipments, char production resulted in about 79.6%. Trompowsky *et al.* (2005), in controlled conditions, demonstrated that two eucalypt species, *E. saligna* and *E. grandis*, had a maximum production of char, 65%, at 300°C carbonization temperature. In that same study, it was also observed that the O/C ratio was higher (0.3) at 300°C and at 500°C it was reduced to 0.06, resulting in a material that contained 90% C in its total mass. The presence of oxygen in higher proportion may indicate higher reactivity of the analyzed char.

Hamer *et al.* (2004) investigated the "priming" effect of

pyrogenic C and the mineralization of glucose. According to these authors the mineralization of the pyrogenic C was stimulated by the addition of glucose to the soil and, vice-versa, the mineralization of glucose was stimulated by the presence of the pyrogenic C. The authors supposed that the effect of the carbonized material occurred because of its high superficial area that served as medium for microbial growth and consequently activity rather than because of higher availability of C since pyrogenic C, being highly aromatic, is not readily available as energy source. Hence, charcoal, however is relatively inert and highly stable in the soil, depending on the conditions of its formation and on the transformations it undergoes, it may have the capacity to contribute to enhance the chemical, physical and consequently biological properties of soils.

The light fraction of SOM. There are only a few studies on the contribution of pyrogenic C to the light fraction of SOM in TPI. This fraction is separated on the basis of the density of soil components after their total dispersion. Glaser *et al.* (2000) determined pyrogenic C in three density fractions: $d < 2.0 \text{ g cm}^{-3}$ (light fraction), $d = 2.0\text{-}2.4 \text{ g cm}^{-3}$ (medium fraction) and $d > 2.4 \text{ g cm}^{-3}$ (mineral matrix of the soil to which organic C is associated by chemical and physical mechanisms) in TPI and adjacent non-anthropogenic Ferralsols. The observed that the contribution of pyrogenic C to the soil mass was higher in TPI in each separated fraction, being the largest in the medium fraction. In the Ferralsols the distribution of the pyrogenic C between the fractions was more balanced.

It is possible that the quantities of pyrogenic C were super estimated by the determination of benzenecarboxylic acid with gas chromatography after partial oxidation of the pyrogenic C with nitric acid (Derenne and Largeau 2001). Nevertheless, their results showed the expressed presence of pyrogenic C in the light fraction of SOM of TPIs indicating that the largest proportion of pyrogenic C in TPI is of particulate nature and part of it is physically not protected by soil structure, however, it can be also found in physically or chemically (organo-mineral complexes) protected forms.

The light fraction is one of the most labile ones with a high pre-disposition to microbial decomposition, especially the free light fraction that is not physically protected by soil structure. In relation to chars, and specifically in the case of TPI, there is no exact information about physical protection in its particulate form. There are indications, as mentioned before, that it occurs in protected and non-protected forms as well. The pyrogenic C in particulate form, however, has a much longer half-life in the soil than non-pyrogenic particulate SOM, if not lost by erosion. The fact that it is more preserved even in the particulate form is another indication of its higher recalcitrance. However the forms of protection of pyrogenic C and its chemical structure is not fully understood, it can be confirmed, with great certainty, that its higher stability is due to its molecular structure and morphology that varies according to its origin (Poirier *et al.* 2000). Pyrogenic C that was formed from lingo-celulosic materials tends to have micro-structure with basic structural units arranged in a little oriented manner. Formed from materials that have low oxygen concentration like plant waxes or hydrocarbonates as fossil fuels they feature micro-structure where the basic structural units are arranged in a concentric manner (Poirier *et al.* 2000; Derenne and Largeau 2001). The basic structural units, of nanometric scale, are made of condensed polyaromatic structures with low H:C ratio that are highly resistant to biotic and abiotic degradation.

More recently, Liang *et al.* (2010) found that the presence of pyrogenic C (black C) led to more rapid incorporation of freshly added organic matter into aggregate and organo-mineral fractions, the physically and chemically protected organic matter, respectively.

Humic substances. The study of the humified organic matter has recently gain new importance due to its role in

Table 1 Variability and distribution of the fulvic acid, HA and humin fractions within a Terra Preta site (Fazenda Roda de Fogo, municipality of Rio Preto da Eva, Amazonas State) (Soares 2007).

Depth, cm	Fulvic acids			HAs			Humin		
	C, mg g ⁻¹	V, %	CV	C, mg g ⁻¹	V, %	CV	C, mg g ⁻¹	V, %	CV
0-5	11.47	5043	0.62	5.31	759	0.52	4.56	176	0.29
5-10	12.04	4981	0.59	6.54	938	0.47	5.69	193	0.24
10-20	10.82	3537	0.55	9.57	1135	0.35	4.73	19	0.09
20-40	10.00	2390	0.49	8.39	580	0.29	3.48	13	0.10
40-60	7.85	3519	0.76	4.81	1127	0.70	1.95	21	0.24
60-80	7.74	3636	0.78	2.82	1370	1.31	2.15	71	0.39
80-100	7.55	4347	0.87	1.42	404	1.41	2.39	99	0.42

V: Variance, CV: Covariance

Table 2 Correlation among fertility attributes, sum of bases (SB) and cation exchange capacity (CEC) of a Terra Preta (Fazenda Hatahara, municipality of Iraduba, Amazonas State) and soil organic matter fractions (Cunha 2005).

Soil fertility attribute	AF	AH	HUM	Total of fractions (AF+AH+HUM)	Corg	Ctot	Cres
SB	R	-0.01 ^{ns}	0.66**	0.29 ^{ns}	0.45 ^{ns}	0.72**	0.10 ^{ns}
CEC	R	0.28 ^{ns}	0.85*	0.56 ^{ns}	0.69**	0.81*	-0.18 ^{ns}

AF: fulvic acids, AH: HAs, HUM: humin, Total of fractions: sum of C in the three humic fractions (chemical fractions), Corg: organic C measured by the modified Walkley-Black method (Embrapa 1997), Ctot: total carbon measured by dry combustion, Cres: Ctot-Corg, SB: sum of bases, CEC: cation exchange capacity, r: coefficient of correlation (Pearson's), ns: not significant at p<0.01, *: significant at p<0.0001, **: significant at p<0.005, ***: significant at p<0.01.

environmental protection. The disequilibrium caused by the elimination of the natural vegetation and the introduction and inappropriate practice of agricultural activities promoted soil degradation and diminished its capacity to sustain life in long term (Mielniczuk *et al.* 2003). The humic substances are among the most important SOM fractions, having fundamental role in the sustainability of soil functions. The largest proportion of humic substances represent a pool of SOM that has small transformation or degradation rate or is recalcitrant and makes 70-80% SOM (Stevenson 1994). These have important role in C accumulation and to retain and make available plant nutrients.

For these reasons the distribution of C among the fractions of humic substances and their molecular characteristics in TPI will be discussed in detail further in the text, always compared with non-anthropogenic soils.

The distribution of C among humic substances fractions in TPI

The most abundant humic fraction is the humin, both in TPI and non-anthropogenic soils under tropical climate. The TPI, however, has a larger proportion of C in the HA fraction (Fig. 1) than in the fulvic acid fraction, in this being different from non-anthropogenic soils. Hence, in TPI, those humic fractions (HAs and humins) that are conventionally considered to be chemically more stable in function of the presence of aromatic groups and benzene rings, higher degree of polymerization and lower susceptibility to microbial decomposition predominate. In Fig. 1 it is also presented that TPI has a great proportion of C (~50% in the whole soil profile) that is not oxidizable by the Walkley-Black method (Nelson and Sommers 1996). This C is chemically stable and consequently has longer persistence in the soil.

The variability of the distribution of the three humic fractions (fulvic acids, HAs and humin) of the humic substances within one TPI site is however great (Table 1), and the fulvic acid fraction has the largest and the humin the smallest variability of all. This variability, similarly to those observed in the case of the fertility properties of TPI, is probably due to the proper anthropogenic origin of these soils that is the cause of the uneven and irregular deposition of the carbonized material within these sites.

Composition and physical-chemical properties of the HA fraction of Terra Preta

In the profile of TPI (Fig. 1) total and organic C showed significant correlations to CEC and with the sum of basic cations (Table 2). Among the chemical fractions of SOM




Fig. 1 Anthropogenic dark earth (Terra Preta) profile at Hatahara Farm, Iraduba, Amazonas State. Concentration of carbon in the chemical fractions of soil organic matter (n = 3). Error bars indicate standard deviation. FA: fulvic acids, HA: HAs, Total of fractions: HA+FA+Humin.


only the HA fraction correlated with these fertility parameters. These data demonstrate the importance of the HA fraction in conditioning some soil fertility attributes in the studied TPIs. Also suggest that to be able to better understand the effect of SOM on the fertility of TPI it is important to investigate, among other aspects, the molecular composition and physical-chemical characteristics of the HA fraction.

To demonstrate the principal differences between the HAs of TPI and non-anthropogenic soils, the respective physical-chemical and molecular properties, that define their function in the soil, will be discussed in the following. The properties that determine the function and transformations of the HAs are their stability, reactivity and susceptibility to mineralization. The stability of HAs in relation to decomposition and leaching is important for the accumulation and persistence of organic C in the soil. The reactivity of HAs is responsible for the interaction of these molecules with other soil components via the formation of organo-metallic complexes whose linkages between the mineral and organic matrix of the soil favor organic C accumulation and stabilization. Via adsorption and desorption reactions occurs the retention and availability of plant nutrients. Via mineralization of HAs, the liberation of nutrients occurs and also the diminution of organic C concentration in the soil. The stability of HAs is determined, principally, by their aro-

Table 3 Indices that describe composition and physical-chemical properties of HAs in Terra Preta (TPI) and adjacent non-anthropogenic soils (Non-TPI) of the Amazon (Cunha 2005).

Solos	TGI	IA	Total acidity, cmol _c kg ⁻¹	Carboxylic acidity, cmol _c kg ⁻¹	O:H	E ₄ :E ₆	FOR, Spin g ⁻¹	C:N	A ₄ :A ₁	I ₄₈₅ :I ₄₀₀
TPI	3.2a	0.86a	673a	445a	7.42a	4.2b	5.75a	12.2a	3.067a	2.109a
Non-TPI	2.3b	0.77b	575b	320b	4.76b	6.0a	2.11b	8.8b	1.713b	1.510b


TPI: Terra Preta under natural forest and cultivated (N=18), Non-TPI: Non-anthropogenic soil, adjacent to TPI under forest (N=4). The numbers followed by different letters within a column are different at $p \leq 0.05$ by Tukey's test. TGI: Thermogravimetric index, IA: Index of aromaticity obtained by infrared spectroscopy (DRIFT), Total acidity: Concentration of all acidic functional groups on molecular surfaces, Carboxylic acidity: Concentration of carboxylic groups, O:H: Atomic ratio Oxygen: Hydrogen, E₄:E₆: Obtained by UV-Visible spectroscopy, indicate condensed aromatic rings, FOR: Obtained by electron paramagnetic resonance spectroscopy (EPR), used to describe humification degree, C:N: atomic ratio Carbon: Nitrogen, A₄:A₁ e I₄₈₅:I₄₀₀: Indices obtained by fluorescence spectroscopy, describe degree of humification of humic substances.

**Fig. 2** Properties that determine function and transformation of humic substances in soil.

matic and hydrophobic structures. Their chemical reactivity is determined by the concentration of acidic functional groups on the molecular surfaces. The susceptibility of HAs to mineralization depends, greatly, on the quantity of hydrophilic aliphatic structures in their molecular associations (Fig. 2).

The principal differences between the HAs of TPI and non-anthropogenic soils, is presented in Tables 3 and 4. The methods used to obtain the indices presented in the respective tables are listed in Table 5 (Cunha 2005). The localization of the respective TPIs and adjacent non-anthropogenic soils is described in Fig. 3.

The humification process results in ever-increasing molecules or molecular associations, increasing the degree of condensation and concentration of free organic radicals. The different humification indices that were obtained via degradative, basically chemical, and non-degradative physical, principally spectroscopic techniques, methods indicated higher humification degree and higher concentration of aromatic poly-condensed structures in HAs extracted from TPIs. The HAs of TPI are more resistant to thermo-degradation, compared to the HAs of adjacent soils, that is expressed by the thermogravimetric index (TGI). This property indicates that the HAs of TPI have greater proportion of stable structural elements such as condensed aromatic nuclei. The index of aromaticity (IA), obtained by infrared spectroscopy, and the distribution of C among the structural elements of humic substances determined by ¹³C nuclear magnetic resonance spectroscopy (Tables 3, 4) confirmed a greater proportion of aromatic structures in the HAs of TPI. The E₄/E₆ ratio that has an inverse relation to the condensed aromatic rings was smaller in TPI HAs than in those of adjacent soils. The concentration of the free organic radicals (FOR), data obtained by electronic paramagnetic resonance (EPR), is also frequently utilized to estimate the degree of humification. The higher the concentration of FOR in the HA fraction, the higher is the degree of humification, and HAs of TPI showed a higher concentration of FOR compared to adjacent soils. Additionally, the humification indices obtained by fluorescence spectroscopy (A₄:A₁ and I₄₈₅:I₄₀₀) also indicated higher degree of humification in TPI. The C:N ratio indicate the degree of incorporation of nitro-

**Fig. 3** Localization (▲) of the Amazonian anthropogenic dark earths and adjacent soils described in Table 5.

gen, in this case, into the structure of humic substances and also the degree of humification. High C:N ratios normally signify low levels of N incorporation and low degree of humification. In the case of TPI, however, the C:N ratio of the HAs falls within the range considered "normal" for stable organic compounds in soils that is between 10 and 12 (Stevenson 1994). It means little secondary mineralization of SOM (Rossel *et al.* 1989). The low (8.8) C:N ratio obtained for the non-anthropogenic soils suggests lower biological stability and the possible presence of secondary mineralization of SOM. The higher total acidity and the presence of carboxylic radicals together with the higher O:H ratio of the HAs of TPI indicate the higher concentration of acidic functional groups that are responsible for the chemical reactivity and, consequently, for the interactions of HAs with other soil components.

Hence, the HAs of TPI are more stable and at the same time have higher chemical reactivity than those in adjacent, non-anthropogenic, soils. Through their physical-chemical and molecular structural properties the humic substances formed in TPI facilitate organic C accumulation and higher fertility and production potential, so much desired in soils under tropical climate where soils without this organic matter are often obstacles to sustainable agricultural production.

The origin of the molecular structural characteristics of the HAs in Terra Preta

The edaphic and climatic conditions that control humification processes are similar for TPI and adjacent soils. The difference is in the higher concentration of char or pyrogenic C in TPIs. The pyrogenic C is considered inert and little susceptible to decomposition but, as mentioned before, it also undergoes some degradation and transformation in the soil. Trompowsky *et al.* (2005), using ¹³C direct polarization magic angle spinning nuclear magnetic resonance (DP/MAS NMR) on HAs extracted from charcoal made from eucalypt wood showed that C was present in two prin-

Table 4 Distribution of ^{13}C among carbon containing chemical structures of the humic acids of Terra Preta and adjacent non-anthropogenic soils of the Amazon obtained by nuclear magnetic resonance spectroscopy (^{13}C VACP/MAS NMR).

Soils	Alkyl	Methoxyl	O-Alkyl	Aromatic	Carboxyl	Carbonyl	Aliphatic	Aromatic
TPI	23 b	10 b	18 b	29 b	11 a	7 a	64 b	36 b
Non-TPI	29 a	12 a	26 a	22 a	11 a	2 b	75 a	25 a

TPI: Terra Preta under natural forest and cultivated (N=18); Non-TPI: Non-anthropogenic soil, adjacent to TPI under forest (N=4); The numbers followed by different letters within a column are different at $P \leq 0.05$ by Tukey's test. Source: Cunha (2005).

Table 5 List of techniques and methods used to obtain humification indices and reactivity, quantify and describing the properties of soil organic matter, of humic substances and, specifically, of the HAs (Cunha 2005), discussed in the text.

Technique/Method	Purpose/Index/Property	Specification
Modified Walkley-Black method Embrapa (1997)	Quantification of organic C in soil	Oxidation of SOM by potassium dichromate
Benites <i>et al.</i> (2003) modified from Yeomans and Bremner (1988)	Quantification of humic substances	Oxidation of SOM by potassium dichromate
Method recommended by IHSS (International Humic Substances Society) (Swift 1996)	Extraction and separation of humic substances	Operational separation of humic substances based on solubility in aqueous solutions of acid and base
Thermogravimetry	Thermogravimetric index (TGI) (Benites <i>et al.</i> 2005b)	Ratio of the weight loss of HAs in the temperature ($^{\circ}\text{C}$) ranges (350-650) and (105-305)
Fourier Transform Infrared Spectroscopy with diffuse reflectance (DRIFT)	Index of aromaticity (IA) (Freixo <i>et al.</i> 2002)	Ratio of the absorbance of the bands (3057-3055 cm^{-1}) and (2934-2928 cm^{-1})
Ba(OH) ₂	Total acidity and carboxylic acidity (Schnitzer and Gupta 1965)	Total acidity = (titrated volume for blank – titrated volume for sample) × N of HCl × 1000/mass of the sample (mg) Carboxylic acidity = (titrated volume for sample – titrated volume for blank) × N of NaOH × 1000/mass of the sample (mg)
Elemental analysis by high temperature combustion (Dumas method; Nelson and Sommer 1996)	O:H and C:N ratios	
UV-Visible Spectroscopy	E ₄ /E ₆ ratio (Kononova 1966)	Ratio of absorbance at 465 and 665 nm
Electron paramagnetic Resonance Spectroscopy (EPR)	Free Organic Radicals (FOR)	Obtained by the integration of the signal of the first derivative of the free radical, compared to a standard (KCl) of known concentration of spins (3×10^{15} spins g^{-1}), using synthetic rubi as secondary standard (Singer 1959; Martin-Neto <i>et al.</i> 1994)
Fluorescence spectroscopy	A ₄ :A ₁ e I ₄₈₅ :I ₄₀₀ indices	A ₄ :A ₁ : ratio of absorbance at (665-585 nm) and (425-345 nm) (Zsolnay <i>et al.</i> 1999). I ₄₈₅ :I ₄₀₀ : ratio of absorbance at 480 nm and 400 nm (Kalbitz <i>et al.</i> 1999)
Nuclear Magnetic Resonance Spectroscopy with Cross Polarization, Variable Amplitude Magic Angle Spinning (^{13}C VACP/MAS NMR)	Table 3 (Knicker <i>et al.</i> 2005a, 2005b; Dick <i>et al.</i> 2005)	

principal structures: aromatic (aryl-C, condensed and deficient in H, at 130 ppm) and carboxylic (-COO⁻, at 169 ppm) groups and a significant part of the carboxylic groups was chemically linked to aromatic structures. As the condensed aromatic structures are highly resistant to degradation the carboxylic groups associated to them are also expected to be resistant and persistent once in the soil.

Benites *et al.* (2005b) examined, using ^{13}C variable angle cross polarization magic angle spinning nuclear magnetic resonance (VACP/MAS NMR), HAs extracted from a soils that were naturally (no anthropogenic effect) enriched in pyrogenic C. The principal structures identified in these HAs were, similarly to the spectra of HAs extracted from charcoals, C in aromatic structures (130 ppm) and in carboxylic groups (169 ppm) probably linked to the aromatic components. In this material the contribution of C in alkyl structures (0-45 ppm) was, however, greater than in the HA extracted from charcoals.

Novotny *et al.* (2007) compared the spectra, also using ^{13}C NMR spectroscopy, of HAs extracted from TPI and from an adjacent Ferralsol. The HA fraction of TPI presented two principal peaks, at 130 ppm, correspondent to aryl-C and at 169 ppm, corresponding to C in carboxylic groups, similarly to the HAs extracted from eucalypt charcoals and from the soil naturally enriched in pyrogenic C. There was a smaller contribution of aliphatic structures such as C in O-alkyl structures of carbohydrates (60-90 ppm), C in methoxyl groups of lignin (45-60 ppm) and alkyl-C (0-45 ppm). The HAs of the non-anthropogenic soil (Ferralsol) consisted of, in basis of the spectrum, significant proportions of C in aliphatic structures, specifically at 0-45 ppm (alkyl-C) and at 45-60 ppm (methoxyl). The presence of O-aryl C (140-

160 ppm), together with methoxyl-C, indicate higher contribution of lignin to the structure of the HAs in the non-anthropogenic soil. In the anthropogenic dark earths the peak of the O-aryl C was smaller and the contribution of aryl-C (poly-condensed aromatic structures) greater. Additionally, the carboxylic groups that were present in TPI HAs, in great proportion were associated with poly-aromatic condensed structures. Hence, the HAs of TPI, besides showing a certain degree of recalcitrance, also had C in functional groups that represent chemical reactivity.

Using data obtained by ^{13}C VACP/MAS NMR and multivariate analysis, Novotny *et al.* (2007) verified that the HAs in TPI had C in their structures that are of pyrogenic origin. This signifies that the HAs of TPI are in fact originated, at least partially, from the pyrogenic C (charcoal) that was incorporated, accumulated and transforming in the soil during long periods of time. This way, charcoal directly contributed to the enhancing of the chemical and physical properties and, consequently, for the formation of more fertile and productive soils.

FINAL CONSIDERATIONS

The organic matter of the Amazonian anthropogenic dark earths has a key role in the formation of the chemical, physical and biological properties of these soils. Because of its high concentration, and principally due to its chemical-physical properties, TPIs feature unique fertility properties and so agricultural potential in humid tropical environments where soils are frequently obstacles of sustainable production.

The anthropogenic dark earths are rich in humified and

stable organic matter that, at the same time gives higher fertility to the soil, making viable the existence of an environment that is favorable to cation exchange reactions, with organic and inorganic colloids, and the complexation of metallic ions, among others. For having stable organic matter, in the TPI predominate processes that favor organic C accumulation. However, the effect of the pyrogenic C and of the fractions of SOM in TPI that inherited some properties of the pyrogenic C on the biological processes in the soil and on the proper accumulation of C is still basically unknown. The mechanisms by which pyrogenic C affects plant growth are also still mainly unknown.

The anthropogenic dark earths of the Amazon, due to their favorable properties for agricultural use, are usually utilized by local communities. The promotion of the agricultural utilization of TPIs, these being national patrimony of Brazil (IPHAN 1988), is not desirable. However, using scientific investigation, important information can be obtained about the functioning of these soils that could be used, as model (Novotny *et al.* 2009; Renard *et al.* in press), in the development of sustainable soil fertility management systems for TPIs already under utilization and for the development of new soil management systems for tropical soils for the benefit of, principally, small producers, family farms and local communities of the Amazon and other tropical regions.

REFERENCES

- Almendros G, González-Vila FJ, Martín F, Fründ R, Lüdemann HD (1992) Solid state NMR studies of fire-induced changes in the structure of humic substances. *The Science of the Total Environment* **117-118**, 63-74
- Beckman EH, Souza LF, Coelho MR, Martins GC, Madari BE (2005) Distribuição vertical e horizontal de substâncias húmicas em terra preta de índio, município Rio Preto da Éva, Amazonas. In: *Anais do XXX Congresso Brasileiro de Ciência do Solo*, SBSCS, Recife-PB (CD-Rom)
- Benites VM, Madari BE, Machado PLOA (2005a) Matéria orgânica do solo. In: Wadt PGS (Ed) *Manejo do Solo e Recomendação de Adubação para o Estado do Acre*, Embrapa Acre, Rio Branco-AC, pp 93-120
- Benites VM, Mendonça ES, Schaefer CEGR, Novotny EH, Reis EL, Ker JC (2005b) Properties of black soil humic acids from high altitude rocky complexes in Brazil. *Geoderma* **127**, 104-113
- Benites VM, Madari BE, Machado PLOA (2003) Extração e fracionamento quantitativo de substâncias húmicas do solo: Um procedimento simplificado de baixo custo. Embrapa Solos, Rio de Janeiro-RJ, pp 1-7 (Comunicado Técnico, 16)
- Bird MI, Moyo C, Veenendaal EM, Lloyd J, Frost P (1999) Stability of elemental carbon in savanna soil. *Global Biogeochemical Cycles* **13**, 923-932
- Birk JJ, Teixeira WG, Neves EG, Glaser B (2011) Faeces deposition on Amazonian Anthrosols as assessed from 5b-stanols. *Journal of Archaeological Science* **38**, 1209-1220
- Costa ML, Kern DC (1999) Geochemical signatures of archaeological sites with black earth soils in Amazon region. *Journal of Geochemical Exploration* **66**, 369-385
- Cunha TJF (2005) Ácidos húmicos de solos escuros da Amazônia (Terra Preta de Índio). PhD thesis, Universidade Federal Rural de Rio de Janeiro, Rio de Janeiro, 118 pp
- Derenne S, Largeau C (2001) A review of some important families of refractory macromolecules: Composition, origin, and fate in soils and sediments. *Soil Science* **166**, 833-847
- Dick DP, Gonçalves CN, Dalmolin RSD, Knicker H, Klamt E, Kögel-Knabner I, Simões ML, Martin-Neto L (2005) Characteristics of soil organic matter of different Brazilian Ferralsols under native vegetation as a function of soil depth. *Geoderma* **124** (3-4), 319-333
- Empresa Brasileira de Pesquisa Agropecuária-Embrapa (1997) *Manual de Métodos de Análise de Solos*, Centro Nacional de Pesquisa de Solos, Embrapa Solos, Rio de Janeiro-RJ, 212 pp (Série Documentos, 1).
- Empresa Brasileira de Pesquisa Agropecuária-Embrapa (1999) *Sistema Brasileiro de Classificação de Solos*, Centro Nacional de Pesquisa de Solos, Embrapa Produção de Informação, Brasília-DF, Embrapa Solos, Rio de Janeiro-RJ, 412 pp
- Fearnside PM, Graça PML, Filho NL, Rodrigues FJA, Robinson JM (1999) Tropical forest burning in Brazilian Amazônia: Measurement of biomass loading, burning efficiency and charcoal formation at Altamira, Pará. *Forest Ecology and Management* **123**, 65-79
- Fearnside PM, Newton L, Fernandes FM (199X) Rainforest burning and the global carbon budget: Biomass, combustion efficiency, and charcoal formation in the Brazilian Amazon. *Journal of Geophysical Research* **98**, 16733-16743
- Freixo AA, Canellas LP, Machado PLOA (2002) Propriedades espectrais da matéria orgânica leve-livre e leve-livre intra-agregado de dois latossolos sob plantio direto e preparo convencional. *Revista Brasileira de Ciência do Solo* **26**, 445-453
- Glaser B, Balashov E, Haumaier L, Guggenberger G, Zech W (2000) Black carbon in density fractions of anthropogenic soils of the Brazilian Amazon region. *Organic Geochemistry* **31**, 669-678
- Glaser B, Lehmann J, Zech W (2002) Ameliorating physical and chemical properties of highly weathered soil in the tropics with charcoal – a review. *Biology and Fertility of Soils* **35**, 219-230
- Golchin A, Clarke P, Baldock JA, Higashi T, Skjemstad JO, Oades JM (1997) The effects of vegetation and burning on the chemical composition of soil organic matter in a volcanic ash soil as shown by ¹³C NMR spectroscopy: I whole soil and HA fraction. *Geoderma* **76**, 155-174
- Graça PMLA, Fearnside PM, Cerri CC (1999) Burning of Amazonian Forest in Ariquemes, Rondônia, Brazil: Biomass, charcoal formation and burning efficiency. *Forest Ecology and Management* **120**, 179-191
- Hamer U, Merschner B, Brodowski S, Amelung W (2004) Interactive priming of black carbon and glucose mineralization. *Organic Geochemistry* **35**, 823-830
- Instituto do Patrimônio Histórico e Artístico Nacional – IPHAN (1988) Coletânea da legislação de proteção ao patrimônio cultural. Iphan, Manaus-AM, pp 199-225
- IUSS Working Group WRB (2006) World Reference Base for Soil Resources 2006. FAO, Rome
- Kalbitz K, Geyer W, Geyer S (1999) Spectroscopic properties of dissolved humic substances – a reflection of land use history in a fen area. *Biogeochemistry* **47**, 219-238
- Kämpf N, Kern DC (2005) O solo como registro da ocupação humana pré-histórica na Amazônia. *Tópicos em Ciência do Solo* **4**, 277-320
- Kämpf N, Woods WI, Sombroek WG, Kern DC, Cunha TJF (2003) Classification of Amazonian dark earths and other ancient anthropic soils. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 77-102
- Kern DC (1996) Geoquímica e pedogequímica de sítios arqueológicos com Terra Preta na Floresta Nacional de Caxiuanã (Portel-Pará). PhD thesis, Universidade Federal do Pará, Belém-PA, 124 pp
- Kern DC (1988) Caracterização pedológica de solos com Terra Preta Aqueológica na Região de Oriximiná, Pará. MSc thesis, Universidade Federal do Rio Grande do Sul, Porto Alegre-RG, 232 pp
- Kern DC, Kämpf N (1989) Antigos assentamentos indígenas na formação de solos com Terra Preta Arqueológica na região de Oriximiná, Pará. *Revista Brasileira de Ciência do Solo* **13**, 219-225
- Kishimoto S, Sugiura G (1985) Charcoal as a soil conditioner. In: *Anais Symposium on Forest Products Research - International Achievements and the Future*, April 26, 1985, National Timber Research Institute, Pretoria, South Africa, pp 22-26
- Knicker H, Totsche KU, Almendros G, González-Vila FJ (2005a) Condensation degree of burnt peat and plant residues and the reliability of solid-state VACP MAS ¹³C NMR spectra obtained from pyrogenic humic material. *Organic Geochemistry* **36**, 1359-1377
- Knicker H, González-Vila FJ, Polvillo O, González JA, Almendros G (2005b) Fire-induced transformation of C- and N- forms in different organic soil fractions from a Dystric Cambisol under a Mediterranean pine forest (*Pinus pinaster*). *Soil Biology and Biochemistry* **37**, 701-718
- Kononova MM (1996) *Soil Organic Matter*, Pergamon Press, Oxford, 544 pp
- Kuhlbusch TAJ, Crutzen PJ (1995) Toward a global estimate of black carbon in residues of vegetation fires representing a sink for aTerra Mulataospheric CO₂ and a source of O₂. *Global Biogeochemical Cycles* **9**, 491-501
- Kumada K (1965) Studies on the color of HAs. Part 1: On the concepts of humic substances and humification. *Soil Science and Plant Nutrition* **11** (4), 11-16
- Lal R (2000) Physical management of soils of the tropics: Priorities for the 21st century. *Soil Science* **165** (3), 191-207
- Lehmann J, Kern DC, German LA, McCann J, Martins GC, Moreira A (2003) Soil fertility and production potential. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 105-124
- Liang B, Lehmann J, Sohi SP, Thies JE, O'Neill B, Trujillo L, Gaunt J, Solomon D, Grossman J, Neves EG, Luizão FJ (2010) Black carbon affects the cycling of non-black carbon in soil. *Organic Geochemistry* **41**, 206-213
- Madari B, Benites VM, Cunha TJF (2003) The effect of management on the fertility of Amazonian dark earths. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 407-432
- Martin-Neto L, Andriulo AE, Tragheta DG (1994) Effects of cultivation on ESR spectra of organic matter from soil fractions of a mollisol. *Soil Science* **157**, 365-372
- McCann JM, Woods WI, Meyer DW (2001) Organic matter and anthrosols in Amazônia: interpreting the American legacy. In: Rees RM, Ball BC, Campbell DC, Watson CA (Eds) *Sustainable Management of Soil Organic Matter*, CAB International, Wallingford, UK, pp 180-189
- Mielniczuk J, Bayer C, Vezzani FM, Lovato T, Fernandes FF, Debarba L

- (2003) Manejo de solo e culturas e sua relação com os estoques de carbono e nitrogênio do solo. *Tópicos em Ciência do Solo* 3, 209-248
- Neves EG, Petersen JB, Bartone RN, Silva CA** (2003) Historical and socio-cultural origins of Amazonian dark earths. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 29-50
- Novotny EH, Azevedo ER, Bonagamba TJ, Cunha TJF, Madari BE, Benites VM, Hayes MHB** (2007) Studies of the compositions of humic acids from Amazonian dark earth soils. *Environmental Science and Technology* 41, 400-405
- Novotny EH, Hayes MHB, Madari BE, Bonagamba TJ, Azevedo ER, Souza AA, Song G, Nogueira CM, Mangrich AS** (2009) Lessons from the Terra Preta de Índios of the Amazon Region for the utilisation of charcoal for soil amendment. *Journal of the Brazilian Chemical Society* 20, 1003-1010
- Pabst E** (1992) Critérios de distinção entre Terra Preta e Latossolo na Região de Belterra e os seus significados para a discussão pedogenética. *Bol. Mus. Par. Emílio Goeldi. Série Antropol.* 7, 5-19
- Poirier N, Derenne S, Rouzaud J-N, Largeau C, Mariotti A, Balesdent J, Maquet J** (2000) Chemical structure and sources of the macromolecular, resistant, organic fraction isolated from a forest soil (Lacadée, south-west France). *Organic Geochemistry* 31 (9), 813-827
- Renard D, Iriarte J, Birk JJ, Rostain S, Glaser B, McKey D** (in press) Ecological engineers ahead of their time: The functioning of pre-Columbian raised-field agriculture and its potential contributions to sustainability today. *Ecological Engineering* VOL. PAGES
- Rossel RA, Angrilo AE, Schnitzer M, Crespo MB, Miglierina AM** (1989) HAs properties of na Argjudoll soil under two tillage systems. *Sci. Total Environ.* 81-82, 391-400
- Ruivo MLP, Arroyo-Kalin MA, Schaefer CER, Costi HT, Arcanjo SHS, Lima HN, Pulleman MM, Creutzberg D** (2003) The use of micromorphology for the study of the formation and properties of Amazonian dark earths. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 243-254
- Schmidt MWI, Noack AG** (2000) Black carbon in soils and sediments: analysis, distribution, implications, and current challenges. *Global Biogeochemical Cycles* 14, 777-793
- Schmidt MWJ, Skjemstad E, Gehrt E, Koel-Knabner I** (1999) Charred organic carbon in German chernozemic soils. *European Journal of Soil Science* 50, 351-365
- Schnitzer M, Gupta UC** (1965) Determination of acidity in soil organic matter. *Soil Science Society of America Proceedings* 27, 274-277
- Schulze DG, Nagel JL, Van Scoyoc GE, Henderson TL, Baumgardner MF, Stott DE** (1993) Significance of organic matter in determining soil colors. In: Bigham JM, Ciolkosz EJ (Eds) *Soil Color*, SSSA Special Publication Number 31, SSSA, Madison, pp 71-90
- Singer LS** (1959) Synthetic ruby as a secondary standard for the measurement of intensities in electron paramagnetic resonance. *Journal of Applied Physics* 30, 1463-1464
- Soares R** (2007) Agregação e distribuição da matéria orgânica em solos de Terra Preta de Índio da Amazônia Central. MSc thesis, Universidade Federal Fluminense, Niterói, 120 pp
- Sollins P, Homann P, Caldwell BA** (1996) Stabilization and destabilization of soil organic matter: Mechanisms and controls. *Geoderma* 74, 65-105
- Sombroek WG** (1966) *Amazonas Soils. A Reconnaissance of the Soils of the Brazilian Amazon Region*, Centre for Agricultural Publication and Documentation, Wageningen, 292 pp
- Stevenson FJ** (1994) *Humus Chemistry: Genesis, Composition, Reactions* (2nd Edn), Wiley, New York, 496 pp
- Swift RS** (1996) Organic matter characterization. In: Sparks DL, Page AL, Helmke PA, Loeppert RH, Soltanpour PN, Tabatabai MA, Johnston CT, Sumner ME (Eds) *Methods of Soil Analysis (Part 3) Chemical Methods*, SSSA Book Series, 5, SSSA/ASA, Madison, pp 1011-1020
- Teixeira WG, Martins GC** (2003) Soil physical characterization. In: Lehmann J, Kern DC, Glaser B, Woods WI (Eds) *Amazonian Dark Earths. Origin, Properties, Management*, Kluwer Academic Publishers, Dordrecht, pp 271-286
- Trompowsky PM, Benites VM, Madari BE, Pimenta AS, Hockaday WC, Hatcher PG** (2005) Characterization of humic like substances obtained by chemical oxidation of eucalyptus charcoal. *Organic Geochemistry* 36, 1480-1489
- Tryon EH** (1948) Effects of charcoal on certain physical, chemical, and biological properties of forest soils. *Ecol. Monogr.* 18, 81-115
- Woods WI, McCann JM** (2001) El origen y persistencia de las tierras negras de la Amazonia. In: Hiraoka M, Mora S (Eds) *Desarrollo Sostenible en la Amazonia*, PUBLISHER. CITY, pp 23-30
- Yeomans JC, Bremner JM** (1988) A rapid and precise method for routine determination of organic carbon in soil. *Communications in Soil Science and Plant Analysis* 19, 1467-1476
- Zech W, Haumaier L, Hempfling R** (1990) Ecological aspects of soil organic matter in tropical land use. In: Mccarthy P, Clapp CE, Malcolm RL, Bloom PR (Eds) *Humic Substances in Soil and Crop Sciences: Selected Readings*, SSSA/ASA, Madison, pp 187-202
- Zsolnay A, Baigar E, Jimenez M, Steinweg B, Saccomandi F** (1999) Differentiating with fluorescence spectroscopy the sources of dissolved organic matter in soils subjected to drying. *Chemosphere* 38, 45-50