Natural variation of arsenic fractions in soils of the Brazilian Amazon

Ediu Carlos da Silva Júnior^a, Gabriel Caixeta Martins^b, Lúcia Helena de Oliveira Wadt^c, Kátia Emídio da Silva^d, Roberval Monteiro Bezerra de Lima^d, Karine Dias Batista^e, Marcelino Carneiro Guedes^f, Raimundo Cosme de Oliveira Junior^g, André Rodrigues Reis^h, Guilherme Lopes^a, Michele Duarte de Menezes^a, Martin R. Broadleyⁱ, Scott D. Youngⁱ, Luiz Roberto Guimarães Guilherme^{a,*}

^a Department of Soil Science, Federal University of Lavras, Lavras, MG CEP 37200-000, Brazil

^b Vale Institute of Technology, Belém, PA CEP 66055-090, Brazil

^c Embrapa Rondônia, Porto Velho, RO CEP 76815-800, Brazil

^d Embrapa Amazônia Ocidental, Manaus, AM CEP 69010-970, Brazil

^e Embrapa Roraima, Boa Vista, RR CEP 69301-970, Brazil

^f Embrapa Amapá, Macapá, AP CEP 68903-419, Brazil

^g Embrapa Amazônia Oriental, Belém, PA CEP 66095-903, Brazil

^h São Paulo State University (UNESP), School of Sciences and Engineering, Rua Domingos da Costa Lopes 780, Tupã, SP CEP 17602-496, Brazil

ⁱ School of Biosciences, University of Nottingham, Sutton Bonington Campus, Loughborough LE12 5RD, UK

HIGHLIGHTS

- Sena Madureira presents As concentration in soils higher than the prevention level.
- Average extractable As (soluble + available) corresponds to 0.24% of the total.
- Available As represents >70% of the extractable As, on average.
- Arsenic fractions are mainly influenced by soil pH and are highly correlated with Al^{3+} .

GRAPHICAL ABSTRACT

ARTICLE INFO

Article history:

Received 31 December 2018

Received in revised form 26 May 2019

Accepted 29 May 2019

Available online 03 June 2019

Editor: Prosun Bhattacharya

ABSTRACT

Arsenic (As) in native soils of the Amazon rainforest is a concern due to its likely origin from the Andean rivers, which transport loads of sediments containing substantial amounts of trace elements coming from the cordilleras. Yet, unveiling soil As baseline concentrations in the Amazon basin is still a need because most studies in Brazil have been performed in areas with predominantly high concentrations and cannot express a real baseline value for the region. In this study, 414 soil samples (0–20, 20–40 and 40–60 cm layers) were collected from different sites throughout the Amazon basin - including native Amazon rainforest and minimally disturbed areas - and used to determine total and extractable (soluble + available) As concentrations along with relevant soil physicochemical properties. Descriptive statistics of the data was performed and Pearson correlation supported

* Corresponding author.

E-mail address: Guilherm@dcs.ufra.br (L.R.G. Guilherme).

Keywords:

Arsenic
Extractable As
Prevention level
Carcinogenic
Baseline
Amazon

by a Principal Component Analysis (PCA) provided an improved understanding of where and how As concentrations are influenced by soil attributes. Total As concentration ranged from 0.98 to 41.71 mg kg⁻¹ with values usually increasing from the topsoil (0–20 cm) to the deepest layer (40–60 cm) in all sites studied. Considering the proportional contribution given by each fraction (soluble and available) on extractable As concentration, it is noticeable that KH₂PO₄-extractable As represents the most important fraction, with >70% of the As extracted on average in all the sites studied. Still, the extractable fractions (soluble + available) correspond to ~0.24% of the total As, on average. Total, available, and soluble As fractions were strongly and positively correlated with soil Al³⁺. The PCA indicated that soil pH in combination with CEC might be the key factors controlling soil As concentrations and the occurrence of each arsenic fraction in the soil layers.

© 2019 Elsevier B.V. All rights reserved.

1. Introduction

Arsenic (As) occurs naturally in rocks, soil and water, with higher levels usually found close to sites with anthropogenic pollution, with special attention to groundwater in several countries in the world, with concentration levels exceeding the WHO drinking water guideline value of 10 µg L⁻¹ (WHO, 2001; Chowdhury et al., 2018). Contamination has been detected in ground and surface water in 15 of the 20 Latin American countries in 2011, and much of the As contamination described during the last 20 years in this region were published much later or never in the international literature (Bundschuh et al., 2012).

The overall mean value of total As for different soils worldwide is estimated as 6.83 mg kg⁻¹ (Kabata-Pendias, 2011), which is very close to the mean value found for Brazilian oxisols (i.e., 5.92 mg kg⁻¹) (Campos et al., 2007). Yet, As is found more abundantly in argillaceous sediments (up to 13 mg kg⁻¹). Anionic forms, AsO₄³⁻, HAsO₄²⁻, and H₂AsO₃²⁻, predominate in soil solution (Kabata-Pendias and Mukherjee, 2007).

Exposure to As has a great impact on human health across the world (Kapaj et al., 2006; Meliker et al., 2007; Chatterjee et al., 2018). In the environment As often raises a great concern because of its high toxicity to diverse human body systems (e.g. integumentary, nervous, respiratory, cardiovascular, hematopoietic, immune, endocrine, hepatic, renal, reproductive) as well as to its capability of causing different types of cancer (skin, lung, and bladder) (Abdul et al., 2015; Gamboa-Loira et al., 2017). In fact, this naturally occurring metalloid is considered a well-established “Class I” human carcinogen by the International Agency of Research on Cancer (IARC) (Minatel et al., 2018).

Arsenic contamination originating from geogenic sources occurs in many regions worldwide, such as Bangladesh and India (Arco-Lázaro et al., 2018; Rahman et al., 2018). It is estimated that over 200 million people are chronically exposed to As concentrations at or above the World Health Organization (WHO) threshold across the world (Naujokas et al., 2013). In addition to the natural occurrence of arsenic, there are many other anthropogenic sources that may increase soil arsenic concentration, for example: mining, smelting, application of wastes, animal manures, As-bearing pesticides, wood preservation, irrigation using contaminated water, coal fly ash, glass manufacturing, wastewater sludge, pharmaceutical waste, livestock dips and smelting activities to phosphate fertilizers (Punshon et al., 2017).

Several studies have clearly shown that the adverse effects of arsenic are not only dependent on the total concentration of this element, but rather on its reactive fractions (Wang et al., 2018). For As, this fractionation may include soluble and exchangeable, bound to carbonates, iron and manganese oxides, organic matter, sulphides and residual fraction among others. Those fractions govern arsenic bioavailability and mobility in soils (Ho et al., 2013; Shahmoradi et al., 2017; Liu et al., 2018). Additionally, soil physicochemical properties such as pH, redox potential, clay and organic matter (OM) content and the presence of metal (Fe, Mn and/or Al) oxides and/or competing ions such as phosphates highly influence arsenic bioavailability and mobility (Sharma et al., 2010; Biswas et al., 2014).

Many sequential extraction procedures (SEP) can be used to extract chemical fractions of trace elements in soil (Wan et al., 2017; He et al., 2013). These methods classify trace elements in soil into different operationally-defined fractions with increasing metal binding strength (Wan et al., 2017). Soil chemical extractions for soluble and available As can help understand the element's dynamic chemical behavior and its possible entrance into the food chain.

Many extractants can be used for this purpose, for example, the soluble fraction can be extracted using water (water-soluble) (Wenzel et al., 2001; Baroni et al., 2004; Jiang et al., 2014; Wan et al., 2017). This fraction contains water-soluble species such as free arsenate, and weakly adsorbed species, representing the most mobile and potentially the most bioavailable As species in the environment (Wan et al., 2017). Additionally, the soluble fraction can be also extracted using inorganic salts such as KCl (Kim et al., 2003) and KNO₃ (Alam et al., 2007).

Another relevant fraction of As - the available fraction - is not well defined in the literature in terms of method of extraction, but would be better classified as the As retained in the solid phase interacting with the soil solution and occupying exchangeable sites as diffuse ion or as outer-sphere complexes (Van Herreweghe et al., 2003) and can be extracted by using Na₂HPO₄ (Kim et al., 2003), MgCl₂ (Jiang et al., 2014), CH₃COOH (Wan et al., 2017), (NH₄)₂SO₄ (Wenzel et al., 2001) and KH₂PO₄ (Thinh et al., 2018) among others.

In Brazil, soil quality criteria for some contaminants were defined by CONAMA in the normative 420/2009, which sets screening values based on total concentration of selected elements. For arsenic, the established threshold for the so-called “prevention level” in soil was 15 mg As kg⁻¹ and is primarily based on phytotoxicity tests and ecological risk assessment. Besides this, investigation/intervention values derived from human health risk assessments were set as 35 mg As kg⁻¹ for agricultural, 55 mg As kg⁻¹ for residential, and 150 mg As kg⁻¹ for industrial areas (Conama, 2009). These thresholds were established for total concentrations and in some situations can be very restrictive, especially in soils with low bioavailability. This justifies investing more effort to assess As availability in soils, especially in tropical ecosystems, which are known for their oxidic characteristics that render arsenic less bioavailable (Campos et al., 2007, 2013a).

Increased As levels have been observed in soils worldwide and also in Brazil, especially due to mining activities (Bundschuh et al., 2012). Hypothetically, we could expect high concentrations of As in soils nearby the Amazon basin because of the sediments coming through the Amazon river (and its tributaries) from the Andes, which are expected to be naturally enriched in selected volatile trace elements (e.g., Se and Hg) (Fadini and Jardim, 2001; Silva Junior et al., 2017). However, there is a lack of comprehensive information about background levels of As in soils from the Amazon basin to prove this assumption since the more recent studies have focused only in specific states of Eastern Amazon (Fernandes et al., 2018). Therefore it is necessary to perform a more inclusive study concerning the occurrence/distribution of As in different Amazon agroecosystems in order to provide information for the local population about possible health issues involved in the

use of soil and water resources for living, building and growing crops in this region.

To the best of our knowledge, no research data regarding the determination of soluble and available soil As in the Brazilian Amazon region have been published so far. This is relevant, as such information could support public policies for decision makers and help guide future research in these sites in order to provide a better planning for a sustainable use of land and natural resources by the local population, while ensuring also human health and food security.

In this context, this study aims to: i) characterize and compare baseline levels of As in different sites in the Amazon region; ii) study the dynamics of extractable As fractions (soluble and available fractions) with depth, i.e., in three different layers (0–20, 20–40 and 40–60 cm) of soils occurring in agroecosystems with Brazil nuts plantations; and, iii) assess the influence of the main soil physicochemical attributes for determining the soil As concentrations and the occurrence of each arsenic fraction in the soils.

2. Materials and methods

2.1. Study approach

This study involved the following steps: (a) soil survey to collect baseline data for As in 9 target sites; (b) description of the natural variation of As and soil physicochemical attributes in each site; and, (c) correlation and PCA analysis to support inference of soil

environmental controls for As variability in the soil. The methods used to reach our goals are described in the following sections.

2.2. Sampling sites and characterization of the study area

Soil samples were collected in nine sites belonging to the Amazon Biome in Brazil. Each site correspond to a Brazil nut stand, identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato Grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracaraí, state of Roraima). In each site, samples of 15 collection points were sampled in the native Amazon rainforest, except for AR (Itacoatiara), where samples of 18 points were collected in the Aruanã farm, which is a large Brazil nuts plantation. Samples were collected between 2014 and 2017. The chosen target sites for this study were the same ones used by [Silva Junior et al. \(2017\)](#) in their survey assessing the natural variation of selenium in Brazil nuts and soils from the Amazon region, to which we have added four additional areas. Sampling sites and detailed geographic information are provided in [Fig. 1](#) and Table S1.

Soil samples were collected at the depths of 0–20, 20–40, and 40–60 cm at a distance of 3 m from the trunk of selected Brazil nuts trees. These samples were collected at four points and mixed to compose a single sample for each depth. The sampling method is represented schematically according to [Silva Junior et al. \(2017\)](#).

Fig. 1. Map of the Brazilian Amazon (within South America) (A) with visualization of the Andes mountain range and Amazon sub-regions (B) as well as indicating the location of the nine studied sites (C) identified by: Sena Madureira (SM) and Xapuri (XP)-state of Acre; Porto Velho (RO)-state of Rondônia; Anori (ANO) and Aruanã farm, Itacoatiara (AR)-state of Amazonas; Itaúba (MT)-state of Mato Grosso; Caracaraí (RR)-state of Roraima; Santarém (PA)-state of Pará and Laranjal do Jari (AP)-state of Amapá. Relevant locations referred in the manuscript for data comparison (e.g., Cerrado biome as well as the State of Minas Gerais) are also shown in the map. DEM - Digital elevation model. Source: U.S. Geological Survey's Center for Earth Resources Observation and Science (EROS).

2.3. Soil physicochemical characterization

Soil samples were taken to the Soil Science Department of the Federal University of Lavras, Minas Gerais-Brazil, where they were grinded, sieved at <2 mm, dried at room temperature, and subjected to chemical and physical (texture) analysis. The soil pH was determined potentiometrically in water (pH H₂O) and KCl (1 mol L⁻¹) in the ratio 1:2.5 soil/solution. The exchangeable cations (Ca²⁺, Mg²⁺ and Al³⁺) were obtained by 1 mol L⁻¹ KCl extractant. Sulfur (S-SO₄²⁻) was extracted as sulfate by monocalcium phosphate with acetic acid and the result converted to S. Available P, Na, K, and micronutrients (Fe, Zn, Mn, and Cu) were obtained by Mehlich-1 (0.05 mol L⁻¹ HCl + 0.0125 mol L⁻¹ H₂SO₄; in ratio 1:10 soil/solution). Potential acidity (H + Al) was estimated indirectly through SMP pH (Shoemaker et al., 1961). Exchangeable aluminum (Al³⁺) was determined by titration with sodium hydroxide (0.025 mol L⁻¹ NaOH). P was determined colorimetrically, K by flame emission photometry, S-SO₄²⁻ by turbidimetry, and Ca, Mg, Zn, Fe, Mn, and Cu were determined by atomic absorption spectrometry (AAS) (Raij and Quaggio, 1983). The organic matter content was determined by oxidation of the organic matter in the soil with a potassium dichromate solution in the presence of sulfuric acid (Carter and Gregorich, 2006). Sum of bases (SB), cation exchange capacity at pH 7 (T), effective cation exchange capacity (t), base saturation (V%), and aluminum saturation (m) were calculated based on the results of the chemical analysis. The particle size analysis (clay, silt, and sand) was performed by the "Bouyoucos" method, as described by Embrapa (1997).

2.4. Total arsenic determination

Total As analyses was performed in air-dried and <2 mm sieved soil samples. A portion of each soil sample was finely grinded (<0.45 mm) using an agate ball mill (Retch PM 400 mill). Soil samples (~0.2 g DW) were fully digested in 70% HF, 70% HNO₃ and 60% HClO₄ (TAG; Fisher Scientific UK Ltd), using perfluoroalkoxy (PFA) digestion vessels and a 48-place teflon-coated graphite digester block (Model A3, Analysco Ltd., Chipping Norton, UK) (Chilimba et al., 2011).

The digester block runs with two different programs, a mix of reagents and a few steps for 3 days as can be observed with further details in Table 1. After the digestion, the block was turned off and the tubes were left to cool down completely for about 30 min. Digested samples were diluted to 50 mL into volumetric flasks using Milli-Q water and stored in 5% HNO₃ at room temperature in universal sample bottles, pending elemental analysis. Before the analysis in the ICP-MS, the extracts were diluted using 1 mL + 9 mL of Milli-Q water.

Each digestion batch (n = 70) included a standard reference material (n = 3) (NIST 2711a; Montana Soil II, National Institute of Standards and Technology, Gaithersburg, MD, USA), which contains 107 ± 5 mg kg⁻¹ As. The average recovery in the standard reference material was 101% (n = 21, SD = 6.61) for As. In addition, 3 blank samples were used in each batch for quality control purposes and for calculating limits of detection. The limit of detection (LOD) was established using 21 blank extracts for soil samples following the overall procedure. The values were calculated with three times the standard deviation (SD)

of the blanks used in each batch (n = 3) using the formula: SD (3 blanks) × 3 × final volume extract (mL) ÷ sample weight (g). The average LOD was 0.029 mg kg⁻¹ for total soil As.

2.5. Soluble and available As in soil (extractable As)

Soluble and available As analyses were performed in sequence, according to the modified method described by Cai et al. (2002). First, for the soluble fraction, 2 g of air-dried and <2 mm sieved soil was added into a 50 mL polycarbonate centrifuge tube. Next 10 mL of 0.01 mol L⁻¹ KNO₃ was added to each tube and the suspensions were shaken for 2 h using a rotary shaker (20 rpm). Soil suspensions were then centrifuged for 30 min at 3500 rpm, filtered to <0.22 µm using a Millex syringe driven filter unit (Millipore, Cork, Ireland) and stored at room temperature prior to analysis of As by ICP-MS. In the second step, for the available fraction, an extraction with 10 mL of 0.016 M KH₂PO₄ was performed in the same tubes from the previous step. The suspensions were shaken for 1 h in the rotary shaker (20 rpm) and then centrifuged for 30 min at 3500 rpm. The supernatant was filtered using <0.22 µm syringe filter and put in a universal tube (used for analysis in the ICP-MS). For the ICP-MS analysis, 4.8 mL of the supernatant collected from KNO₃ and KH₂PO₄ extractions were added into an ICP tube containing 0.2 mL of 50% HNO₃ for As analysis. The limits of detection (LOD) for soluble and available soil As were calculated as previously mentioned, and corresponded to 0.047 and 0.056 µg kg⁻¹, respectively.

2.6. Analytical determination of As

The analytical determinations of As were performed using Inductively Coupled Plasma-Mass Spectrometry (ICP-MS, X-Series II, Thermo Fisher Scientific) in the Environmental geochemistry analytical suite from the University of Nottingham-UK. Samples were introduced at 1 mL min⁻¹ through a concentric glass Venturi nebuliser and Peltier-cooled (3 °C) spray chamber. Instrumental drift was corrected using internal standards (⁷²Ge, ¹⁰³Rh and ¹⁸⁵Re). Data for total As concentration in soils are reported on a dry weight basis (DW) and expressed in mg kg⁻¹. Data for soluble and available As are expressed in µg kg⁻¹.

2.7. Statistical analysis

A descriptive analysis of the concentrations of total As, as well as for soluble and available As was carried out using Box plots in order to display data distribution. The data were subjected to Pearson's correlation analysis, which was performed between total, soluble and available As concentrations and other physicochemical soil properties. All the graphics were performed using the software R 3.4.4 (R Development Core Team, 2018).

Analysis of variance was performed to assess how the log-transformed As concentration in soils varied across different sites. After verifying the significance through ANOVA, the mean values for As concentration were compared among sites for the three layers studied using Tukey's HSD test with the package emmeans v 2.23 (Lenth, 2016) in R 3.4.4 (R Development Core Team, 2018). The same approach was used to compare soluble and available As in soils, but in this case using generalized least squares (GLS) for the data log-transformed, in order to allow the variance to be estimated separately for each site (Cleasby and Nakagawa, 2011).

A principal component analysis (PCA) was performed to find out the main soil attributes influencing As concentration in the soil according to the degree of variance explained and to confirm the previous relationships obtained by Pearson correlation analyses. Because the distribution of As concentrations and fractions was not normal, the data were log-transformed, before analysis to meet the assumption of normality required for the Pearson correlation and PCA, except for soil pH, because it follows a normal distribution in soil data.

Table 1
HF digestion steps with digester block.

Day	Program	Reagent	Volume (mL)	Steps
1°	1 ^a	HNO ₃	2	1) 80 °C, 8 h;
		HClO ₄	1	2) 100 °C, 2 h;
2°	2	HF	2.5	1) 120 °C, 1 h;
				2) 140 °C, 3 h;
				3) 160 °C, 4 h;
3°	–	HNO ₃	2.5	4) 50 °C, 1 h.
		Milli-Q H ₂ O	2.5	

^a Left overnight with the program turned on.

3. Results and discussion

3.1. Soil physicochemical properties, element contents and associations

The chemical and textural attributes of soil varied widely among the sites studied, reflecting a great variability in soil characteristics throughout the Amazon basin that may be related to diversity of parent material and soil types. Most of collected samples show acidic soils (mean pH ranging from 3.75 to 5.5) which in fact reflects the well-known weathered soils from this region according to [Alvarez et al. \(1999\)](#). The OM content in the topsoil is classified as moderate in most sites (average OM content ranging from 1.16% to 2.63%) according to [Alvarez et al. \(1999\)](#) (Table 2). The moderate contents of OM can be explained by the high temperature and precipitation levels (humidity) in the region, i.e., the most important climatic factors controlling the dynamic of ecological processes ([Da Silva et al., 2018](#)), which reflects not only in the biomass production but also in decomposition rates of the litter in the soil.

The dominant soil texture was clay (AR, RO and PA) and sandy loam (MT and AP). Such high percentage of clay content facilitates As retention, especially onto kaolinite, as well as Fe and Al oxyhydroxide surfaces that predominate in the clay fraction of the highly developed soils of the Amazon region, which is subjected to strong weathering conditions ([De Souza et al., 2018](#)) (Table 2).

Soil cation exchange capacity (CEC) ranged from 2.96 to 25.2 $\text{cmol}_c \text{dm}^{-3}$ demonstrating a great variability among the sites studied. The soils from ANO showed the highest mean values of CEC (22.6 $\text{cmol}_c \text{dm}^{-3}$). For the others sampled sites, CEC values were all below 13 $\text{cmol}_c \text{dm}^{-3}$ with the lowest values observed in the sites RR and AP (5.33 and 4.28 $\text{cmol}_c \text{dm}^{-3}$, respectively). This wide variation in soil CEC indicates that besides the predominance of dystrophic and highly weathered soils with secondary minerals, there are still soils with 2:1 clay minerals, composing less weathered soils with high activity in the clay fraction in this region (Table 2).

Extractable S and P in the present study exhibited considerable variation among the sites with mean values ranging from 5.29 to 47.9 mg kg^{-1} (S) and from 0.52 to 3.72 mg kg^{-1} (P). In general, the content of available P was moderate in most sites, where also the OM contents are usually higher than in deeper layers, except in the sites MT and RO in southern Amazon, which contained available P values below 2 mg kg^{-1} (Table 2). The content of extractable S on the other hand presented higher values in the sites AR and RO (Table 2).

Iron (Fe) usually plays an important role in controlling the concentration of trace elements in soil solution because of its involvement in adsorption. The variation of available Fe among the sites was also quite outstanding with mean values ranging from 58.5 mg kg^{-1} (AP) to 489.7 mg kg^{-1} (ANO) (Table 2). We observed that soil pH is the major factor explaining the concentration of this element in soil

solution, since the site with lowest Fe content presents also the highest pH (site AP). On the other hand, available Fe might not reflect the concentration of iron (hydr) oxides that explains the retention of anions in soil.

3.2. Total As concentration in soil

Total As concentrations ranged from 0.98 to 41.7 mg kg^{-1} , with 75% of the soil samples showing values lower than 6.93 mg kg^{-1} , which is close to the mean value of total As reported for different soils worldwide (i.e., 6.83 mg kg^{-1}) ([Kabata-Pendias, 2011](#)). The sites with highest baseline As concentration were SM followed by RO and XP. On the other hand, the sites with lowest As concentrations were AP and RR, according to the Tukey HSD test (Fig. 2).

Interestingly, total As concentration in the western Amazon (SM, XP and RO) presented the highest values and a wide variation not only among the layers but also within the three layers evaluated. The total As concentration in the western Amazon showed median concentrations increasing from the topsoil to the deepest layers as follow: SM (15.9 < 19.0 < 26.5 mg kg^{-1}) and XP (5.96 < 7.20 < 9.34 mg kg^{-1}) for the layers 0–20, 20–40 and 40–60 cm, respectively. A wide variation for arsenic concentration within layers was observed especially at the deepest layer (40–60 cm), which also presented the highest As concentrations for each site studied in the western Amazon: SM (17.3 to 41.7 mg kg^{-1}), XP (5.69 to 17.8 mg kg^{-1}), and RO (5.83 to 23.6 mg kg^{-1}) (Fig. 2).

There is no nationwide As baseline level established by the Brazilian legislation to be used in terms of quality reference values (QRVs) for soils. This is also true for soils of the Brazilian Amazon region. However, a QRV for As in soils from the Eastern Amazon was proposed by [Fernandes et al. \(2018\)](#). These authors observed total As concentration ranging from 0.07 to 7.75 mg kg^{-1} in the 0–20 cm layer and, considering the 90th percentile of the sample population after eliminating anomalies (cumulative frequency distribution), 2.7 mg kg^{-1} was suggested as the QRV. Given the heterogeneity in the pedogeoclimatic conditions in the region, the 90th percentile seems to best represent the QRVs because of the wide variances for the values observed, as suggested by [Conama \(2009\)](#).

In the present study 57% of the samples from the 0–20 cm soil layer show As concentrations higher than the QRV for the 90th percentile proposed by [Fernandes et al. \(2018\)](#), which reinforces the difficulty of establishing a single QRV for a very large region as the Brazilian Amazon. In addition, the medians for total As concentrations in the three layers (0–20, 20–40 and 40–60 cm) from the site SM are higher than the As prevention level (15 mg kg^{-1}) established by the Brazilian Normative CONAMA 420, for soils nationwide ([Conama, 2009](#)) (Fig. 2). This is indicative that the current Brazilian prevention values need to be validated using representative soils from the Amazon basin since this

Table 2

Physicochemical properties of the topsoil (0–20 cm) from samples collected at different sites in the Amazon region.

Site	n	pH (H ₂ O)	OM (%)	CEC ($\text{cmol}_c \text{dm}^{-3}$)	S (mg kg^{-1})	P (mg kg^{-1})	Fe (mg kg^{-1})	Clay (%)	Silt	Sand
SM	15	4.53 ± 0.34	1.42 ± 0.35	6.92 ± 1.74	5.29 ± 1.17	3.72 ± 1.27	174.9 ± 86.3	21.6 ± 4.70	22.5 ± 2.38	55.9 ± 4.85
XP	15	4.36 ± 0.32	1.62 ± 0.55	7.21 ± 2.25	8.09 ± 5.16	3.08 ± 0.75	190.1 ± 83.9	18.4 ± 5.37	32.3 ± 10.2	49.3 ± 10.5
MT	15	4.19 ± 0.16	2.32 ± 0.40	7.10 ± 2.52	7.32 ± 2.00	1.46 ± 0.32	324.3 ± 69.6	19.5 ± 2.61	3.86 ± 1.85	76.6 ± 2.82
AR	18	4.26 ± 0.09	2.63 ± 0.32	7.97 ± 1.34	16.5 ± 7.06	2.56 ± 0.61	183.0 ± 48.6	77.6 ± 3.46	10.1 ± 2.37	12.3 ± 2.56
ANO	15	4.00 ± 0.35	2.03 ± 0.70	22.6 ± 10.9	9.55 ± 4.97	2.53 ± 0.85	489.7 ± 134.1	27.2 ± 5.82	56.7 ± 8.05	16.1 ± 11.9
AP	15	5.49 ± 0.34	1.16 ± 0.34	4.28 ± 0.67	9.49 ± 1.60	2.25 ± 0.72	58.5 ± 22.6	19.9 ± 2.60	3.40 ± 1.92	76.7 ± 3.24
RR	15	4.46 ± 0.16	2.43 ± 0.46	5.33 ± 0.73	10.1 ± 1.64	2.38 ± 0.28	177.2 ± 50.2	44.1 ± 5.97	8.13 ± 3.54	47.7 ± 6.12
RO	15	3.75 ± 0.22	2.40 ± 0.37	12.6 ± 3.63	18.0 ± 8.88	1.87 ± 0.37	196.2 ± 70.8	51.7 ± 9.78	28.1 ± 12.5	20.2 ± 7.70
PA ^a	15	4.41 ± 0.26	2.20 ± 0.42	9.24 ± 2.24	10.6 ± 3.51	2.25 ± 0.43	242.4 ± 62.5	52.0	8.0	40.0

Values of soil physicochemical attributes are mean ± standard deviation (SD).

Each site is identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracaraí, state of Roraima).

^a Texture in PA was derived from one single analysis performed by a composite sample resulted of subsamples from 5 spots in the site.

Fig. 2. Raw data for total As concentrations in soils from different sites in the Amazon region for the layers 0–20, 20–40 and 40–60 cm. Each site correspond to a Brazil nut stand, identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracarái, state of Roraima). The letter on top of each state indicates the result of Tukey's HSD test ($p = 0.05$) on the log-transformed As concentration, following a significant ANOVA ($p < 0.001$).

region may present soils that may be naturally enriched with As. All other studied sites presented median As concentrations below 6 mg kg^{-1} - a value close to the average As content reported by Campos et al. (2007) for 17 Brazilian oxisols (i.e., 5.92 mg kg^{-1}) - with the lowest median concentrations observed in the site AP (1.30 , 1.53 and 1.71 mg kg^{-1} for 0–20, 20–40 and 40–60 cm respectively) (Fig. 2).

The sites from the central and eastern portion of the Amazon basin (ANO, AR, PA and AP) presented low As concentrations in all the three layers evaluated. They also presented low variation among and within the layers, as observed in the medians concentrations in the site AR (5.07 , 5.33 and 5.41 mg kg^{-1} for 0–20, 20–40 and 40–60 cm respectively) (Fig. 2).

The sites located in the northern and southern portion of the Amazon, which are RR (medians of 1.39 , 1.70 and 1.80 mg kg^{-1}) and MT (medians of 2.56 , 2.99 and 3.03 mg kg^{-1}) also presented low total As concentration for the three layers evaluated compared to the other sites studied, and no meaningful variation among and within the layers was noticed (Fig. 2).

Campos et al. (2013b) determined background levels of As in representative soils from the states of Goiás and Minas Gerais (from 2000 to 2500 km far away from the studied sites), which belongs to Cerrado

biome (Brazil) in three sub-regions and observed concentrations similar to those found in the southern Amazon in the present study, with mean concentrations in a decreasing order as follow: East of state of Goiás (3.29 mg kg^{-1}) > Triângulo Mineiro (state of Minas Gerais) (2.18 mg kg^{-1}) > Northeast of Minas Gerais (0.62 mg kg^{-1}). The highest As concentrations were observed in profiles of plintossols and cambisols from the East of Goiás. In another study, Campos et al. (2013a) observed levels of As ranging between 0.14 mg kg^{-1} (for a gley soil) and 9.3 mg kg^{-1} (for an oxisol) in soils from Minas Gerais. Those concentrations are usually found in non-contaminated soils ($<10 \text{ mg kg}^{-1}$) (Adriano, 2001; Campos et al., 2007, 2013a; Bundschuh et al., 2012) and are similar to the ones reported in the present study.

In a different scenario far away from the Amazon region, in the Iron Quadrangle, southeast region of Brazil, within the state of Minas Gerais, the presence of As is associated with primary gold deposits where it is contained predominantly in S-bearing minerals such as arsenopyrite and pyrite (Borba et al., 2003). Also in the iron quadrangle, Alves and Rietzler (2015) observed high As concentrations in soils from mining areas in Nova Lima (13.2 in the control site and 489 mg kg^{-1} in the mine stream) and Santa Bárbara (82.3 in the control site and 1329 mg kg^{-1} nearby the mine plant). Out of the Iron Quadrangle, but still in Minas Gerais, natural soil As concentrations found in a private

natural reserve (native Cerrado vegetation) located nearby a gold-mining area ranged from 25 to 35 mg kg⁻¹ (Ono et al., 2012), which is also greater than the prevention level established by Normative 420, for Brazilian soils (Conama, 2009). These studies emphasize the importance of the parent material in total arsenic concentration in soils and make us believe that the main source of arsenic, in the present study, is geogenic.

3.3. Extractable soil As (soluble and available)

Extractions for soluble and available As showed the highest concentrations for the site SM which is in accordance with the results observed for total As. For the soluble fraction the concentrations ranged from <0.047 (MT) to 16 µg kg⁻¹ (SM). Available As varied from 2.14 (AR) to 236.5 µg kg⁻¹ (SM) (Fig. 3).

Observing the mean concentrations for soluble As, it is noteworthy that in SM the values are increasing from the topsoil to the deepest layers. On the other hand, in the sites AR, ANO, PA, RO and AP, soluble As decreases from the topsoil (0–20 cm) to the deepest layers (20–40 and 40–60 cm). The sites RR and AP presented the lowest average

concentration for soluble As (Fig. 3). The site with highest pH (AP) also presented the lowest median values for soluble As (Fig. 3). This effect is confirmed by the negative correlation between soluble As and pH in all the three layers studied (Table 3).

The average concentration for available As in SM in the layers 0–20, 20–40 and 40–60 cm (45.3, 68.6 and 117 µg kg⁻¹) are at least 4-fold higher than those observed in all other sites studied. There was no meaningful variation among the sites for available As.

The observed behavior of soluble and available As in these soils may be partly explained by their interactions with available P (Table 2), which showed positive correlation with both fractions in all the three layers studied (Table 3). This is expected as a greater content of P in oxidic soils (which retain phosphate strongly) make As much more available and soluble due to competitive adsorption on positively-charged sites (Campos et al., 2013a).

Considering the proportional contribution given by each fraction (soluble and available) to the extractable As concentration, it is noticeable that available As represents the major fraction with >70% of the As being extracted by KH₂PO₄ on average in all the sites studied.

Fig. 3. Raw data for concentration of As available and soluble for 9 sites in the Amazon region. Each site correspond to a Brazil nut stand, identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracará, state of Roraima). The letter on top of each state indicates the result of Tukey's HSD test ($p = 0.05$) on the log-transformed As concentration, following a significant ANOVA ($p < 0.001$) using generalized least squares.

Table 3
Pearson's correlation coefficients (r) for log transformed physicochemical variables versus total, soluble and available arsenic concentrations from different sites in the Amazon region in the layers 0–20 cm, 20–40 and 40–60 cm.

Soil attribute	0–20 cm			20–40 cm			40–60 cm		
	As	As-KNO ₃	As-KH ₂ PO ₄	As	As-KNO ₃	As-KH ₂ PO ₄	As	As-KNO ₃	As-KH ₂ PO ₄
pH	−0.43**	−0.33**	−0.14	−0.44**	−0.48**	−0.17*	−0.46**	−0.36**	−0.17
OM	0.06	0.27**	−0.12	−0.02	0.11	−0.19*	−0.12	−0.15	−0.23**
CEC	0.34**	0.26**	0.15	0.33**	0.35**	0.22*	0.39**	0.39**	0.30**
P	0.27**	0.39**	0.43**	0.31**	0.25**	0.32**	0.29**	0.23**	0.31**
S	0.03	0.08	−0.23*	0.14	0.10	−0.14	0.06	−0.05	−0.19*
Al	0.53**	0.40**	0.41**	0.52**	0.55**	0.44**	0.56**	0.60**	0.47**
Fe	0.08	0.09	−0.01	−0.06	0.09	−0.15	−0.15	−0.19*	−0.29**
Clay	0.15	0.26**	−0.06	0.04	0.10	−0.21*	0.01	0.0	−0.29**
Silt	0.55**	0.31**	0.40**	0.57**	0.45**	0.37**	0.56**	0.51**	0.40**

Soil attributes: total As (HF digestion); soluble arsenic (KNO₃-extractable As); available arsenic (KH₂PO₄-extractable As); pH (pH in H₂O); P (available phosphate); S (extractable S-SO₄^{2−}); CEC (cation exchange capacity); OM (organic matter); Al (exchangeable aluminum); Fe (available iron); clay and silt (%).

* Significant correlation ($p < 0.05$).

** Significant correlation ($p < 0.01$). The correlation includes all the sites studied except the site “Santarém/PA” because the data for texture was obtained in only one point.

Generally, the contribution given by the soluble fraction in all the sites reduces from the topsoil to the deepest layer, as follows: 25% > 18% > 16% (0–20, 20–40 and 40–60 cm respectively) (Fig. 4). The site with

the greatest contribution given by available As was SM with 92%, 94% and 94% for the 0–20, 20–40 and 40–60 soil layers. On the other hand, the sites with smallest contribution given by available As were RR in

Fig. 4. Relative contribution in % from soluble and available As in soils from different sites in the Amazon region. Each site is identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracaraí, state of Roraima).

the layer 0–20 cm (67.7%), PA in the layer 20–40 cm (75.7%) and AR in the layer 40–60 cm (77.5%) (Fig. 4).

Arsenic availability was also assessed using tri-distilled water, ammonium sulfate, ammonium phosphate, ammonium oxalate + oxalic acid, a mixture of organic acids, Mehlich-1, and USEPA 3051A in artificially contaminated soils (entisol and oxisol) by Melo et al. (2012). The extracting solutions tested in the experiment were equally efficient in assessing plant availability of As in the soils, however there was greater concentration of As extracted from the clayey entisol when compared with the medium-textured oxisol due to differences between the adsorptive capacities of these soils. Besides validating extractants for plant-available As, these authors emphasized the capacity of tropical soils, with variable charge, to adsorb As, especially those soils rich in clay (Melo et al., 2012).

3.4. Relationship between soil As and physicochemical attributes

Total As concentration (total As) correlated positively ($p < 0.05$) with CEC, available P, K, exchangeable Al, and silt in the three layers studied and also with available Zn in the 20–40 and 40–60 cm layers. Total As correlated negatively ($p < 0.05$) with pH, exchangeable Ca, available Cu in the three layers studied and also with exchangeable Mg and available Mn in the 0–20 cm layer and with available B and sand in the 0–20 and 20–40 cm layers. These results indicate that, in tropical soil conditions, increasing pH values will decrease total As content either directly because of its influence on colloidal charges (i.e., causing a decrease in positive charges) or indirectly influencing the availability and retention of competing anions such as phosphate and sulfate (considering that soils are open systems and available forms can be removed, thereby reducing total levels) (Table 3).

Soluble As correlated positively ($p < 0.05$) with CEC, available P, exchangeable Al and silt in the three layers evaluated and also with OM and clay in the 0–20 cm layer and with available Zn and K in the

40–60 cm layer. Soluble As correlated negatively ($p < 0.05$) with pH, exchangeable Ca and available Cu in the three layers studied and also with exchangeable Mg, available B and sand in the 0–20 and 20–40 cm layers following by Fe in the 40–60 cm and available Mn in the 0–20 and 40–60 cm layers (Table 3).

On the other hand, available As correlated positively ($p < 0.05$) with available P, K, Zn, exchangeable Al and silt in all the three layers studied and also with CEC in the 20–40 and 40–60 cm layers and with available Mn in the 20–40 cm layer following by sand in the 40–60 cm layer. Available As correlated negatively ($p < 0.05$) with OM and clay in the 20–40 and 40–60 cm, and also with pH in the 20–40 cm, with extractable S in the 0–20 and 40–60 cm layers, Fe in the 40–60 cm layer and available Cu in the 0–20 cm layer (Table 3).

In the present study, an important factor influencing As bioavailability that negatively affected total and soluble As was soil pH as can be observed in the Fig. 5. As pointed out by Zhang et al. (2017), numerous studies have shown that soil pH is an important factor controlling arsenic geochemistry, with adsorption of As decreasing as pH increases. This is attributed to the higher amounts of negative charges on soil components that generate variable charge, such as 1:1 clays, along with less amounts of positive charges on Fe and Al oxides. Such pH influence on charge generation is pronounced in tropical soils, since these are variable charge soils (Fontes et al., 2001).

According to Huang et al. (2006), NaH_2PO_4 -extractable As correlates well with total As concentrations found in edible parts of various crops and soils collected from paddy rice fields in suburban areas of Fukang province in southeast China, showing the usefulness of this extractant to assess the bioavailability of As in soils. These authors also reported that As availability decreased significantly with increasing silt size (0.02–0.002 mm) and free iron content, but increased significantly with soil pH and organic matter content. Those findings are not supported by results observed in the present study for the silt fraction, which showed positive correlation with

Fig. 5. Major mechanism at the soil colloid-soil solution interface controlling arsenic availability in tropical soils.

total, soluble, and available As in soil samples for all the three layers studied (Table 3).

Similarly to what was observed in the present study (20–40 and 40–60 cm layers), Melo et al. (2012) observed that As availability (expressed by concentration of As in plants) was negatively correlated with clay content. The main reason for finding negative correlations between available As and clay content, especially in subsurface, is because of the particular composition and characteristics of the clay fraction in tropical soils, which presents significant amounts of Fe and Al oxides and oxyhydroxides as well as high ZPC (Fontes et al., 2001). According to McBride (1994), chemical adsorption of As occurs mainly in Fe and Al oxides, as well as in amorphous aluminosilicates, with small quantities being retained by clay silicates and organic substances. For this reason, the mobility and bioavailability of As is generally lower in clayey soils with variable charge.

It is noteworthy that the highest correlation coefficients were observed between exchangeable Al (Al^{3+}) and total, available, and soluble As in all the three layers studied, which were all positive ($p < 0.05$) (Table 3). The mechanism by which high Al^{3+} is associated with high As concentrations may be related to the preferential retention of Al in the negative sites of variable-charge minerals (including clay minerals, Fe and Al oxides and oxyhydroxides), thus creating a bridge for binding As through formation of surface ternary complexes, which has been previously reported (Oliveira et al., 2015).

Given that As (as arsenate) and P (as phosphate) present similar chemical behavior in soils, available P has an antagonistic effect upon As retention, as P can compete with As for anionic sorption sites (Sharma et al., 2011). This explains the positive correlation observed between the soluble and available As fractions and the available P in soil samples of all the three layers studied (Table 3). Several studies show evidence supporting a strong relationship between As availability in soil and soil available P as a consequence of the chemical and structural similarities between their anionic forms (Punshon et al., 2017; Arco-Lázaro et al., 2018; Chowdhury et al., 2018). Jiang et al. (2014) observed significant positive correlation between soil available P and exchangeable As. The authors also stated that the main factors influencing As uptake by rice in that study included soil pH, total organic carbon, as well as soil available P and Fe. In this case, soil pH affected As uptake indirectly via impact on available P and Fe. Wang et al. (2018) conducted a pot experiment to study the effect of P application on the lability and phytoavailability of As in the soil and observed that application of exogenous phosphate induced soil arsenic mobilization, thereby increasing arsenic lability due to a competition effect. Finally, a positive correlation found between available P and total As in all soil layers is probably related to the fact that positively-charged colloidal surfaces that adsorb P in tropical agroecosystems are also relevant for retaining As in the soil system.

Extractable sulfur in this study affected negatively the amount of available As in the soil, in the layers 0–20 and 40–60 cm. The competition between S and As, mainly in the 40–60 cm layer, where sulfur could have more affinity with the binding sites in tropical soils is relevant if we consider that As in soil solution cannot overcome S in the sites available, due to a concentration effect, i.e., S in solution is generally higher than As in solution in most soils. Therefore the decreasing of available As in a long term is certain when the concentration of S is higher. But this is not the behavior often showed in experiments of shorter term where the addition of S displace As from the binding sites and increase its concentration in soil solution. In fact, Song et al. (2015) studying the effect of sulfate on As mobility in contaminated soils observed that addition of sulfate significantly enhanced As availability. Moreover, in a study aiming to evaluate the competitive adsorption of arsenate/phosphate and arsenate/sulfate in Brazilian soils, Campos et al. (2013a) observed that the addition of phosphate and sulfate decreased arsenate adsorption and consequently increased the concentration of As in soil solution and its availability.

The negative correlation between OM and available As in the 20–40 and 40–60 cm layers observed in the present study (Table 3), suggest that OM can be considered a trap and should be able to alleviate As mobility in soil by formation of covalent bindings, especially in OM-rich soils which is the case of some points sampled in these sites studied (Langner et al., 2012).

The adsorption capacity of As by soils provides an indication of the system's buffer power, since many soil attributes (such as texture, Fe and Al oxide content, among others) interfere with the buffer capacity and, consequently, influence leaching losses (Campos et al., 2013a). Although we did not find any relevant correlation among Fe and As fractions in soils from the present study, a great number of studies in the literature shows otherwise. In fact, a close coupling between the biogeochemical cycles of iron and arsenic in both oxidizing and reducing environment has been well established (Dixit and Hering, 2003). The mechanism that is presumably involved in this interaction is that both As(III) and As(V) adsorb to iron (hydr) oxides through, in part, formation of inner-sphere complexes (Tufano et al., 2008). Wang et al. (2012), studying As retention in muddy sediments observed that As retention was strongly coupled to Fe and S, and correlation analysis showed that total and extractable As correlated positively and well with total Fe and HCl-extractable Fe.

Fig. 6 shows the gradient of influence of the soil chemical variables and correlation groups of the samples for each site studied in the Amazon region. The model including the two principal components accounted for 52.7% of the total variance explained in the three layers studied (29.8% for PC1 and 22.9% for PC2).

The first component (PC1) was negatively dominated by P, S, K, total As, available As, soluble As, CEC, OM, Al and Fe and positively dominated by Zn, Mn, Cu, B, Mg, Ca and pH, which accounted for 29.8% of the variance. The second (PC2) was negatively dominated by S and positively dominated by Fe, OM, Al, AS, CEC, soluble As, available As, K and P, representing 22.9% of the total variance in the data (Fig. 6).

The soil attribute with the biggest contribution for PC1 is soil pH, followed by Ca and Mg, i.e., in terms of acidity and the presence of basic cations, soils from the Amazon are very diverse, reflecting a heterogeneous mineralogy and capacity to adsorb anions as can be observed by the groupings of sites formed along the axis of these attributes (Fig. 6 B). The soil attribute that contributed the most for PC2 is Zn, followed by K and P, which are influencing the grouping of the samples from the site XP.

Total, soluble, and available As are positively correlated with K, P, CEC, OM, Al and Fe and also present a major influence in the samples of the sites ANO, XP and SM. The close correlation between the As fractions and CEC and OM suggest that these soil properties are important factors to retain these As forms in the soil. With regard to CEC, unless there is a formation of a ternary complex in the interface soil-solution with an intermediate cation in the system, we believe that the correlation between CEC and As fractions is indirectly influenced by soil organic matter. The extractable S was separated from the other soil chemical attributes in the third quadrant and showed no relationship with total and soluble As, and only a weak relationship with available As (negative correlation in the 0–20 and 40–60 cm layers) (Table 3) and therefore represented a major influence for the samples from the sites AR and PA (Fig. 6).

The site AP forms a separated group in the border of first and fourth quadrant, which indicates that this site present its chemistry distinct from the others, mainly because of the higher values of pH, Ca, Mg, B, Mn, and Cu. On the other hand, this site receives lower impact from the total, soluble and available As fractions because of its lower As concentrations compared with the other sites (Fig. 6). In summary, soil pH in combination with CEC and OM might be the key factors that influence total As concentrations in soil for the conditions of the present study.

Fig. 6. Principal Component Analysis (PCA) of log-transformed data for A) Vectors with soil chemical attributes and total, soluble and available soil As concentrations and B) sampling points distributed from 9 different sites of the Amazon region identified by: SM (Sena Madureira, state of Acre); XP (Xapuri, state of Acre); ANO (Anori, state of Amazonas); AR (Aruanã farm-Itacoatiara, state of Amazonas); AP (Laranjal do Jari, state of Amapá); MT (Itaúba, state of Mato grosso); PA (Santarém, state of Pará); RO (Porto Velho, state of Rondônia) and RR (Caracará, state of Roraima). The symbols represent each soil depth as follow – circle: 0–20 cm; cross: 20–40 cm and triangle: 40–60 cm. Soil chemical attributes: S (extractable $S-SO_4^{2-}$ in $mg\ kg^{-1}$); available Mn, Zn, Fe, Cu, B, K, and P ($mg\ kg^{-1}$); exchangeable Mg, Ca, and Al in $cmol_c\ dm^{-3}$; CEC (cation exchange capacity in $cmol_c\ dm^{-3}$); OM (organic matter in %); pH (pH in H_2O); total As ($mg\ kg^{-1}$); soluble As ($\mu g\ kg^{-1}$); available As ($\mu g\ kg^{-1}$). Graphic was designed using the software PAST (Hammer et al., 2001).

4. Conclusions

Soil pH and CEC are indirectly the main chemical attributes influencing the dynamic of As fractions in soils from the sites studied in the Amazon basin. In a smaller magnitude but not less important, the competing anions P and S affect soluble as well as available As in the soil.

The highest variable correlating with total, soluble and available As is soil Al^{3+} in the three layers studied. The mechanisms by which this element affect As mobility and retention needs further investigation, but might be related to formation of ternary complexes.

Our study raises the need to complement the information about the background levels of As in soils from the Amazon region using representative soils that reflects the natural concentrations of each sub-region in order to provide guideline values for a better use of soil and water resources in the region.

Acknowledgements

The authors are grateful to CNPq, CAPES and FAPEMIG from Brazil for financial support to the research project and especially CAPES for

granting the doctorate scholarship for ECSJ. This study was financed in part by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – Brasil (CAPES) – Finance Code 001.

GCM would like to thank for his Post-Doc scholarship provided by CAPES (Grant number 88887.160998/2017-00).

ARR also thanks the National Council for Scientific and Technological Development (Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq) for the research fellowship (Grant number 309380/2017-0).

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.scitotenv.2019.05.446>.

References

- Abdul, K.S.M., Jayasinghe, S.S., Chandana, E.P.S., Jayasumana, C., De Silva, P.M.C.S., 2015. Arsenic and human health effects: a review. *Environ. Toxicol. Pharmacol.* 40, 828–846.

- Adriano, D.C., 2001. Trace Elements in Terrestrial Environments: Biogeochemistry, Bioavailability, and Risks of Metals. 2nd. ed. Springer-Verlag New York, New York, USA.
- Alam, M.G.M., Tokunaga, S., Stagnitti, F., 2007. Removal of arsenic from contaminated soils using different salt extractants. *J. Environ. Sci. Heal. Part A* 42, 447–451.
- Alvarez, V.V.H.R.F., Novais, N.F., Barros, R.B., Cantarutti, A., Lopes, S., Alvarez, V.V.H., 1999. Interpretation of soil analysis results. In: Ribeiro, A.C., PTG, Guimarães (Eds.), *Recommendations for Use of Amendments and Fertilizers in Minas Gerais, Brazil*, 5th ed, pp. 25–32 Viçosa. (In Portuguese).
- Alves, R.H., Rietzler, A.C., 2015. Efeitos tóxicos de arsênio em eisenia andrei em exposição a solos do entorno de minerações de ouro. *Rev. Bras. Cienc. do Solo* 39, 682–691.
- Arco-Lázaro, E., Pardo, T., Clemente, R., Bernal, M.P., 2018. Arsenic adsorption and plant availability in an agricultural soil irrigated with As-rich water: effects of Fe-rich amendments and organic and inorganic fertilisers. *J. Environ. Manag.* 209, 262–272.
- Baroni, F., Boscagli, A., Di Lella, L.A., Protano, G., Riccobono, F., 2004. Arsenic in soil and vegetation of contaminated areas in southern Tuscany (Italy). *J. Geochemical Explor.* 81, 1–14.
- Biswas, A., Petter, J., Neidhardt, H., Halder, D., Kundu, A.K., Chatterjee, D., Berner, Z., 2014. Role of competing ions in the mobilization of arsenic in groundwater of Bengal basin: insight from surface complexation modeling. *Water Res.* 55, 30–39.
- Borba, R.P., Figueiredo, B.R., Matschullat, J., 2003. Geochemical distribution of arsenic in waters, sediments and weathered gold mineralized rocks from Iron Quadrangle, Brazil. *Environ. Geol.* 4, 39–52.
- Bundschuh, J., Litter, M.I., Parvez, F., Román-Ross, G., Nicolli, H.B., Jean, J.S., Liu, C.W., López, D., Armienta, M.A., Guilherme, L.R.G., Cuevas, A.G., Cornejo, L., Cumbal, L., Toujaguez, R., 2012. One century of arsenic exposure in Latin America: a review of history and occurrence from 14 countries. *Sci. Total Environ.* 429, 2–35.
- Cai, Y., Cabrera, J.C., Georgiadis, M., Jayachandran, K., 2002. Assessment of arsenic mobility in the soils of some golf courses in South Florida. *Sci. Total Environ.* 291, 123–134.
- Campos, M.L., Guilherme, L.R.G., Lopes, R.S., Antunes, A.S., Marques, J.J.G. de S. e M., Curi, N., 2007. Teor e capacidade máxima de adsorção de arsênio em Latossolos brasileiros. *Rev. Bras. Ciência do Solo* 31, 1311–1318.
- Campos, M.L., Guilherme, L.R.G., Antunes, A.S., Borges, K.S.C., 2013a. Teor de arsênio e adsorção competitiva arsênio/fosfato e arsênio/sulfato em solos de Minas Gerais, Brasil. *Ciência Rural* 43, 985–991.
- Campos, M.L., Guilherme, L.R.G., Marques, J.J.G. de S.e.M., Curi, N., Araújo, A.S.A., Miquelluti, D.J., Lopes, C., Spiazzi, F.R., 2013b. Teores de arsênio e cádmio em solos do bioma Cerrado. *Rev. Bras. Cienc. do Solo* 37, 281–286.
- Carter, M.R., Gregorich, E.G., 2006. *Soil Sampling and Methods of Analysis*. 2^o. ed. CRC Press. Canadian Society of Soil Science (1224 p).
- Chatterjee, D., Adak, S., Banerjee, N., Bhattacharjee, P., Bandyopadhyay, A.K., Giri, A.K., 2018. Evaluation of health effects, genetic damage and telomere length in children exposed to arsenic in West Bengal, India. *Mutat. Res. - Genet. Toxicol. Environ. Mutagen.* 836, 82–88.
- Chilimba, A.D.C., Young, S.D., Black, C.R., Rogerson, K.B., Ander, E.L., Watts, M.J., Lammel, J., Broadley, M.R., 2011. Maize grain and soil surveys reveal suboptimal dietary selenium intake is widespread in Malawi. *Sci. Rep.* 1–9.
- Chowdhury, M.T.A., Deacon, C.M., Steel, E., Imamul Huq, S.M., Paton, G.I., Price, A.H., Williams, P.N., Meharg, A.A., Norton, G.J., 2018. Physiographical variability in arsenic dynamics in Bangladeshi soils. *Sci. Total Environ.* 612, 1365–1372.
- Cleasby, I.R., Nakagawa, S., 2011. Neglected biological patterns in the residuals. *Behav. Ecol. Sociobiol.* 65, 2361–2372.
- Conama, 2009. Normative No. 420 of December 28th, 2009. Valores Orientadores de Qualidade do Solo Quanto a Presença de Substâncias Químicas, Conselho Nacional de Meio Ambiente, 2009. Accessed Dec 03, 2018 from <http://www2.mma.gov.br/port/conama/legiabre.cfm?codlegi=620>.
- Da Silva, W.B., Périco, E., Dalzochio, M.S., Santos, M., Cajaiba, R.L., 2018. Are litterfall and litter decomposition processes indicators of forest regeneration in the neotropics? Insights from a case study in the Brazilian Amazon. *For. Ecol. Manag.* 429, 189–197.
- De Souza, J.J.L., Fontes, M.P.F., Gilkes, R., da Costa, L.M., de Oliveira, T.S., 2018. Geochemical signature of Amazon tropical rainforest soils. *Rev. Bras. Cienc. do Solo* 42, 1–18.
- Dixit, S., Hering, J.G., 2003. Comparison of arsenic (V) and arsenic (III) sorption onto iron oxide minerals: implications for arsenic mobility. *Environ. Sci. Technol.* 37, 4182–4189.
- Empresa Brasileira de Pesquisa Agropecuária – Embrapa, 1997. *Manual de métodos de análise de solo*. 2.ed. Rio de Janeiro. (212p).
- Fadini, P.S., Jardim, W.F., 2001. Is the Negro River Basin (Amazon) impacted by naturally occurring mercury? *Sci. Total Environ.* 275, 71–82.
- Fernandes, A.R., Souza, E.S. de, de Souza Braz, A.M., Birani, S.M., Alleoni, L.R.F., 2018. Quality reference values and background concentrations of potentially toxic elements in soils from the eastern Amazon, Brazil. *J. Geochemical Explor.* 190, 453–463.
- Fontes, M.P.F., Camargo, O.A. de, Sposito, G., 2001. Eletroquímica das partículas coloidais e sua relação com a mineralogia de solos altamente intemperizados. *Sci. Agric.* 58, 627–646.
- Gamboa-Loira, B., Cebrián, M.E., Franco-Marina, F., López-Carrillo, L., 2017. Arsenic metabolism and cancer risk: a meta-analysis. *Environ. Res.* 156, 551–558.
- Hammer, Ø., Harper, D.A.T., Ryan, P.D., 2001. *PAST: Paleontological statistics software package for education and data analysis*. *Palaeontol. Electron.* 4 (1), 9. http://palaeo-electronica.org/2001_1/past/issue1_01.htm.
- He, Q., Ren, Y., Mohamed, I., Ali, M., Hassan, W., Zeng, F., 2013. Assessment of trace and heavy metal distribution by four sequential extraction procedures in a contaminated soil. *Soil Water Res.* 8, 71–76.
- Ho, H.H., Swennen, R., Cappuyns, V., Vassilieva, E., Van Gerven, T., Van Tran, T., 2013. Speciation and mobility of selected trace metals (As, Cu, Mn, Pb and Zn) in sediment with depth in Cam River-Mouth, Haiphong, Vietnam. *Aquat. Geochemistry* 19, 57–75.
- Huang, R., Gao, S., Wang, W., Staunton, S., Wang, G., 2006. Soil arsenic availability and the transfer of soil arsenic to crops in suburban areas in Fujian Province, Southeast China. *Environ. Process.* 368, 531–541.
- Jiang, W., Hou, Q., Yang, Z., Zhong, C., Zheng, G., Yang, Z., Li, J., 2014. Evaluation of potential effects of soil available phosphorus on soil arsenic availability and paddy rice inorganic arsenic content. *Environ. Pollut.* 188, 159–165.
- Kabata-Pendias, A., 2011. *Trace Elements in Soils and Plants*. 4th ed. CRC Press LLC, USA.
- Kabata-Pendias, A., Mukherjee, A.B., 2007. *Trace Elements From Soil to Human*. Springer-Verlag, Berlin.
- Kapaj, S., Peterson, H., Liber, K., Bhattacharya, P., 2006. Human health effects from chronic arsenic poisoning-a review. *J. Environ. Sci. Heal.* 41, 2399–2428.
- Kim, K., Kim, J., Davis, A.P., 2003. Stabilization of available arsenic in highly contaminated mine tailings using iron. *Environ. Sci. Technol.* 37, 189–195.
- Langner, P., Mikutta, C., Kretzschmar, R., 2012. Arsenic sequestration by organic sulphur in peat. *Nat. Geosci.* 5, 66–73.
- Lenth, R.V., 2016. Least-squares means: the R package lsmeans. *J. Stat. Softw.* 69, 1–33.
- Liu, L., Li, W., Song, W., Guo, M., 2018. Remediation techniques for heavy metal-contaminated soils: principles and applicability. *Sci. Total Environ.* 633, 206–219.
- McBride, M.B., 1994. *Environmental Chemistry of Soils*. Oxford University, New York.
- Meliker, J.R., Wahl, R.L., Cameron, L.L., Nriagu, J.O., 2007. Arsenic in drinking water and cerebrovascular disease, diabetes mellitus, and kidney disease in Michigan: a standardized mortality ratio analysis. *Environ. Health* 6, 1–11.
- Melo, É.E.C., Guilherme, L.R.G., Nascimento, C.W.A., Penha, H.G.V., 2012. Availability and accumulation of arsenic in oilseeds grown in contaminated soils. *Water Air Soil Pollut.* 223, 233–240.
- Minatel, B.C., Sage, A.P., Anderson, C., Hubaux, R., Marshall, E.A., Lam, W.L., Martinez, V.D., 2018. Environmental arsenic exposure: from genetic susceptibility to pathogenesis. *Environ. Int.* 112, 183–197.
- Naujokas, M.F., Anderson, B., Ahsan, H., Vasken Aposhian, H., Graziano, J.H., Thompson, C., Suk, W.A., 2013. The broad scope of health effects from chronic arsenic exposure: update on a worldwide public health problem. *Environ. Health Perspect.* 121, 295–302.
- Oliveira, L.K. De, Melo, C.A., Goveia, D., Lobo, A., Aurora, M., Hernández, A., Fernandes, L., Rosa, A.H., 2015. Adsorption/desorption of arsenic by tropical peat: influence of organic matter, iron and aluminium. *Environ. Technol.* 36, 149–159.
- Ono, F.B., Guilherme, L.R.G., Penido, E.S., Carvalho, G.S., Hale, B., Toujaguez, R., Bundschuh, J., 2012. Arsenic bioaccessibility in a gold mining area: a health risk assessment for children. *Environ. Geochem. Health* 34, 457–465.
- Punshon, T., Jackson, B.P., Meharg, A.A., Warczak, T., Scheckel, K., Guerinot, M. Lou, 2017. Understanding arsenic dynamics in agronomic systems to predict and prevent uptake by crop plants. *Sci. Total Environ.* 581–582, 209–220.
- R Development Core Team, 2018. *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria (Version 3.4.4).
- Rahman, M.A., Rahman, A., Khan, M.Z.K., Renzaho, A.M.N., 2018. Human health risks and socio-economic perspectives of arsenic exposure in Bangladesh: a scoping review. *Ecotoxicol. Environ. Saf.* 150, 335–343.
- Raij, B.Van., Quaggio, J.A., 1983. Métodos de análise de solo para fins de fertilidade (Boletim Técnico, 81). Instituto Agrônomo, Campinas (31 p).
- Shahmoradi, S., Afyuni, M., Hajabbasi, M.A., 2017. Speciation, fractionation and plant availability of arsenic as induced by sorbents mixed with soil of Zarshuran (Iran). *Int. J. Environ. Sci. Technol.* 14, 767–776.
- Sharma, P., Ofner, J., Kappler, A., 2010. Formation of binary and ternary colloids and dissolved complexes of organic matter, Fe and As. *Environ. Sci. Technol.* 44, 4479–4485.
- Sharma, P., Rolle, M., Kocar, B.D., Fendorf, S., Kappler, A., 2011. Influence of natural organic matter on As transport and retention. *Environ. Sci. Technol.* 45, 546–553.
- Shoemaker, H.E., McLean, E.O., Pratt, P.F., 1961. Buffer methods for determining lime requirement of soils with appreciable amounts of extractable aluminum. *Soil Science Society of American Proceedings*, Madison 25, 274–277.
- Silva Junior, E.C., Wadt, L.H.O., Silva, K.E., Lima, R.M.B., Batista, K.D., Guedes, M.C., Carvalho, G.S., Carvalho, T.S., Reis, A.R., Lopes, G., Guilherme, L.R.G., 2017. Natural variation of selenium in Brazil nuts and soils from the Amazon region. *Chemosphere* 188, 650–658.
- Song, Y., Wang, S., Jia, Y., Yuan, Z., Wang, X., Gómez, M.A., 2015. Effects of nutrient and sulfate additions on As mobility in contaminated soils: a laboratory column study. *Chemosphere* 119, 902–909.
- Thinh, N., Van, Osanai, Y., Adachi, T., Thai, P.K., Nakano, N., Ozaki, A., Kuwahara, Y., Kato, R., Makio, M., Kurosawa, K., 2018. Chemical speciation and bioavailability concentration of arsenic and heavy metals in sediment and soil cores in estuarine ecosystem, Vietnam. *Microchem. J.* 139, 268–277.
- Tufano, K.J., Reyes, C., Saltikov, C.W., 2008. Reductive processes controlling arsenic retention: revealing the relative importance of iron and arsenic reduction. *Environ. Sci. Technol.* 42, 8283–8289.
- Van Herreweghe, S., Swennen, R., Vandecasteele, C., Cappuyns, V., 2003. Solid phase speciation of arsenic by sequential extraction in standard reference materials and industrially contaminated soil samples. *Environ. Pollut.* 122, 323–342.
- Wan, X., Dong, H., Feng, L., Lin, Z., Luo, Q., 2017. Comparison of three sequential extraction procedures for arsenic fractionation in highly polluted sites. *Chemosphere* 178, 402–410.
- Wang, S., Xu, L., Zhao, Z., Wang, S., Jia, Y., Wang, H., Wang, X., 2012. Arsenic retention and remobilization in muddy sediments with high iron and sulfur contents from a heavily contaminated estuary in China. *Chem. Geol.* 314–317, 57–65.

- Wang, J., Zeng, X., Zhang, H., Li, Y., Zhao, S., Su, S., Bai, L., Wang, Y., Zhang, T., 2018. Effect of exogenous phosphate on the lability and phytoavailability of arsenic in soils. *Chemosphere* 196, 540–547.
- Wenzel, W.W., Kirchbaumer, N., Prohaska, T., Stingeder, G., Lombi, E., Adriano, D.C., 2001. Arsenic fractionation in soils using an improved sequential extraction procedure. *Anal. Chim. Acta* 436, 309–323.
- WHO, 2001. Environmental Health Criteria 224, Arsenic and Arsenic Compounds. Inter-organization Programme for the Sound Management of Chemicals. Geneva.
- Zhang, L., Qin, X., Tang, J., Liu, W., Yang, H., 2017. Review of arsenic geochemical characteristics and its significance on arsenic pollution studies in karst groundwater, Southwest China. *Appl. Geochem.* 77, 80–88.