

## Review Article

# Genomics and Genetic Transformation in *Arachis*

Ana Cristina Miranda Brasileiro\*, Ana Claudia Guerra Araujo,  
Soraya Cristina Leal-Bertioli, Patricia Messenberg Guimarães

EMBRAPA Genetic Resources and Biotechnology, Brazil

## \*Corresponding author

Ana Cristina Miranda Brasileiro, Embrapa Genetic Resources and Biotechnology, Parque Estação Biológica (PqEB) – PO Box 02372, 70.770-917 - Brasília – DF, Brazil, Tel: + 55. 613448-4901; Fax: +55. 613340-3658; Email: ana.brasileiro@embrapa.br

Submitted: 07 April 2014

Accepted: 30 August 2014

Published: 02 September 2014

ISSN: 2333-6668

Copyright

© 2014 Miranda Brasileiro et al.

OPEN ACCESS

## Abstract

This review focus on *Arachis* functional and structural genomic studies, and their application in peanut improvement via plant transformation, that constitutes one of the most promising strategies to advance the introgression and modulation of important traits which can be translated into crop gain.

## Keywords

- Arachis
- Functional genomics
- Transgenic
- Genome
- Peanut

## ABBREVIATIONS

Mya: Million Years Ago; BAC: Bacterial Artificial Chromosome; LTR: Long Terminal Repeat; ESTs: Expressed Sequenced Tags; NGS: Next Generation Sequencing; SSH: Suppression Subtractive Hybridization; qRT-PCR: Quantitative Reverse Transcription PCR; DDPCR: Differential Display PCR; NCBI: National Center for Biotechnology Information; RGAs: Resistance Gene Analogs; NBS-LRR: Nucleotide Binding Site-Leucine Rich Region; HR: Hypersensitive Response; TFs: Transcription Factors; miRNA: Micro RNA; SSRs: Simple Sequence Repeats; GM: Genetically Modified; TE: Transpiration Efficiency; iRNA: Interference RNA

## INTRODUCTION

The main purpose of functional genomics is to assess, in parallel, the gene function of a large number, if not all, of genes contained in a given genome, by making use of the information provided by structural genomics [1]. For that, high-throughput methods are applied to both, the identification of which protein is encoded by each gene and, in a broader sense, deciphering the role of each gene in the overall function of the cell and organism.

Functional genomics is especially important in crops that lack a reference genome or with limited genomic information and a narrow genetic base, relying mostly in alien alleles for improvement of important traits. This is particularly true for peanut (*Arachis hypogaea* L.), as most sources of alleles related to the adaptability to adverse environments are frequently harbored by the cross-incompatible *Arachis* wild species.

The recent understanding on the genome structure of cultivated and wild *Arachis* has brought new insights for gene occurrence and evolution in the genome context. In parallel, functional genomics has elucidated expression profiling and regulation of *Arachis* genes in a global genome network context. Therefore, within a “genome to gene” approach, structural and functional genomic data integration will allow not only gene discovery, but the understanding of gene function, modulation

and their importance for the control of agronomical traits. This knowledge will facilitate and broaden the use of the wild *Arachis* extensive genetic diversity for peanut improvement, by either introgression using interspecific hybridization or transgenic approaches.

This review focus on *Arachis* functional and structural genomic studies, and their application in peanut improvement via plant transformation, that constitutes one of the most promising strategies to advance the introgression and modulation of important traits which can be translated into crop gain.

## The genus *arachis*

The genus *Arachis* has 80 described species [2] that includes *A. hypogaea*, the commercial peanut, also known as groundnut. It is a major oil seed crop, grown extensively in Africa, Asia and America, with a world annual production of around 38 million tons [3]. Besides its role in the confectionery industry, peanut represents a rich source of human food and animal livestock, containing 48% of oil, including unsaturated oil [4], 26% of protein, and 3% of fiber, added to calcium, thiamine and niacin (reviewed by [5]).

*A. hypogaea* is an allotetraploid species (AABB genome;  $2n = 4x = 40$ ) [6], most probably the result from a single hybridization event between two wild diploid species (*A. duranensis* and *A. ipaënsis*), followed by a rare and spontaneous chromosome doubling [7-9]. Due to its tetraploid character, peanut has been reproductively isolated from other wild species, which are mostly diploid, thus showing a limited genetic variability [10,11].

Wild diploid *Arachis* species ( $2n = 2x = 20$ ), which are in origin exclusive to South America [2], are more genetically diverse than the cultivated species [12]. Also, these species have been selected during evolution in a range of environments [12,13] and under innumerable abiotic and biotic stresses, constituting a rich source

of allele diversity potentially useful for genomic and genetic approaches targeting peanut improvement [14-17].

On the other hand, peanut, as other cultivated crops, is particularly susceptible to a number of constraints. Among infectious diseases, there are those caused by fungi, viruses, nematodes, bacteria, phytoplasm and the plant parasite *Alecta vogelli*. Susceptibility to some abiotic stresses such as drought, salinity, high temperature, herbicide toxicity and lack of nutrients are also included (reviewed by [18]).

Amid the pathogens, fungi are the largest group affecting the peanut crop, with the pathogenic species distributed in almost 50 different genera. Early (*Cercospora arachidicola*) and late (*Phaeoisariopsis personata*) leaf spots, along with peanut rust (*Puccinia arachidis*) are the main leaf diseases, responsible for production losses of almost 70%. Another important peanut fungus is *Aspergillus* spp., mostly *A. flavus* and *A. parasiticus* that produce, as secondary metabolites, a group of toxic compounds denominated aflatoxins during peanut seed development. Aflatoxins can cause necrosis of liver, cirrhosis and carcinoma in several animals and humans, and the preharvest aflatoxin contamination is considered a major economic problem for the peanut industry [19]. Other important diseases in peanut are caused by viruses, mainly PeMV (Peanut mottle virus) and GRV (Groundnut rosette virus). Nematodes, such as the root-knot (*Meloidogyne arenaria*, *M. javanica* and *M. hapla*) and root-lesion (*Pratylenchus brachyurus* and *P. coffeae*), are also major pathogens to the crop.

Concerning abiotic stresses, it should be recalled that peanut is cultivated mostly in semi-arid tropics (70% of its production), where water is usually a limiting factor. Therefore, drought and its associated problems, such as aflatoxin contamination, salinity and high temperatures, are considered the main abiotic stress for this crop.

Besides environmental and agronomic issues for peanut production, seeds and peanut-containing products are one of the major responsible for food allergenic reactions in children and adults worldwide. Some part of the world population is affected by reactions to peanut major allergens Arah1-9 (reviewed by [20]), which, in hypersensitive individuals, can cause death. The production of an allergen free or hypoallergenic peanut is one of the main goals for crop improvement.

### Arachis genome structure

Peanut is an underrepresented crop in terms of genome sequencing and structural genomics studies comparing to model plants or economically important crops, such as *Arabidopsis*, rice and soybean. Fortunately, a recent international joint initiative (The International Peanut Genome Initiative – IPGI; <http://www.peanutbioscience.com>) was launched to sequence the entire genome from *A. hypogaea* and its diploid parental *A. duranensis* and *A. ipaënsis*. This effort will increase the genome structural knowledge of *Arachis* genus as a whole.

Different studies have revealed important aspects of the peanut genome structure, such as its large size at 2.8 Gb [21]. *A. duranensis* and *A. ipaënsis* are estimated to have a similar size, with is almost half (1.3-1.4 Gb) of the peanut genome [22,23].

Cytological analysis indicated that the peanut karyotype is equivalent to the *A. duranensis* (A genome, having a small pair of chromosomes, ‘the A chromosomes’) [24] and *A. ipaënsis* (B genome, having symmetric karyotypes, but without ‘A chromosomes’) [22,23]. This has also been shown at molecular [25,26] and phylogenetic levels [11,27].

The A and B subgenomes of peanut have a high diploidized character, with no frequent rearrangements evidence in the tetraploid level, as it is only rarely that chromosomes from both genomes pair during meiosis, suggesting that A and B subgenomes might have diverged molecularly during evolution. Indeed, the estimated date of this divergence is 3.5 Mya [11,28]. Despite the indicative of molecular divergence, both A and B gene-spaces are likely to be ordered into ten conserved blocks, with genic sequences showing high identity [29]. From the diploid genetic maps [30,31], it is apparent that the order of molecular markers in the A and B genomes is mostly collinear, with only a few major rearrangements distinguishing them [32,33].

Regardless its recent origin, high transposon activities are detectable in the peanut genome. Indeed, the repetitive fraction of the peanut genome was estimated by the renaturation kinetics in around 64% [34]. Also, there are differences in the frequency and distribution of repetitive components in both subgenomes of peanut, as shown by *in situ* hybridizations probed with the retroelements FIDEL [35] and Matita [28].

As fundamental tools for structural genomic studies, BAC libraries of *A. duranensis* and *A. ipaënsis* were built aiming to exploit the ancestors’ peanut genomes and also for the characterization and isolation of genes of agronomical importance [36]. Analysis of 1.26 Mb of 12 genomic sequences from *A. duranensis* BAC clones [33], each clone containing genomic regions, revealed several repetitive sequences and 20 complete LTR retroelements, confirming that many of the repetitive DNA elements diverge in its origin and distribution (review by [37]). Most abundant LTR was FIDEL, a Ty3-*gypsy* type [35] with its truncated copies or solo LTRs, named Feral (non-autonomous retroelements). FIDEL and Feral were followed by Pipa, also a non-autonomous element, and a probable parasite on the retrotransposon named Pipoka. These four elements together, covered most of the analysed sequences, along with other less frequent LTRs such as Gordo, Mico and Curu [33]. Nonetheless, LTR retrotransposons further studies will certainly uncover a much larger world of retrotransposons than it is now known.

The repetitive sequences seem to occupy a substantial proportion of the peanut genome, mostly with some recent LTR retrotransposons and their non-autonomous elements, solo LTRs and nested retroelements [33]. This retroelements abundance is critical for the chromosomes structure and genome evolution. But also, it represents a difficulty for the upcoming sequencing and assembling of *Arachis* whole genome sequences, as well as for accurate genes annotation. A greater integration between *Arachis* structural and comparative genomics [29] will certainly help the unraveling of the occurrence, evolution, function and control of target genes.

### Arachis functional genomics for gene discovery

Efforts aiming peanut improvement have taken advantage

of the *Arachis* wild relatives, as the crosses of *A. monticola* with *A. hypogaea* [38], and the introgression of segments from *A. cardenasii* to the cultivars COAN and Nematan, resulting in resistance to root-knot nematode [39-41]. Nevertheless, the concern on the occurrence of an unpredictable gene drag has hindered the development of new cultivars using the inter specific hybridization approach.

Functional genomics and biotechnological techniques can be important allies for the strengthening of peanut breeding programs, as they enable the discover and characterization of genes of agronomic importance through deep analysis of the transcriptome, and their direct transference to chosen cultivars by plant transformation. Thus the transgenic approach circumvents the unsought genomic arrest and can prove to be very useful for the development of peanut varieties with high field performance and productivity.

Nevertheless, despite the considerable improvement on peanut genetic and genomic tools [42,43], few were the genes identified and isolated and even less those incorporated in peanut cultivars. As a matter of fact, only in the last decade, most transcriptome resources and gene discovery tools became available for *Arachis* [44].

The emerging of high-throughput sequencing technologies, which have granted an unprecedented methodological easiness and a dramatic decrease on costs, has allowed a substantial increase on *Arachis* functional genomics. The development of experimental approaches to assign gene functions in the genus has followed the trends of those applied for other more studied crops. It started with the building of ESTs (Expressed Sequence Tags) databanks for wild and cultivated species, followed by their rapid amplification due to the escalation on the production of expressed sequences by NGS technologies (Next Generation Sequencing). The contribution of some of these methods for the elucidation of the *Arachis* genome is discussed below.

### **cDNA libraries sequencing (ESTs) and microarrays**

The use of functional genomics for discovering genes which could confer new interesting traits in cultivated peanut is not an easy task due to several factors such as polyploidy, large genome size and abundance of repetitive sequences [33,45,46]. Therefore, the use of diploid wild relatives as source of genetic variability has a strategic role as they can offer new unexploited genes useful for the improvement of peanut cultivars.

Peanut, and its wild relatives, still have relatively little transcriptome information available when compared to major crops or model species. So far, the identification and studies on genes of agronomic interest have been quite limited to peanut: from the 254,541 ESTs currently available on public databases, 178,532 are from *A. hypogaea*; 35,291 from *A. duranensis*; 32,291 from *A. ipaënsis*; 6,264 from *A. stenosperma* and 750 from *A. magna*

Most of the initial functional studies in *Arachis* have concentrated on the generation of cDNA libraries and their subsequent partial (ESTs) or complete (full length cDNA) sequencing, which expressed sequences were used for select directly candidate genes or to produce oligonucleotide

microarrays. Some of these studies aimed at identifying plant genes preferentially expressed in specific organs or plant tissues. Wu et al. [47] used subtraction at mRNA level to identify uniquely expressed genes in peanut immature pod and leaf, whilst Li et al. [48] sequenced full-length cDNA libraries to compare the expression of acetyl-CoA carboxylase, an enzyme involved in fatty acid biosynthesis, in developing seeds and other tissues.

Others intended to analyze in parallel, the spatial (organ/tissue) distribution of the expression of multiple genes, as Bi et al. [49] and Payton et al. [50]. The first used microarrays to show diverse expression patterns of several genes in peanut seeds, leaves, stems, roots, flowers, and gynophores. The latter used ESTs available in public databanks as templates for probe design to create an oligonucleotide microarray containing unique probes from peanut. This microarray enabled the analysis of transcript levels in different tissues and organs aiming the identification of peanut genes found to be pod-specific and correlated to seed storage proteins, desiccation, oil production and cellular defense.

Oligonucleotide microarrays produced from peanut EST sequences were also developed by Chen et al. [51] to investigate the impact of artificial selection on expression diversity. By comparing gene expression profiles in pod and leaf of five widespread cultivars in southern China, they were able to show that differential expression owing to artificial selection, which probably led to low-expression polymorphisms in both tissues and cultivars, has contributed to expand the narrow genetic diversity in peanut.

Some other studies, using cDNA libraries or microarrays, have concentrated on the analysis of gene expression during incompatible and compatible interactions between peanut and its pathogens. These analyses included the spatial and temporal aspects of gene expression, for the identification of candidate resistance genes against fungi, nematodes and virus. This strategy was used to identify genes involved in resistance to late leaf spot (*P. personata*) in contrasting peanut genotypes [52], and to identify differentially regulated genes in response to aflatoxin-producing fungus *A. parasiticus* and drought stress [53,54]. Nobile et al. [55] used suppression subtractive hybridization (SSH) and differential screening of cDNA macroarrays to identify, in a resistant peanut genotype, unique ESTs involved in the early stages of *P. personata* pathogenesis. Differently expressed genes were also identified in peanut seeds infected with *A. parasiticus* [56], and in leaf tissues collected under natural occurrence of TSWV and leaf spot diseases (*C. arachidicola* and *P. personata*) [57]. Guo et al. [58] produced another oligonucleotide microarray representing over 40% of the protein coding genes in the peanut genome. They identified 22 putative *Aspergillus*-resistance genes that were constitutively up-expressed in the resistant cultivar which were homologous to corn and soybean genes, previously shown to confer resistance to fungal infection.

Gene discovery through functional genomics strategies were also used in less studied peanut diseases, such as those caused by nematodes and bacteria. Huang et al. [59] identified differentially expressed genes, including those involved in ethylene and jasmonic acid signal transduction pathways, from both peanut leaves and roots challenged with the bacterial pathogen *Ralstonia solanacearum* [59]. For the identification of



resistance genes against the root-knot nematode *M. arenaria* in peanut, Tirumalaraju et al. [60] constructed SSH libraries from challenged roots from near-isogenic lines from nematode-resistant and susceptible cultivars, NemaTAM and Florunner, respectively. A number of genes related to nematode infection response was identified and their transcriptional regulation pattern further analyzed via qRT-PCR (quantitative Reverse Transcription PCR).

Different genotypes of cultivated and wild *Arachis* harbor diverse mechanisms of response and adaptation to drought which might include the expression of a distinct set of genes that trigger changes in specific biochemical pathways. To gain a better understanding of peanut drought stress responses at molecular level, several studies have been carried out aiming the genomic analysis of stress-responsive genes. Differential Display PCR (DDPCR) and ESTs analyses were used to evaluate the transcriptome of different peanut genotypes and successfully identified drought stress responsive genes [61-63]. Su et al. [64] carried out an EST sequencing project using peanut seed cDNA library to isolate a Late Embryogenesis Abundant (LEA) gene, which is known to be accumulated in vegetative tissues in response to drought, salinity, and cold stress. Ranganayakulu et al. [65] analyzed the expression profile of drought tolerance candidate genes obtained from a subtracted cDNA library by macroarray and reverse Northern techniques. This data indicated that the adjustment in transcript levels of drought responsive genes in peanut was due to several key regulatory genes, such as dehydrins, Transcription Factors (TFs) and heat shock proteins, which are able to operate under successive levels of water loss process to confer desiccation tolerance. Recently, Dang et al. [66] compared TFs expression between contrasting peanut genotypes under drought conditions, and identified some highly induced transcripts containing DNA binding domains, which are also present in Ethylene Responsive Factors (ERF) family of TFs, and have already been related to drought stress.

Over the past years, transcriptome analysis by generation of ESTs played key role in providing resources for *in silico* gene discovery and annotation, as shown by Kumari et al. [67]. In this work, peanut ESTs already available in databases, coupled with a number of bioinformatics tools, were used to identify drought stress responsive genes in peanut. The strategy enabled the selection of putative candidate genes belonging to different families associated with drought stress responses, and also the 3D protein structure prediction of some of these genes. In the same way, Mondal et al. [68] have identified ESTs closely linked to rust (*P. arachidis*) Resistance (R) gene in peanut, by mining 5,184 *A. hypogaea* ESTs from NCBI database.

Resistance Gene Analogs (RGAs) account for the largest number of known disease R genes in plants. Although a great number of RGAs have been identified in peanut genome [69-71] only a small portion has been proved to be expressed. Radwan et al. [72] isolated several expressed RGAs from peanut which were available in public databases and developed R gene candidate (RGC) markers. This strategy has also been successfully applied by Liu et al. [73] that identified 385 putative RGAs from the public available peanut and wild *Arachis* ESTs database corresponding to six different classes of known plant disease R genes. These

peanut-expressed RGAs encoded for NBS-LRR proteins, protein kinases, LRR-PK/transmembrane proteins, toxin reductases, LRR-domain containing proteins and for TM-domain containing proteins. Both studies identified genes that can be used for future peanut functional genomics and genetic engineering researches.

Concerning *Arachis* wild species, the number of functional genomics studies has been quite restricted. A highly root-knot nematode resistant species, *A. stenosperma*, was used to produce the first wild *Arachis* ESTs databank, which comprised *M. arenaria* challenged and non-challenged infected roots [74]. Genes potentially related to *A. stenosperma* nematode resistance mediated by Hypersensitive Response (HR) were subsequently validated by Northern blot and DNA macroarrays [75]. More recently, Morgante et al. [76] extended the survey for *M. arenaria* resistance related genes in *A. stenosperma* by analyzing the expression profiles of 19 genes selected from this and other *Arachis* species using qRT-PCR. Sixteen genes were significantly differentially expressed in *A. stenosperma* infected roots, either involved in the HR displayed by this species [77] or production of secondary metabolites related to pathogen defense.

Besides constituting an important source of pathogen R genes, wild *Arachis* species show different transpiration behavior compared to cultivated peanut under gradual drought stress [78], and might constitute also important sources of genes related to mechanisms of drought tolerance. Therefore, the transcriptome of *A. magna* roots subjected to gradual water stress was analyzed by the construction of two SSH cDNA libraries [79,80]. *In silico* analysis revealed 13 candidate genes differentially expressed in response to drought stress that were subsequently validated by qRT-PCR.

## NEXT GENERATION SEQUENCING (NGS)

In the last decade, the advent of new technologies of mass parallel sequencing of nucleic acids known as NGS (Next Generation Sequencing) increased by several orders of magnitude the amount of sequences produced with unprecedented sequencing data for a number of plant species [81-83]. The use of NGS technologies has proven to be a powerful strategy, particularly in functional genomics, as it enables a comprehensive transcriptome analysis of induced changes in gene expression, allows the prediction of the roles and interactions of individual or correlated genes, and helps the elucidation of more complex signaling pathways activated in response to external stimuli [84]. Moreover, even when fully sequence genomes are not yet available, as in the case of peanut, *de novo* transcriptome assembly from NGS data can still be performed for the measurement of both absolute and differential expression. In fact, as previous studies have shown, there is a strong concordance between results obtained from microarrays and NGS data, with NGS showing higher sensitivity and dynamic range, coupled with lower technical variation [85]. For *Arachis*, with relatively few genomic resources available, the use of these new NGS technologies has enabled an enormous leap on the amount of data and genomic resources produced (molecular markers, genetic maps and candidate genes).

Chen et al. [86] used 454 sequencing technology (pyrosequencing) for the analysis of the aerial and subterranean pod transcriptome in peanut. Genes differentially regulated

in each tissue were identified, which are involved in embryo abortion in the aerial developing pod. RNA-seq technology was also used to profile the gene expression patterns from seeds in different developmental stages of high- and low-oil peanut varieties [87]. Over 59,000 unique sequences were produced and some were identified as being involved in lipid metabolism. This dataset allows a further understanding of the molecular basis of lipid biosynthesis in this crop, with identification of seven possible metabolic pathways involved in the accumulation of oil during seed development. Zhang et al. [88] employed RNA-seq technology to analyze the transcriptome of immature seeds of three peanut varieties, with different oil contents.

Wu et al. [89] also used RNA-seq technology to produce a comprehensive Unigene database that represents the whole genetic content of the Spanish peanut type. In both studies, a large number of Unigenes was assembled, and thousands of SSR markers identified. This data has the potential to facilitate gene discovery and functional genomics studies elucidating the complex response mechanisms of this allotetraploid species.

Likewise, the role of microRNAs (miRNA) in the cell, by affecting negatively gene expression at the post-transcriptional levels, has also been investigated in peanut by deep NGS sequencing [Zhao et al. [90] This study identified novel and conserved miRNA families from peanut which might play an important role in plant growth, development, and response to environmental stress.

NGS technology has also been applied for the identification of genes related to resistance to stresses in wild peanut relatives. Guimaraes et al [91] applied mass 454 technology sequencing on transcriptome analysis of *A. stenosperma*, which harbors resistances to a number of pests, challenged with the late leaf spot fungus (*P. personata*) and to *A. duranensis*, which shows conservative transpiration responses, submitted to water limited conditions. Almost 18,000 Unigenes were generated for *A. stenosperma* and 22,000 for *A. duranensis*, and 214 EST-SSRs polymorphic primers were identified. These databases of wild *Arachis* includes transcripts induced in both biotic and abiotic stresses, which constitute a rich source for gene discovery and molecular markers development in this underexploited wild germplasm. Transcriptome analysis of roots and developing seeds of two *A. duranensis* genotypes by NGS approach was used to construct a high-density genetic map based on *de novo* generated EST databases [92].

The availability of *Arachis* transcriptome data can lead the genus into a post-genomic era. However, in spite of the great numbers of expressed sequences and candidate genes identified in *Arachis*, few genes had their expression validated and their potential role in the control of traits of interest unraveled. In this scenario, elucidating genes biological functions is the main bottleneck and a challenge to the scientific community.

For that, *in vitro* gene validation using accurate gene-by-gene methods (eg. qRT-PCR) followed by studies on its modulation and phenotype associated effects in transgenic model plants are essential to untangle gene function.

### **Peanut genetic transformation**

In recent years, the development and commercialization of

Genetically Modified (GM) crops harbouring traits of agronomic interest is worldwide adopted (exceed 175 million hectares in 2013 [93]) and impacted significantly agriculture and food chain [94]. However, GM technology is yet restricted to few major crops, due to its high cost of development, risk assessment and commercialization.

The first transgenic plant of cultivated peanut was obtained in 1993 [95] and since then, different protocols have been described based on existing plant regeneration systems. Most of the protocols exploits the genotype-independent biolistic approach, using preferentially embryonic tissues as target [96,97]. Genotypes of the three major varietal-groups of peanut (Runner, Virginia and Spanish) have been successfully transformed using this system [98]. Furthermore, faster and improved protocols have been described using *Agrobacterium*-mediated transformation, increasing peanut transformation efficiency [99,100]. However, as a genotype-dependent approach, the majority of *Agrobacterium*-mediated transgenic peanut are only within Spanish and Valencia groups. In general, regardless of the system, peanut transformation efficiency is low and regeneration of transgenic plant is time-consuming. Therefore, the major limitations to produce transgenic peanut remain the difficulties for plant regeneration by tissue culture and selection of transgenic events.

Despite all limitations, transgenic peanut expressing genes that modulate different traits such as resistance to virus, insect and fungus, drought tolerance and grain quality have been developed by several research groups, particularly in United States, China and India. Nevertheless, none of these events are commercially available yet. These transgenic plants are under evaluation at different containment levels: *in vitro*, greenhouse and field conditions [97,98]. Only after that, these transgenes can be introgressed and used as sources of new alleles for peanut breeding programs.

### **Herbicide resistance**

The first successful transgenic peanut expressing a trait of agronomic interest was achieved in 1994 by introducing the *bar* gene to confers herbicide resistance [101]. The *bar* gene encodes the enzyme phosphinotricin acetyltransferase, isolated from *Streptomyces hygrosopicus*, widely used to confer full resistance to the active ingredient phosphinotricin of many broad-spectrum herbicides. The expression *bar* gene conferred to transgenic peanut resistance towards commercial formulations of this ingredient without plant damage. The further introduction of Bcl-xL gene in transgenic peanut also conferred tolerance to the herbicide paraquat, a bipyridilium herbicide [102]. Bcl-xL gene is a negative regulator of programmed cell death in humans and has a high potential to increase tolerance to biotic and abiotic stresses in plants. However, high levels of Bcl-xL gene expression seem to be deleterious for plant cells and not allow a proper growth and development of transgenic peanut.

### **Virus resistance**

TSWV is one of the most aggressive pathogens to several economically important crops, including peanut. As a strategy to prevent peanut infection by TSWV, the gene encoding a nucleocapsid protein N (N-gene) of TSWV strain isolated from

lettuce (*Lactuca sativa*) [101,103,104] and peanut [105] was introduced in *sense* and *antisense* directions, respectively, in peanut aiming to silence the expression of the virus protein. Three years of field evaluation have shown that N-gene silencing induced resistance to the virus (symptomless or delay in symptom development) and perhaps, to other tospoviruses in transgenic progenies [105,106]. Another N-gene of the genus *Tospovirus* (Peanut bud necrosis virus; PBNV) was also introduced and expressed in the *sense* direction in peanut [107]. However, greenhouse and field analyses of transgenic progeny indicated partial and non-durable resistance to PBNV.

Likewise, the expression of the gene coding for the coat protein of Peanut Stripe Virus (PStV) induced high levels of resistance in transgenic peanut lines challenged with PStV in greenhouse [108]. PStV is a potyvirus that imposes constraints on peanut production in Southeast Asia and China. Recently, Mehta et al., [109] expressed the coat protein of Tobacco Streak Virus (TSV) to successfully develop virus tolerant transgenic peanuts. TSV belongs to the genus *Ilarvirus* and is associated with the peanut stem necrosis disease, a new threat to peanut yield. The expression of genes coding for coat or nucleocapsid proteins or any other key virus protein for engineering resistance in plants is based on the concept of pathogen-derived resistance, by which plants expressing gene or sequence from the pathogen are protected against its deleterious effects, reviewed by [110].

### Insect resistance

In 1997, Singit et al., [111] had demonstrated the efficiency of chimeric *cryIA(c)* endotoxin gene of the soil bacterium *Bacillus thuringiensis* for the control of lesser cornstalk borer (*Elasmopalpus lignosellus*) larvae on transgenic peanut plants. This insect pest occurs widely in the western hemisphere, causing serious damage to economically important crops, including peanut. More recently, transgenic peanut plants expressing the chimeric protein of the  $\delta$ -endotoxin Cry1EC showed effective protection against the larvae of tobacco cutworm (*Spodoptera litura*) in leaf feeding bioassays [112]. Cry1EC is a synthetic  $\delta$ -endotoxin modified from *B. thuringiensis* for high-level expression in plants. It has insecticidal properties against *S. litura*, an insect pest that causes serious damage to peanut crops in South India, with yield losses up to 71%.

### Fungal resistance

Transgenic peanut plants with enhanced resistance to fungal diseases have been obtained by over expression of antifungal genes, in particular those encoding Pathogenesis-Related (PR) proteins. Over expression of chitinase, a tobacco (*Nicotiana tabacum*) PR protein was successfully used to achieve resistance to *C. arachidicola*, the causal agent of early leaf spot disease in peanut [113]. Recently, peanut plants over expressing *Rchit*, a chitinase gene from rice (*Oryza sativa*), displayed a broad range resistance to the three most economically important fungal pathogens for peanut: the foliar fungi that caused late leaf spot and leaf rust diseases (*P. personata* and *P. arachidis*, respectively) and the aflatoxin-producing fungus *A. flavus*, that dramatically affects the peanut quality making it unfit for consumption [114]. Overexpression of chitinase from different organisms in transgenic plants as a strategy to confer resistance to fungal

diseases has been widely described in the literature (reviewed by [115]). Chitinase antifungal activity seems to be related to the degradation of chitin hyphae, slowing of fungus growth, and pathogen elicitor release, that induces defence response signalling in plants.

Transgenic peanut plants also displayed increased resistance to leaf spot diseases (*P. personata* and *C. arachidicola*) by overexpressing another class of PR protein, a defensin from mustard (*Brassica juncea*) [116]. Defensins are cysteine-rich antimicrobial peptides which protects seeds against pathogen attack by altering fungal membrane permeability and/or inhibiting macromolecules biosynthesis [117]. The co-expression of two other classes of PR proteins, an osmotin-like from black nightshade (*Solanum nigrum*) and an antifungal protein-2 from radish (*Raphanus sativus*) also conferred resistance to *P. personata* in transgenic peanut [118]. Resistance to *A. flavus* and *C. arachidicola* was also obtained by overexpression of another class of PR protein, a  $\beta$  1-3 Glucanase from tobacco [119]. As chitinase, glucanase has also been widely used for improving fungal disease resistance in several plant species (reviewed by [115]). The defence protein glucanase catalyses the hydrolysis of glucan, an important cell wall component of most filamentous fungi and, when overexpressed, enhances fungal resistance in host. Transgenic peanut plants expressing a rice chitinase and/or an alfalfa (*Medicago sativa*) glucanase evaluated in fields over three years period demonstrated consistent increased resistance to Sclerotinia blight compared to the susceptible cultivar [120]. Sclerotinia blight is caused by the soil-borne fungus *Sclerotinia minor* and is one of the most destructive diseases of peanut.

In addition to PR proteins, other antifungal proteins have been introduced in transgenic peanut plants, such as oxalate oxidase from barley (*Hordeum vulgare*) that conferred Sclerotinia blight resistance [121]. Evaluation of these plants over five years in naturally infested fields with *S. minor* has shown a high level of resistance [122]. Oxalate oxidase catalyzes the degradation of oxalic acid that induces early oxidative burst in plant cell and can trigger the activation of plant defense response mechanism and the production of antimicrobial metabolites [117]. Thus, overexpression of oxalate oxidase can reduce the predisposition of cells to fungal infection, preventing the disease development.

The gene encoding nonheme chloroperoxidase (*cpo-p*) from *Pseudomonas pyrrocinia* was introduced in peanut and inhibited *A. flavus* hyphal growth [123]. Although the exact mechanism is not determined, *cpo-p* is described as a potent candidate gene to reduce preharvest aflatoxin contamination.

### Drought tolerance

The first report describing drought-tolerant peanut transgenic plants was in 2007, with the expression of the gene coding the transcription factor DREB1A, under the control of an abiotic stress inducible promoter (rd29A), both from *Arabidopsis thaliana* [124]. DREB1A is a transcription factor that binds to the dehydration-responsive element (DRE) located in the promoter region of stress-responsive genes, thereby activates its expression in response to water deficit. The DREB1A-expressing peanuts showed similar phenotype to well-watered control plants and an increased transpiration efficiency (TE) when exposed to soil



drying conditions. TE is an important component of water use efficiency and of plant performance under drought conditions. However, subsequent studies of antioxidative responses of those transgenic lines were unable to provide an explanation for its higher TE performance [125]. Recent analyzes, in a lysimeter system that mimics field conditions, showed that rooting growth under water-limited conditions is higher in transgenic lines than in controls, particularly in deep soil layers, leading to an increase in water extraction [126].

Transgenic peanut plants expressing isopentenyltransferase (*ipt*) gene isolated from *Agrobacterium tumefaciens*, under the control of the drought-inducible promoter SARK from bean (*Phaseolus vulgaris*), showed also increased tolerance to drought [127]. IPT is a key enzyme in the biosynthesis of cytokinins and plays a critical role in the control of cytokinin levels. Increased cytokinin levels leads to hormone homeostasis changing and robust root development, which can enhanced the water absorption, even under water-limited conditions. Transgenic plants evaluated in field for two years produced significantly higher yields than controls in dry land conditions.

### Salinity tolerance

Banjara et al. [128] demonstrated that the overexpression of AtNHX1 gene could confer high salt tolerance in peanut transgenic plants. AtNHX1 antiporter protein is the most abundant vacuole Na<sup>+</sup>/H<sup>+</sup> antiporter in *A. thaliana*. Its overexpression increases the ability to sequester sodium into vacuoles, thus reducing its toxicity in the cytosol, favouring water uptake by root cells, and improving tissue water retention under stress. Peanut plants expressing AtNHX1 gene driven by 35S promoter displayed better growth and performance when compared to controls, at levels up to 150 mM NaCl.

### Drought and salinity tolerance

Although drought and salinity tolerance are complex multigenic traits, a same single gene introduced into several crops, including peanut, can displayed simultaneously, tolerance to multiple abiotic stresses [129]. Indeed, overexpression of AtNHX1 gene driven by 35S promoter provided to transgenic peanut plants not only enhanced tolerance to high salt concentrations but also to water deprivation [130]. Transgenic lines also accumulated more salt and proline in leaves under salt and drought stress conditions. Likewise, overexpression of AVP1 also increased simultaneously salinity and drought tolerance in transgenic peanuts [131]. AVP1 encodes an H<sup>+</sup> pyrophosphatase with a proton pump activity on vacuole membrane and its overexpression can increase secondary transporters activity, such as the AtNHX1 Na<sup>+</sup>/H<sup>+</sup> antiporter, thereby decreasing vacuoles water potential leading to increased tolerance. AVP1-expressing transgenic peanuts plants outperformed wild-types under reduced irrigation or high saline conditions. Recently, the expression of *mtlD* gene from *Escherichia coli* in peanut plants transgenic lines demonstrated an enhanced tolerance to salinity and drought [132]. The bacterial *mtlD* gene encodes the enzyme Mannitol 1-Phosphatase Dehydrogenase that converts mannitol 1-phosphate to mannitol which is accumulated in transgenic tissues.

### Grain quality

Arah2 and Arah6 (conglutins) are peanut seed storage proteins recognized as allergens by more than 90% of hyper sensible individuals, eliciting IgE-mediated response and causing severe reactions. Therefore, strategies to reduce peanut seed allergens are one of priorities of peanut grain quality improvement and food industry. Gene silencing using the RNA interference (RNAi) strategy allowed the reduction of Arah2 and Arah6 content in transgenic peanut seeds [133,134]. Further studies of RNA-silenced transgenic lines showed changes in the allergen seed content including Ara h10 (oleosin 1), 13-lipoxygenase and Ahy-3 (arachin), as well as the decrease of conarachin [135].

Although there is a considerable number of genes already transferred to peanut by plant transformation, the only example of peanuts expressing genes/sequences isolated from the genetic pool of the genus *Arachis* is Arah-silencing plants. This is mainly due to the previous restricted knowledge on their genomes and genes availability. The recent increase on wild genomes data will hopefully enable that the full potential of these species, as resistance and tolerance genes donors, can be fulfilled.

The integration of genomic studies, along with their potential biotechnological applications (e.g molecular markers, genes and promoters, transgenic plants), constitutes an important alternative for the development of cultivars with traits of interest to both, the producer and the consumer.

### ACKNOWLEDGEMENTS

This work was funded by the host institution, FAP/DF; CAPES and CNPq (Brazil) and Generation Challenge Program (GCP-CGIAR

### REFERENCES

1. Bouchez D, Höfte H. Functional genomics in plants. *Plant Physiol.* 1998; 118: 725-732.
2. Valls J, Simpson C. New species of *Arachis* L. (Leguminosae) from Brazil, Paraguay and Bolivia. *Bonplandia* (Argentina). 2005; 14: 35-64.
3. USDA-FAS. Major Oilseeds: World Supply and Distribution (Commodity View). 2013.
4. Holbrook C, Stalker H. Peanut breeding and genetic resources. *Plant Breeding Rev.* 2003; 22: 297-355.
5. Sarvamangala C, Gowda MVC, Varshney RK. Identification of quantitative trait loci for protein content, oil content and oil quality for groundnut (*Arachis hypogaea* L.). *Field Crops Res.* 2011; 122: 49-59.
6. Smartt J, Stalker H. Speciation and cytogenetics in *Arachis*. *Peanut Sci Technol.* 1982: 21-49.
7. Kochert G, Halward T, Branch WD, Simpson CE. RFLP variability in peanut (*Arachis hypogaea* L.) cultivars and wild species. *Theor Appl Genet.* 1991; 81: 565-570.
8. Stalker H, Simpson C. Germplasm resources in *Arachis*. *Adv Peanut Sci.* 1995; 14-53.
9. Grabielle M, Chalup L, Robledo G, Seijo G. Genetic and geographic origin of domesticated peanut as evidenced by 5S rDNA and chloroplast DNA sequences. *Plant Syst Evol.* 2012; 298: 1151-1165.
10. Tang RH, Zhuang WJ, Gao GQ, He LQ, Han ZQ, Shan SH, et al.

- Phylogenetic relationships in genus *Arachis* based on SSR and AFLP markers. *Agri Sci China*. 2008; 7: 405-414.
11. Moretzsohn MC, Gouvea EG, Inglis PW, Leal-Bertioli SC, Valls JF, Bertioli DJ. A study of the relationships of cultivated peanut (*Arachis hypogaea*) and its most closely related wild species using intron sequences and microsatellite markers. *Ann Bot*. 2013; 111: 113-126.
  12. Krapovickas A, Gregory W. Taxonomia del genero *Arachis* (Leguminosae). *Bonplandia*. 1994; 8:1-186.
  13. Freitas FO, Moretzsohn MC, Valls JF. Genetic variability of Brazilian Indian landraces of *Arachis hypogaea* L. *Genet Mol Res*. 2007; 6: 675-684.
  14. Nelson SC, Simpson CE, Starr JL. Resistance to *Meloidogyne arenaria* in *Arachis* spp. *Germplasm. J Nematol*. 1989; 21: 654-660.
  15. Leal-Bertioli SC, José AC, Alves-Freitas DM, Moretzsohn MC, Guimarães PM, Nielen S, et al. Identification of candidate genome regions controlling disease resistance in *Arachis*. *BMC Plant Biol*. 2009; 9: 112.
  16. Upadhyaya HD, Sharma S, Dwivedi SL. *Arachis*. *Wild Crop Relatives: Genomic and Breeding Resources: Kole C, editor. In: Springer Berlin Heidelberg; 2011; 1-19.*
  17. Stalker HT, Tallury SP, Ozias-Akins P, Bertioli D, Leal-Bertioli SCM. The value of diploid peanut relatives for breeding and genomics. *Peanut Sci*. 2013; 40: 70-88.
  18. Pio-Ribeiro G, Pereira GA, Moraes SA, Filho PdAM. Principais doenças do amendoim e seu controle. In: Santos RC, Freire RMM, Lima LM, editors. *O Agronegócio do Amendoim no Brasil. Brasília -DF: Embrapa; 2013; 585.*
  19. Klich MA. *Aspergillus flavus*: the major producer of aflatoxin. *Mol Plant Pathol*. 2007; 8: 713-722.
  20. Koppelman SJ, Hefle SL, Taylor SL, de Jong GA. Digestion of peanut allergens Ara h 1, Ara h 2, Ara h 3, and Ara h 6: a comparative in vitro study and partial characterization of digestion-resistant peptides. *Mol Nutr Food Res*. 2010; 54: 1711-1721.
  21. Greilhuber J. Intraspecific variation in genome size in angiosperms: identifying its existence. *Ann Bot*. 2005; 95: 91-98.
  22. Seijo G, Lavia GI, Fernández A, Krapovickas A, Ducasse DA, Bertioli DJ, Moscone EA. Genomic relationships between the cultivated peanut (*Arachis hypogaea*, Leguminosae) and its close relatives revealed by double GISH. *Am J Bot*. 2007; 94: 1963-1971.
  23. Seijo JG, Lavia GI, Fernández A, Krapovickas A, Ducasse D, Moscone EA. Physical mapping of the 5S and 18S-25S rRNA genes by FISH as evidence that *Arachis duranensis* and *A. ipaensis* are the wild diploid progenitors of *A. hypogaea* (Leguminosae). *Am J Bot*. 2004; 91: 1294-1303.
  24. Husted L. Cytological studies on the peanut, *Arachis*. II. Chromosome number, morphology and behavior, and their application to the problem of the origin of the cultivated forms. *Cytologia*. 1936; 7: 396-422.
  25. Halward T, Stalker H, Larue E, Kochert G. Genetic variation detectable with molecular markers among unadapted germplasm resources of cultivated peanut and related wild species. *Genome*. 1991; 34: 1013-20.
  26. Milla SR, Isleib TG, Stalker HT. Taxonomic relationships among *Arachis* sect. *Arachis* species as revealed by AFLP markers. *Genome*. 2005; 48: 1-11.
  27. Burow MD, Simpson CE, Faries MW, Starr JL, Paterson AH. Molecular biogeographic study of recently described B- and A-genome *Arachis* species, also providing new insights into the origins of cultivated peanut. *Genome*. 2009; 52: 107-119.
  28. Nielen S, Vidigal BS, Leal-Bertioli SC, Ratnaparkhe M, Paterson AH, Garsmeur O, D'Hont A. Matita, a new retroelement from peanut: characterization and evolutionary context in the light of the *Arachis* A-B genome divergence. *Mol Genet Genomics*. 2012; 287: 21-38.
  29. Bertioli DJ, Moretzsohn MC, Madsen LH, Sandal N, Leal-Bertioli SC, Guimarães PM, Hougaard BK. An analysis of synteny of *Arachis* with *Lotus* and *Medicago* sheds new light on the structure, stability and evolution of legume genomes. *BMC Genomics*. 2009; 10: 45.
  30. Moretzsohn MC, Barbosa AV, Alves-Freitas DM, Teixeira C, Leal-Bertioli SC, Guimarães PM, Pereira RW. A linkage map for the B-genome of *Arachis* (Fabaceae) and its synteny to the A-genome. *BMC Plant Biol*. 2009; 9: 40.
  31. Moretzsohn MC, Leoi L, Proite K, Guimarães PM, Leal-Bertioli SC, Gimenes MA, Martins WS. A microsatellite-based, gene-rich linkage map for the AA genome of *Arachis* (Fabaceae). *Theor Appl Genet*. 2005; 111: 1060-1071.
  32. Burow MD, Simpson CE, Starr JL, Paterson AH. Transmission genetics of chromatin from a synthetic amphidiploid to cultivated peanut (*Arachis hypogaea* L.). broadening the gene pool of a monophyletic polyploid species. *Genetics*. 2001; 159: 823-837.
  33. Bertioli DJ, Vidigal B, Nielen S, Ratnaparkhe MB, Lee TH, Leal-Bertioli SC, Kim C. The repetitive component of the A genome of peanut (*Arachis hypogaea*) and its role in remodelling intergenic sequence space since its evolutionary divergence from the B genome. *Ann Bot*. 2013; 112: 545-559.
  34. Dhillon SS, Rake AV, Miksche JP. Reassociation Kinetics and Cytophotometric Characterization of Peanut (*Arachis hypogaea* L.) DNA. *Plant Physiol*. 1980; 65: 1121-1127.
  35. Nielen S, Campos-Fonseca F, Leal-Bertioli S, Guimarães P, Seijo G, Town C, Arrial R. FIDEL-a retrovirus-like retrotransposon and its distinct evolutionary histories in the A- and B-genome components of cultivated peanut. *Chromosome Res*. 2010; 18: 227-246.
  36. Guimarães PM, Garsmeur O, Proite K, Leal-Bertioli SC, Seijo G, Chaine C, Bertioli DJ. BAC libraries construction from the ancestral diploid genomes of the allotetraploid cultivated peanut. *BMC Plant Biol*. 2008; 8: 14.
  37. Bertioli D, Araujo A, Nielen S, Heslop-Harrison P, Guimaraes P, Schwarzacher T, et al. An overview of peanut genome structure. In: Varshney R, editor. *Genetics, Genomics and Breeding of Peanuts: in press*. 2014.
  38. Isleib T, Wynne J. Use of plant introductions in peanut improvement. In: Shands H, editor. *Use of Plant Introductions in Cultivar Development Volume 2*. Madison: Crop Science Society of America; 1992; 75-116.
  39. Simpson CE, Starr JL. Registration of 'COAN' peanut appreciation is expressed to the Texas Peanut Producers Board for their generous support of this research from 1988 to the present time. Registration by CSSA. *Crop Sci*. 2001; 4: 918.
  40. Simpson CE, Starr JL, Church GT, Burow MD, Paterson AH. Registration of 'NemaTAM' peanut registration by CSSA. *Crop Sci*. 2003; 43: 1561-.
  41. Starr JL, Morgan ER, Simpson CE. Management of the peanut root-knot nematode, *Meloidogyne arenaria*, with host resistance. *Online Plant Health Progress* 2002.
  42. Pandey MK, Monyo E, Ozias-Akins P, Liang X, Guimarães P, Nigam SN, et al. Advances in *Arachis* genomics for peanut improvement. *Biotechnol Adv*. 2012; 30: 639-651.
  43. Varshney RK, Mohan SM, Gaur PM, Gangarao NV, Pandey MK, Bohra A, Sawargaonkar SL. Achievements and prospects of genomics-assisted breeding in three legume crops of the semi-arid tropics. *Biotechnol Adv*. 2013; 31: 1120-1134.



44. Varshney RK, Glaszmann JC, Leung H, Ribaut JM. More genomic resources for less-studied crops. *Trends Biotechnol.* 2010; 28: 452-460.
45. Burrow MD, Selvaraj MG, Upadhyaya H, Ozias-Akins P, Guo B, Bertoli DJ, et al. Genomics of peanut, a major source of oil and protein. *Genomics of Tropical Crop Plants: Springer*; 2008; 421-40.
46. Stalker HT, Weissinger AK, Milla-Lewis S, Holbrook CC. Genomics: An evolving science in peanut. *Peanut Sci.* 2009; 36: 2-10.
47. Wu N, Matand K, Williams S. A novel mRNA level subtraction method for quick identification of target-orientated uniquely expressed genes between peanut immature pod and leaf. *Biol Proced Online.* 2010; 12: 44-55.
48. Li M-J, Xia H, Zhao C-Z, Li A-Q, Li C-S, Bi Y-P, et al. Isolation and characterization of putative Acetyl-CoA Carboxylases in *Arachis hypogaea* L. *Plant Mol Biol Rep.* 2010; 28: 58-68.
49. Bi YP, Liu W, Xia H, Su L, Zhao CZ, Wan SB, Wang XJ. EST sequencing and gene expression profiling of cultivated peanut (*Arachis hypogaea* L.). *Genome.* 2010; 53: 832-839.
50. Payton P, Kottapalli KR, Rowland D, Faircloth W, Guo B, Burrow M, Puppala N. Gene expression profiling in peanut using high density oligonucleotide microarrays. *BMC Genomics.* 2009; 10: 265.
51. Chen X, Hong Y, Zhang E, Liu H, Zhou G, Li S, et al. Comparison of gene expression profiles in cultivated peanut (*Arachis hypogaea*) under strong artificial selection. *Plant Breeding.* 2012; 131: 620-30.
52. Luo M, Dang P, Bausher MG, Holbrook CC, Lee RD, Lynch RE, Guo BZ. Identification of Transcripts Involved in Resistance Responses to Leaf Spot Disease Caused by *Cercosporidium personatum* in Peanut (*Arachis hypogaea*). *Phytopathology.* 2005; 95: 381-387.
53. Luo M, Liang XQ, Dang P, Holbrook CC, Bausher MG, Lee RD, et al. Microarray-based screening of differentially expressed genes in peanut in response to *Aspergillus parasiticus* infection and drought stress. *Plant Sci.* 2005; 169: 695-703.
54. Liang XQ, Holbrook CC, Lynch RE, Guo BZ. beta-,3-Glucanase Activity in Peanut Seed (*Arachis hypogaea*) is Induced by Inoculation with *Aspergillus flavus* and Copurifies with a Conglutin-Like Protein. *Phytopathology.* 2005; 95: 506-511.
55. Nobile PM, Lopes CR, Barsalobres-Cavallari C, Quecini V, Coutinho LL, Hoshino AA, et al. Peanut genes identified during initial phase of *Cercosporidium personatum* infection. *Plant Sci.* 2008; 174: 78-87.
56. Guo B, Chen X, Dang P, Scully BT, Liang X, Holbrook CC, Yu J. Peanut gene expression profiling in developing seeds at different reproduction stages during *Aspergillus parasiticus* infection. *BMC Dev Biol.* 2008; 8: 12.
57. Guo B, Chen X, Hong Y, Liang X, Dang P, Brennehan T, Holbrook C. Analysis of Gene Expression Profiles in Leaf Tissues of Cultivated Peanuts and Development of EST-SSR Markers and Gene Discovery. *Int J Plant Genomics.* 2009; 2009: 715605.
58. Guo B, Fedorova ND, Chen X, Wan CH, Wang W, Nierman WC, Bhatnagar D. Gene expression profiling and identification of resistance genes to *Aspergillus flavus* infection in peanut through EST and microarray strategies. *Toxins (Basel).* 2011; 3: 737-753.
59. Huang J, Yan L, Lei Y, Jiang H, Ren X, Liao B. Expressed sequence tags in cultivated peanut (*Arachis hypogaea*): discovery of genes in seed development and response to *Ralstonia solanacearum* challenge. *J Plant Res.* 2012; 125: 755-769.
60. Tirumalaraju SV, Jain M, Gallo M. Differential gene expression in roots of nematode-resistant and -susceptible peanut (*Arachis hypogaea*) cultivars in response to early stages of peanut root-knot nematode (*Meloidogyne arenaria*) parasitization. *J Plant Physiol.* 2011; 168: 481-92.
61. Devaiah KM, Bali G, Athmaram TN, Basha MS. Identification of two new genes from drought tolerant peanut up-regulated in response to drought. *Plant Growth Regul.* 2007; 52: 249-58.
62. Govind G, Harshavardhan VT, Patricia JK, Dhanalakshmi R, Senthil Kumar M, Sreenivasulu N, et al. Identification and functional validation of a unique set of drought induced genes preferentially expressed in response to gradual water stress in peanut. *Mol Genet Genomics.* 2009; 281: 591-605.
63. Guo BZ, Yu J, Holbrook CC, Lee RD, Lynch RE. Application of differential display RT-PCR and EST/microarray technologies to the analysis of gene expression in response to drought stress and elimination of aflatoxin contamination in corn and peanut. *Toxin Rev.* 2003; 22(2-3): 287-312.
64. Su L, Zhao CZ, Bi YP, Wan SB, Xia H, Wang XJ. Isolation and expression analysis of LEA genes in peanut (*Arachis hypogaea* L.). *J Biosci.* 2011; 36: 223-228.
65. Ranganayakulu GS, Chandraobulreddy P, Thippeswamy M, Veeranagamallaiah G, Sudhakar C. Identification of drought stress-responsive genes from drought-tolerant groundnut cultivar (*Arachis hypogaea* L. cv K-134) through analysis of subtracted expressed sequence tags. *Acta Physiol Plant.* 2012; 34: 361-77.
66. Dang PM, Chen CY, Holbrook CC. Evaluation of five peanut (*Arachis hypogaea*) genotypes to identify drought responsive mechanisms utilising candidate-gene approach. *Funct Plant Biol.* 2013; 40: 1323-33.
67. Kumari A, Kumar A, Wany A, Prajapati GK, Pandey DM. Identification and annotation of abiotic stress responsive candidate genes in peanut ESTs. *Bioinformatics.* 2012; 8: 1211-1219.
68. Mondal S, Badigannavar AM, D'Souza SF. Development of genic molecular markers linked to a rust resistance gene in cultivated groundnut (*Arachis hypogaea* L.). *Euphytica.* 2012; 188: 163-73.
69. Zheng Y-x, Li C-j, Liu Y, Yan C-x, Zhang T-t, Zhuang W-j, et al. Cloning and characterization of a NBS-LRR resistance gene from peanut (*Arachis hypogaea* L.). *J Agr Sci (1916-9752).* 2012; 4.
70. Yuksel B, Estill JC, Schulze SR, Paterson AH. Organization and evolution of resistance gene analogs in peanut. *Mol Genet Genomics.* 2005; 274: 248-263.
71. Bertoli DJ, Leal-Bertoli SC, Lion MB, Santos VL, Pappas G Jr, Cannon SB, Guimarães PM. A large scale analysis of resistance gene homologues in *Arachis*. *Mol Genet Genomics.* 2003; 270: 34-45.
72. Radwan OE, Ahmed TA, Knapp SJ. Phylogenetic analyses of peanut resistance gene candidates and screening of different genotypes for polymorphic markers. *Saudi J Biol Sci.* 2010; 17: 43-49.
73. Liu Z, Feng S, Pandey MK, Chen X, Culbreath AK, Varshney RK, Guo B. Identification of expressed resistance gene analogs from peanut (*Arachis hypogaea* L.) expressed sequence tags. *J Integr Plant Biol.* 2013; 55: 453-461.
74. Proite K, Leal-Bertoli SC, Bertoli DJ, Moretzsohn MC, da Silva FR, Martins NF, Guimarães PM. ESTs from a wild *Arachis* species for gene discovery and marker development. *BMC Plant Biol.* 2007; 7: 7.
75. Guimarães PM, Brasileiro AC, Proite K, Araujo AC, Leal-Bertoli SC, Pic-Taylor A, et al. A study of gene expression in the nematode resistant wild peanut relative, *Arachis stenosperma*, in response to challenge with *Meloidogyne arenaria*. *Trop Plant Biol.* 2010: 1-10.
76. Morgante CV, Brasileiro ACM, Roberts PA, Guimarães LA, Araujo ACG, Fonseca LN, et al. A survey of genes involved in *Arachis stenosperma*

- resistance to *Meloidogyne arenaria* race 1. *Funct Plant Biol.* 2013; 40(12): 1298-309.
77. Proite K, Carneiro R, Falcao R, Gomes A, Leal-Bertioli S, Guimaraes P, et al. Post-infection development and histopathology of *Meloidogyne arenaria* race 1 on *Arachis* spp. *Plant Pathology.* 2008; 57: 974-80.
78. Leal-Bertioli SCM, Bertioli DJ, Guimaraes PM, Pereira TD, Galhardo I, Silva JP, et al. The effect of tetraploidization of wild *Arachis* on leaf morphology and other drought-related traits. *Environ Exp Bot.* 2012; 84: 17-24.
79. Morgante CV, Guimarães PM, Martins AC, Araújo AC, Leal-Bertioli SC, Bertioli DJ, et al. Reference genes for quantitative reverse transcription-polymerase chain reaction expression studies in wild and cultivated peanut. *BMC Res Notes.* 2011; 4: 339.
80. Brasileiro A, Morgante C, Leal-Bertioli S, Araujo A, Silva A, Martins A, et al. Identificação de genes associados à resposta ao estresse hídrico em espécies silvestres de *Arachis*. *Heringeriana.* 2012; 6: 35-6.
81. Morozova O, Marra MA. Applications of next-generation sequencing technologies in functional genomics. *Genomics.* 2008; 92: 255-264.
82. Varshney RK, Roorkiwal M, Nguyen HT. Legume genomics: From genomic resources to molecular breeding. *Plant Genome.* 2013; 6.
83. Edwards D, Batley J, Snowden RJ. Accessing complex crop genomes with next-generation sequencing. *Theor Appl Genet.* 2013; 126: 1-11.
84. Kim KH, Kang YJ, Kim DH, Yoon MY, Moon JK, Kim MY, Van K. RNA-Seq analysis of a soybean near-isogenic line carrying bacterial leaf pustule-resistant and -susceptible alleles. *DNA Res.* 2011; 18: 483-497.
85. Oshlack A, Robinson MD, Young MD. From RNA-seq reads to differential expression results. *Genome Biol.* 2010; 11: 220.
86. Chen X, Zhu W, Azam S, Li H, Zhu F, Li H, Hong Y. Deep sequencing analysis of the transcriptomes of peanut aerial and subterranean young pods identifies candidate genes related to early embryo abortion. *Plant Biotechnol J.* 2013; 11: 115-127.
87. Yin D, Wang Y, Zhang X, Li H, Lu X, Zhang J, Zhang W. De novo assembly of the peanut (*Arachis hypogaea* L.) seed transcriptome revealed candidate unigenes for oil accumulation pathways. *PLoS One.* 2013; 8: e73767.
88. Zhang J, Liang S, Duan J, Wang J, Chen S, Cheng Z, Zhang Q. De novo assembly and characterisation of the transcriptome during seed development, and generation of genic-SSR markers in peanut (*Arachis hypogaea* L.). *BMC Genomics.* 2012; 13: 90.
89. Wu N, Matand K, Wu H, Li B, Li Y, Zhang X, He Z. De novo next-generation sequencing, assembling and annotation of *Arachis hypogaea* L. Spanish botanical type whole plant transcriptome. *Theor Appl Genet.* 2013; 126: 1145-1149.
90. Zhao CZ, Xia H, Frazier TP, Yao YY, Bi YP, Li AQ, Li MJ. Deep sequencing identifies novel and conserved microRNAs in peanuts (*Arachis hypogaea* L.). *BMC Plant Biol.* 2010; 10: 3.
91. Guimarães PM, Brasileiro AC, Morgante CV, Martins AC, Pappas G, Silva OB Jr, Togawa R. Global transcriptome analysis of two wild relatives of peanut under drought and fungi infection. *BMC Genomics.* 2012; 13: 387.
92. Nagy ED, Guo Y, Tang S, Bowers JE, Okashah RA, Taylor CA, Zhang D. A high-density genetic map of *Arachis duranensis*, a diploid ancestor of cultivated peanut. *BMC Genomics.* 2012; 13: 469.
93. James C. *Global Status of Commercialized Biotech/GM Crops*: 2013. Ithaca: ISAAA, 2013.
94. Cressley D. Transgenics: A new breed. *Nature.* 2013; 497: 27-29.
95. Ozias-Akins P, Schnall JA, Anderson WF, Singsit C, Clemente TE, Adang MJ, et al. Regeneration of transgenic peanut plants from stably transformed embryogenic callus. *Plant Sci.* 1993; 93: 185-94.
96. Chu Y, Bhattacharya A, Wu C, Knoll J, Ozias-Akins P. Improvement of peanut (*Arachis hypogaea* L.) transformation efficiency and determination of transgene copy number by relative quantitative real-time PCR. *In Vitro Cell Dev Biol -Plant.* 2013; 49: 266-75.
97. Ozias-Akins P, Gill R. Progress in the development of tissue culture and transformation methods applicable to the production of transgenic peanut. *Peanut Sci.* 2001; 28: 123-31.
98. Holbrook CC, Ozias-Akins P, Chu Y, Guo B. Impact of molecular genetic research on peanut cultivar development. *Agronomy.* 2011; 1: 3-17.
99. Bhatnagar M, Prasad K, Bhatnagar-Mathur P, Narasu ML, Waliyar F, Sharma KK. An efficient method for the production of marker-free transgenic plants of peanut (*Arachis hypogaea* L.). *Plant Cell Rep.* 2010; 29: 495-502.
100. S Sharma KK, Anjaiah V V. An efficient method for the production of transgenic plants of peanut (*Arachis hypogaea* L.) through *Agrobacterium tumefaciens*-mediated genetic transformation. *Plant Sci.* 2000; 159: 7-19.
101. Brar GS, Cohen BA, Vick CL, Johnson GW. Recovery of transgenic peanut (*Arachis hypogaea* L.) plants from elite cultivars utilizing ACCELL@ technology. *Plant J.* 1994; 5: 745-53.
102. Chu Y, Deng XY, Faustinelli P, Ozias-Akins P. Bcl-xL transformed peanut (*Arachis hypogaea* L.) exhibits paraquat tolerance. *Plant Cell Rep.* 2008; 27: 85-92.
103. Li Z, Jarret R, Demski J. Engineered resistance to tomato spotted wilt virus in transgenic peanut expressing the viral nucleocapsid gene. *Transgenic Res.* 1997; 6: 297-305.
104. Yang H, Singsit C, Wang A, Gonsalves D, Ozias-Akins P. Transgenic peanut plants containing a nucleocapsid protein gene of tomato spotted wilt virus show divergent levels of gene expression. *Plant Cell Rep.* 1998; 17: 693-9.
105. Magbanua Z, Wilde HD, Roberts J, Chowdhury K, Abad J, Moyer J, et al. Field resistance to Tomato spotted wilt virus in transgenic peanut (*Arachis hypogaea* L.) expressing an antisense nucleocapsid gene sequence. *Mol Breeding.* 2000; 6: 227-36.
106. Yang H, Ozias-Akins P, Culbreath AK, Gorbet DW, Weeks JR, Mandal B, et al. Field evaluation of Tomato spotted wilt virus resistance in transgenic peanut (*Arachis hypogaea*). *Plant Dis.* 2004; 88: 259-64.
107. Rao SC, Bhatnagar-Mathur P, Kumar PL, Reddy AS, Sharma KK. Pathogen-derived resistance using a viral nucleocapsid gene confers only partial non-durable protection in peanut against peanut bud necrosis virus. *Arch Virol.* 2013; 158: 133-143.
108. Higgins CM, Hall RM, Mitter N, Cruickshank A, Dietzgen RG. Peanut stripe potyvirus resistance in peanut (*Arachis hypogaea* L.) plants carrying viral coat protein gene sequences. *Transgenic Res.* 2004; 13: 59-67.
109. Mehta R, Radhakrishnan T, Kumar A, Yadav R, Dobaria JR, Thirumalaisamy PP, et al. Coat protein-mediated transgenic resistance of peanut (*Arachis hypogaea* L.) to peanut stem necrosis disease through *Agrobacterium*-mediated genetic transformation. *Indian J Virol.* 2013; 24: 205-213.
110. Collinge DB, Jørgensen HJ, Lund OS, Lyngkjær MF. Engineering pathogen resistance in crop plants: current trends and future prospects. *Annu Rev Phytopathol.* 2010; 48: 269-291.
111. Singsit C, Adang MJ, Lynch RE, Anderson WF, Wang A, Cardineau G, Ozias-Akins P. Expression of a *Bacillus thuringiensis* cryIA(c) gene in transgenic peanut plants and its efficacy against lesser cornstalk borer. *Transgenic Res.* 1997; 6: 169-176.

112. Tiwari S, Mishra DK, Singh A, Singh PK, Tuli R. Expression of a synthetic cry1EC gene for resistance against *Spodoptera litura* in transgenic peanut (*Arachis hypogaea* L.). *Plant Cell Rep.* 2008; 27: 1017-1025.
113. Rohini VK, Sankara Rao K. Transformation of peanut (*Arachis hypogaea* L.) with tobacco chitinase gene: variable response of transformants to leaf spot disease. *Plant Sci.* 2001; 160: 889-898.
114. Prasad K, Bhatnagar-Mathur P, Waliyar F, Sharma K. Overexpression of a chitinase gene in transgenic peanut confers enhanced resistance to major soil borne and foliar fungal pathogens. *J Plant Biochem Biotechnol.* 2013; 22: 222-33.
115. Ceasar SA, Ignacimuthu S. Genetic engineering of crop plants for fungal resistance: role of antifungal genes. *Biotechnol Lett.* 2012; 34: 995-1002.
116. Swathi Anuradha T, Divya K, Jami SK, Kirti PB. Transgenic tobacco and peanut plants expressing a mustard defensin show resistance to fungal pathogens. *Plant Cell Rep.* 2008; 27: 1777-1786.
117. Punja ZK. Genetic engineering of plants to enhance resistance to fungal pathogens: a review of progress and future prospects. *Can J Plant Pathol.* 2001; 23: 216-35.
118. Vasavirama K, Kirti PB. Increased resistance to late leaf spot disease in transgenic peanut using a combination of PR genes. *Funct Integr Genomics.* 2012; 12: 625-634.
119. Sundaresha S, Manoj Kumar A, Rohini S, Math SA, Keshamma E, Chandrashekar SC, et al. Enhanced protection against two major fungal pathogens of groundnut, *Cercospora arachidicola* and *Aspergillus flavus* in transgenic groundnut over-expressing a tobacco  $\beta$  1-3 glucanase. *Eur J Plant Pathol.* 2010; 126: 497-508.
120. Chenault KD, Melouk HA, Payton ME. Field reaction to Sclerotinia blight among transgenic peanut lines containing antifungal genes. *Crop Sci.* 2005; 45: 511-5.
121. Livingstone DM, Hampton JL, Phipps PM, Grabau EA. Enhancing resistance to Sclerotinia minor in peanut by expressing a barley oxalate oxidase gene. *Plant Physiol.* 2005; 137: 1354-1362.
122. Partridge-Telenko DE, Hu J, Livingstone DM, Shew BB, Phipps PM, Grabau EA. Sclerotinia blight resistance in Virginia-type peanut transformed with a barley oxalate oxidase gene. *Phytopathology.* 2011; 101: 786-793.
123. Niu C, Akasaka-Kennedy Y, Faustinelli P, Joshi M, Rajasekaran K, Yang H, et al. Antifungal activity in transgenic peanut (*Arachis hypogaea* L.) conferred by a nonheme chloroperoxidase gene. *Peanut Sci.* 2009; 36: 126-32.
124. Bhatnagar-Mathur P, Devi MJ, Reddy DS, Lavanya M, Vadez V, Serraj R, et al. Stress-inducible expression of At DREB1A in transgenic peanut (*Arachis hypogaea* L.) increases transpiration efficiency under water-limiting conditions. *Plant Cell Rep.* 2007; 26: 2071-2082.
125. Bhatnagar-Mathur P, Devi MJ, Vadez V, Sharma KK. Differential antioxidative responses in transgenic peanut bear no relationship to their superior transpiration efficiency under drought stress. *J Plant Physiol.* 2009; 166(11): 1207-17.
126. Vadez V, Rao JS, Bhatnagar-Mathur P, Sharma KK. DREB1A promotes root development in deep soil layers and increases water extraction under water stress in groundnut. *Plant Biol (Stuttg).* 2013; 15: 45-52.
127. Qin H, Gu Q, Zhang J, Sun L, Kuppu S, Zhang Y, et al. Regulated expression of an isopentenyltransferase gene (IPT) in peanut significantly improves drought tolerance and increases yield under field conditions. *Plant Cell Physiol.* 2011; 52: 1904-1914.
128. Banjara M, Zhu L, Shen G, Payton P, Zhang H. Expression of an Arabidopsis sodium/proton antiporter gene (AtNHX1) in peanut to improve salt tolerance. *Plant Biotechnol Rep.* 2012; 6: 59-67.
129. Sun L, Hu R, Shen G, Zhang H. Genetic engineering peanut for higher drought-and salt-tolerance. *Food Nutr Sci.* 2013; 4: 1.
130. Asif MA, Zafar Y, Iqbal J, Iqbal MM, Rashid U, Ali GM, Arif A. Enhanced expression of AtNHX, in transgenic groundnut (*Arachis hypogaea* L.) improves salt and drought tolerance. *Mol Biotechnol.* 2011; 49: 250-256.
131. Qin H, Gu Q, Kuppu S, Sun L, Zhu X, Mishra N, et al. Expression of the Arabidopsis vacuolar H<sup>+</sup>-pyrophosphatase gene AVP1 in peanut to improve drought and salt tolerance. *Plant Biotechnol Rep.* 2013; 7: 345-55.
132. Bhauso TD, Thankappan R, Kumar A, Mishra GP, Dobarja JR, Rajam M. Over-expression of bacterial mtd gene confers enhanced tolerance to salt-stress and water-deficit stress in transgenic peanut (*Arachis hypogaea*) through accumulation of mannitol. *Aust J Crop Sci.* 2014; 8: 413-21.
133. Chu Y, Faustinelli P, Ramos ML, Hajduch M, Stevenson S, Thelen JJ, Maleki SJ. Reduction of IgE binding and nonpromotion of *Aspergillus flavus* fungal growth by simultaneously silencing Ara h 2 and Ara h 6 in peanut. *J Agric Food Chem.* 2008; 56: 11225-11233.
134. Dodo HW, Konan KN, Chen FC, Egnin M, Viquez OM. Alleviating peanut allergy using genetic engineering: the silencing of the immunodominant allergen Ara h 2 leads to its significant reduction and a decrease in peanut allergenicity. *Plant Biotechnol J.* 2008; 6: 135-145.
135. Stevenson SE, Chu Y, Ozias-Akins P, Thelen JJ. Validation of gel-free, label-free quantitative proteomics approaches: applications for seed allergen profiling. *J Proteomics.* 2009; 72: 555-566.

Cite this article

Miranda Brasileiro AC, Guerra Araujo AC, Leal-Bertioli SC, Guimarães PM (2014) Genomics and Genetic Transformation in *Arachis*. *Int J Plant Biol Res* 2(3): 1017.