



Effect of hydrogen peroxide on thawed ovine sperm motility

M.S. Maia^{1,3}, S.D. Bicudo², L. Rodello²

¹Brazilian Agricultural Research Corporation (EMBRAPA), Semi-Arid Research Center, Petrolina, PE, Brazil.

²São Paulo State University, Faculty of Veterinary Medicine and Animal Science, Botucatu, SP, Brazil.

Abstract

Oxidative stress, resulting from excessive levels of ROS in semen, has a negative impact on functional parameters and sperm fertility. In this study we examined the influence of the oxidative stress induced by hydrogen peroxide (H₂O₂) in ovine sperm motility after thawing and catalase (CAT) ability to preserve sperm motility. Semen was incubated at 37°C with 100 µM H₂O₂, 1 U catalase + 100 µM H₂O₂ or no treatment, for 30 min. Immediately after adding treatments, sperm motility was determined by computer-assisted semen analysis (CASA). Incubation with H₂O₂ led to a significant (P < 0.05) decrease in motility parameters whereas catalase prevented a decline in motility secondary to oxidative stress. After 30 min of incubation with H₂O₂, total motility (12.0% vs. control 73.0%, H₂O₂ +CAT 70.0%), progressive motility (0.0% vs. control 19.0%, H₂O₂ +CAT 19.0%) and rapid motility (1.0% vs. control 43.0%, H₂O₂ +CAT 40.0%) decreased significantly (P < 0.05), whereas percentage of static cells increased (84.0% vs. control 18.0%, H₂O₂ +CAT 20.0%). We conclude that H₂O₂ causes damage to ovine sperm motility and that catalase is able to avoid detrimental effect of H₂O₂ on sperm motility.

Keywords: antioxidants, catalase, oxidative stress, ROS.

Introduction

Oxidative stress is determined by the balance between the generation and degradation of reactive oxygen species (ROS) within a tissue. Sperm and seminal plasma possess a number of enzymes and low-molecular weight antioxidants that scavenge ROS in order to prevent possible cellular damage (Aitken, 1995; Halliwell and Gutteridge, 1999).

The action of ROS on sperm function is concentration-dependent. Low concentrations of ROS, such as superoxide anion (O₂⁻) and hydrogen peroxide (H₂O₂) are needed for sperm capacitation, hyperactivation of motility, acrosome reaction and sperm-oocyte fusion (De Lamirande and Gagnon, 1993; Aitken, 1995; Baumber *et al.*, 2003; O'Flaherty *et al.*, 2003). On the other hand, excessive levels of ROS due to overproduction and/or inadequate antioxidant defenses leads to oxidative stress, resulting in a negative impact on functional parameters and fertility of sperm (Aitken,

1995; Armstrong *et al.*, 1999; Peris *et al.*, 2007).

The ROS most commonly generated by spermatozoon are: the superoxide anion, hydrogen peroxide and hydroxyl radical (Aitken, 1995). Several studies have shown that H₂O₂ exogenously added or produced by sperm, is toxic to mammalian spermatozoon causing damage to spermatid cell, including inhibition of motility and decline in energy metabolism (Armstrong *et al.*, 1999; O'Flaherty *et al.*, 1999; Baumber *et al.*, 2000, 2002; Bilodeau *et al.*, 2001, 2002; Garg *et al.*, 2009). Once produced, the H₂O₂ is removed by antioxidant enzymes such as catalase, glutathione peroxidase and peroxidases that catalyze their reduction to O₂ and water (Halliwell and Gutteridge, 1999).

Catalase has been used successfully in extenders in order to improve the antioxidant capacity of semen and preserve sperm motility after thawing (Krzyzosiak *et al.*, 2000; Baumber *et al.*, 2002; Bilodeau *et al.*, 2002; Maia *et al.*, 2009).

In light of the relationship between ROS generation and spermatid function, we examined the effect of the addition of exogenous H₂O₂ on frozen/thawed ram sperm motility during incubation for 30 min and the catalase ability to preserve sperm motility in semen incubated with H₂O₂.

Materials and Methods

Chemicals were purchased from SIGMA-ALDRICH, Inc. (St Louis, MO, USA) except hydrogen peroxide, which was obtained from Merck Millipore (Darmstadt, Germany) and Orvus® Es Paste (sodium lauryl sulfate), which was purchased from Procter & Gamble (Cincinnati, OH, USA).

Semen collection and processing

Semen was collected from sixteen adult (24-32 months of age) Santa Inês rams (*Ovis aries*) with an artificial vagina and the study was carried out using one ejaculate per male. Immediately after collection, ejaculates were evaluated for volume, concentration (Neubauer chamber) and motility (using CASA). Semen was diluted to 400 x 10⁶ cells/ml with a Tris-egg yolk extender, containing 250.25 mM Tris-hydroxymethyl-aminomethane, 79.71 mM monohydrated citric acid, 9.99 mM glucose, 20.0% (v/v) egg yolk, 1.0% (v/v)

³Corresponding author: marcianemaia@yahoo.com.br

Phone: +55(84)3232-5875; Fax: +55(84)3232-5868

Received: February 7, 2014

Accepted: June 13, 2014



Orvus ES Paste, 100 mg/100 ml streptomycin sulfate and 100 000 IU Potassic Penicillin (Fraction A, pH: 6.91; osmolarity: 308 mOsm) and 14.0% glycerol (v/v) in the Fraction B (Maia *et al.*, 2009). Dilution was performed at 32°C in a two-step process. The extended semen was filled in 0.25 French straws, sealed with polyvinyl alcohol and then cooled and frozen in an automatic freezer (Tetakon® -TK 3000, TK Tecnologia em Congelação Ltda, Uberaba, MG, Brazil) at a rate of -0.25°C/min from 32 to 5°C and -20°C/min from 5 to -120°C.

Thawing was carried out in a water bath at 42°C for 20 s. After thawing, straws were wiped and their content was poured in a micro tube, and then split in three aliquots for treatments: control; oxidant treatment (H₂O₂) and catalase/oxidant treatment (H₂O₂+CAT). In the control treatment, 60 µl semen (24 x 10⁶ cells/ml) was diluted in 440 µl Tris-citric acid buffer solution (pH 7.4, Maia *et al.*, 2010); the oxidant treatment constituted of semen sample, 60 µl, 435 µl Tris-citric acid buffer solution and 5 µl of 10 mM H₂O₂ (100 µM; Bilodeau *et al.*, 2001); the catalase/ oxidant treatment constituted of semen sample, 60 µl, 415 µl Tris-citric acid buffer solution, 5 µl of 10 mM H₂O₂ (100 µM) and 20 µl of 25 U/ml catalase solution (2 U/ml; Bilodeau *et al.*, 2002). The hydrogen peroxide (Perhydrol 30% H₂O₂ PA- Merck) and catalase (Catalase from bovine liver, SIGMA, C-9322) were prepared just prior to being added to the incubation medium.

Samples were then incubated at 37°C for 30 min, in order to induce oxidative stress, and sperm motility was determined immediately after adding of the treatments (time 0) and then at 15 min intervals.

Evaluation of sperm motility using CASA

Motility was assessed using a computer assisted sperm analysis (CASA). A sample of 10 µl of semen was placed in a pre-warmed MAKLER chamber (37°C) and transferred to the CASA analyzer (Hamilton-Thorn Biosciences IVOS 12, Beverly, MA, USA). For each semen sample, three fields were selected at random and various motility parameters including total motility (TM; %), progressive motility (PM; %), sperm with rapid motility (RAPID; %), static cells (STATIC; %), average path velocity (VAP; µm/s), curvilinear velocity (VCL; µm/s), straight-line velocity (VSL; µm/s), amplitude of lateral head displacement (ALH; µm), beat-cross frequency (BCF; Hz), straightness (STR; %) and linearity (LIN; %; Maia *et al.*, 2009) were evaluated.

The settings used for ovine spermatozoa were negative phase-contrast optics, at a recording rate of 60 frames/s, minimum contrast 60, minimum cell size five pixels, cell intensity 55, non motile head size 2, non motile head intensity 50, progressive VAP threshold value 75 µm/s, slow cells VAP cutoff 21.9 µm/s, slow cells VSL cutoff 6.0 µm/s, threshold STR 80%.

Progressive cells were those exhibiting a VAP of 75 µm/s and an STR of 80%. For each sperm sample, three fields were selected at random and assessed to generate data from at least 450 sperm/sample.

Detection of hydrogen peroxide

The presence of hydrogen peroxide in a sample was assessed using the HRPO-dependent oxidation of phenol red to a yellow derivate which concentration was measured by its absorbance at 610 nm (Maia *et al.*, 2010).

Statistical analysis

One ejaculate from each of the 16 rams was used in the 3 treatments (control, H₂O₂ and H₂O₂+CAT) and 3 incubation times (0; 15 and 30 min) totaling n = 144. Analyses of variance were performed employing a randomized block design. Each male was considered a block and treatments and time were random variables blocked within rams. The linear model for ANOVA was:

$$Y_{ijk} = \mu + \text{Trat}_i + \text{Time}_j + I_{ij} + \text{Male}_k + \varepsilon_{ijk}$$

Where Y is a motility parameter; μ the overall mean, Trat is the effect of treatment; Time is the effect of incubation time; I, effect of interaction (Trat x time); Male, effect of ram and ε the error component. Mean values that were different at P < 0.05 were tested using Duncan's test. The values for sperm motility parameters were expressed as the mean \pm SEM.

Results

The effect of H₂O₂ on sperm motility during incubation is shown in Table 1. Significant differences were observed in motility parameters after exposure to H₂O₂. During incubation, catalase prevented the decrease in all motility parameters.

Total motility (TM) in H₂O₂ group was significantly lower (P < 0.05) than in control from the beginning of incubation and after 15 min incubation compared to the H₂O₂/CAT group. At 15 min incubation a significant reduction in total motility (P < 0.05) in H₂O₂/CAT group was observed compared to control, whereas at 30 min no difference was found between the two groups.

Progressive motility was the parameter most affected by exposure to H₂O₂. After 15 min incubation a significant reduction in progressive motility (P < 0.05) was observed in H₂O₂ group compared to the other groups. After 30 min of incubation, no sperm with progressive motility was observed in this group.

After 15 min incubation, the percentage of sperm with rapid motility was significantly lower (P < 0.05) in H₂O₂ than in the control and H₂O₂+CAT group.

Exposure to H₂O₂ rapidly led to a significant increase in the percentage of static cells when compared with control and H₂O₂+CAT treatment. At 30 min incubation the percentage of static sperm in control, H₂O₂



and H₂O₂+CAT groups were 18 ± 2.6%, 84 ± 2.6% and 20 ± 2.6% respectively. The other motility parameters (VAP, VSL, VCL, ALH, BCF, STR and LIN) decreased significantly after 15 min of incubation in H₂O₂ group compared to the other groups.

After incubation of the sample for 30 min,

the concentration of H₂O₂ detected in control was 7.4 ± 2.8 nmol/ml, in H₂O₂ group 19.6 ± 5.7 nmol/ml and in H₂O₂+CAT group was 6.6 ± 2.6 nmol/ml. The concentration of H₂O₂ was greater (P < 0.05) in H₂O₂ group than in samples from control or H₂O₂+CAT, that were similar.

Table 1. Effect of hydrogen peroxide (H₂O₂) on frozen-thawed sperm motility during incubation at 37°C for 30 min.

Parameter	Treatment	Incubation time (min)		
		0	15	30
TM (%)	Control	71 ± 2.5 ^{Aa}	75 ± 2.5 ^{Aa}	73 ± 2.5 ^{Aa}
	H ₂ O ₂	59 ± 2.5 ^{Bb}	24 ± 2.5 ^{Cc}	12 ± 2.5 ^{Bd}
	H ₂ O ₂ + CAT	64 ± 2.5 ^{Aa}	66 ± 2.5 ^{Ba}	70 ± 2.5 ^{Aa}
PM (%)	Control	22 ± 1.0 ^{Aa}	23 ± 1.0 ^{Aa}	19 ± 1.0 ^{Ab}
	H ₂ O ₂	20 ± 1.0 ^{Ab}	1.0 ± 0.6 ^{Bc}	0.4 ± 0.1 ^{Bc}
	H ₂ O ₂ + CAT	20 ± 1.0 ^{Ab}	20 ± 1.0 ^{Ab}	19 ± 1.0 ^{Ab}
RAPID (%)	Control	39 ± 2.2 ^{Aa}	46 ± 2.2 ^{Aa}	43 ± 2.2 ^{Aa}
	H ₂ O ₂	32 ± 2.2 ^{Ab}	3.0 ± 2.2 ^{Bc}	0.9 ± 0.2 ^{Bc}
	H ₂ O ₂ + CAT	38 ± 2.2 ^{Ab}	39 ± 2.2 ^{Ab}	40 ± 2.2 ^{Ab}
STATIC (%)	Control	17 ± 2.6 ^{Ba}	16 ± 2.6 ^{Ca}	18 ± 2.6 ^{Ba}
	H ₂ O ₂	31 ± 2.6 ^{Ac}	70 ± 2.6 ^{Ab}	84 ± 2.6 ^{Aa}
	H ₂ O ₂ + CAT	27 ± 2.6 ^{Ac}	24 ± 2.6 ^{Bc}	20 ± 2.6 ^{Bd}
VAP (µm/s)	Controle	84.5 ± 1.6 ^{ABa}	87.0 ± 1.6 ^{Aa}	84.1 ± 1.6 ^{Aa}
	H2O2	80.2 ± 1.6 ^{Bb}	46.0 ± 1.6 ^{Bc}	45.0 ± 1.6 ^{Bc}
	H2O2 + CAT	87.0 ± 1.6 ^{Aa}	83.3 ± 1.6 ^{Aa}	81.5 ± 1.6 ^{Aa}
VSL (µm/s)	Controle	62.0 ± 1.3 ^{ABb}	62.5 ± 1.3 ^{Ab}	59.3 ± 1.3 ^{Ab}
	H2O2	60.4 ± 1.3 ^{Bb}	32.0 ± 1.3 ^{Bc}	25.2 ± 1.3 ^{Bd}
	H2O2 + CAT	65.0 ± 1.3 ^{Aa}	62.0 ± 1.3 ^{Aab}	59.0 ± 1.3 ^{Ab}
VCL (µm/s)	Controle	161.0 ± 3.1 ^{Aa}	160.1 ± 3.1 ^{Aa}	159.0 ± 3.1 ^{Aa}
	H2O2	151.4 ± 3.1 ^{Aa}	112.0 ± 3.1 ^{Bb}	101.0 ± 3.1 ^{Bc}
	H2O2 + CAT	161.4 ± 3.1 ^{Aa}	154.0 ± 3.1 ^{Aa}	152.4 ± 3.1 ^{Aa}
ALH (µm)	Controle	8.1 ± 0.5 ^{Aa}	7.7 ± 0.5 ^{Aa}	7.7 ± 0.5 ^{Aa}
	H2O2	7.5 ± 0.5 ^{Aa}	4.4 ± 0.5 ^{Bb}	3.1 ± 0.5 ^{Bb}
	H2O2 + CAT	7.6 ± 0.5 ^{Aa}	7.4 ± 0.5 ^{Aa}	7.2 ± 0.5 ^{Aa}
BCF (Hz)	Controle	38.0 ± 0.7 ^{Aa}	36.0 ± 0.7 ^{Aa}	36.0 ± 0.7 ^{Bb}
	H2O2	37.0 ± 0.7 ^{Ab}	31.4 ± 0.7 ^{Cc}	31.5 ± 0.7 ^{Cc}
	H2O2 + CAT	37.4 ± 0.7 ^{Ab}	37.0 ± 0.7 ^{Ba}	36.5 ± 0.7 ^{Ba}
STR (%)	Controle	69 ± 1.3 ^{Aa}	70 ± 1.3 ^{Aa}	68 ± 1.3 ^{Aa}
	H2O2	71 ± 1.3 ^{Aa}	68 ± 1.3 ^{Bb}	55 ± 1.3 ^{Bc}
	H2O2 + CAT	72 ± 1.3 ^{Aa}	72 ± 1.3 ^{Aa}	70 ± 1.3 ^{Aa}
LIN (%)	Controle	38 ± 1.2 ^{Aa}	39 ± 1.2 ^{Aa}	38 ± 1.2 ^{Aa}
	H2O2	40 ± 1.2 ^{Aa}	30 ± 1.2 ^{Bb}	27 ± 1.2 ^{Bb}
	H2O2 + CAT	40 ± 1.2 ^{Aa}	40 ± 1.2 ^{Aa}	39 ± 1.2 ^{Aa}

Values with different superscript letters in the same row (lower case letters) or column (capital letters) differ significantly at P < 0.05. Data is expressed as mean ± S.E of 16 replicates (one ejaculate from each of the sixteen rams).



Discussion

Our findings support the evidences that hydrogen peroxide is toxic to mammalian spermatozoon. The exposure to exogenous H_2O_2 affected adversely all sperm motility parameters, like what was observed in previous reports with various species (O'Flaherty *et al.*, 1999; Bilodeau *et al.*, 2001; 2002; Peris *et al.*, 2007; Garg *et al.*, 2009; Du Plessis *et al.*, 2010). This study also confirmed the beneficial effect of catalase on sperm survival during incubation under oxidative stress (Aitken *et al.*, 1993; Armstrong *et al.* 1999; Baumber *et al.*, 2000; Bilodeau *et al.*, 2002; Fernández-Santos *et al.*, 2008) indicating that H_2O_2 exerts a direct cytotoxic effect on spermatozoa.

Previous reports showed that catalase on concentration ranging from 1 - 4000 U/ml prevents the decrease of sperm motility in semen under oxidative stress, induced by H_2O_2 exogenous or by ROS generate systems (Aitken *et al.*, 1993; Armstrong *et al.*, 1999; Baumber *et al.*, 2000; Bilodeau *et al.*, 2002; Fernández-Santos *et al.*, 2008). In our study, the addition of only 2 U/ml of catalase to the incubation medium was sufficient to prevent the loss of sperm motility caused by H_2O_2 . This could be partly explained because during the catalytic reaction to eliminate two molecules of H_2O_2 , free catalase is regenerated and molecular oxygen is released (Chance *et al.*, 1979; Bartosz, 2005). Thus the enzyme may continue to react and a single molecule of catalase is capable of decomposing thousands of molecules of H_2O_2 . According to Bartosz (2005) the number of peroxide molecules decomposed per CAT molecule per second is 3.5×10^6 . Moreover, the catalase is most effective when the concentration of H_2O_2 is highly elevated which may have favored the catalase activity in our study. The catalytic reaction in which H_2O_2 react with catalase and Compound I does not obey Michaelis-Menten Kinetics; the rate of substrate decomposition increases linearly with hydrogen peroxide concentration over a wide concentration range (Bartosz, 2005). According to Chance *et al.* (1979) catalase is especially effective as a "safety valve" for dealing with much H_2O_2 . The catalase activity like H_2O_2 scavenger was observed in our study, in sperm treated with H_2O_2 and catalase, the H_2O_2 levels remain similar to untreated sperm.

The damaging effect caused by 100 μM H_2O_2 (100 nmol/ml) on sperm motility during incubation in the diluents without catalase was very fast, 15 min after its addition to the extender; the same was observed by Bilodeau *et al.* (2001, 2002) and Garg *et al.* (2009). Possibly, this occurred due to the amount of H_2O_2 added to the system, which was much greater than the concentrations generated spontaneously by ram sperm, after cryopreservation (7.39 nmol/ml). However, after 30 min of incubation, the amount of hydrogen peroxide detected in the H_2O_2 group was only about 20% of original concentration added to the system (100

nmol/ml), possibly reflecting a scavenging effect of spermatozoa and seminal plasma. Nevertheless, this concentration was sufficient to affect sperm motility. It has been established that the decline in sperm motility, after treatment with H_2O_2 , occurs in a dose-dependent manner (Armstrong *et al.*, 1999; Peris *et al.*, 2007). Thus, the amount of H_2O_2 added to the system was enough to cause a severe oxidative stress to sperm, resulting in a rapid decline (after 15 min) in all motility parameters.

The parameter most affected by H_2O_2 was the progressive motility, such as that detected in human sperm by Calamera *et al.* (2001) and du Plessis *et al.* (2010). Moreover, Calamera *et al.* (2001) observed that although progressively immovable, the spermatozoa were alive and showing no signs of major membrane damage. Thus, the mechanism by which H_2O_2 affects sperm motility is still unclear. Some studies (Armstrong *et al.*, 1999; Bilodeau *et al.*, 2002) attributed the loss of sperm motility to the fact that the H_2O_2 inhibits sperm ATP production by different ways. However, Calamera *et al.* (2001) reported that the loss of motility was accompanied by a parallel increase in sperm ATP concentration. Therefore, depletion of ATP may not be the cause of the decrease of sperm movement, instead, the decrease in the use or consumption by progressively immotile spermatozoa.

Furthermore, the inactivation of the intracellular scavenger systems may also be involved in decline in sperm motility, after treatment with H_2O_2 . According to Krzyzosiak *et al.* (2000) the effect of H_2O_2 on sperm motility during incubation without catalase could be related to the inactivation of glutathione peroxidase/reductase system. This system is present in ovine semen (Bucak *et al.*, 2008; Marti *et al.*, 2008). Since biological membranes are highly permeable to H_2O_2 , and glutathione peroxidase is present in the mitochondria (Chance *et al.*, 1979), probably, when the intracellular concentration of H_2O_2 increased, intracellular glutathione (GSH) began to go down as well as the sperm motility. Therefore, the loss of sperm motility caused by H_2O_2 could be because no antioxidant enzyme that reduces H_2O_2 was present in the media.

On the other hand, in the semen incubated with H_2O_2 and catalase there was a time lag before the effect of catalase on motility was detected. Possibly, this occurred because the catalase reacted with H_2O_2 only after most GSH had been consumed, since glutathione peroxidase has a higher affinity for H_2O_2 (Bartosz, 2005) than does catalase. In erythrocyte Mendiratta *et al.* (1998) observed that with increasing fluxes of H_2O_2 coming into the cells, intracellular GSH decreased as the amount of H_2O_2 increased, reaching the catalase and that GSH need not be substantially depleted before H_2O_2 reaches catalase. In this way, the increase in sperm motility, arising from the effect of catalase, compared to the H_2O_2 group was only observed after 15 min of incubation.



In conclusion, the deleterious effect of H₂O₂ on sperm motility can be counteracted by adding catalase to the extender. The finding that H₂O₂ is cytotoxic to ovine spermatozoon suggests the importance of the addition of systems for H₂O₂ scavenge, like catalase, in the semen cryopreservation extender.

Acknowledgment

The authors are grateful to the Foundation of Research Support of São Paulo State (FAPESP) for funding this work.

References

- Aitken RJ, Buckingham D, Harkiss D.** 1993. Use of a xanthine oxidase free radical generation system to investigate the cytotoxic effect of reactive oxygen species on human spermatozoa. *J Reprod Fertil*, 97:441-450.
- Aitken RJ.** 1995. Free radicals, lipid peroxidation and sperm function. *Reprod Fertil Dev*, 7:659-668.
- Armstrong JS, Rajasekaran M, Chamulitrat W, Gatti P, Hellstrom WJ, Sikka SC.** 1999. Characterization of reactive oxygen species induced effects on human spermatozoa, and energy metabolism. *Free Radic Biol Med*, 26:869-880.
- Bartosz G.** 2005. Superoxide dismutase and catalase. In: Grune T (Ed.). *Reaction and Process*. Berlin: Springer. pp. 109-149. (The Handbook of Environmental Chemistry, pt. 2/20).
- Baumber J, Ball BA, Gravance CG, Medina V, Davies-Morel MCG.** 2000. The effect of reactive oxygen species on equine sperm motility, viability, acrossomal integrity, mitochondrial membrane potential and membrane lipid peroxidation. *J Androl*, 21:895-902.
- Baumber J, Vo A, Sabeur K, Ball BA.** 2002. Generation of reactive oxygen species by equine neutrophils and their effect on motility of equine spermatozoa. *Theriogenology*, 57:1025-1033.
- Baumber J, Sabeur K, Vo A, Ball BA.** 2003. Reactive oxygen species promote tyrosine phosphorylation and capacitation in equine spermatozoa. *Theriogenology*, 60:1239-1247.
- Bilodeau JF, Blanchette S, Gagnon C, Sirard MA.** 2001. Thiols prevent H₂O₂ - mediated loss of sperm motility in cryopreserved bull semen. *Theriogenology*, 56:275-286.
- Bilodeau JF, Blanchette S, Cormier N, Sirard MA.** 2002. Reactive oxygen species-mediated loss of bovine sperm motility in egg yolk Tris extender: protection by pyruvate, metal chelators and bovine liver or oviductal fluid catalase. *Theriogenology*, 57:1105-1122.
- Bucak MN, Atessahin A, Yüce A.** 2008. Effect of antioxidants and oxidative stress parameters on ram semen after the freeze-thawing process. *Small Rumin Res*, 75:128-134.
- Calamera JC, Fernandez PJ, Buffone MG, Acosta AA, Doncel GF.** 2001. Effects of long-term in vitro incubation of human spermatozoa: functional parameters and catalase effect. *Andrologia*, 33:79-86.
- Chance B, Sies H, Boveris A.** 1979. Hydroperoxide metabolism in mammalian organs. *Physiol Rev*, 59:527-605.
- De Lamirande E, Gagnon C.** 1993. Human sperm hyperactivation and capacitation as parts of an oxidative process. *Free Radic Biol Med*, 14:255-265.
- Du Plessis SS, McAllister DA, Luu A, Savia J, Agarwal A, Lampiao F.** 2010. Effects of H₂O₂ exposure on human sperm motility parameters, reactive oxygen species levels and nitric oxide levels. *Andrologia*, 42:206-210.
- Fernández-Santos MR., Domínguez-Rebolledo AE, Esteso MC, Garde JJ, Martínez-Pastor F.** 2008. Catalase supplementation on thawed bull spermatozoa abolishes the detrimental effect of oxidative stress on motility and DNA integrity. *Int J Androl*, 32:353-359.
- Garg A, Kumaresan A, Ansari MR.** 2009. Effects of hydrogen peroxide (H₂O₂) on fresh and cryopreserved buffalo sperm function during incubation at 37°C in vitro. *Reprod Domest Anim*, 44:907-912.
- Halliwell B, Gutteridge JMC.** 1999. *Free Radicals in Biology and Medicine*, 3rd ed. Oxford: Oxford University Press. 936 pp.
- Krzyzosiak J, Evenson D, Pitt C, Jost L, Molan P, Vishwanath R.** 2000. Changes in susceptibility of bovine sperm to in situ DNA denaturation, during prolonged incubation at ambient temperature under conditions of exposure to reactive oxygen species and nuclease inhibitor. *Reprod Fertil Dev*, 12:251-261.
- Maia MS, Bicudo SD, Azevedo HC, Sicherle CC, Sousa DB, Rodello L.** 2009. Motility and viability of ram sperm cryopreserved in a Tris-egg yolk extender supplemented with anti-oxidants. *Small Rumin Res*, 85:85-90.
- Maia M da S, Bicudo SD, Sicherle CC, Rodello L, Gallego ICS.** 2010. Lipid peroxidation and generation of hydrogen peroxide in frozen-thawed ram semen cryopreserved in extenders with antioxidants. *Anim Reprod Sci*, 122:118-123.
- Martí E, Martí JI, Muiño-Blanco T, Cebrián-Pérez JA.** 2008. Effect of the cryopreservation process on the activity and immunolocalization of antioxidant enzymes in ram spermatozoa. *J Androl*, 29:459-467.
- Mendiratta S, Qu Z-C, May JM.** 1998. Erythrocyte defenses against hydrogen peroxide: the role of ascorbic acid. *Biochim Biophys Acta*, 1380:389-395.
- O'Flaherty C, Beorlegui N, Beconi MT.** 1999. Reactive oxygen species requirements for bovine sperm capacitation and acrosome reaction. *Theriogenology*, 52:289-301.
- O'Flaherty C, Beorlegui N, Beconi MT.** 2003. Participation of superoxide anion in the capacitation of cryopreserved bovine sperm. *Int J Androl*, 26:109-114.
- Peris SI, Bilodeau J-F, Dufour M, Bailey J.** 2007. Impact of cryopreservation and reactive oxygen species on DNA integrity, lipid peroxidation, and functional parameters in ram sperm. *Mol Reprod Dev*, 74:878-892.