

Short Communication

THE STINGLESS BEE SPECIES, *SCAPTOTRIGONA* AFF. *DEPILIS*, AS A POTENTIAL INDICATOR OF ENVIRONMENTAL PESTICIDE CONTAMINATIONANNELISE DE SOUZA ROSA,*† ROBBIE I'ANSON PRICE,‡ MARIA JULIANA FERREIRA CALIMAN,† ELISA PEREIRA QUEIROZ,†
BETINA BLOCHTEIN,§ CARMEN SÍLVIA SOARES PIRES,|| and VERA LUCIA IMPERATRIZ-FONSECA†

†Department of Entomology, University of São Paulo, Ribeirão Preto, Brazil

‡Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland

§Department of Biology, Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre, Brazil

||Embrapa, Brasília, Brazil

(Submitted 22 February 2015; Returned for Revision 13 March 2015; Accepted 19 March 2015)

Abstract: Neonicotinoids have the potential to enter the diet of pollinators that collect resources from contaminated plants. The species *Scaptotrigona* aff. *depilis* (Moure, 1942) can be a useful indicator of the prevalence of these chemicals in the environment. Using high-performance liquid chromatography–mass spectrometry, the authors devised a protocol for neonicotinoid residue extraction and detected the presence of neonicotinoids in the bee bodies. Thus, the authors consider this species to be a potential indicator of environmental contamination. *Environ Toxicol Chem* 2015;34:1851–1853. © 2015 SETAC

Keywords: Biomonitoring Environmental toxicology Insecticide Imidacloprid Neonicotinoid

INTRODUCTION

Neonicotinoid pesticides are registered for use in more than 120 countries and account for 24% of the global pesticide market. This means neonicotinoids affect a wide array of plants, animals, and environments [1]. The potential harm they cause insect populations has been studied heavily [2]. As a systemic insecticide, compounds within neonicotinoids can enter plant cells and be present in substrates that are attractive to insects. As such, they may be ingested in nectar and pollen by foraging bees [3,4]. Even without robust evidence, the potential presence of neonicotinoid compounds in pollinator diets has been linked to pollinator declines [5].

In Brazil, the native stingless bee species *Scaptotrigona* aff. *depilis* (Moure, 1942) visits a wide range of floral resources [6]. This bee species is likely to come into contact with plants that have been treated with pesticides; thus, we chose to investigate the incidence of the 2 most widely used neonicotinoids using this species. Previous studies using honey bees have shown that the expression of detoxification, behavior, immunity, and nutrition genes is altered in individuals exposed to pesticides [7]. This suggests that honey bees can metabolically combat the incidence of pesticides in their bodies; indeed, the same could potentially occur in stingless bees. Against this background, the present study aimed to detect neonicotinoid chemicals in stingless bees and monitor the time it takes for detoxification of bees from imidacloprid.

MATERIAL AND METHODS

We collected *S. aff. depilis* foragers that were returning to the colony after foraging. These were collected at the entrance of 5 colonies at the meliponary at São Paulo University, Ribeirão Preto campus. Colonies were assigned letters: A, B, C, D, and E.

One hundred bees were sampled from each colony. At the same time, we collected samples of bees from 2 natural nests (colonies F and G), approximately 3 km away from the 5 colonies. Given the flight range (903 m) of this species, we would not expect the foraging areas of our 2 groups to overlap [8]. Bee samples were frozen immediately after returning to the laboratory. Thereafter, we monitored the presence of imidacloprid and thiamethoxam in colonies of *S. aff. depilis* maintained in the meliponary.

We established the following protocol as the method to extract residue in workers from the *S. aff. depilis* species. A sample of 20 bees was put in a 50-mL Falcon tube with 10 mL of acetone. The sample was ground using a tissue homogenizer for 1 min and 30 s. It was then placed in a centrifuge for 15 min at 4000 rpm and 4° C, and the supernatant was removed. This was filtered using a 0.20- μ m filter, transferred to a test tube, and set aside to evaporate in compressed air. Then, 250 μ L of Mobile Phase (50% MeOH:H₂O [v/v] + 0.5% CH₃COOH:H₂O [v/v]) was added to the tube, and the samples were placed in the spectrometer using high-performance liquid chromatography–mass spectrometry. The liquid chromatography system consisted of an LC10AD pump and a CTO-10AS column oven from Shimadzu. The separation of the chemicals was performed on a C18 column (Agilent ZORBAX Rapid Resolution Eclipse Plus C18 Analytical 4.6 \times 150 mm 5- μ ; Agilent Technologies). Detection was obtained with a Quattro Micro LC triple quadrupole mass spectrometer (Micromass) fitted with an electrospray interface source and tandem mass operated in the positive ion mode.

Detection was carried out using the internal standards imidacloprid and thiamethoxam, both obtained from Sigma Aldrich (PESTANAL, analytical standard, Fluka). The instrument conditions were adapted from Kamel [9]—retention time: imidacloprid 3.89 s and thiamethoxam 3.39 s; primary transition molecules and collision energy (in electron volts; eV): imidacloprid 256>175, 23eV, and thiamethoxam 292>132, 19eV; secondary transition molecules and collision energy:

* Address correspondence to annesouzar@gmail.com.

Published online 10 July 2015 in Wiley Online Library (wileyonlinelibrary.com).

DOI: 10.1002/etc.2998

imidacloprid 256>209, 21eV, and thiamethoxam 292>211, 31eV.

RESULTS

Imidacloprid was detected in the samples obtained from colonies A, B, and C (Figure 1). This indicated that these colonies were suitable for further study. Thiamethoxam was found in 1 sample (colony D), and 1 sample was free from contamination of both neonicotinoids (colony E). The bees collected from wild nests 3 km outside the meliponary (colonies F and G) did not show contamination with the neonicotinoids. Additional samples were collected from colonies A, B, and C. Twenty bees from the newly collected samples were tested immediately for the presence of neonicotinoids and once more showed contamination with imidacloprid. We used bees from the remainder of the sample to monitor the time before the

neonicotinoid compounds were no longer present in the bees. We confined the bees in boxes of 30 individuals and kept them in a greenhouse with noncontaminated syrup (sucrose and water, 1:1). Two confinement times were used: individuals from colonies A and B were confined for 24 h, and individuals from colonies A, B, and C were confined for 96 h. At the end of these isolation periods, the bees were collected and frozen. Bee samples from colonies A and B (confined for 24 h) had peaks revealing imidacloprid was still present. Individuals from colonies A, B, and C (confined for 96 h) did not present the chemical anymore.

DISCUSSION

The present study is the first to extract and detect pesticide residues in *S. aff. depilis*. This procedure can be considered for application and adaptation for other species of stingless bee.




Figure 1. (A–C) Chromatograms showing the initial detection of imidacloprid in 3 of the 5 colonies: colony A (A), colony B (B) and colony C (C). Each chromatogram is followed by information on the primary (256>209) or secondary (256>175) transition molecules of imidacloprid. The x-axis represents the retention time of the imidacloprid molecules (in seconds) and the y-axis represents the signal intensity (in percentage).

Faria et al. [6] reported that this species visits a wide range of floral resources inside the São Paulo University campus by analyzing the pollen collected by *S. aff. depilis*. Within a 500-m radius of the meliponary, Aleixo et al. [10] surveyed 289 species of plants, among which 194 offer resources to bees. These flowers could have been contaminated and are within reach of the foraging workers of *S. aff. depilis*, which have an estimated flight radius of 903 m [8].

Notably, after 96 h of confinement, bees that were previously contaminated by imidacloprid showed no sign of contamination. Because the chemicals were removed in a short period of time, *S. aff. depilis* could be a sensitive indicator of environmental contamination. Furthermore, it is likely that the concentration at which the active substance was ingested is a key factor in the rate it is eliminated [11]. If this is the case, this species would enable researchers to quantify the current level of environmental pesticide contamination using several isolation periods. Part of our future plan is to use the stingless bee diversity on campus. It is home to at least 14 species of stingless bee in wild nests [12]; this variety offers us the opportunity to use several species when analyzing pesticide contamination in the environment. We will also continue to develop the procedures established in the present study to devise a method to quantify residue. This will enable us to deduce the concentrations of the chemicals ingested using the time to elimination. It is likely that the presence of neonicotinoids in the diet of stingless bees has nonlethal effects [13]; thus, bee mortality might not be observed initially. Sublethal effects, situations in which the insecticide does not kill individuals immediately but affects long-term colony fitness, have been demonstrated in bees [14,15]; in fact, studies have attributed the use of neonicotinoids at sublethal concentrations to colony collapse disorder [16,17]. There is a possibility that neonicotinoid chemicals are accumulated on the colonies on campus. Bees play an important role in their environment; therefore, a study that investigates the long-term effects of neonicotinoid ingestion could help us understand the environmental consequences of using neonicotinoid pesticides.

Finally, our results enable us to consider using *S. aff. depilis* as an indicator of environmental pesticide contamination. Further research using the techniques developed in the present study will help monitor toxicity levels in stingless bee species with the potential to create proposals that use stingless bee contamination as an environmental indicator.

Acknowledgment—The authors thank the Faculdade de Filosofia, Ciências e Letras de Ribeirão Preto from the University of São Paulo, for providing the area to collect bees. We also thank the Faculdade de Ciências Farmacêuticas (University of São Paulo) for their help with the chemical analysis. Finally, thanks to Conselho Nacional de Desenvolvimento Científico e Tecnológico for financial support.

Data availability—All data are found within the present study.

REFERENCES

- Jeschke P, Nauen R, Schindler M, Elbert A. 2010. Overview of the status and global strategy for neonicotinoids. *J Agric Food Chem* 59:2897–2908.
- Thompson HM. 2010. Risk assessment for honey bees and pesticides—Recent developments and ‘new issues’. *Pest Manag Sci* 66:1157–1162.
- Rortais A, Arnold G, Halm MP, Touffet-Briens F. 2005. Modes of honeybees’ exposure to systemic insecticides: Estimated amounts of contaminated pollen and nectar consumed by different categories of bees. *Apidologie* 36:71–83.
- Blacquiere T, Smaghe G, Van Gestel CA, Mommaerts V. 2012. Neonicotinoids in bees: A review on concentrations, side-effects and risk assessment. *Ecotoxicology* 21:973–992.
- Laycock I, Cotterell KC, O’Shea-Wheller TA, Cresswell JE. 2014. Effects of the neonicotinoid pesticide thiamethoxam at field-realistic levels on microcolonies of *Bombus terrestris* worker bumble bees. *Ecotoxicol Environ Saf* 100:153–158.
- Faria LBD, Aleixo KP, Garófalo CA, Imperatriz-Fonseca VL, Silva CID. 2012. Foraging of *Scaptotrigona aff. depilis* (Hymenoptera, Apidae) in an urbanized area: Seasonality in resource availability and visited plants. *Psyche: A Journal of Entomology* 2012:1–12.
- Schmehl DR, Teal PE, Frazier JL, Grozinger CM. 2014. Genomic analysis of the interaction between pesticide exposure and nutrition in honey bees *Apis mellifera*. *J Insect Physiol* 71:177–190.
- Lichtenberg EM, Imperatriz-Fonseca VL, Nieh JC. 2010. Behavioral suites mediate group-level foraging dynamics in communities of tropical stingless bees. *Insectes Soc* 57:105–113.
- Kamel A. 2010. Refined methodology for the determination of neonicotinoid pesticides and their metabolites in honey bees and bee products by liquid chromatography–tandem mass spectrometry (LC-MS/MS). *J Agr Food Chem* 58:5926–5931.
- Aleixo KP, Faria LB, Groppo M, Castro MMN, Silva CI. 2014. Spatiotemporal distribution of floral resources in a Brazilian city: Implications for the maintenance of pollinators, especially bees. *Urban For Urban Green* 13:689–696.
- Suchail S, De Sousa G, Rahmani R, Belzunces LP. 2004. *In vivo* distribution and metabolism of 14C-imidacloprid in different compartments of *Apis mellifera* L. *Pest Manag Sci* 60:1056–1062.
- de Freitas GS, de Assis AF, de Souza CCM, de Sousa J, Soares AEE. 2009. O doce lar das abelhas indígenas sem ferrão. *Mensagem Doce* 100:15–20.
- Cresswell JE. 2011. A meta-analysis of experiments testing the effects of a neonicotinoid insecticide (imidacloprid) on honey bees. *Ecotoxicology* 20:149–157.
- Tomé HVV, Martins GF, Lima MAP, Campos LAO, Guedes RNC. 2012. Imidacloprid-induced impairment of mushroom bodies and behavior of the native stingless bee *Melipona quadrifasciata anthidioides*. *PLoS ONE* 7:e38406.
- Sandrock C, Tanadini M, Tanadini LG, Fauser-Misslin A, Potts SG, Neumann P. 2014. Impact of chronic neonicotinoid exposure on honeybee colony performance and queen supersedure. *PLoS ONE* 9:e103592.
- Chensheng LU, Warchol KM, Callahan RA. 2012. In situ replication of honey bee colony collapse disorder. *Bulletin of Insectology* 65: 99–106.
- Chensheng LU, Warchol KM, Callahan RA. 2014. Sub-lethal exposure to neonicotinoids impaired honey bees winterization before proceeding to colony collapse disorder. *Bull Insectology* 67:125–130.