



SETAC Latin America 11th Biennial Meeting

7–10 September 2015 | Buenos Aires, Argentina | sla2015.setac.org

THE ROLE OF SCIENCE IN ENVIRONMENTAL DECISION-MAKING



SETAC Latin America
11th Biennial Meeting
Organizing Committee

Chairs

Pedro Carriquiriborde
CONICET-UNLP
La Plata, Buenos Aires, Argentina

Gustavo M. Somoza
CONICET-UNSAM
Chascomús, Buenos Aires,
Argentina

Members

Horacio Asprea
SAyDS
Buenos Aires, Argentina

Afonso Celso Dias Bainy
Universidade Federal de Santa
Catarina, Florianopolis, Brazil

Ricardo Barra
Universidad de Concepción
Concepción, Chile

Julie Brodeur
CONICET-INTA
Castelar, Buenos Aires, Argentina

Sandra Demichelis
Universidad Nacional de Lanús
Lanús, Buenos Aires, Argentina

Damián Marino
CONICET-UNLP
La Plata, Buenos Aires, Argentina

Karina Miglioranza
CONICET-UNMdP
Mar del Plata, Buenos Aires,
Argentina

Sebastián Reynaldi
Universidad Nacional
Medellín, Colombia

Déborah Tasat
Universidad de San Martín
San Martín, Buenos Aires,
Argentina

Abstract Book

Buenos Aires 2015

M.J. Damonte, M.B. Poliserpi, INTA; M.F. D'Andrea, J.C. Brodeur, CONICET INTA

Se evaluó en campos de soja los efectos de la aplicación del insecticida ENGEO® (Lam: Lambdacialotrina 10,6% p/v + Tia: Tiametoxam 14,1% p/v) sobre *Leptodactylus latrans*, *Leptodactylus latinasus* e *Hypsiboas pulchellus*. El muestreo se realizó 15 días antes y 2 y 4/5 días después de la aplicación del insecticida en tres lotes de soja de Bs. As. (L1, L2 y L3). Además se muestreó la reserva "El Destino" como sitio control. En todos los sitios y en las diferentes fechas se capturaron ranas y se tomaron muestras de suelo y agua. Se evaluó la condición corporal y biomarcadores de exposición a xenobióticos (ChE: colinesterasas y GST: glutatión-S-transferasa) o de estrés oxidativo (CAT: catalasa y GSH: glutatión reducido). Las muestras de suelo y agua no contenían Tia ni Lam antes de la aplicación. Después de la aplicación, se detectó Engeo en el suelo de L2 (Tia 190 ppb, lam 231 ppb) pero no en los otros dos lotes, probablemente debido a la gran cobertura vegetal de estos sitios. La condición corporal de *L. latinasus* estuvo reducida con respecto al control en todos los lotes siendo significativa en L3. No hubo diferencias entre los sitios cultivados y control para *L. latrans*. *H. pulchellus* presentó un significativo aumento de la condición corporal. La actividad de ChE fue significativamente reducida dos días después de la aplicación para *L. latinasus* en L1 y L2 e *H. pulchellus* solo L2, recuperando sus niveles basales 4 días después. Lo mismo ocurre con CAT, dos días después de la aplicación, sus valores se reducen significativamente para *L. latinasus* en L1 y L2. Los niveles de actividad de GST fueron significativamente menores para *L. latrans* en L1 y L3 con respecto al control, tanto antes como después de la aplicación. Este trabajo es el segundo en reportar una inhibición de la GST en *L. latrans* de zonas agrícolas de la región pampeana. La mayor cantidad de alteraciones encontradas en *L. latinasus* podrían deberse a que es una especie exclusivamente terrestre, estando más expuesta a la aplicación de plaguicidas. En este estudio, se demuestra la importancia de muestrear inmediatamente después de la aplicación de este formulado.

PT61. Analisis of the take-back program for empty pesticide containers in the municipality of Bom Repouso, mg, brasil

M. Rodrigues Silva, Universidade de Sao Paulo / Escola de Engenharia de São Carlos; M.T. Nunes, Escola de Engenharia de São Carlos USP / Hidraulica e Saneamento

Brazil is the first in the ranking of take - back program for empty pesticide containers comparing the returns with the amount of pesticides placed on market since 2011, reaching a percentage of 96% of return in 2013. The Brazilian take-back program was originally designed targeting large rural areas, which may not apply to small rural communities, generally more isolated and with poor infrastructure, that must require large storage and transport costs. Therefore, it is still very common to burn and bury empty pesticides containers in this areas, such as storing it in inappropriate places or leaving it in the field. Thus, this research aimed to analyze the efficiency of empty pesticide containers reverse logistics process in small Brazilian agricultural counties, taking as a case study the municipality of Bom Repouso, MG, Brazil, which is characterized by the production of strawberry and potato on properties with family labor and intensive use of pesticides. Therefore, it was made a comparative analysis of the amount of packaging sold in the city and the amount returned to the place where occurs the management of the empty containers that it's under INPEV's responsibility, called Central de Pousos Alegres, in MG, Brazil, between the years 2012 and 2014. We compared the total amount of returned packaging with the sales for the seven best-selling pesticides in the city. Considering the requirement of the law that farmers must return empty pesticides containers in a maximum of one year after purchase, it was taken a comparison between two periods, from the year 2012 to 2013 and 2013 to 2014. In both periods analyzed, the amount of packaging sold, considering only the seven most traded products, it was much higher than the returned packaging, reaching a difference of up to 67% between the product with the highest sales volume and the total amount of returned packages. Besides, there was a decrease of the total quantity of empty pesticides containers returned from 2013 to 2014, from 156,29 kg to 62,76 kg. Hard storage for empty pesticide containers and transportation costs were the main points highlighted by dealers as reasons given by producers for not returning and, consequently, improperly dispose such waste. It is believed that simple solution as the itinerant collection, can help to improve this scenario and ensure proper disposal of empty pesticides containers. Key-words: pesticides, small rural communities, waste handling, waste recovery.

PT62. Efficiency of Biobeds for pesticide disposal: Ecotoxicity evaluation using *Collembola* reproduction tests

L. Camargo, Soil Department; O. Filho; L.C. Filho, UDESC / Soil Department; L. Gebler, Embrapa; J.C. Niemeyer, Universidade Federal de Santa Catarina

Pesticide waste disposal from washing the equipment consist in a source of contamination for agricultural soils. An alternative to this problem was founded in some countries: the discard is done in reactors filled with straw, peat and soil mixed, called Biobeds. In Brazil, this sort of effluent is not remarkable. In South of Brazil, Mancozeb and Chlorpirifos are used in apple cultures, but their waste is discharge in soil and effects on soil organisms are unknown. An experiment with Biobeds was installed on the Experimental Station of Embrapa, at Vacaria City, in the apple harvest between 2013-2014, aiming to determine the toxicity into a biobed and in a local natural soil – Oxisol, during the harvest. The reactors were water boxes with biomix (Biobed) or Oxisol. The boxes received 50 L of effluent for each application in the crop. Samples of biomix and soils were collected on time 0 (without pesticide), 90, 150, 210 and 240 days after the first pesticide application, and ecotoxicity reproduction tests were conducted with *Collembola* (*Folsomia candida*) following ISO 11267 (2014). The number of juveniles for each treatment (Biomix or Oxisol) was compared for every time, with t-test. The collembolas not survive in any treatment for two first samplings. Only the last one (240 days) there was reproduction, and this not diverged statically from control, suggesting a toxicity reduction throughout time. Whereas, when compared Biomix against Oxisol, reproduction rates for the first one was upper than second. Although absence of toxicity in natural soil reactors after a period, in a natural system, discharge a substance which have an unknown toxicity and behavior, could impact soil organisms promoting a dissipation from that area, increasing the contamination already high, due lack of the pesticide dissipation. The study points to the need to know the toxicity of this waste, so common and generated in high amounts by agricultural activities and still without specific legislation in Brazil.

PT63. Assessing risk in the age of global trade: a comparison of regulatory bee requirements on plant protection products in South America, USA and Europe

M. Delgado, M. Faupel, A. Hoffarth, Rifcon GmbH

Are the regulations governing the registration of plant protection products (PPPs) in developing countries just copied from their counterparts in the industrialized nations? Are the current and emerging agricultural superpowers in South America (e.g. Brazil) adapting their regulatory standard to the agricultural markets of the USA and the European Union? What about issues relevant to environmental, consumer and farmer protection in the countries of origin: are standards the same as in the USA or the EU? We take a closer look at the registration process of different regions across the globe (South America, U.S. and Europe). Taking the risk assessment of bees as the basis for a comparison, we identify the specific regional data requirements and risk assessment approaches for registration of PPPs. We then discuss the implications of South America's eco-climatic variability and agricultural diversity with respect to the assessment of potential risk associated with PPPs on bees. Finally, we identify global trends in regulatory requirements and discuss the advantages and possible drawbacks of setting common approaches to evaluate risk and for streamlining the registration process.

PT64. Biomarkers of oxidative damage and its relationship with risk factors in a locality environmentally exposed to agrochemicals

L. Odetti, Universidad Nacional del Litoral / Cát Toxicología Farmacología y Bioquímica Legal Fac Bioquímica y Cs Biológicas Santa Fe; M. Dechiara, Universidad Nacional del Litoral, Santa Fe, Argentina. / Cát Toxicología Farmacología y Bioquímica Legal Fac Bioquímica y Cs Biológicas; G. Poletta, University of the Litoral / Cát Toxicología Farmacología y Bioq Legal Fac Bioquímica y Cs Biológicas; D. Marino, Centro de Investigaciones del Medio Ambiente Universidad Nacional de La Plata / Centro de Investigaciones del Medio Ambiente; M. Simoniello, Universidad Nacional del Litoral / Toxicología Farmacología y Bioq Legal

In Santa Fe province, located in the center of Argentina, has been historically one of the main agricultural centers. The production of the main crops (soybean, sunflower, corn, wheat and sorghum) significantly increased in recent years. The aim of this study was to evaluate oxidative damage generated by environmental exposure to agrochemical mixtures in people living in Santo Domingo, Santa Fe, Argentina and to compare them with healthy non-exposed people from Santa Fe city. The study involved 72 subjects, 44 from Santo Domingo (rural area) and 28 healthy donors from Santa Fe city (urban area). Samples were taken at two different moments, in September (low pesticide spraying) and in March (post-period of high spray). The activity of Catalase (CAT), Superoxide dismutase