

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/296688896>

Short Communication: Soil carbon pools in different pasture systems

Article in SPANISH JOURNAL OF AGRICULTURAL RESEARCH · March 2016

DOI: 10.5424/sjar/2016141-7939

CITATIONS

0

READS

65

4 authors, including:



[R.F.V. Carneiro](#)

Universidade Federal de Alfnas

24 PUBLICATIONS 119 CITATIONS

[SEE PROFILE](#)



[Ademir Araujo](#)

Universidade Federal do Piau

106 PUBLICATIONS 868 CITATIONS

[SEE PROFILE](#)

All content following this page was uploaded by [Ademir Araujo](#) on 05 June 2016.

The user has requested enhancement of the downloaded file. All in-text references [underlined in blue](#) are added to the original document and are linked to publications on ResearchGate, letting you access and read them immediately.



Soil carbon pools in different pasture systems

Francisco M. Cardozo Jr.¹, Romero F. V. Carneiro¹, Luiz F. C. Leite² and Ademir S. F. Araujo³

¹ Federal University of Piauí, Agricultural Science Center, Pos-Graduation Program in Animal Science, Teresina, PI, 64049-550, Brazil. ² Embrapa Mid-North, Av. Duque de Caxias, SN, Teresina, PI, 64000-000, Brazil. ³ Federal University of Piauí, Agricultural Science Center, Soil Quality Lab., Teresina, PI, 64000-000, Brazil.

Abstract

The aim of this study was to assess the carbon pools of a tropical soil where the native forest was replaced with different pasture systems. We studied five pasture production systems, including four monoculture systems with forage grasses such as *Andropogon*, *Brachiaria*, *Panicum*, and *Cynodon*, and an agroforestry system as well as a native vegetation plot. Greater availability of fulvic acid was detected in the agroforestry system as compared with that in the other systems. Higher lability of C was detected in the *Andropogon* system during the dry and rainy seasons and during the dry season in *Cynodon*. During the dry season, all pastures systems showed deficits in the net removal of atmospheric CO₂. The structure and practices of the agroforestry system enables more carbon to be sequestered in the soil as compared with the monoculture pasture, suggesting that it is an important practice to mitigate climatic change and to improve soil quality.

Additional key words: humic substances; carbon management; agroforestry system

Abbreviations used: AE (alkaline extract); AFS (agroforestry system); AND (Andropogon); BRA (Brachiaria); C-FAF (C-fulvic acid fraction); C-HAF (C-humic acid fraction); C-HF (C-humin fraction); CMI (carbon management index); CPI (carbon pool index); CYN (Cynodon); HI (humification index); LC (labile organic carbon); L (lability); LI (lability index); NLC (non-labile organic carbon); NV (native vegetation); PAN (Panicum); SOM (soil organic matter); TOC (total organic carbon)

Citation: Cardozo, F. M. Jr.; Carneiro, R. F. V.; Leite, L. F. C.; Araujo, A. S. F. (2016). Short communication: Soil carbon pools in different pasture systems. Spanish Journal of Agricultural Research, Volume 14, Issue 1, e11SC01. <http://dx.doi.org/10.5424/sjar/2016141-7939>.

Received: 29 Apr 2015. **Accepted:** 08 Feb 2016

Copyright © 2016 INIA. This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial (by-nc) Spain 3.0 Licence, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Funding: The authors received no specific funding for this work

Competing interests: The authors have declared that no competing interests exist.

Correspondence should be addressed to Ademir Araujo: asfarauj@yahoo.com.br

Introduction

The pasture ecosystem is characterized by interactions between plants, animals, soil, climate, and management practices implemented by the farmer. Meat and milk productions are important economic activities in Brazil, and the sustainable practices of these production systems contribute to efficient carbon cycling, thereby improving soil quality (Carvalho *et al.*, 2014; Soussana & Lemaire, 2014). Soils are a carbon pool for terrestrial ecosystems; thus, they are relevant with regard to environmental problems associated with global warming and deforestation (Bao *et al.*, 2015).

Conversion from native forest to livestock grazing has decreased soil C storage (Carvalho *et al.*, 2014) and has been the most common land use change in Brazil (Soussana & Lemaire, 2014). It is estimated that Brazil releases more than 1.090 Mt CO₂/year into the

atmosphere through deforestation, burning grasslands, and enteric fermentation by cattle (Bustamante *et al.*, 2012).

Loss of organic carbon can be minimized by managing soil to reduce disturbances and maximize forage productivity using fertilizers and by integrating pastures and trees, which promotes benefits for the animals, increases productivity, litter inputs, nutrients cycling, and water infiltration (Murgeitio *et al.*, 2011). Therefore, the evaluation of the impact of agricultural systems on pasture, through the quantification of total organic carbon (TOC) stocks and humic fractions, as well as assessing carbon lability, is important to maintain soil quality (Yang *et al.*, 2012).

Specifically in pastures, organic input from vegetation and animal activities can contribute to increase the organic C content and consequently cause an impact on C pools (Lopes *et al.*, 2010). As it may vary accord-

ing to different pasture system, we hypothesized that the management method applied for different pasture systems could influence the soil carbon pool. In this context, the objective of this study was to assess the C pools of a tropical soil where the native forest was replaced with different pasture systems.

Material and methods

This study was conducted as a long-term experiment on pasturelands belonging to the Animal Science Department, Agriculture Science Center, Federal University of Piauí, Brazil (05°05'21" S, 42°48'07" W; 74 m asl). The climate is tropical with two seasons: rainy (January to May) and dry (June to December). The mean of precipitation is 1,300 mm/yr. The soil is a Haplic Acrisol. The experimental area presents plots with the following pasture system: a) *Andropogon gayanus* Kunth (AND) [plots without liming and chemical fertilization; production of 2.1 tons/ha (dry weight); 2.21% N and a C/N ratio of 21]; b) *Brachiaria brizantha* (BRA) [plots annually fertilized with 120, 180, and 100 kg/ha urea, triple superphosphate, and potassium chloride, respectively; production of 4.35 tons/ha (dry weight); 0.91% N and a C/N ratio of 37]; c) *Panicum maximum* (PAN) [plots annually fertilized with 70, 80, and 50 kg/ha urea, super triple phosphate, and potassium chloride, respectively; production of 3.0 tons/ha (dry weight); 1.22% N and a C/N ratio of 31]; d) *Cynodon dactylon* (CYN) [plots annually fertilized with 75, 30, and 30 kg/ha urea, super triple phosphate, and potassium chloride, respectively; production of 1.3 tons/ha (dry weight); 1.37% N and a C/N ratio of 36.9]; e) agroforestry system (AFS) [plots composed of grass (*A. gayanus* Kunth) and trees (*Mimosa* sp. and *Thiloua glaucocarpa* Benth); production of 7.4 tons of plant litter (dry weight)/ha]; and f) native vegetation (NV) (plots composed of native plant species, including *Cenostigma macrophyllum*, *Tabebuia serratifolia*, *Hymenaea courbaril*, *Orbignya phalerata*, *Combretum leprosum*, *Guarea kunthiana*, and *Lecythis pisonis*; production of 9.5 tons of plant litter (dry weight)/ha).

Soil sampling was carried out in March (rainy season) and September (dry season) 2014. Soil samples were obtained from three transects from each plot (three points per transect) at a depth of 0–20 cm. The soil samples were ground and passed through a 0.21-mm sieve to determine TOC by wet combustion using a mixture of potassium dichromate and sulfuric acid under heating (Yeomans & Bremner, 1988). Labile organic carbon (LC) was quantified by wet oxidation with 0.33 M KMnO_4 , as described by Blair *et al.* (1995). Non-labile carbon (NLC), that is equivalent to

non-oxidized carbon by KMnO_4 , was calculated as a difference ($\text{NLC} = \text{TOC} - \text{LC}$). Based on the difference between TOC-forest (reference) and TOC systems, a carbon pool index was created (CPI) and calculated as $\text{CPI} = \text{TOC-system}/\text{TOC-forest}$.

According to changes in the proportion of LC (*i.e.*, $L = \text{LC}/\text{NLC}$) in the soil, a lability index (LI) was calculated as $\text{LI} = L \text{ system}/L \text{ reference}$. These two indices were used to calculate the carbon management index (CMI) using the following expression: $\text{CMI} = \text{CPI} \times \text{LI} \times 100$ (Blair *et al.*, 1995). C-CO₂ emission or sequestration rate was estimated for 0–20 cm depth, using native vegetation as a reference (TOC stocks native vegetation – TOC stocks management systems/number of years). A conversion factor of C to CO₂ of 3.67 (molar mass of CO₂/molar mass of C) was used.

Soil humic substances (humic acids, fulvic acids, and humin) were extracted and fractionated using the method recommended by the International Humic Substances Society, as described by Swift (1996). The carbon content of the fulvic acid (C-FAF), humic acid (C-HAF), and humin (C-HF) fractions was measured using the dichromate oxidation method (Yeomans & Bremner, 1988). The ratios of C-HAF by C-FAF and the alkali soluble fractions ($\text{C-FAF} + \text{C-HAF} = \text{AE}$) by H ($\text{AE}/\text{C-HF}$) were calculated to characterize the humified fraction of soil organic matter (SOM). Additionally, the humification index (HI) was calculated using the following formula to estimate the proportion of humified organic matter in relation to TOC content: $\text{HI} = (\text{C-FAF} + \text{C-HAF} + \text{C-HF}) / \text{TOC} \times 100$.

Data were analyzed using one-way analysis of variance, and means were compared using the least significant difference (LSD) values calculated at the 5% level of significance. All analyses were performed using STATISTICA 7.1 (StatSoft).

Results and discussion

The C-HAF was lower than the C-FAF during the wet and dry seasons in all systems (Table 1). The lower C-HAF found in all systems may have occurred because this humic fraction can easily migrate in soil with high porosity (Martins *et al.*, 2009). Therefore, in the evaluated areas the soil porosity may facilitate the movement of C-HAF through the soil horizons. The C-HAF values in plots under *Brachiaria*, *Panicum* and *Cynodon* were similar to those in native vegetation during the dry season. Greater C-FAF availability was detected in the agroforestry system during the rainy and dry seasons because this plot showed higher TOC and Ca⁺² content, which is a favorable condition for the complexation of this C pool in the soil (Barros *et al.*,

Table 1. Carbon content from humic acid (HAF), fulvic acid (FAF), and humin (HF) fractions, HAF:FAF ratio (HAF/FAF), alkaline extract and HF ratio (HAF+FAF/HF), and humification index (HI) under different pastures and during two seasons (rainy and dry). AND - *Andropogon gayanus*; BRA - *Brachiaria brizantha*; PAN - *Panicum maximum*; CYN - *Cynodon dactylon*; AFS – Agroforestry system; NV - native vegetation.

	Rainy season						Dry season					
	AND	BRA	PAN	CYN	AFS	NV	AND	BRA	PAN	CYN	AFS	NV
HAF (g/kg)	1.1 b	0.7 c	0.7 c	0.5 c	1.3 a	1.2 ab	1.30 b	0.70 c	0.60 c	0.70 c	1.50 a	0.80 c
FAF (g/kg)	2.2 b	2.1 c	2.0 c	2.0 c	3.0 a	2.7 b	2.80 a	1.90 b	2.00 b	1.80 b	3.20 a	2.10 b
HF (g/kg)	5.3 b	4.9 c	4.8 c	5.0 c	7.2 a	4.3 d	6.00 c	6.10 b	6.20 bc	6.30 bc	5.90 c	7.00 a
HAF/FAF	0.50 a	0.33a	0.35 a	0.25 a	0.43 a	0.44 a	0.46 a	0.36 a	0.30 a	0.38 a	0.46 a	0.38 a
(HAF+FAF)/HF	0.62 b	0.57 b	0.56 b	0.50 b	0.59 b	0.90 a	0.68 b	0.42 c	0.41 c	0.39 c	0.79 a	0.41 c
HI (%)	94.5 a	95.0 a	94.9 a	94.8 a	96.6 a	95.3 a	95.9a	88.7b	95.7a	95.6a	96.3a	94.2a
TOC (g/kg)	9.1 b	8.1 b	7.9 b	7.9 b	11.9 a	8.6 b	10.5 b	9.8 c	9.1 c	11.0 a	10.5 b	9.1c

Means followed by the same letter in each row and season do not differ statistically from each other at $p < 0.05$ (LSD test).

2012). The high carbon content from FAF in this system may have stimulated microbial diversity because that fraction is more easily used as an energy source by soil microorganisms, generating more negative charges and improving nutrient cycling and ecosystem productivity (Moraes *et al.*, 2011).

The C-HF was higher than the C-HAF and C-FAF in all evaluated plots during both seasons, suggesting a greater stability of humin to mineralization than of humic and fulvic acids (Barros *et al.*, 2012). The C-HF was higher in the native vegetation (dry season) and agroforestry system (rainy season), indicating the presence of more stable humus, low degradation, and strong stimulus to soil microbial activity (Barros *et al.*, 2012). The presence of lignin derived from plant residues increased also the humin in soil (Carvalho *et al.*, 2014). This increase in humin in soil is resulting from the loss of oxidative C and, at the same time, an increase in the C stable (Moraes *et al.*, 2011). Therefore, the permanent inputs of organic C from herbaceous plants and

trees in the agroforestry system may indicate a high potential for nutrient cycling and increased fertility.

No significant difference in the HA:FA ratio was observed between areas. The ratio (HAF + FAF)/HF in the native vegetation (rainy season) and agroforestry system (dry season) was higher than in the pasture systems, indicating that these systems contain more chemically stable organic C, for which the turnover time is approximately 2,000 years (Chan *et al.*, 2001). The humification index did not differ between systems, which was likely because of the low proportion of labile C found in all plots. These results are indicative of soils with low organic matter input; however, with potential to stimulate the microbial growth in these ecosystems because of the complexity of their organic molecules (Bausenwein *et al.*, 2008).

Higher lability of C was detected in the soil with *Andropogon* during the dry and rainy seasons, whereas soil with *Cynodon* showed higher lability of C during the dry season (Table 2). All plots showed a

Table 2. Labile carbon (LC), labile carbon: total organic carbon ratio (LC/TOC), non-labile carbon (CNL), lability (L), lability index (LI), carbon pool index (CPI), carbon management index (CMI), and carbon stock (C stock) under different pastures and during two seasons (rainy and dry). AND - *Andropogon gayanus*; BRA - *Brachiaria brizantha*; PAN - *Panicum maximum*; CYN - *Cynodon dactylon*; AFS – Agroforestry system; NV - native vegetation.

	Rainy season						Dry season					
	AND	BRA	PAN	CYN	AFS	NV	AND	BRA	PAN	CYN	AFS	NV
LC (g/kg)	0.96 a	0.86 b	0.74 c	0.69 d	0.83 b	0.60 d	0.87 a	0.78 b	0.73 c	0.89 a	0.76 b	0.67 d
LC/TOC (%)	10.50 a	10.60 a	9.30 b	8.70 b	6.90 c	6.90 c	8.20 b	7.90 c	8.02 b	9.60 a	6.90 d	6.30 d
CNL (g/kg)	8.14 b	7.24 b	7.16 b	7.21 b	11.07 a	8.00 b	9.63 cd	9.02 c	8.37 d	8.31 cd	10.24 a	9.83 b
L	0.11 a	0.10 b	0.10 c	0.09 c	0.07 d	0.07 d	0.09 a	0.08 b	0.08 b	0.10 a	0.07 c	0.06 c
LI	1.57 a	1.57 b	1.42 c	1.28 c	1.00 d	–	1.50 a	1.33 b	1.33 b	1.66 a	1.10 c	–
CPI	1.05 b	0.94 b	0.91 b	0.91 b	1.38 a	–	1.00 a	0.93 b	0.86 b	0.87 b	1.04 a	–
CMI	164.8a	147.5 b	129.2 d	117.3 e	138.8 c	–	150.0 a	123.6 b	111.3 b	143.1 a	114.4 b	–
TOC (g/kg)	9.1 b	8.1 b	7.9 b	7.9 b	11.9 a	8.6 b	10.5 b	9.8 c	9.1 c	9.2 c	11.0 a	10.5 b
C stock (Mg C/ha)	25.9 b	20.7 c	21.9 c	24.3 b	32.8 a	25.4 b	27.3 b	23.5 d	23.6 d	25.7 c	30.5 a	29.4 b

Means followed by the same letter in each row and season do not differ statistically from each other at $p < 0.05$ (LSD test).




Figure 1. Emission and sequestration rate of C-CO₂. Means with similar small letters in the dry season (red bars) and capital letters in the rainy season (blue bars), do not differ significantly according to LSD test ($p < 0.05$). Plots: AND, *Andropogon gayanus*; BRA, *Brachiaria brizantha*; PAN, *Panicum maximum*; CYN, *Cynodon dactylon*; AFS, Agroforestry system. Positive values indicate C sequestration and negative values indicate C loss.

CMI < 100 (Table 2), indicating high plant input and minimal soil disturbance (Leite *et al.*, 2014). According to Blair *et al.* (1995), a CMI < 100 indicates a strong negative impact of management practices on a soil ecosystem. Specifically, the agroforestry system showed a higher CMI than in the *Brachiaria*, *Panicum*, and *Cynodon*. The absence of trees and different pasture management practices reduce the C management index over time, reflecting a decrease in the potential to restore pre-existing carbon stocks (Leite *et al.*, 2014).

The agroforestry system showed highest carbon stock values and high C sequestration rate for the rainy and dry seasons, whereas *Brachiaria*, *Panicum*, and *Cynodon* showed the lowest C stock and a high C loss (Fig. 1). It means that in multicropping systems, such as agroforestry system, the presence of pioneer tree species and the lesser removal of plant residues contribute significantly to mitigate the loss of C and decrease greenhouse gas emissions. Also, agroforestry systems have higher potential to build up and sequester C in soils because of the increased rates of organic matter addition and retention (Lenka *et al.*, 2012).

In conclusion, conversion of native vegetation to pasture system causes changes in C pools, increasing CO₂ emissions into the atmosphere. The agroforestry system has the potential to sequester more carbon in the soil than the pasture system, and it may be an alternative to produce forage for animal production.

References

Bao X, Li Q, Hua J, Zhao T, Liang W, 2015. The interactive effects of elevated ozone and wheat cultivars on soil microbial community composition and metabolic diversity.

- Appl Soil Ecol 87: 11-18. <http://dx.doi.org/10.1016/j.apsoil.2014.11.003>
- Barros KRM, Lima HV, Canellas LP, Kern DC, 2012. Fracionamento químico da matéria orgânica e caracterização física de Terra Preta de Índio. Rev Cienc Agrar 55: 44-51. <http://dx.doi.org/10.4322/rca.2012.037>
- Bausenwein U, Gattinger A, Langer U, Embacher A, Hartmann HP, Sommer M, Munch JC, Schloter M, 2008. Exploring soil microbial communities and water extractable organic matter: availability and interactions in an integratedly managed arable soil. J Appl Soil Ecol 140: 67-77. <http://dx.doi.org/10.1016/j.apsoil.2008.03.006>
- Blair GJ, Lefroy RDB, Lisle L. 1995. Soil carbon fractions based on their degree of oxidation, and the development of a carbon management index for agricultural systems. Aust J Agr Res 46: 1459-1466. <http://dx.doi.org/10.1071/AR951459>
- Bustamante MMC, Nobre CA, Smeraldi R, 2012. Estimating greenhouse gas emissions from cattle raising in Brazil. Clim Ch 115: 559-577. <http://dx.doi.org/10.1007/s10584-012-0443-3>
- Carvalho JLN, Raucci GS, Frazao LA, Cerri CEP, Bernoux M, Cerri CC, 2014. Crop pasture rotation: A strategy to reduce soil greenhouse gas emissions in the Brazilian Cerrado. Agr Ecosyst Environ 183:167-175. <http://dx.doi.org/10.1016/j.agee.2013.11.014>
- Chan KY, Bowman A, Oates A, 2001. Oxidizable organic carbon fractions and soil quality changes in a paleustalf under different pasture leys. Soil Sci 166: 61-67. <http://dx.doi.org/10.1097/00010694-200101000-00009>
- Leite LFC, Iwata BF, Araujo ASF, 2014. Soil organic matter pools in a tropical savanna under agroforestry system in northeastern Brazil. Rev Árvore 38: 1-8. <http://dx.doi.org/10.1590/S0100-67622014000400014>
- Lenka NK, Choudhury PR, Sudhishri S, Dass A, Patnaik US, 2012. Soil aggregation, carbon build up and root zone soil moisture in degraded sloping lands under selected agroforestry based rehabilitation systems in eastern India. Agr Ecosyst Environ 150: 54-62. <http://dx.doi.org/10.1016/j.agee.2012.01.003>
- Lopes MM, Salviano, AAC, Araujo ASF, Nunes LAPL, Oliveira ME, 2010. Changes in soil microbial biomass and activity in different Brazilian pastures. Span J Agric Res 8: 1253-1259. <http://dx.doi.org/10.5424/sjar/2010084-1411>
- Martins EL, Coringa JES, Weber OLS, 2009. Carbono orgânico nas frações granulométricas e substâncias húmicas de um Latossolo Vermelho. Amarelo distrófico-LVAd sob diferentes agrossistemas. Acta Amaz 39: 655-660. <http://dx.doi.org/10.1590/S0044-59672009000300021>
- Moraes GM, Xavier FA, Mendonca E, Filho JA, Oliveira JN, 2011. Chemical and structural characterization of soil humic substances under agroforestry and conventional systems. Rev Bras Ci Solo 35: 1597-1608. <http://dx.doi.org/10.1590/S0100-06832011000500014>
- Murgeitio E, Calle Z, Uribe F, Calle A, Solorio B, 2011. Native trees and shrubs for the productive rehabilitation of tropical cattle ranching lands. Forest Ecology and Management 261: 1654-1663. <http://dx.doi.org/10.1016/j.foreco.2010.09.027>
- Soussana JF, Lemaire G, 2014. Coupling carbon and nitrogen cycles for environmentally sustainable intensification of

- grasslands and crop-livestock systems. *Agr Ecosyst Environ* 190: 9-17. <http://dx.doi.org/10.1016/j.agee.2013.10.012>
- Swift RS, 1996. Organic matter characterization. In: *Methods of soil analysis*; Sparks DL *et al.* (eds). pp. 1011-1020. Soil Sci Soc Am, Madison, WI, USA.
- Yang X, Ren W, Sun B, Zhang S, 2012. Effects of contrasting soil management regimes on total and labile soil organic carbon fractions in a loess soil in China. *Geoderma* 177: 49-56. <http://dx.doi.org/10.1016/j.geoderma.2012.01.033>
- Yeomans C, Bremner JM, 1988. A rapid and precise method for routine determination of organic carbon in soil. *Comm. Soil Sci Plant Anal* 19: 1467-1476. <http://dx.doi.org/10.1080/00103628809368027>