

## ***Leptocybe invasa* (HYMENOPTERA: EULOPHIDAE) ON *Eucalyptus* SEEDLINGS IN SANTA CATARINA STATE, BRAZIL**

### ***Leptocybe invasa* (HYMENOPTERA: EULOPHIDAE) EM MUDAS DE EUCALIPTO NO ESTADO DE SANTA CATARINA, BRASIL**

Leonardo Rodrigues Barbosa<sup>1</sup> Angelo Peruffo Rodrigues<sup>2</sup> Amanda Rodrigues de Souza<sup>3</sup> Barbara de Oliveira Poretz<sup>4</sup> Carlos Frederico Wilcken<sup>5</sup> José Cola Zanuncio<sup>6</sup>

#### **ABSTRACT**

The blue gum chalcid, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae), native from Australia, can severely damage eucalyptus (Myrtaceae) plants. This insect was found on *Eucalyptus grandis* seedlings in May 2015, in Treze de Maio municipality in Santa Catarina state, Brazil.

**Keywords:** blue gum chalcid; invasive species; nurseries; spread.

#### **RESUMO**

A vespa-da-galha, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae), originária da Austrália, pode danificar severamente plantas de eucalipto (Myrtaceae). Este inseto foi registrado em mudas de *Eucalyptus grandis* em maio de 2015 no município de Treze de Maio, Estado de Santa Catarina, Brasil.

**Palavras-chave:** vespa-da-galha-do-eucalipto; espécies exóticas; mudas; dispersão.

#### **INTRODUCTION**

In 2016, Santa Catarina state had 116,240 ha of *Eucalyptus* plantations, occupying the 13<sup>th</sup> position of the eucalyptus production in Brazil (INDÚSTRIA BRASILEIRA DE ÁRVORES, 2017).

Insect pests have been reported damaging forest plantations, including *Eucalyptus* in several countries (DOOLEY; SIX; POWELL, 2015; FRANÇOIS; THIFFAULT, 2015). Gall-inducers of the Hymenoptera are pests for this plant (CARVALHO-FERNANDES et al., 2016). The feeding habits of gall-inducers change the anatomy of host plant tissues in response to damage (DIAS et al., 2013).

The blue gum chalcid, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae) is a gall wasp and important pest of *Eucalyptus* spp. (Myrtaceae) (MENDEL et al., 2004). This insect, native from Australia, was detected in the Mediterranean Basin and the Middle East in 2000 (MENDEL et al., 2004) and has spread to 29 countries (ZHENG et al., 2014).

In Brazil, *L. invasa* was found on eucalyptus seedlings in nurseries and on a *Eucalyptus camaldul-*

1 Engenheiro Agrônomo, Dr., Pesquisador da Empresa Brasileira de Pesquisa Agropecuária - Embrapa Florestas, Estrada da Ribeira, Km 111, CEP 83411-000, Colombo (PR), Brasil. leonardo.r.barbosa@embrapa.br

2 Engenheiro Agrônomo, Mestrando do Departamento de Zoologia, Universidade Federal do Paraná, Rua dos Funcionários, 1540, Cabral, CEP 81531-990, Curitiba (PR), Brasil. aperufforodrigues@gmail.com

3 Engenheira Agrônoma, Dra., Professora Substituta do Departamento de Proteção Vegetal, Faculdade de Ciências Agronômicas - Universidade Estadual Paulista "Júlio de Mesquita Filho", Rua José Barbosa de Barros, 1780, Fazenda Lageado, CEP 18610-307, Botucatu (SP), Brasil. agroamandarodrigues@yahoo.com.br

4 Engenheira Florestal, MSc., Doutoranda do Departamento de Proteção Vegetal, Faculdade de Ciências Agronômicas - Universidade Estadual Paulista "Júlio de Mesquita Filho", Rua José Barbosa de Barros, 1780, Fazenda Lageado, CEP 18610-307, Botucatu (SP), Brasil. bah\_poretz@hotmail.com

5 Engenheiro Agrônomo, Dr., Professor do Departamento de Proteção Vegetal, Faculdade de Ciências Agronômicas - Universidade Estadual Paulista "Júlio de Mesquita Filho", Rua José Barbosa de Barros, 1780, Fazenda Lageado, CEP 18610-307, Botucatu (SP), Brasil. cwilcken@fca.unesp.br

6 Engenheiro Florestal, Dr., Professor do Departamento de Entomologia/BIOAGRO, Universidade Federal de Viçosa, Avenida P.H. Rolfs, s/n, Centro, CEP 36571-900, Viçosa (MG), Brasil. zanuncio@ufv.br

Recebido para publicação em 2/03/2017 e aceito em 24/08/2017

*lensis* Dehnh x *Eucalyptus grandis* W. Hill ex Maiden hybrid in the northeast of Bahia state in 2008 (COSTA et al., 2008). It has spread to Rio Grande do Sul (GARLET et al., 2013), Minas Gerais (FERNANDES et al., 2014), Goiás (PEREIRA et al., 2014), Espírito Santo, Maranhão, Mato Grosso, Mato Grosso do Sul, Pará, Paraná, Pernambuco, São Paulo, Sergipe, and Tocantins states (WILCKEN et al., 2015).

*Leptocybe invasa* adults are between 1.1 to 1.4 mm long. Their body is dark brown, with a blue-green metallic shine, mainly on the head and mesosoma, with a yellowish color at the basal antenna joints and brown legs (MENDEL et al., 2004). The life cycle (egg to adult) of *L. invasa* lasted 132.6 days on *Eucalyptus camaldulensis* in a greenhouse in Israel (MENDEL et al., 2004). The insect development period was 59.5 days on *Eucalyptus tereticornis* Smith in India (KAVITHA-KUMARI et al., 2010) and 46 days on *E. camaldulensis* in Thailand (SANGTONGPRAOW; CHARERNSON; SIRIPATANADILOK, 2011).

*Leptocybe invasa* females oviposit on the epidermis of the leaf petiole midrib, causing leaf senescence and gall formation on growing branch parenchyma. These galls may stunt young plants growth causing deformations on stems and leaves (MENDEL et al., 2004; NYEKO; MUTITU; DAY, 2009). This may reduce productivity (COSTA et al., 2008) especially of susceptible genotypes. Thus, efficient strategies to manage *L. invasa* need to be developed (ZHENG et al., 2014) focusing on plant resistance and biological control.

*Eucalyptus* genotypes resistant to *L. invasa* were identified in Laos, Thailand, Vietnam (OATES et al., 2012), Kenya, South Africa and Uganda (NYEKO et al., 2010). Tolerance and/or resistance of *Eucalyptus* clones and species make them potentially suitable for planting in areas with different *L. invasa* infestation levels (NYEKO; MUTITU; DAY, 2009; DITTRICH-SCHRÖDER et al., 2012). *Eucalyptus camaldulensis*, *Eucalyptus grandis* and *Eucalyptus tereticornis* were more susceptible to this pest in Vietnam (THU; DELL; BURGESS, 2009). *Eucalyptus dunnii* Maiden, *Eucalyptus nitens* (Deane & Maiden) Maiden, *Eucalyptus saligna* Smith x *Eucalyptus urophylla*, *Eucalyptus smithii* (R.T. Baker) and *Eucalyptus urophylla* S.T. Blake (DITTRICH-SCHRÖDER et al., 2012) and *Eucalyptus benthamii* Maiden & Cambridge and *Corymbia citriodora* (Hook.) K.D. Hill and L.A.S. Johnson (WILCKEN et al., 2015) had low or no infestation by *L. invasa*.

Biological control is an effective strategy to manage insect pests such a *L. invasa* (KIM et al., 2008; KULKARNI et al., 2010). Eulophidae parasitoids are natural enemies of eucalyptus pests (PEREIRA et al. 2008a; 2008b), including *Selitrichodes neseri* Kelly & La Salle in South Africa (KELLY et al., 2012) and Brazil (SOUZA, 2016; MASSON et al., 2017; DE SOUZA et al., 2018) and *Quadrastichus mendeli* Kim & La Salle and *Selitrichodes kryceri* Kim & La Salle were introduced in Israel, for the biological control of *L. invasa* (KIM et al., 2008). The search for natural *L. invasa* enemies identified the Hymenoptera parasitoids *Aprostocetus* sp. and *Megastigmus* sp. (Torymidae), *Parallelaptera* sp. (Mymaridae) and *Telenomus* sp. (Scelionidae) (GUPTA; POORANI, 2009). In Brazil, the native species *Megastigmus brasiliensis* Doğanlar, Zaché & Wilcken (Hymenoptera: Torymidae) is associated with *L. invasa* (DOĞANLAR; ZACHÉ; WILCKEN, 2013).

The aim of this study was to report the occurrence of *L. invasa* in Santa Catarina state, expanding the distribution of this pest in Brazil.

## MATERIALS AND METHODS

The presence of *L. invasa* on *Eucalyptus grandis* seedlings was confirmed based on the characteristic gall symptoms normally found on the central leaf veins, the petioles and green stems. This assessment was made during May 2015 in a forest nursery in Treze de Maio municipality in Santa Catarina state, Brazil (28° 35' S latitude × 49° 10' W longitude and 169 m above sea level). The average minimum and maximum temperatures in Santa Catarina state during the autumn season were 15.36 °C and 24.93 °C, respectively, with 1,444 mm average annual rainfall (WREGGE et al., 2011).

*Eucalyptus* seedlings were screened for symptoms of this insect in Araranguá, Iporã do Oeste, Jacinto Machado, Palma Sola, São João do Oeste, São José do Cedro, São Miguel do Oeste, and Urussanga municipalities in the Santa Catarina state.

## RESULTS AND DISCUSSION

The blue gum chalcid was recorded on *Eucalyptus grandis* seedlings in Treze de Maio, Santa Catarina state, Brazil. *Eucalyptus grandis* seedlings in full sun were more infested by *L. invasa* than those predominantly in the shade. Galls induced by the pest occurred mostly on the central leaf veins, the petioles and the green stems of these plants (Figure 1). Approximately 10% of nursery seedlings were infested and subsequently eliminated. The rest of the seedlings were treated with chemicals registered for this pest in Brazil.



FIGURE 1: *Eucalyptus grandis* seedlings infested by *Leptocybe invasa* (Hymenoptera: Eulophidae) in Treze de Maio municipality, Santa Catarina state, Brazil.

FIGURA 1: Mudanças de *Eucalyptus grandis* infestadas por *Leptocybe invasa* (Hymenoptera: Eulophidae) no município de Treze de Maio, no Estado de Santa Catarina, Brasil.

*Eucalyptus grandis* seedlings with galls from Treze de Maio were sent to the Embrapa Forests in Colombo, Paraná state, Brazil. The insects that emerged were identified as *L. invasa* based on its antenna, forewing, gaster, head and mesosoma (MENDEL et al., 2004).

In addition to detecting *L. invasa* in Treze de Maio, this insect was also found in eight other municipalities of Santa Catarina state, Brazil, which shows the presence of this insect on opposite sides of the state (Figure 2). This demonstrates that this insect may be present throughout the State.




FIGURE 2: The blue gum chalcid, *Leptocybe invasa* (Hymenoptera: Eulophidae), distribution in Santa Catarina State, Brazil.

FIGURA 2: Distribuição da vespa-da-galha, *Leptocybe invasa* (Hymenoptera: Eulophidae) no Estado de Santa Catarina, Brasil.

Damage by *L. invasa* reduces plant growth and may cause plant death in severe cases. The presence of galls may be a diagnostic indicator for *L. invasa* presence (WILCKEN et al., 2015). Forest nurseries growing eucalyptus seedlings need periodic monitoring by direct visual or indirect assessment through yellow sticky traps. Chemical control with the systemic and contact insecticides such as Imidacloprid, Thiamethoxam and Fipronil are recommended for application on eucalyptus seedlings in forest nurseries and it is currently the most used *Leptocybe invasa* management strategy (WILCKEN et al., 2015).

Biological control is the major strategy in integrated management programs for this pest with importing and rearing of the parasitoid *Selitrichodes neseri* in Brazil and South Africa. In Thailand, the native parasitoid *Megastigmus thitipornae* Doğanlar & Hassan (Hymenoptera: Torymidae) is being studied (SANGTONGPRAOW; CHARERNSOM, 2013). In Israel, *Selitrichodes kryceri* and *Quadrastichus mendeli* are successfully used to manage *L. invasa* (KIM et al., 2008). Finding a suitable host to mass rear its natural enemies is the major question in developing biological control programs for *L. invasa* (FAVERO et al., 2013).

## CONCLUSION

The rapid spread and damage caused by *L. invasa* in various Brazilian states shows the need to develop efficient strategies to manage this insect pest.

## ACKNOWLEDGEMENTS

To Health Protection Vegetable Integrated Agricultural Development Company of Santa Catarina (CIDASC) for sending seedlings attacked. To “Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq)”, “Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG)” and “Programa Cooperativo sobre Proteção Florestal - Instituto de Pesquisas e Estudos Florestais” (PROTEF - IPEF)” for the financial support. Dr. Phillip John Villani (The University of Melbourne, Australia) revised and corrected the English language used in this manuscript.

## REFERENCES

- CARVALHO-FERNANDES, S. P. et al. Diversity of insect galls associated with coastal shrub vegetation in Rio de Janeiro, Brazil. **Anais da Academia Brasileira de Ciências**, Rio de Janeiro, v. 88, n. 3, p. 1407-1418, 2016.
- COSTA, V. A. et al. *Eucalyptus* gall wasp, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae) in Brazil: New forest pest reaches the New World. **Revista de Agricultura**, Piracicaba, v. 83, n. 2, p. 136-139, 2008.
- DE SOUZA et al. Longevity and survival of *Leptocybe invasa* (Hymenoptera: Eulophidae), an invasive gall inducer on *Eucalyptus*, with different diets and temperatures. **PeerJ**, v. 6, n. 1, p. 5265, 2018.
- DIAS, G. C. et al. Developmental pathway from leaves to galls induced a sap-feeding insect on *Schinus polygamus* (Cav.) Cabrera (Anacardiaceae). **Anais da Academia Brasileira de Ciências**, Rio de Janeiro, v. 85, n. 1, p. 187-200, 2013.
- DITTRICH-SCHRÖDER, G. et al. Diversity in *Eucalyptus* susceptibility to the gall-forming wasp *Leptocybe invasa*. **Agricultural and Forest Entomology**, Malden, v. 14, n. 4, p. 419-427, 2012.
- DOĞANLAR, M.; ZACHÉ, B.; WILCKEN, C. F. New species of *Megastigmus* (Hymenoptera: Torymidae: Megastigminae) from Brazil. **Florida Entomologist**, Lutz, v. 96, n. 1, p. 196-199, 2013.
- DOOLEY, E. M.; SIX, D. L.; POWELL, J. A. A comparison of mountain pine beetle (Coleoptera: Curculionidae, Scolytinae) productivity and survival in lodgepole and whitebark pine after a region-wide cold weather event. **Forest Science**, Bethesda, v. 61, n. 2, p. 235-246, 2015.
- FAVERO, K. et al. *Tenebrio molitor* (Coleoptera: Tenebrionidae) as an alternative host for the parasitoid *Trichospilus diatraeae* (Hymenoptera: Eulophidae). **Revista Colombiana de Entomología**, Bogotá, v. 39, n. 1, p. 47-48, 2013.
- FERNANDES, B. F. et al. *Leptocybe invasa* (Hymenoptera: Eulophidae), an exotic pest of *Eucalyptus*, in Minas Gerais State, Brazil. **Florida Entomologist**, Lutz, v. 97, n. 2, p. 824-826, 2014.
- FRANÇOIS, H.; THIFFAULT, N. Vegetation management, stock type, and scarification effects on white pine weevil incidence and early Norway spruce growth in Québec, Canada. **Forest Science**, Bethesda, v. 61, n. 5, p. 966-972, 2015.
- GARLET, J. et al. *Leptocybe invasa* em *Eucalyptus* sp. no estado do Rio Grande do Sul, Brasil. **Ciência Rural**, Santa Maria, v. 43, n. 12, p. 2175-2177, 2013.
- GUPTA, A.; POORANI, J. Taxonomic studies on a collection of Chalcidoidea (Hymenoptera) from India with new distribution records. **Journal of Threatened Taxa**, Coimbatore, v. 1, n. 5, p. 300-304, 2009.
- INDÚSTRIA BRASILEIRA DE ÁRVORES. **Indústria Brasileira de Árvores**. São Paulo: IBÁ, 2017. Disponível em: <[http://iba.org/images/shared/Biblioteca/Iba\\_RelatorioAnual2017.pdf](http://iba.org/images/shared/Biblioteca/Iba_RelatorioAnual2017.pdf)>. Acesso em: 30 ago. 2017.
- KAVITHA-KUMARI, N. et al. Biology of eucalyptus gall wasp, *Leptocybe invasa* Fisher and La Salle (Hymenoptera: Eulophidae). **Karnataka Journal of Agricultural Sciences**, Dharwad, v. 23, n. 1, p. 211-212, 2010.
- KELLY, J. et al. *Selitrichodes neseri* n. sp., a new parasitoid of the eucalyptus gall wasp *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae: Tetrastichinae). **Zootaxa**, Auckland, v. 3333, n. 1, p. 50-57, 2012.
- KIM, I.-K. et al. Taxonomy, biology, and efficacy of two Australian parasitoids of the eucalyptus. **Zootaxa**, Auckland, v. 1910, n. 1, p. 1-20, 2008.
- KULKARNI, H. et al. Release and recovery of parasitoids in *Eucalyptus* against gall wasp, *Leptocybe invasa* (Hymenoptera: Eulophidae) under green house. **Karnataka Journal of Agricultural Sciences**, Dharwad, v. 23, n. 1, p. 91-92, 2010.
- MASSON, M. V. et al. *Selitrichodes neseri* (Hymenoptera: Eulophidae) recovered from *Leptocybe invasa* (Hymenoptera: Eulophidae) galls after initial release on *Eucalyptus* (Myrtaceae) in Brazil, and data on its biology. **Florida Entomologist**, Lutz, v. 100, n. 2, p. 589-593, 2017.
- MENDEL, Z. et al. Taxonomy and biology of *Leptocybe invasa* gen. & sp. n. (Hymenoptera: Eulophidae), an invasive gall inducer on *Eucalyptus*. **Australian Journal of Entomology**, Canberra, v. 43, n. 1, p. 101-

113, 2004.

NYEKO, P. et al. Variations in *Leptocybe invasa* (Hymenoptera: Eulophidae) population intensity and infestation on *Eucalyptus* germplasms in Uganda and Kenya. **International Journal of Pest Management**, London, v. 56, n. 2, p. 137-144, 2010.

NYEKO, P.; MUTITU, K. E.; DAY, R. K. *Eucalyptus* infestation by *Leptocybe invasa* in Uganda. **African Journal of Ecology**, Malden, v. 47, n. 3, p. 299-307, 2009.

OATES, C. N. A hypothetical *Eucalyptus grandis* defense model against *Leptocybe invasa* based on transcriptome sequencing. **South African Journal of Botany**, Pietermaritzburg, v. 79, n. 1, p. 204, 2012.

PEREIRA, F. F. et al. New record of *Trichospilus diatraeae* as a parasitoid of the eucalypt defoliator *Thyrinteina arnobia* in Brazil. **Phytoparasitica**, Heidelberg, v. 36, n. 3, p. 304-306, 2008a.

PEREIRA, F. F. et al. Species of Lepidoptera defoliators of *Eucalyptus* as new host for the parasitoid *Palmistichus elaeisis* (Hymenoptera: Eulophidae). **Brazilian Archives of Biology and Technology**, Curitiba, v. 51, n. 2, p. 259-262, 2008b.

PEREIRA, J. M. et al. Record of the *Leptocybe invasa* in the state Goiás, Brazil. **Ciência Rural**, Santa Maria, v. 44, n. 10, p. 1721-1724, 2014.

SANGTONGPRAOW, B.; CHARERNSOM, K. Evaluation of parasitism capacity of *Megastigmus thitipornae* Doganlar & Hassan (Hymenoptera: Torymidae), the local parasitoid of *Eucalyptus* gall wasp, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae). **Kasetsart Journal: Natural Science**, Bangkok, v. 47, n. 2, p. 191-204, 2013.

SANGTONGPRAOW, B.; CHARERNSON, K.; SIRIPATANADILOK, S. Longevity, fecundity and development time of *Eucalyptus* gall wasp, *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae) in Kanchanaburi Province, Thailand. **Thai Journal of Agricultural Science**, Bangkok, v. 44, n. 3, p. 155-163, 2011.

SOUZA, A. R. **Aspectos biológicos de *Leptocybe invasa* (Hymenoptera: Eulophidae) e de seu parasitóide *Selitrichodes neseri* (Hymenoptera: Eulophidae) e levantamento de vespas galhadoras em florestas de eucalipto na Austrália**. 2016. 75 f. Tese (Doutorado) - Universidade Estadual Paulista "Júlio de Mesquita Filho", Faculdade de Ciências Agrônômicas, Botucatu, 2016.

THU, P. Q.; DELL, B.; BURGESS, T. I. Susceptibility of 18 eucalypt species to the gall wasp *Leptocybe invasa* in the nursery and young plantations in Vietnam. **Science Asia**, Langkawi, v. 35, n. 1, p. 113-117, 2009.

WILCKEN, C. F. et al. A vespa-da-galha-do-eucalipto, *Leptocybe invasa* Fisher & La Salle. In: VILELA, E. F.; ZUCCHI, R. A. (Org.). **Pragas introduzidas no Brasil: insetos e ácaros**. Piracicaba: FEALQ, 2015. p. 835-844.

WREGE, M.S. et al. **Atlas Climático da Região Sul do Brasil: Estados do Paraná, Santa Catarina e Rio Grande do Sul**. Pelotas: Embrapa Clima Temperado; Colombo: Embrapa Florestas, 2011. 336 p.

ZHENG, X. L. et al. A review of invasive biology, prevalence and management of *Leptocybe invasa* Fisher & La Salle (Hymenoptera: Eulophidae: Tetrastichinae). **African Entomology**, Pretoria, v. 22, n. 1, p. 68-79, 2014.