



Agronomic Performance of Sunflower Hybrids in Brazilian Savannah Region

Flávio Carlos Dalchiavon^{1*}, Marcos Birck¹, Diogo Stasiak¹, Rosivaldo Hiolanda¹ and Claudio Guilherme Portela de Carvalho²

¹*Department of Agronomy, Federal Institute of Education, Science and Technology of Mato Grosso/IFMT, 78360-000, Campo Novo do Parecis - MT, Brazil.*

²*Brazilian Agricultural Research Corporation, Embrapa Soybean, 86001-970, Londrina - PR, Brazil.*

Authors' contributions

This work was carried out in collaboration between all authors. The authors of note 1 elaborated the study, collected the data, performed the statistical analysis and wrote the manuscript. Author of note 2 elaborated the study, performed the statistical analysis and wrote the manuscript. All authors compile the literature review and approve the final manuscript.

Article Information

DOI: 10.9734/JEAI/2018/44619

Editor(s):

(1) Dr. Marco Aurelio Cristancho, Professor, National Center for Coffee Research, CENICAFÉ, Colombia.

Reviewers:

(1) Habu Saleh Hamisu, National Horticultural Research Institute, Nigeria.

(2) Parmeshwar Sahu, Indira Gandhi Agriculture University, India.

(3) Silvane Isabel Brand, Luiz de Queiroz College of Agriculture, University of São Paulo, Brazil.

Complete Peer review History: <http://www.sciencedomain.org/review-history/27228>

Original Research Article

Received 23 August 2018

Accepted 08 November 2018

Published 15 November 2018

ABSTRACT

The establishment of sunflower crop in the Brazilian savannah region depends, among other factors, on the use of cultivars with adaptation to the growing regions and adequate characteristics, such as drought tolerance and resistance to main diseases and early cycle. The objective of this work was to evaluate the agronomic performance of sunflower cultivars sown in the second summer crop in Campo Novo do Parecis (Mato Grosso), the main sunflower producing region in the Brazilian savannah, in 2014 and 2015. The experimental design was a randomised complete block with eight treatments (AGUARÁ 04, AGUARÁ 06, CF 101, GNZ NEON, HELIO 251, MG 305, SYN 045 and SYN 3950HO) and four replications. Sowing was carried out on March 7, 2014, and March 3, 2015. Plant height, head diameter, harvest index, 1000-achene mass, achene productivity, oil content and oil yield were assessed. Data were submitted to analysis of variance and the Scott-Knott test at 5%

*Corresponding author: E-mail: flavio.dalchiavon@cnp.ifmt.edu.br;

probability ($P < .05$). The hybrids AGUARÁ 04, SYN 3950HO, SYN 045, CF 101 and MG 305 stand out for grain and oil yields. AGUARÁ 04 and CF 101 presented the lowest plant height and the highest harvest rate. The hybrid SYN 3950HO has associated high grain and oil yields.

Keywords: Grain yield; Helianthus annuus L; oil yield; varietal competition.

1. INTRODUCTION

Sunflower (*Helianthus annuus* L.) achenes can be used to extracted high-quality oil for human consumption and high protein meal for the production of animal feed [1,2,3], besides the plant serve as ornamentation [4]. Sunflower oil may be rich in linoleic (traditional sunflower) or oleic (high oleic sunflower) fatty acids that favour the reduction of the LDL-cholesterol (bad cholesterol) fraction and, consequently, the risk of cardiovascular disease [5]. Also, oil with high oleic acid content is less susceptible to oxidative changes during refining, storage and frying.

In the 2015/2016 harvest, the sunflower was the third oilseed in achene production (40.57 million tons) and the fourth in oil production (15.5 million tons) in the world [6]. Despite the relevance, the Brazilian cultivation area is still reduced. In this harvest, it was 51.4 thousand ha, 80% of which was established in the savannah region, mainly in Campo Novo do Parecis (Mato Grosso) with an area of over 20 thousand ha [7].

In Brazilian savannah region, the sunflower is grown in the double summer cropping from February / March due to the occurrence of suitable rainfall (500 to 700 mm) and temperature (20 to 28°C) conditions for its cultivation [8]. Currently, soybean is usually sown in October and November and harvested between January and March and maize is the most sown crop in the second crop in a total area of approximately 10.0 million hectares [7]. Sunflower cultivation may be an option for crop rotation with maize.

To increase the area of sunflower cultivation in the country, it is relevant to have information on the agronomic behaviour of hybrids in the producing regions to increase the producer profitability and to facilitate the cultivation practices [9]. In Brazil, this evaluation is being carried out by the Sunflower Trial Network, coordinated by Embrapa, which counts with the participation of public and private companies [10,11,12,13]. In this network, hybrids are assessed for two years under different soil and climatic conditions. Thus, the evaluation of

cultivars (best hybrids evaluated in this network) under the same growing conditions can facilitate the choice of the best cultivar by the producer.

The objective of this work was to evaluate the agronomic performance of sunflower cultivars sown in the summer double cropping in Campo Novo do Parecis (Mato Grosso), the main sunflower producing region in the Brazilian savannah.

2. MATERIALS AND METHODS

2.1 Field Trials

The trials were carried out in Campo Novo do Parecis (Mato Grosso), in a system of summer double cropping in succession to soybean in 2014 and 2015. In the last five years, the area was cultivated in the soy/corn succession system. The geographical coordinates of the area are latitude 13°40'37" S, longitude 57°47'30" W and altitude of 564 m. The soil is classified as typical dystrophic Red Latosol (Typic Tropudox) [9].

The soil textural characterisation in the 0-0.20 m depth layer is 506, 134 and 360 g kg⁻¹ for clay, silt and sand, respectively. The initial characterization of fertility (0-0.20 m) showed the following values: pH (CaCl₂) = 5.7; OM = 26 g dm⁻³; P (resin) = 5.9 mg dm⁻³; K, Ca, Mg and H + Al = 1.5; 32; 11 and 40 mmol_c dm⁻³, respectively; with BS = 54.8%.

The local climate is the type AW according to Köppen, the dry season taking place from May to September and the rainy season from October to April [14], as indicated in the historical series of the years from 2003 to 2010 (Fig. 1). However, the precipitation height was 570 and 820 mm (Figs. 2A,B), values sufficient to satisfy the water demand of the crop, since this requires accumulated precipitation of 500 to 700 mm, regularly distributed throughout its cycle [8]. The mean values for maximum, average and minimum temperatures were 30.3; 23.2 and 18.9 °C (2014) and 30.6; 23.5 and 19.5 °C (2015), respectively.

Seeding was carried out on March 7, 2014 and March 3, 2015. In fertilisation, a total of 267 kg ha⁻¹ of NPK 10-30-20 + 45 kg ha⁻¹ of Potassium Chloride were applied - 26.7 kg ha⁻¹ of N, 80 kg ha⁻¹ of P₂O₅ and 80 kg ha⁻¹ of K₂O, according to the interpretation of soil analysis and crop demand [15]. The desiccation together with the application of 2 kg ha⁻¹ of boron (boric acid) was carried out on the same day of sowing. At 32 DAE, nitrogen fertilisation was performed at a dose of 50 kg ha⁻¹ of N (urea). Other cultural

practices were made to allow optimum plant development [15].

2.2 Agronomic Traits

The agronomic traits were evaluated in 10 plants randomly sampled in the useful area: achene yield (AY, kg ha⁻¹) obtained from all plants (R.9; physiological maturity) of the useful area and corrected for 11% moisture; oil content (OC %) predicted by spectroscopy [16]; oil





Fig. 1. Rainfall and thermal averages in the experimental area between 2003 and 2010 (Campo Novo do Parecis, Mato Grosso)






Fig. 2. Rainfall and thermal averages in the experimental area between March and June of 2014 (a) and of March and June of 2015 (b) (Campo Novo do Parecis, Mato Grosso)

yield (OY, kg ha^{-1}) calculated by the product between the oil content (%) and achene yield (kg ha^{-1}) / 100; plant height (PH, cm) measured from the soil base to the plant apex, in R5.5 (50% of the disk flowers have completed or are in anthesis); head diameter (HD; cm), measured in R9 (physiological maturity); harvest index (HI), determined by dividing the achene mass by the head mass and 1000-achene mass (1000AM) obtained by counting and weighing samples with a thousand achenes.

2.3 Experimental Design and Statistic

The experimental design was a randomised complete block with four replications. Each plot (replicate) consisted of four rows 6.5 m long and spaced from 0.45 m. The two external rows of each plot (borders) were discarded, as were 0.5 m from each end of the two central rows, leaving a useful area of 4.5 m^2 .

The treatments were the hybrids (commercial cultivars) AGUARÁ 04, AGUARÁ 06, CF 101, GNZ NEON, HELIO 251, MG 305, SYN 045 and SYN 3950HO. These hybrids are simple and traditional (their oils are rich in linoleic acids), except SYN 3950HO which is high oleic. The analysis of variance and Scott-Knott test ($P < .05$) were carried out by using SISVAR software program [17] and taking the average of two years (2014 and 2015). The homogeneity of residual variances obtained in the individual analysis was verified. In this test, variances were considered

as homogeneous when the ratio between the larger and the smaller residual mean square was smaller than 7.0 [18].

3. RESULTS AND DISCUSSION

3.1 Achene and Oil Yields

The producer must take into account the achene and oil yields in choosing the best hybrid sunflower. There were significant differences ($P = .01$) between hybrids grown in the Brazilian summer double cropping for oil content and oil yield, but not for achene yield (Table 1). The non-distinction between the hybrids for this trait may be due to them being commercial cultivars with wide adaptability to the edaphoclimatic conditions of Brazilian savannah [10,11,12,13].

The hybrids did not differ in the achene yield ($P > .05$), even when this difference was above 300 kg ha^{-1} (Table 2). The AGUARÁ 06 yield ($1368.3 \text{ kg ha}^{-1}$) was, for example, 22% higher than that of GNZ NEON ($1671.3 \text{ kg ha}^{-1}$). The highest achene yield was obtained in 2014 (mean of $2086.6 \text{ kg ha}^{-1}$ in 2014 and $1090.6 \text{ kg ha}^{-1}$ in 2015), a year with the lowest rainfall (Figs. 2A, B), reflecting in higher leaf health due to the lower incidence of *Alternaria stem helianthi* and *Sclerotinia sclerotiorum*. These are the main diseases of the culture in the Brazilian savannah region and that drastically reduces the photosynthetically active leaf area [15].

Table 1. Analysis of variance for agronomic traits of sunflower hybrids grown in the second summer crop (Campo Novo do Parecis, Mato Grosso, mean of data collected in 2014 and 2015)

Trait	F	C.V. (%) ^a	GM ^b
Plant height (cm)	48.1**	2.1	170.8
Head diameter (cm)	4.9**	4.2	15.4
Harvest index	8.9**	6.1	0.61
1000-achene mass (g)	16.4**	5.7	44.8
Achene yield (kg ha ⁻¹)	1.9	9.4	1588.6
Oil content (%)	57.2**	1.8	42.1
Oil yield (kg ha ⁻¹)	3.8**	9.7	671.5

** significant at $P < .01$ probability; ^a C.V. = coefficient of variation; ^b GM = general mean.

Table 2. Mean values for achene yield (AY), oil content (OC) and oil yield (OY) of sunflower hybrids grown in the second summer crop (Campo Novo do Parecis, Mato Grosso, mean of data collected in 2014 and 2015)

Hybrid	AY (kg ha ⁻¹)	OC (%)*	OY (kg ha ⁻¹)
AGUARÁ 04	1670.5	45.0 a	755.8 a
AGUARÁ 06	1368.3	40.9 c	562.0 b
CF 101	1559.0	44.2 a	694.5 a
GNZ NEON	1671.3	37.6 d	629.5 b
HELIO 251	1624.3	38.4 d	626.8 b
MG 305	1535.5	43.4 b	666.5 a
SYN 045	1652.5	42.6 b	708.3 a
SYN 3950HO	1627.3	44.5 a	728.5 a

Means followed by the same letter, in the column, do not significantly differ from each other by the Scott-Knott test at $P < .01$ probability

The hybrids AGUARÁ 04, SYN 3950HO and CF 101 showed the highest oil contents, with values between 44.2 and 45.0% (Table 2), while GNZ NEON (37.6%) and HELIO 251 (38.4%) showed the lowest contents. The mean oil content of the hybrids was 42.1%. The desirable oil content set by the industry is above 40%. Some industries provide subsidies to farmers who produce achenes with values higher than this index [10,11,16,12,13], since oil is the main commercial product of the sunflower crop. However, the hybrid with higher achene yield will not always result in higher oil yield. Therefore, the higher the subsidy, the greater the preference of producers for hybrids with higher oil contents and oil yields.

In this study, as there was no statistical difference between hybrids for achene yield, those with the highest oil yield were those with the highest oil content, such as AGUARÁ 04 (755.8 kg ha⁻¹), SYN 3950HO (728.5 kg ha⁻¹), SYN 045 (708.3 kg ha⁻¹), CF 101 (694.5 kg ha⁻¹) and MG 305 (666.5 kg ha⁻¹) (Table 2). Of which, the hybrid SYN 3950 is the only high oleic. [16] also reported hybrids grown in Brazilian savannah with high achene and oil yields associated with oil quality.

The mean of hybrid oil yields was 671.5 kg ha⁻¹ (Table 1), higher than the value highlighted by [19], who worked with two hybrids in Lavras (Minas Gerais) (savannah region) and obtained a mean of 565.4 kg ha⁻¹. On the other hand, Grunvald et al. [10] worked with several hybrids in the Sunflower Trial Network, between 2004 and 2007 in Brazilian savannah region, and obtained mean of oil productivity from 861.0 to 1022.0 kg ha⁻¹, with achene productivity ranging from 2021.0 to 2245.0 kg ha⁻¹ and oil contents ranging from 43.8 to 45.5%. These values show that other regions favourable to sunflower cultivation, in addition to Campo Novo do Parecis (Mato Grosso).

3.2 Other Agronomic Traits

Head size can be considered an indicator of the plant development, as well as the achene yield, even in situations of environmental stress [8]. Heldwein et al. [20] reported that both head size and 1000-achene mass have a direct implication on the achene yield. In the present study, the hybrids AGUARÁ 06 (16.6 cm), AGUARÁ 04 (16.0 cm), HELIO 251 (15.7 cm) and MG 305 (15.6 cm) showed the highest means for head size (Tables 1 and 2). The overall mean was

Table 3. Mean values for plant height (PH), head diameter (HD), harvest index (HI) and 1000-achene mass (1000AM) of sunflower hybrids grown in the second summer crop (Campo Novo do Parecis, Mato Grosso, mean of data collected in 2014 and 2015)

Hybrid	PH* (cm)	HD (cm)	HI	1000 GM (g)
AGUARÁ 04	166.8 b	16.0 a	0.69 a	40.7 c
AGUARÁ 06	179.7 a	16.6 a	0.59 b	38.3 c
CF 101	147.7 c	14.9 b	0.65 a	42.0 c
GNZ NEON	181.5 a	14.8 b	0.50 c	46.6 b
HELIO 251	162.1 b	15.7 a	0.60 b	41.1 c
MG 305	165.4 b	15.6 a	0.63 a	48.1 b
SYN 045	184.4 a	14.4 b	0.64 a	53.9 a
SYN 3950HO	178.7 a	15.2 b	0.61 b	47.4 b

*Means followed by the same letter, in the column, do not significantly differ from each other by the Scott-Knott test at $P < .01$ probability

15.4 cm. On the other hand, the hybrids SYN 045, MG 305 (48.1 cm), SYN 3950HO (47.4) and GNZ NEON (46.6) showed the highest 1000-achene mass (53.9 g) (Table 3). Thus, the hybrids that stood out for head size showed a lower 1000-achene mass, so that there was no significant difference ($P > .05$) between hybrids for achene yield (Table 1).

In addition to grain and oil yields, plant height (PH) was relevant for sunflower crop. The overall mean of the hybrids for plant height was 170.8 cm (Table 1). Those of smaller height were CF 101 (147.7 cm), HELIO 251 (162.1 cm), MG 305 (165.4 cm) and AGUARÁ 04 (166.8 cm) (Table 3). [21] reported an average of 170.0 cm for CF 101 in the North of Minas Gerais (savannah region), which shows an environmental effect for this characteristic. In addition to facilitating cultural practices, lower-sized reduces the lodge and stem breakage under unfavourable climatic conditions and minimises losses in mechanised harvesting [22,8].

The values of the hybrid harvest indexes were also shown in Table 3. The higher the value, the greater the commercial mass of the head, increasing the interest of the industry since this requires the achenes for the industrial processing [9]. Hybrids AGUARÁ 04, CF 101, SYN 045 and MG 305 showed the highest harvest indexes, ranging from 0.63 to 0.69, while GNZ NEON showed the lowest index (0.5), indicating that most of the energy translocated to heads of this hybrid was used to form the floral receptacle that has no commercial value, and not to form the achenes. This result corroborated by [9] which evaluated the agronomic performance of SYN 045 in the Campo Novo do Parecis (Mato Grosso) and reported a harvest index of 0.66.

4. CONCLUSION

For the summer second crop of Brazilian savannah region, the hybrids AGUARÁ 04 and CF 101 associate high achene and oil yields with lower-sized and good harvest index and the hybrid SYN 3950HO associates high achene and oil yields.

ACKNOWLEDGEMENTS

GPF (Phytochemical Research Group) of the IFMT - Campus Campo Novo do Parecis. FAPEMAT (Process N°. 460160/2014 and Process N°. 157050/2014). CNPq (Process 402022 / 2014-9 and Process 313036/2014-4) for fellowships.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Balbinot Jr AA, Backes RL, Souza AM de. Performance of sunflower cultivars in three planting dates in north plateau of Santa Catarina state. *Sci Agrar*. 2009;10(2):127-133. Portuguese.
- Brightenti AM. Control of volunteer soybean plants in sunflower crop. *Pesqui Agropecu Trop*. 2015;45(3):274-281. Portuguese.
- Azerbaijan MMLC, Farias ME de, CF Pine de, Teixeira SB, Ávila GE, Deuner C, Borella J, Deuner S. Antioxidant Defense in Sunflower against Drought. *Journal of Experimental Agriculture International*. 2018;25(1):1-9.

4. Souza, RN de, Paz VP da S, Gonçalves KS, Soares TM, Azevedo Neto AD, Gheyi HR. Treated domestic effluents: An option for cultivation of ornamental sunflower in a hydroponic system. *Journal of Experimental Agriculture International*. 2018;24(6):1-11.
5. Miller JF, Zimmerman DC, Vick BA. Genetic control of high oleic acid content in sunflower oil. *Crop Sci*. 1987;27(5):923-926.
DOI:<http://dx.doi.org/10.2135/cropsci1987.0011183X002700050019x>
6. United States - Department of agriculture - Foreign agricultural service. *Oilseeds: world markets and trade*. Washington. 2017;36.
7. Conab - National Supply Company - Monitoring of harvest: grains. Joint monthly, March 2017. Brasília, National Supply Company. 2017;172. Accessed 25 April 2017.
Available:<http://www.conab.gov.br>
8. Castro C de, Leite RMVB de C. Main aspects of sunflower production in Brazil. *OCL* 2018;25(1):D104.
DOI:<https://doi.org/10.1051/ocli/2017056>
9. Dalchiavon FC, Montanari R, Andreotti M, Dallacort R, Souza MFP. Relationship between sunflower productivity and soil's chemical properties by geo-statistical techniques. *Afr J Agric Res*. 2015;10(35):3525-3532.
DOI:<http://dx.doi.org/10.5897/AJAR2014.9472>
10. Grunvald AK, Carvalho CGP de, Oliveira ACB de, Andrade CA de B. Adaptability and stability of sunflower genotypes in Central Brazil. *Pesqui. Agropecu. Bras*. 2008;43(11):1483-1493.
DOI:<http://dx.doi.org/10.1590/S0100204X2008001100006>. Portuguese.
11. Porto WS, Carvalho CGP de, Pinto RJB, Oliveira MF de, Oliveira ACB de. Evaluation of sunflower cultivar for central Brazil. *Sci Agric*. 2008;65(2):139-144.
DOI:<http://dx.doi.org/10.1590/S010390162008000200005>.
12. Carvalho CGP de, Ozawa EKM, Amabile RF, Godinho V de PC, Gonçalves SL, Ribeiro JL, Seifert A. L. Adaptability and stability of sunflower genotypes imidazolinone resistant in second crop cultivation. *Brazilian Journal of Agricultural Sciences*. 2015;10(1):1-7.
DOI:<http://dx.doi.org/10.5039/agraria.v10i1a3804>. Portuguese.
13. Dalchiavon FC, Carvalho CGP de, Amabile RF, Godinho VPC, Ramos NP, Anselmo JL. Agronomic traits and their correlations in sunflower hybrids adapted to second crop. *Pesqui. Agropecu. Bras*. 2016;51(11):1806-1812.
DOI:<http://dx.doi.org/10.1590/S0100-204X2016001100002>. Portuguese.
14. Dallacort R, Martins JA, Inoue MH, Freitas PSL, Coletti AJ. Rain distribution in Tangará da Serra, mid-northern Mato Grosso State, Brazil. *Acta Scientiarum Agronomy*. 2011;33(2):193-200.
Portuguese.
15. Oliveira ACB de, Rosa APSA da. Practical guide to sunflower cultivation. Brasília: Embrapa. 2013;53.
16. Grunvald AK, Carvalho CPG de, Leite RS, Mandarino JMG, Andrade CA de B, Scapim CA. Predicting the oil contents in sunflower genotype seeds using near-infrared reflectance (NIR) spectroscopy. *Acta Scientiarum Agronom*. 2014a;36(2):233-237.
DOI:<http://dx.doi.org/10.4025/actasciagron.v36i2.17677>
17. Ferreira DF. Sisvar: A computer statistical analysis system. *Science and Agrotechnology*. 2011;35(6):1039-1042.
DOI:<http://dx.doi.org/10.1590/S1413-70542011000600001>
18. Pimentel Gomes F. Course of experimental statistics. 13 ed. São Paulo: USP-Esalaq. 1985;468.
19. Silva MDL, Faria MA de, Morais AR de, Andrade GP, Lima EM de C. Growth and productivity of sunflower cultivated in late summer and with different water depths. *Brazilian Journal of Agricultural and Environmental Engineering*. 2007;11(5):482-488.
DOI:<http://dx.doi.org/10.1590/S1415-43662007000500006>. Portuguese.
20. Heldwein AB, Loose LH, Lucas DDP, Hinnah FD, Bortoluzzi MP, Maldaner IC. Yield and growth characteristics of sunflower sown from August to February in Santa Maria, RS. *Brazilian Journal of Agricultural and Environmental Engineering*. 2014;18(9):908-913.
DOI:<http://dx.doi.org/10.1590/1807-1929/agriambi.v18n09p908-913>. Portuguese.

21. Nobre DAC, Resende JCF, Junior DSSB, Costa CA. Agronomic performance of sunflower genotypes in the north of Minas Gerais, Brazil. Magazine Agro@ambiente On-line. 2012;6(2):140-147.
DOI:<http://dx.doi.org/10.18227/1982-8470ragro.v6i2.781>. Portuguese.
22. Pivetta LG, Guimarães VF, Fioreze SL, Pivetta LA, Castoldi G. Evaluation of sunflower hybrids and the relationship between productive and qualitative parameters. Cienc Agron. 2012;43(3):561-568. Portuguese.

© 2018 Dalchiavon et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

*The peer review history for this paper can be accessed here:
<http://www.sciencedomain.org/review-history/27228>*