



# Draft Genome Sequence of Vancomycin-Resistant *Enterococcus faecium* UEL170 (Sequence Type 412), Isolated from a Patient with Urinary Tract Infection in a Tertiary Hospital in Southern Brazil

Eliandro Reis Tavares,<sup>a</sup> Lucas Fernando da Silva,<sup>b</sup> Alexandre Tadachi Morey,<sup>c</sup> Admilton Gonçalves de Oliveira,<sup>a,b</sup> Sergio Paulo Dejato da Rocha,<sup>a,b</sup> Renan Augusto Ribeiro,<sup>d</sup>  Mariangela Hungria,<sup>b,d</sup> Isabella Ramos Trevizani Thihara,<sup>a</sup> Marcia Regina Eches Perugini,<sup>e</sup> Lucy Megumi Yamauchi,<sup>a,b</sup> Sueli Fumie Yamada-Ogatta<sup>a,b</sup>

<sup>a</sup>Department of Microbiology, Universidade Estadual de Londrina (UEL), Londrina, Paraná, Brazil

<sup>b</sup>Postgraduate Program of Microbiology, Department of Microbiology, UEL, Londrina, Paraná, Brazil

<sup>c</sup>Federal Institute of Rio Grande do Sul, Campus Canoas, Canoas, Rio Grande do Sul, Brazil

<sup>d</sup>Soil Biotechnology Laboratory, EMBRAPA Soja, Londrina, Paraná, Brazil

<sup>e</sup>Department of Pathology and Clinical and Toxicological Analysis, UEL, Londrina, Paraná, Brazil

**ABSTRACT** *Enterococcus faecium* is a leading cause of health care-associated infections, with specific lineages circulating in hospital settings worldwide. Here, we report the draft genome sequence of the multidrug-resistant and biofilm-producing *E. faecium* UEL170, sequence type 412 (ST412), isolated from an inpatient with a urinary tract infection. This strain is a member of clonal complex 17 (CC17), a globally hospital-associated clone.

*Enterococcus faecium* can be found as a harmless colonizer of the gastrointestinal tract of humans and animals. However, as an opportunistic pathogen, this bacterium is an important causative agent of health care-associated infections (HAI) worldwide (1–3). This scenario may be associated with *E. faecium*'s remarkable capacity to acquire resistance to many antimicrobials and the extraordinary plasticity of its genome, which contributes to bacterial adaptation in different environments (4, 5). Molecular epidemiological studies revealed the existence of specific genogroups, including a hospital-associated *E. faecium* subpopulation, collectively known as clonal complex 17 (CC17) (6, 7). This clonal complex has been widely detected in hospital settings around the world (7, 8), including in Brazil (2, 9–11). In particular, the multidrug-resistant *E. faecium* sequence type 412 (ST412) that belongs to CC17 has been disseminated in various Brazilian hospitals (9–11).

University Hospital of Londrina (UHL) is a teaching hospital and a major referral center in the north of Paraná, Brazil, for the Sistema Único de Saúde, a government public health system. In this institution, vancomycin-resistant enterococci (VRE) that were detected in surveillance rectal swab cultures from patients housed in intensive care units were first reported in 2002; the first outbreak of VRE (predominantly *E. faecium* [VRE<sub>fm</sub>]) occurred in 2005 (12), and it remains among the leading Gram-positive bacteria isolated from HAI (13). Here, we describe the draft genome of the VRE<sub>fm</sub> UEL170 strain, which was isolated from an inpatient with a urinary tract infection in 2007. This strain is phenotypically resistant to ampicillin, ciprofloxacin, erythromycin, streptomycin, tetracycline, teicoplanin, and vancomycin (1); it is a strong biofilm-producing bacterium on abiotic surfaces (1), and it belongs to ST412 (14).

For DNA isolation, 5 CFU was inoculated into brain heart infusion (BHI) broth (HiMedia, Brazil), and the culture was incubated for 24 h at 37°C. The cells were

**Citation** Tavares ER, da Silva LF, Morey AT, de Oliveira AG, da Rocha SPD, Ribeiro RA, Hungria M, Thihara IRT, Perugini MRE, Yamauchi LM, Yamada-Ogatta SF. 2019. Draft genome sequence of vancomycin-resistant *Enterococcus faecium* UEL170 (sequence type 412), isolated from a patient with urinary tract infection in a tertiary hospital in southern Brazil. *Microbiol Resour Annot* 8:e01365-18. <https://doi.org/10.1128/MRA.01365-18>.

**Editor** David Rasko, University of Maryland School of Medicine

**Copyright** © 2019 Tavares et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Sueli Fumie Yamada-Ogatta, [ogatta@uel.br](mailto:ogatta@uel.br).

E.R.T. and L.F.D.S. contributed equally to this article.

**Received** 30 October 2018

**Accepted** 16 January 2019

**Published** 14 February 2019

harvested by centrifugation ( $2,600 \times g$  for 10 min at  $10^{\circ}\text{C}$ ) and washed twice with 0.15 phosphate-saline buffer at pH 7.2. Genomic DNA was extracted using the DNeasy blood and tissue kit (Qiagen, Germany) according to the manufacturer's standard protocols and processed on a MiSeq platform using a MiSeq reagent v3 kit (600 cycles; Illumina, Brazil) at Embrapa Soja in Londrina, Brazil. Shotgun sequencing with the paired-end sequence strategy allowed a genome coverage of approximately 550-fold, which resulted in a total of 6,422,034 reads (300 bp each). The genome was assembled by the *de novo* strategy using the A5-miseq assembly pipeline. The data were filtered and trimmed to a Phred score of  $>20$  using the quality assessment tool for genome assemblies (QUAST [<http://quast.sf.net>]) according to the default software recommendations (15), generating 227 contigs. The genome was annotated with Rapid Annotation using System Technology (RAST) v2.0 (<http://rast.nmpdr.org>) using FIGfams technology release 70 according to the default software recommendations (16). Assembly of the genome generated a total size of 3,040,378 bp and a mean GC content of 38% with  $N_{50}$  and  $L_{50}$  values of 37,880 bp and 24 bp, respectively. The following acquired antimicrobial resistance and virulence genes were identified by ResFinder v3.0 (17) and VirulenceFinder (18), respectively (<http://cge.cbs.dtu.dk/services>): *aph(3')-III* and *ant(6)-Ia* (aminoglycoside resistance), *erm(B)* (macrolide resistance), *msr(C)* (macrolide, lincosamide, and streptogramin B resistance), *vanA* (glycopeptide resistance), *acm* (collagen-binding adhesin of microbial surface components recognizing adhesive matrix molecule [MSCRAMM] family), *efaAfm* (endocarditis-specific antigen A adhesin), and *espfm* (enterococcal surface protein). The availability of this genome sequence may help in exploring new targets for antimicrobial agents, including those exhibiting antibiofilm activity against VREfm.

**Data availability.** A data file containing the raw reads was uploaded to the NCBI Sequence Read Archive (accession number [SRR8362446](https://www.ncbi.nlm.nih.gov/sra/SRR8362446)). The whole-genome shotgun sequence of VREfm UEL170 was deposited at DDBJ/ENA/GenBank under the accession number [RBBX00000000](https://www.ncbi.nlm.nih.gov/nuclseq/RBBX00000000) (BioProject number [PRJNA493784](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA493784), BioSample number [SAMN10142019](https://www.ncbi.nlm.nih.gov/biosample/SAMN10142019)). The version described here is the first version.

## ACKNOWLEDGMENTS

E.R.T. received a research fellowship from Programa Nacional de Pós-Doutorado de Coordenação de Aperfeiçoamento de Pessoal de Nível Superior, Brazil (PNPD-CAPES). I.R.T.T. received a student scholarship from Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq). M.H. and S.F.Y.-O. received research fellowships from CNPq. This study was financed in part by the CAPES-Finance Code 001.

## REFERENCES

- Ruzon FI, de Paula SB, Kanoshiki RL, Pereira-Santos J, Kerbauy G, Kobayashi RK, Yamauchi LM, Perugini MR, Yamada-Ogatta SF. 2010. Virulence determinants in vancomycin-resistant *Enterococcus faecium* vanA isolated from different sources at University Hospital of Londrina, Paraná, Brazil. *J Microbiol* 48:814–821. <https://doi.org/10.1007/s12275-010-0099-5>.
- Sacramento AG, Zanella RC, Esposito F, Costa EAS, de Almeida LM, Pires C, de Brito AC, Mamizuka EM, Cerdeira LT, Lincopan N. 2017. Changed epidemiology during intra and interhospital spread of high-risk clones of *vanA*-containing *Enterococcus* in Brazilian hospitals. *Diagn Microbiol Infect Dis* 88:348–351. <https://doi.org/10.1016/j.diagmicrobio.2017.05.008>.
- Ulrich N, Vonberg RP, Gastmeier P. 2017. Outbreaks caused by vancomycin-resistant *Enterococcus faecium* in hematology and oncology departments: a systematic review. *Heliyon* 3:e00473. <https://doi.org/10.1016/j.heliyon.2017.e00473>.
- de Mello SS, Van Tyne D, Dabul AN, Gilmore MS, Camargo IL. 2016. High-quality draft genome sequence of the multidrug-resistant clinical isolate *Enterococcus faecium* VRE16. *Genome Announc* 4:e00992-16. <https://doi.org/10.1128/genomeA.00992-16>.
- Leong KWC, Cooley LA, Anderson TL, Gautam SS, McEwan B, Wells A, Wilson F, Hughson L, O'Toole RF. 2018. Emergence of vancomycin-resistant *Enterococcus faecium* at an Australian hospital: a whole genome sequencing analysis. *Sci Rep* 8:6274. <https://doi.org/10.1038/s41598-018-24614-6>.
- Willems RJ, Top J, van Santen M, Robinson DA, Coque TM, Baquero F, Grundmann H, Bonten MJ. 2005. Global spread of vancomycin-resistant *Enterococcus faecium* from distinct nosocomial genetic complex. *Emerg Infect Dis* 11:821–828. <https://doi.org/10.3201/1106.041204>.
- Guzman Prieto AM, van Schaik W, Rogers MR, Coque TM, Baquero F, Corander J, Willems RJ. 2016. Global emergence and dissemination of enterococci as nosocomial pathogens: attack of the clones? *Front Microbiol* 7:788. <https://doi.org/10.3389/fmicb.2016.00788>.
- Lee T, Pang S, Abraham S, Coombs GW. 2018. Antimicrobial-resistant CC17 *Enterococcus faecium*: the past, the present and the future. *J Glob Antimicrob Resist*. <https://doi.org/10.1016/j.jgar.2018.08.016>.
- Palazzo IC, Pitondo-Silva A, Levy CE, da Costa Darini AL. 2011. Changes in vancomycin-resistant *Enterococcus faecium* causing outbreaks in Brazil. *J Hosp Infect* 79:70–74. <https://doi.org/10.1016/j.jhin.2011.04.016>.
- Alves GDS, Pereira MF, Bride LL, Nunes APF, Schuenck RP. 2017. Clonal dissemination of vancomycin-resistant *Enterococcus faecium* ST412 in a Brazilian region. *Braz J Infect Dis* 21:656–659. <https://doi.org/10.1016/j.bjid.2017.07.001>.
- Marchi AP, Perdigão Neto LV, Martins RCR, Rizek CF, Camargo CH, Moreno LZ, Moreno AM, Batista MV, Basqueira MS, Rossi F, Amigo U,

- Guimaraes T, Levin AS, Costa SF. 2018. Vancomycin-resistant enterococci isolates colonizing and infecting haematology patients: clonality, and virulence and resistance profile. *J Hosp Infect* 99:346–355. <https://doi.org/10.1016/j.jhin.2017.10.010>.
12. Perugini MR, Nomi SM, Lopes GK, Belei RA, van der Heijden IM, Mostachio AK, Grion C, Couto EB, Jr, Costa SF. 2011. Impact of the reduction of environmental and equipment contamination on vancomycin-resistant *Enterococcus* rates. *Infection* 39:587–593. <https://doi.org/10.1007/s15010-011-0140-6>.
  13. Souza ES, Belei RA, Carrilho CMDM, Matsuo T, Yamada-Ogatta SF, Andrade G, Perugini MRE, Pieri FM, Dessunti EM, Kerbauy G. 2015. Mortality and risks related to healthcare associated infection. *Texto Contexto Enferm* 24:220–228. <https://doi.org/10.1590/0104-07072015002940013>.
  14. Larsen MV, Cosentino S, Rasmussen S, Friis C, Hasman H, Marvig RL, Jelsbak L, Sicheritz-Ponten T, Ussery DW, Aarestrup FM, Lund O. 2012. Multilocus sequence typing of total-genome-sequenced bacteria. *J Clin Microbiol* 50:1355–1361. <https://doi.org/10.1128/JCM.06094-11>.
  15. Gurevich A, Saveliev V, Vyahhi N, Tesler G. 2013. QUAST: quality assessment tool for genome assemblies. *Bioinformatics* 29:1072–1075. <https://doi.org/10.1093/bioinformatics/btt086>.
  16. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
  17. Zankari E, Hasman H, Cosentino S, Vestergaard M, Rasmussen S, Lund O, Aarestrup FM, Larsen MV. 2012. Identification of acquired antimicrobial resistance genes. *J Antimicrob Chemother* 67:2640–2644. <https://doi.org/10.1093/jac/dks261>.
  18. Joensen KG, Scheutz F, Lund O, Hasman H, Kaas RS, Nielsen EM, Aarestrup FM. 2014. Real-time whole-genome sequencing for routine typing, surveillance, and outbreak detection of verotoxigenic *Escherichia coli*. *J Clin Microbiol* 52:1501–1510. <https://doi.org/10.1128/JCM.03617-13>.