

Genetic variability revealed by microsatellite markers in a germplasm collection of *Jatropha curcas* L. in Brazil: an important plant for biofuels

Crop Breeding and Applied Biotechnology
19: 337-346, 2019
Brazilian Society of Plant Breeding.
Printed in Brazil
<http://dx.doi.org/10.1590/1984-70332019v19n3a46>

Romero de Lima Souza¹, Luiz Antônio dos Santos Dias^{1*}, Thais Roseli Corrêa², Eveline Teixeira Caixeta^{1,3}, Erika da Costa Fernandes¹, Dandara Rêgo Muniz¹, Lucas Barbosa de Castro Rosmaninho¹ and Poliane Marcele Ribeiro Cardoso¹

Abstract: We evaluated the genetic variability of a collection of *Jatropha curcas* germplasm, represented by 93 accessions, using microsatellite markers. Among the 60 markers tested, five of them detected polymorphisms, with a total of 11 alleles and mean of 2.2 alleles per loci. These five markers enabled the quantification of genetic variability through estimates of expected ($H_e=0.42$) and observed ($H_o=0.64$) heterozygosity, Shannon-Weaver index ($H'=0.62$), coefficient of inbreeding ($f=-0.44$) and the formation of 11 clusters. Simultaneously, 14 accessions randomly sampled among the 93 and represented by seven plants each, were analyzed with these same five markers to quantify the within and between variability. Most of the genetic variation (92.58%) was contained within the accessions. These analyses revealed, for the first time, expressive genetic variability to be explored in this collection. The accessions UFVJC 05, 07, 12, 18, and 53 presented expressive variability among them with potential for the constitution of a base population for the breeding program.

Keywords: Variability, molecular markers, germplasm bank, breeding.

INTRODUCTION


Oil price volatility, combined with the need to reduce greenhouse gas emissions, has boosted global demand for biofuels. Among the potential species for the production of biofuels, *Jatropha curcas* L. deserves to be highlighted, as its seeds contain high oil content (36.2%) with the best quality (Freitas et al. 2016). These aspects have led to rapid expansion of cultivated areas and demand for improved cultivars (Sorrel et al. 2010).

Despite its great potential, *J. curcas* is still a species undergo domestication, but with considerable variability to be explored. However, information regarding genetic variability and population structure is still limited (Bressan et al. 2012) and breeding programs are rare compared to other oilseed species (Dias et al. 2012, Pecina-Quintero et al. 2014).

The initial phase of a breeding program involving any kind of plant species requires a germplasm collection, functioning as a repository of genes for the future development of varieties. Therefore, the success of an improvement

*Corresponding author:

E-mail: lasdias@ufv.br

 ORCID: 0000-0002-1828-6938

Received: 24 April 2019

Accepted: 07 August 2019

¹ Universidade Federal de Viçosa (UFV),
Departamento de Fitotecnia, 36.571-900,
Viçosa, MG, Brazil

² Universidade Estadual do Maranhão
(UEMA), Departamento de Fitotecnia e Fitos-
sanidade, 65.055-970, São Luís, MA, Brazil

³ Embrapa Café. Current address: UFV, Bio-
agro, 36.571-900, Viçosa, MG, Brazil

program depends on knowledge of the available genetic variability, which will allow the efficient selection of different genotypes to produce hybrids and similar genotypes to produce lines (Pecina-Quintero et al. 2014).

Molecular markers are suitable tools for the characterization of genetic variability in germplasm collections. Microsatellites, also known as Simple Repeated Sequence (SSR) markers, have high reproducibility, codominant inheritance and high polymorphism. Studies involving these markers have been successfully used to characterize the genetic variability of *J. curcas* (Pamidimarri et al. 2008, Sun et al. 2008, Sudheer et al. 2009, Rosado et al. 2010, Wen et al. 2010, Bressan et al. 2012, Na-ek et al. 2011, Pecina-Quintero et al. 2014, Sinha et al. 2015, Santos et al. 2016, Vásquez-Mayorga et al. 2017, Gangapur et al. 2018). In all of these studies, the genetic variability revealed for this species has been considered low.

The present study aimed to quantify the genetic variability present in a germplasm collection of *J. curcas*, composed of 93 accessions, using microsatellite markers. This study is part of the strategy to identify superior genotypes for cultivar development.

MATERIAL AND METHODS

Plant material

The germplasm collection of *J. curcas* of the Federal University of Viçosa (UFV), located in the Araponga Experimental Farm (lat 20° 39' S, long 42° 32' W and alt 823 m asl), in the municipality of Araponga, MG, Brazil is composed of accessions originating from different Brazilian geographic regions and from abroad (Table 1), all propagated by seeds. Currently, this collection is composed of 93 accessions (1504 plants) and is installed in modules of five trials in randomized block design, with four replications and 4-plant plots in 2 x 2 m spacing, with two common controls (Freitas et al. 2011, Freitas et al. 2016).

Samples of young and fully developed leaves from the 93 accessions were collected, each with seven sub-samples (plants), totaling 658 plants sampled. The leaves were wrapped in identified aluminum foil and placed in styrofoam boxes with ice for transportation to the Federal University of Viçosa, where they were stored at -80 °C.

DNA extraction

The DNA extraction process was conducted at the Laboratory of Forest Pathology of the UFV, based on the protocol for eucalyptus, modified from Doyle and Doyle (1990). This modification was due to the maceration of the samples in the following the steps: after removing of the central and secondary veins, the leaves were placed in 2 mL microcentrifuge tubes with metal beads and 700 µL extraction buffer (100 mM Tris-HCl pH 8.0, 20 mM EDTA, 1.4 M NaCl, 2% (w/v) CTAB, 2% (w/v) PVP and 0.4% (v/v) β-mercaptoetanol, the latter separated from the other components). The samples were macerated with TissueLyser II (Qiagen) and incubated in a water bath at 65 °C for 30 minutes. After incubation, 500 µL of chloroform-isoamyl alcohol (24:1) was added to the tubes, which were manually inverted several times. The tubes were then centrifuged at 12,000 rpm for 5 min. The supernatants were transferred to new tubes and the extraction with chloroform was repeated. The supernatants (~ 500 µL) were transferred to fresh tubes and 0.9 volumes (450 µL) of cold isopropanol were added. The precipitated DNA was washed twice with 500 µL of cold 70% and 95% ethanol. The DNA was dried at room temperature for 1 hour and dissolved in 50 µL of TE (10 mM Tris and 1 mM EDTA, pH 8.0) plus RNase (10 µg mL⁻¹) for 2 hours at 37 °C and then stored at -20 °C. DNA quantification was performed using a Thermo Scientific's NanoDrop spectrophotometer and standardized with final concentrations adjusted to 10 ng mL⁻¹.

Microsatellite molecular markers

Sixty pairs of microsatellite primers were tested (see Table 2). All primers used were previously reported in the literature surrounding *J. curcas*, some of them drawn based on microsatellite loci derived from genomic sequences and specific ESTs of the species by Bressan et al. (2012), partial genomic sequences developed initially for cassava by Wen et al. (2010), and a number developed by Sudheer Pamidimarri et al. (2009) for differentiation of *J. curcas* toxic and non-toxic genotypes. PCRs (Polymerase Chain Reaction) were performed in a volume of 20 µL containing 50 ng DNA sample, 1x Taq DNA polymerase buffer, 100 µM of each dNTP, 1.5 mM MgCl₂, 0.2 µM of each primer and 1.0 U Taq DNA polymerase (Life Science). Amplifications were performed in an MJ Research PTC 100 thermocycler with denaturation at 94 °C for 3 minutes, followed by 40 cycles of denaturation at 94 °C for 1 min, annealing for 1 min at the specific

temperature of each primer and extension at 72 °C for 1 min. Final extension was performed at 72 °C for 8 min. PCR products were separated on a 6% denaturing polyacrylamide gel and visualized by silver staining solution, according to Creste et al. (2001).

Table 1. Provenance of 93 accessions of *Jatropha curcas* L. evaluated

Accessions ¹	Location of collection	Accessions	Location of collection
UFVJC 1	Santa Vitória, MG	UFVJC 52	Barbacena, MG
UFVJC 3	Santa Vitória, MG	UFVJC 53	Barbacena, MG
UFVJC 4	Santa Vitória, MG	UFVJC 54	Barbacena, MG
UFVJC 5	João Pinheiro, MG	UFVJC 55	Janaúba, MG
UFVJC 6	João Pinheiro, MG	UFVJC 56	Janaúba, MG
UFVJC 7	João Pinheiro, MG	UFVJC 57	Janaúba, MG
UFVJC 8	João Pinheiro, MG	UFVJC 58	Janaúba, MG
UFVJC 9	João Pinheiro, MG	UFVJC 59	Janaúba, MG
UFVJC 10	João Pinheiro, MG	UFVJC 60	Pompéu, MG
UFVJC 11	João Pinheiro, MG	UFVJC 61	Rio Grande do Sul
UFVJC 12	João Pinheiro, MG	UFVJC 62	Juiz de Fora, MG
UFVJC 13	Tauá, MG	UFVJC 63	Novo Repartimento, PA
UFVJC 14	Ólhos D'água, MG	UFVJC 65	Desconhecido
UFVJC 15	Veredas, MG	UFVJC 66	Desconhecido
UFVJC 16	Jaíba, MG	UFVJC 67	Desconhecido
UFVJC 17	Montalvânia, MG	UFVJC 68	São Luiz, MA
UFVJC 18	Cana Brava II-Montalvânia, MG	UFVJC 70	Ariquemes, RO
UFVJC 19	Cana Brava II-Montalvânia, MG	UFVJC 71	João Pinheiro, MG
UFVJC 20	Poçoões, MG	UFVJC 72	Camboja
UFVJC 22	Ipatinga, MG	UFVJC 73	Camboja
UFVJC 23	Don Lara-Caratinga, MG	UFVJC 74	Camboja
UFVJC 24	Santa Luzia-Caratinga, MG	UFVJC 75	Bomfim, MG
UFVJC 25	Imbé-Caratinga, MG	UFVJC 79	Jordânia, MG
UFVJC 28	Poté, MG	UFVJC 80	Desconhecido
UFVJC 29	Poté, MG	UFVJC 81	Jordânia, MG
UFVJC 30	Poté, MG	UFVJC 82	Jordânia, MG
UFVJC 31	Poté, MG	UFVJC 83	Araras, SP
UFVJC 32	Poté, MG	UFVJC 84	Petrolina, PE
UFVJC 33	Itaipé, MG	UFVJC 85	Jataí, GO
UFVJC 34	Itaipé, MG	UFVJC 86	Jequié, BA
UFVJC 35	Ervália, MG	UFVJC 87	Jequié, BA
UFVJC 36	Serra da Ipiapaba, MG	UFVJC 88	Jequié, BA
UFVJC 37	Janaúba, MG	UFVJC 89	Jitaúna, BA
UFVJC 38	Petrolina, PE	UFVJC 90	Ipiaú, BA
UFVJC 39	Natal, RN	UFVJC 91	Apuarema, BA
UFVJC 40	Formoso, TO	UFVJC 92	Itaitê, BA
UFVJC 41	Jales, SP	UFVJC 93	Itaitê, BA
UFVJC 42	Dourados, MS	UFVJC 94	Andaraí, BA
UFVJC 43	Matozinhos, MG	UFVJC 95	Andaraí, BA
UFVJC 44	São Carlos, SP	UFVJC 96	Mucugê, BA
UFVJC 45	Barra do Bugre, MT	UFVJC 97	Mucugê, BA
UFVJC 46	Barra do Bugre, MT	UFVJC 98	Iraquara, BA
UFVJC 47	Pirajá, SP	UFVJC 99	Iraquara, BA
UFVJC 48	Getulina, SP	UFVJC 100	Souto Soares, BA
UFVJC 49	Bocaiúva, MG	UFVJC 101	Wagner, BA
UFVJC 50	Bocaiúva, MG	UFVJC 102	Desconhecido
UFVJC 51	Rio Pompa, MG		

¹ Accessions in bold were utilized to quantify the within and between variability.

Table 2. SSR primers tested in the evaluation of the genetic variability of *Jatropha curcas* L. accessions

Name	Forward primer	Reverse primer
¹ JCENA 27	CATTTTTTCATCAAGGCCTAC	GTATTTCTCCACACGCAACT
¹ JCENA 41	CTTTCTTACCCCTCATCCTT	AAAGCCAGGCATACCTTGAA
¹ JCENA 47	GCCCGAGTTCTCTATAAGGT	CCAAGAGAAATTAGGAATGC
¹ JCENA 87	<u>ATCTGGAGTGAACCAAAAGA</u>	<u>CACATGGTAAGCATTACAAGC</u>
² JCDS 10	<u>CATCAAATGCTAATGAAAGTACA</u>	<u>CACACCTAGCAAACACTACTTGCA</u>
² JCDS 24	GGATATGAAGTTTCATGGGACAAG	TTCATTGAATGGATGGTTGTAAGG
² JCDS 41	AACACACCATGGGCCACAGGT	TGCATGTGTGCGGGTTTGATTAC
² JCDS 58	TCCATGAAGTTTGCTGGCAAT	AGGTCATCTGGTAAAGCCATACC
² JCDS 66	CCTACGAGTGATTGGATAGTTTCTCA	TCTTCCATCAAGAGTCGTTGGGCA
² JCPS 6	CCAGAAGTAGAATTATAAATAAA	AGCGGCTCTGACATTATGTAC
² JCPS 9	GTACTTAGATCTCTTGTAACTAACAG	TATCTCTTGTTCAGAAATGGAT
² JCPS 20	ACAGCAAGTGCAACAATCTCA	TACTGCAGATGGATGGCATGA
² JCMS 30	GGGAAAGAGGCTCTTTGC	ATGAGTTCACATAAAATCATGCA
³ CESR 0163	AACCACAGGAGTTGGTAATG	GAAAGAAGCAACAGAAATGG
³ CESR 0231	GCTTTAGCAAACCAAGATTC	AATCATCGTTATCGTTGGAC
³ CESR 0235	TGTCACAGTACATGCCACTT	ACCTTCGATTCATTGTGTGC
³ CESR 0290	TCAATCAATCCAACAACAA	CTAAATTCTCAACCAGGTGC
³ CESR 0293	CAAAGAAGCCATTTCTGTTC	TATCATCACAAGGTGCAAA
³ CESR 0303	AAAGCCTAGTGCATTTGAAG	TCATGCTCTTTCTCCACTT
³ CESR 0312	AAGAAACAACAATTGTGCC	TAGAAATCCTGTGGGTTA
³ CESR 0333	ACATCTACAATGGCGATTC	TAATGAATCTGTAGGCCCG
³ CESR 0361	AGGCAGATATGCTGATTAAGTT	AGGCAGATATGCTGATTAAGTT
³ CESR 0370	ATCAATGAACCTGTTGAAGG	TCCAGACTCCAACTTAATCA
³ CESR 0382	ATGACCTACAAGCAAGCACT	CCAATGACACCTCAGAATTT
³ CESR 0386	ATTACTGTGAATGAAGCCCA	ACAGAGCACCCCTAACGATAA
³ CESR 0398	CAATCTTATGCAAGTCTGA	CAAACAGCTAGAATCCCTTG
³ CESR 0399	CAATGCATGGATCATAAGTG	CTCAAGTCAAATCTGGGAAC
³ CESR 0494	GCCAAAGCTTCTCTTAACAA	TAATCACCATACCCACCTTC
³ CESR 0498	GCTCGTTTCATTCATCTCACT	TTAACTGTGAATGCCCTTCT
³ CESR 0718	ATGATGTTAAGGACGAGGTG	ATCGGCTTCTTCTTCTTCT
³ CESR 0719	ATGATGTTAAGGACGAGGTG	GCAAGTCTGGATTGTTCTC
³ CESR 0733	CAAACAGATCAAGTACCCAGA	AAATTGATGGTGGTGATGAT
³ CESR0756	<u>CAGGTTCTGTTCTTCAACT</u>	<u>ATATATGATCCCGACAACAA</u>
³ CESR 0804	CGTATTACCACCAACATCCT	CCATCAATCCACTTATCGTT
³ CESR 0820	CTCGAGCACATGTTTAATGA	TGATGATCAATCAACCACAA
³ CESR 0844	GAAATAGAGACAGAAGGCCGA	AAATGAACAGGGAATTGTTG
³ CESR 0889	GCTCTTGATTGATCCATTT	TGATGATGATGATGATGAGG
³ CESR 0934	GTTATTATCTCGCCACTTG	TTTGATTTGTTTACACGGGA

To be continued...

Variability between and within accessions

To quantify the genetic variability among the 93 accessions, one individual was selected from the seven collected, based on the highest concentration and the best DNA quality. After that, the most polymorphic primers were selected to analyze the variability between and within the accessions. To quantify the genetic variability between and within accessions, DNA from seven plants was extracted from 14 accessions randomly selected from the 93 accessions (Table 2).

Statistical analyses

The markers were coded as codominant, assigning numbers to the alleles. Thus, when a locus presented three alleles, the codes 11, 22, 33 were attributed to the homozygotes and 12, 13, and 23 for the heterozygotes. Popgene software version 1.31 (Yeh et al. 1999) was used to estimate the genetic variability statistics such as allele frequencies, number

Name	Forward primer	Reverse primer
³ CESR 1041	TTGCTGAAGCCCTTTCTAT	CAGTGTGAGATCATAGCGA
³ CESR 1042	TTGGATTCCTATGAACAAC	TTTGTCTGTCGAATCCTCTC
³ CESR 1044	TTGTCSAAGCTAAGGATTTTC	CCATCTTTCTCTCTTTGTG
³ CESR 1050	TTTCCACACATCAGCGGC	ATAAACCTTCAAACGAGCAA
³ CESR 1055	TTTGAGAGGTGGCAATAACT	GTCACAACCGCAATTAG
³ NS. 260	TCAGCTGTATGTTGAGTGAGCA	AGGGAAGGAACACCTCTCCTA
³ NS. 308	GGAAATTGGTTATGCTCTTTCC	CGCATTGGACTTCTACAAA
³ NS. 689	AGGATGATGATGAGACAAGAAGA	CAGACTGGACTTGAACCTTCACT
³ NS. 720	CCATTACTTACACATTGGACTTCTCCT	GGAAATTGGTTATGCTCTTTCC
³ SSRY 4	ATAGAGCAGAAGTGACAGGCG	CTAACGCACACGACTACGGA
³ SSRY 7	TGCTAAGGAAAATTCATTTCAT	TGCTAAGCTGGTCATGCACCT
³ SSRY 107	CCATTTTCTCTTCTCTGTGCA	TGGTTTGAAGTCTATAAAAATCCTT
³ SSRY 112	CGCAAGGTAATCGGAGCTA	ACAATCAAAGGAGTCGTGTAATC
³ SSRY 113	TTTGTGCTGACTGCCACAATA	TCAACAATTGGACTAAGCAGC
³ SSRY 127	CTTCGGCCTTACAAAAGGA	GCTGAACTGCTTTGCCAACT
³ SSRY 133	AGCATGTCATTGCACCAAAAC	CGACTGCATCAGAACAATGC
³ SSRY 146	TTCCCTCGCTAGAATTGTC	CTATTTGACCGTCTTCGCCG
³ SSRY 150	CAATGCAGGTGAAGTGAATACC	AGGGTGTCTTTCAGAGAAAGG
³ SSRY 151	AGTGAAATAAGCCATGTGATG	CCCATAATTGATGCCAGGTT
³ SSRY 153	TTCCAGAAAGACTTCCGTTCA	CTCAACTACTGCTGCTGACTC
³ SSRY 159	CTTATCCTGTCCCCTCCACC	GACAATTGCATAGGAAGCACA
³ SSRY 177	ACCACAAACATAGGCACGAG	CACCCAATTCAACAATTACCA

Primers underlined are the polymorphic used in the study. Superscripts numbers identify the researchers who developed the primers: ¹ Bressan (2012), ² Pamidimarri et al. (2008), ³ Wen et al. (2010).

of alleles (n_a), number of effective alleles (n_e), observed (H_o) and expected (H_e) heterozygosity, Shannon-Weaver index (H'), Nei's (1972) genetic distance of and the coefficient of inbreeding (f). Analysis of molecular variance (AMOVA, Excoffier et al. 1992) and polymorphism information content (PIC, Botstein et al. 1980) were performed using Genes software (Cruz 2013). For the construction of a circular dendrogram, the Mega7 (Kumar et al. 2016) software was used to perform the UPGMA clustering algorithm from Nei's (1972) genetic distance. Interpretation of the dendrogram was conducted taking into consideration high-change points of cluster fusion.

RESULTS

Among the 60 microsatellite markers tested in the 93 accessions, five were able to detect polymorphisms (Table 2, underlined); therefore, these were used in the analysis of genetic variability.

Genetic variability among the 93 accessions

The locus CESR 0756 was the only one among the five analyzed which allowed the detection of three alleles in the 93 accessions. The locus JCENA 87 enabled the detection of less polymorphism according to allele frequency, although two alleles were detected, similar to the JCDS 10, SSRY 107 and SSRY 127 loci. In JCENA 87, the allele A_1 frequency was almost 100%. The five microsatellite loci used to evaluate the accessions generated a total of 11 alleles (n_a), with an average of 2.2 alleles per locus. The number of effective alleles (n_e) ranged from 1.01 to 2.71, with a mean of 1.91 (Table 3). Because the A_2 allele of locus JCENA87 presented a frequency lower than 0.05 ($P < 0.05$) it can be considered to be rare, with occurrence in only one individual of the accession UFVJC18.

To quantify the genetic variability between 93 accessions, estimates of expected (H_e) and observed (H_o) heterozygosity are important. H_o values were found to be ranging from 0.02 to 0.92, with an average of 0.64. H_e values spanned 0.02 to 0.64, with a mean of 0.42 (Table 3), which indicated a possible heterozygous origin of the collection accessions. For all loci (except JCENA 87), H_o was higher than H_e , revealing an excess of heterozygotes relative to that expected when in

Table 3. Diversity statistics and frequency of the alleles A_1 , A_2 and A_3 for five polymorphic SSR loci used in the evaluation of genetic variability among 93 accessions of *Jatropha curcas* L.. The same five loci were also used on 14 accessions, randomly selected among 93 accessions, for evaluating the variability within and between them

Loci	Allelic frequencies and diversity statistics ¹											
	93 accessions									14 accessions		
	A_1	A_2	A_3	na	ne	Ho	He	f	H'	Ho	He	PIC
JCENA 87	0.9946	0.0054	-	2	1.01	0.02	0.02	0.005	0.03	0.00	0.00	0.00
JCDS 10	0.5054	0.4946	-	2	1.99	0.92	0.50	-0.85	0.69	0.30	0.22	0.18
SSRY 107	0.4194	0.5806	-	2	1.94	0.84	0.49	-0.72	0.68	0.79	0.46	0.35
SSRY 127	0.3804	0.6196	-	2	1.89	0.76	0.47	-0.61	0.66	0.13	0.10	0.10
CESR 0756	0.4835	0.2363	0.2802	3	2.71	0.65	0.64	-0.02	1.05	0.41	0.27	0.22
Mean				2.2	1.91	0.64	0.42	-0.44	0.62	0.42	0.25	0.20
Total				11								

¹ Number of alleles (na), Number of effective alleles (ne), Observed (Ho) and Expected (He) heterozygosity, Coefficient of inbreeding (f), Shannon-Weaver index (H') and Polymorphism information content (PIC).

Hardy-Weinberg equilibrium. For genetic improvement purposes, this result demonstrated the presence of heterozygotes able to be explored in this collection.

A parameter of great importance in assessing genetic variability in populations, by measuring the level of homozygosity, is the coefficient of inbreeding (f). It is essential to verify the existence of crossing among related individuals. Here, f values ranged from -0.005 to -0.85 (Table 3). Notably, negative f values are interpreted as null inbreeding, suggesting that there were no crosses between related individuals in the collection.

The Shannon-Weaver index (H') presented values ranging from 0.03 to 1.05, and the mean value found was 0.62 (Table 3). This mean value of 0.62 reveals the existence of high genetic variability in the collected accessions of *J. curcas*, more than sufficient for the continuity of the breeding program.

All 93 accessions were clustered by the UPGMA method (Figure 1) and the formation of the following 11 clusters was observed: cluster I (orange, 23 accessions), cluster II (purple, 15 accessions), cluster III (pink, 10 accessions), cluster VI (blue, 14 accessions), cluster V (yellow, 4 accessions), cluster VII (dark green, 5 accessions), cluster VIII (red, 4 accessions), cluster IX (light green, 6 accessions), cluster X (dark blue, 10 accessions) and two single clusters (X and XI) formed by accessions UFVJC 53 and UFVJC 7, respectively, which stood out from the others and were collected in Barbacena (MG) and João Pinheiro (MG), respectively. The accessions UFVJC 61 and 70 collected in Rio Grande do Sul (RS) and Ariquemes (RO) were grouped in the same cluster II.

Genetic variability within and between 14 accessions

The same five polymorphic SSR loci were used on 14 accessions, randomly selected on 93, for evaluating the variability within and between them. A mean of 1.86 alleles per locus was found. The analysis of polymorphism information content (PIC) allowed quantification of the genetic polymorphism of each locus in the accessions evaluated. The highest PIC value observed was 0.35 for the locus SSRY 107, while the lowest value occurred for the locus JCENA 87 which did not distinguish within and between accessions (Table 3). The mean value of PIC (0.20) indicated a moderate level of polymorphism of the analyzed loci, with the exception of the locus JCENA 87.

Values of He ranged from 0 for the locus JCENA 87 to 0.46 in the locus SSRY 107, with mean of 0.25. Ho ranged from 0 for the locus JCENA 87 to 0.79 at the locus SSRY 107, with an average of 0.42 (Table 3).

The proportion of variability within and between the 14 accessions was evaluated by AMOVA. Most of the genetic variation (92.58%) was within the populations (Table 4). Thus, it can be concluded that there is considerable genetic variability within populations.

DISCUSSION

The characterization of germplasm collections using molecular markers is an important strategy for the success of




Figure 1. Dendrogram based on the UPGMA cluster analysis among 93 accessions of *Jatropha curcas* L. using five microsatellite loci.

breeding programs. Molecular markers have been used in several studies to characterize accessions of *J. curcas*. Sun et al. (2008) used SSR and AFLP markers to characterize 58 accessions from China, and found a low level of polymorphism. Tatikonda et al. (2009) evaluated the genetic variability of 48 accessions from six different regions of China using AFLP markers. Sudheer et al. (2009) used SSR markers to characterize germplasm from India. Santos et al. (2016) evaluated genetic variability using SSR and ISSR markers in 48 accessions from northern Minas Gerais state, Brazil.

The evaluation of genetic variability, by means of microsatellite markers, in 93 accessions from the UFV's *J. curcas* collection was performed for the first time. This rich collection presents accessions from several Brazilian regions and abroad and its genetic evaluation is important for both genetic conservation and development of improved cultivars.

Regarding the genetic variability statistics obtained here, the number of alleles per loci (n_a), ranged from 2 to 3 (mean=2.2), was similar to that found by several other authors. Santos et al. (2016), using 11 SSR markers, found 2 to 5 alleles per locus, whereas Bressan et al. (2012) found 2 to 8 alleles per locus. However, Na-ek et al. (2011) identified only 1.4 alleles per locus, after evaluating 32 accessions. Rosado et al. (2010) found 1 to 2 alleles per locus in accessions in other Brazilian collection, and of the six microsatellite markers selected by them, four were monomorphic. According to Cruz et al. (2011) it is important to have polymorphic loci that have sufficient numbers of alleles to infer the genetic variability of a population in relation to another or its own over time, especially when subjected to evolutionary forces that promote differentiation.

The number of effective alleles (n_e) is a measure that quantifies the alleles with significant frequency in a population. In the present study, n_e ranged from 1.01 to 2.71. This result was superior to those found by Pecina-Quintero et al.

Table 4. Analysis of molecular variance within and between 14 accessions of *Jatropha curcas* L.

Source of variation	df	Sums of Square	Variation (%)	ϕ_{ST}
Between populations	13	3.86	7.42	0.0742
Within populations	82	15.7	92.58	
Total	95	19.56	100.00	

(2014) (ne from 1.06 to 1.25) evaluating genetic variability in nine *J. curcas* populations from Mexico. Wen et al. (2010), in a study of genetic variability of five *J. curcas* populations, found values for ne ranging from 1.45 to 1.68.

In addition to estimating the number of polymorphic loci, other quantitative measures can be adopted, such as H_o and H_e , which allow to infer about the genetic structure of the population. The means of H_o and H_e in the present study ($H_o=0.64$ and $H_e=0.40$) were similar to those found by Santos et al. (2016). Bressan et al. (2012) also reported similar values ($H_o=0.53$ and $H_e=0.66$) for these measures, although with H_e higher than H_o , in a direction contrary to that verified in the present study and by Santos et al. (2016).

The coefficient of inbreeding or fixation index (f) is a parameter of importance in breeding programs that aim at the development of superior cultivars, as it allows measurement of the level of homozygosity in the population. In our study, the values of f varied from -0.01 to -0.85, being close to expected measures, given the genetic nature of the accessions. Vásquez-Mayorga et al. (2017), in evaluating accessions from Costa Rica, found negative values (-0.10) for f , evidencing that there was no crossing between relatives among accessions. Cruz et al. (2011) stated that negative values of the inbreeding coefficient are common when the H_o values are greater than the expected heterozygosity, suggesting an excess of heterozygous loci. The negative f values should be interpreted as estimates of null inbreeding, that is, there was no crossing among related individuals. *J. curcas* is a monoic species, so it is expected that the loci are in the heterozygous state due to the mating system by allogamy, as evidenced by Sun et al. (2008), Rosado et al. (2010) and Wen et al. (2010).

The Shannon-Weaver index (H') has been used in genetic studies as a measure of genetic variability within populations and resembles a genotype richness index. Here, H' values varied from 0.03 to 1.05 (mean of 0.62), which revealed the existence of high genetic variability among our 93 accessions of *J. curcas* collection. Wen et al. (2010) also used this index to verify genotypic richness in 45 accessions. These authors found an average value of 0.55 using SSR markers and suggested a high level of genetic variability to be explored in five *J. curcas* populations.

Among the accessions that showed the greatest genetic diversity, as revealed by the UPGMA algorithm (Figure 1), we highlight: UFVJC 05, 07, 12, 18, and 53, collected in João Pinheiro, Montalvânia and Barbacena (MG). The greater variability present in the accessions collected in Minas Gerais ratifies the study presented by Dias et al. (2012) that considered the State of Minas Gerais as a secondary center of genetic variability in *J. curcas*.

The formation of 11 clusters with the UPGMA algorithm evidences the expressive genetic variability and structuring of the collection. Sun et al. (2008), evaluating 58 accessions using microsatellite markers, found low genetic variability. Rosado et al. (2010) analyzed the genetic variability of 192 accessions by means of RAPD and SSR markers, finding limited variability among the accessions. Na-ek et al. (2011) evaluated 32 plants from different regions of the world with five SSR markers and also recorded low genetic variability. Naresh et al. (2015) assessed genetic variability among 14 accessions from India using RAPD markers and observed considerable variability among accessions.

The collection of seeds led to the formation of our genebank, prioritizing higher number of seeds per accessions, may be the differential of it in terms of expressive variability. Dias and Kageyama (1991) reported that knowledge of the level of genetic variation and its distribution within and between populations is critical. It is possible to better target the breeding strategies to be adopted, in order to maximize genetic gains through the selection cycles. In *J. curcas*, some studies have also detected a greater amount of genetic variability within populations (Bhering et al. 2015, Pioto et al. 2015, Sinha et al. 2015). This high concentration of genetic variation within populations implies sampling more plants per population, as recommended by Dias and Kageyama (1991) and Bhering et al. (2015). This strategy was effectively practiced in our collection, when priority was given to collecting more seeds per accession. The present study confirmed the accuracy of our accession collection process, in which each accession was represented by 16 plants. Previous studies

employing molecular markers evaluated collections with a reduced number of plants per accession. This is possibly the main reason why they have systematically revealed low genetic variability in the species.

CONCLUSION

The five microsatellite markers selected (JCENA 87, JCDS 10, SSRY 107, SSRY 127 and CESR 0756) were able to detect genetic variability among *J. curcas* accessions, and to enable genetic evaluation through diversity statistics.

The accessions UFVJC 05, 07, 12, 18 and 53 exhibited expressive diversity, being able to comprise a base population for breeding.

Our results demonstrated the existence of molecular genetic variability within and between accessions, indicating that our germplasm collection can be used as a base for breeding program.

ACKNOWLEDGMENTS

Thanks are due to CNPq, FAPEMIG and CAPES for financial support.

REFERENCES

- Bhering LL, Peixoto LA, Leite NLSF and Laviola BG (2015) Molecular analysis reveals new strategy for data collection in order to explore variability in *Jatropha*. **Industrial Crops and Products** **74**: 898-902.
- Botstein D, White RL, Skolnick M and Davis RW (1980) Construction of genetic linkage map in man using restriction fragment length polymorphisms. **American Journal of Human Genetics** **32**: 314-331.
- Bressan EA, Scotton DC, Ferreira RR, Jorge EC, Sebbenn AM, Gerald LTS and Figueira A (2012) Development of microsatellite primers for *Jatropha curcas* (Euphorbiaceae) and transferability to congeners. **American Journal of Botany** **99**: 237-239.
- Creste S, Tulmann Neto A and Figueira A (2001) Detection of single sequence repeat polymorphisms in denaturing polyacrylamide sequencing gels by silver staining. **Plant Molecular Biology Reporter** **19**: 299-306.
- Cruz CD, Ferreira FM and Pessoni LA (2011) **Biometria aplicada ao estudo da diversidade genética**. Universidade Federal de Viçosa, Viçosa, 620p.
- Cruz CD (2013) Genes - a software package for analysis in experimental statistics and quantitative genetics. **Acta Scientiarum** **35**: 271-276.
- Dias LAS and Kageyama PY (1991) Variação genética em espécies arbóreas e consequências para o melhoramento florestal. **Agrotrópica** **3**: 119-127.
- Dias LAS, Missio RF and Dias DCFS (2012) Antiquity, botany, origin and domestication of *Jatropha curcas* (Euphorbiaceae), a plant species with potential for biodiesel production. **Genetics and Molecular Research** **11**: 2719-2728.
- Excoffier L, Smouse PE and Quattro JM (1992) Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. **Genetics** **131**: 479-491.
- Freitas RG, Missio RF, Matos FS, Resende MDV and Dias LAS (2011) Genetic evaluation of *Jatropha curcas*: an important oilseed for biodiesel production. **Genetics and Molecular Research** **10**: 1490-1498.
- Freitas RG, Dias LAS, Cardoso PMR, Evaristo AB, Silva MF and Araújo NM (2016) Diversity and genetic parameter estimates for yield and its components in *Jatropha curcas* L. **Genetics and Molecular Research** **15**: 1-10.
- Gangapur DR, Agarwal P and Agarwal PK (2018) Molecular markers for genetic diversity studies in *Jatropha* (*Jatropha curcas* L.). **AIMS Environmental Science** **5**: 340-352.
- Kumar S, Stecher G and Tamura K (2016) MEGA7: Molecular evolutionary genetics analysis version 7.0 for bigger datasets. **Molecular Biology and Evolution** **33**: 1870-1874.
- Na-ek Y, Wongkaew A, Phumichai T, Kongsiri N, Kaveeta R, Reewongchai T and Phumichai C (2011) Genetic diversity of physic nut (*Jatropha curcas* L.) revealed by SSR markers. **Journal of Crop Science and Biotechnology** **14**: 10-110.
- Naresh B, Medipally SR, Dhanasri G and Devi P (2015) Genetic diversity of *Jatropha curcas* L. accessions and screening for high biodiesel production. **International Journal of Pure & Applied Bioscience** **3**: 124-135.
- Nei M (1972) Genetic distance between populations. **American Naturalist** **106**: 283-292.
- Pamidimarri DVNS, Singh S, Mastan SG, Patel J and Reddy MP (2008) Molecular characterization and identification of markers for toxic and non-toxic varieties of *Jatropha curcas* L. using RAPD, AFLP and SSR markers. **Plant Molecular Biology Report** **36**: 1357-1364.
- Pecina-Quintero V, Anaya-López JL, Zamarripa-Colmenero A, Núñez-Colín CA, Montes-García N, Solís-Bonilla JL and Jiménez-Becerril MF (2014) Genetic structure of *Jatropha curcas* L. in Mexico and probable centre of origin. **Biomass & Bioenergy** **60**: 147-155.
- Pioto F, Costa RS, França SC, Gavioli EA, Bertoni BW and Zingaretti SM (2015) Genetic diversity by AFLP analysis within *Jatropha curcas* L. populations in the State of São Paulo, Brazil. **Biomass & Bioenergy** **80**: 316-320.

- Rosado TB, Laviola BG, Faria DA, Pappas MR, Bhering LL, Quirino B and Grattapaglia D (2010) Molecular markers reveal limited genetic diversity in a large germplasm collection of the biofuel crop *Jatropha curcas* L. in Brazil. **Crop Science** **50**: 2372-2382.
- Santos DN, Ferreira JL, Setoaw TA, Caçado GMA, Pasqual M, Londe LCN, Saturnino HM and Vendrame WA (2016) Genetic structure from the oldest *Jatropha* germplasm bank of Brazil and contribution for the genetic improvement. **Anais da Academia Brasileira de Ciências** **88**: 2363-2374.
- Sinha P, Islam MA, Negi MS and Tripathi SB (2015) First identification of core accessions of *Jatropha curcas* from India based on molecular genetic diversity. **Journal of Plant Genetic Resources** **14**: 77-80.
- Sorrell S, Miller R, Bentley R and Speirs J (2010) Oil futures: a comparison of global supply forecasts. **Energy Policy** **38**: 4990-5003.
- Sudheer PDVN, Singh S, Mastan SG, Patel J and Reddy MP (2009) Molecular characterization and identification of markers for toxic and non-toxic varieties of *Jatropha curcas* L. using RAPD, AFLP and SSR markers. **Molecular Biology Reports** **36**: 1357-1364.
- Sun QB, Li LF, Li Y, Wu GJ and Ge XJ (2008) SSR and AFLP markers reveal low genetic diversity in the biofuel plant *Jatropha curcas* in China. **Crop Science** **48**: 1865-1871.
- Tatikonda L, Wani SP, Kannan S, Beerelli N, Sreedevi TK, Hoisington DA, Devi P and Varshney RK (2009) AFLP-based molecular characterization of an elite germplasm collection of *Jatropha curcas* L., a biofuel plant. **Plant Science** **176**: 505-513.
- Vásquez-Mayorga M, Fuchs EJ, Hernández EJ, Herrera F, Hernández J, Moreira I, Arnáez E and Barboza NM (2017) Molecular characterization and genetic diversity of *Jatropha curcas* L. in Costa Rica. **Peer J** **5**: 29-31.
- Wen M, Wang H, Xia Z, Zou M, Lu C and Wang W (2010) Development of EST-SSR and genomic-SSR markers to assess genetic diversity in *Jatropha curcas* L. **BMC Research Notes** **3**: 42-50.
- Yeh FC, Boyle T, Rongca Y, Ye Z and Xiyang JM (1999) **POPGENE. Microsoft window-based freeware for population genetic analysis**. Version 1.31, Manual.