

Inheritance of seed size in watermelon populations

Herencia del tamaño de la semilla en poblaciones de sandía


TIAGO LIMA DO NASCIMENTO^{1, 4}FLÁVIO DE FRANÇA SOUZA²RITA DE CÁSSIA SOUZA DIAS²JOICE SIMONE DOS SANTOS³NATONIEL FRANKLIN DE MELO²

¹ Feira de Santana State University, Fundação de Amparo à Pesquisa da Bahia; CAPES Scholarship Student in Plant Genetic Resources, Department of Biological Sciences, Feira de Santa (Brazil). ORCID Nascimento, T.L.: <https://orcid.org/0000-0002-0916-6344>

² Embrapa, Brazilian Semiarid Agricultural Research Corporation, Genetics and Plant Breeding, Petrolina (Brazil). ORCID Souza, F.F.: <https://orcid.org/0000-0002-9632-2666>; ORCID Dias, R.C.S.: <https://orcid.org/0000-0002-5527-3693>; Melo, N.F.: <https://orcid.org/0000-0001-6888-4090>

³ Federal Institute of Education, Science and Technology of Ceará, Crateus (Brazil). ORCID Santos, J.S.: <https://orcid.org/0000-0003-3597-2021>

⁴ Corresponding author. tiago_lim.a@hotmail.com



Seed phenotype in the studied populations.
Photo: T.L. Nascimento

Last name: NASCIMENTO / SOUZA / DIAS / SANTOS / MELO

Short title: Inheritance of seed size in watermelon

Doi: <https://doi.org/10.17584/rcch.2020v14i1.10521>

Received for publication: 20-01-2020 Accepted for publication: 30-03-2020

ABSTRACT

Seed size is an important agronomic trait and is applicable to different abilities. Small seeds guarantee the greater use of the pulp, while larger seeds facilitate sowing. However, there is little work on the genetic control of this characteristic in watermelon. The objective of this work was to study the seed size inheritance in watermelon populations by crossing contrasting genotypes, seeking to gain information to provide technical support during the selection of seed size for the development of new watermelon genotypes. The seed lengths of six populations, P₁, P₂, F₁, F₂, BC₁ and BC₂, were measured using the GENES software segregating and nonsegregating generations procedure. This trait is controlled by two genes with incomplete dominance. In addition, depending on the populations studied, inheritance for the characteristic in question may behave differently. Nevertheless, the selection of superior individuals within populations can be performed based on this phenotype, which allows the exploitation of these individuals within breeding programs to develop lines or hybrids.

Additional keywords: *Citrullus lanatus*; seed characteristics; crop improvement; selection; incomplete dominance.

RESUMEN

El tamaño de las semillas es un rasgo agronómico importante, pudiendo utilizarse para diferentes habilidades. Mientras que las semillas pequeñas garantizan un mayor aprovechamiento de la pulpa, semillas más grandes facilitan la siembra, sin embargo, existen pocos trabajos sobre el control genético de este carácter en la sandía. El objetivo de este trabajo fue estudiar la herencia del tamaño de las semillas en poblaciones de sandía cruzando genotipos contrastantes, buscando proporcionar soporte técnico durante la selección del tamaño de la semilla en el desarrollo de nuevos genotipos de este fruto. Fue medida la longitud de la semilla de seis poblaciones P₁, P₂, F₁, F₂, RC₁ y RC₂ utilizando el procedimiento generaciones segregantes y no segregantes (GENES), por medio del cual fue posible concluir que la herencia del tamaño de semillas en las poblaciones estudiadas es controlada por dos genes con dominancia incompleta. Además, dependiendo de las

poblaciones estudiadas la herencia del carácter en cuestión puede comportarse de manera diferente. La selección de individuos superiores dentro de las poblaciones puede ser realizada con base en el fenotipo, lo que posibilita la exploración de estos individuos dentro de los programas de mejoramiento para componer líneas o híbridos.

Palabras clave adicionales: *Citrullus lanatus*; características de la semilla; mejoramiento de cultivos; selección; codominancia incompleta.

INTRODUCTION

Watermelon [*Citrullus lanatus* (Thunb.) Matsum and Nakai] is one of the highest yielding fruit crops in Brazil, and according to the 2016 world ranking, the country was ranked as the 4th largest watermelon producer, with 2,090,432 t of fruit, second only to China (79,043,138 t), Turkey (3,928,892 t) and Iran (3,813,850 t) (Fao, 2018). The development of new genotypes with commercial attributes different from those already on the market, such as the size of the seeds, can make the production of Brazilian watermelon more expressive. Moreover, the consumer market has increasingly been demanding nutritious and healthy fruits (Farias *et al.*, 2014).

For fresh consumption, the smaller the seed size, the easier it becomes to consume the fruit, while for the production of fruit and sowing in the field, the larger the seed size is, the greater the production and the easier the planting. Thus, understanding the genetic control of traits is important for the development of genotypes aimed at having the most diverse characteristics (Li *et al.*, 2018). However, it is worth mentioning that information about the genetic behavior of watermelon characteristics is scarce. Among these reports, those studying the size of the seeds, besides being rare, are quite old, which characterizes it as being one of the main obstacles for breeders.

In previous studies, Poole *et al.* (1941), cited by Li *et al.* (2018), described that the genetic control of seed size was controlled by two genes, 's' and 'l', with 's' being epistatic to 'l'. A few years later, the 'Ti' genes for tiny seeds and 'ti' genes for tomato seeds were described (Tanaka *et al.*, 1995; Li *et al.*, 2018). According to Yong *et al.* (2009), watermelon seed size can be classified as giant, large, medium, small, micro and tiny. In addition to seed size, inheritance studies of other characteristics in watermelon have been reported, including fruit yield (Kumar and Wehner, 2013), seed mass (Adjouman *et al.*, 2016), resistance to the PRSV-W virus (Alves *et al.*, 2014), and egusi seed (Gusmini *et al.*, 2004). This suggests that before

performing a characteristic improvement, it is necessary to understand the genetic control mechanisms within the populations of interest.

Thus, the objective of the present work was to study the inheritance of seed size characteristics in watermelon populations through crossing of contrasting genotypes.

MATERIAL AND METHODS

Obtaining the populations

The experiment was conducted from 2017 to 2018 at the Bebedouro Experimental Station at Embrapa Semiárido in Petrolina-PE (Brazil). Two watermelon lines, with contrasting seed size, were used line 31715.001 (“P₁”) (small seed) and lineage 31717.005 (“P₂”) (large seed)—to obtain F₁, F₂, BC₁ and BC₂ populations.

As such, crosses were carried out via controlled hand-pollination (CHP) according to the methods used by Nascimento *et al.* (2018). Regarding the CHP, in the flowering stage, male and female flowers of the lineages and the F₁ hybrids were isolated at preanthesis using aluminum sachet bags. On the following day, the open male flowers were detached from the plants, their petals were removed to expose their anthers, and then pollen grains were gently deposited on the stigmas of the female flowers. After this procedure, the pollinated flowers were identified with cardboard tags containing information on the parental crossings, plant type, and CHP date. Afterward, the flowers were isolated again for a period of 72 h. Between 30 and 35 d after pollination, the fruits were harvested, and their seeds were extracted, washed and left to dry in the shade.

To obtain the F₁ population, between January and April 2017, seeds were sown in polystyrene trays containing proper commercial substrate and then kept in a greenhouse for 12 d. After that period, the seedlings were transplanted to the field.

In the following generation, five seeds from the parents (P₁ and P₂) and from the F₁ population were sown during the months of May to August 2017, and using CHP, the F₁ plants were self-fertilized and simultaneously crossed with the parents, P₁ and P₂, which produced the seeds of F₂, BC₁ and BC₂ populations. All populations obtained were then evaluated in the field, from May to August 2018, under open pollination.

Fertilization was carried out according to a soil analysis and as recommended by Mendes *et al.* (2010), with 30 kg ha⁻¹ of N, 80 kg ha⁻¹ of P₂O₅, and 30 kg ha⁻¹ of K₂O, plus 15 kg ha⁻¹ of ZnSO₄ and 10 kg ha⁻¹ of CuSO₄. Fertilizers were top-dressed with irrigation water

containing 50 kg ha⁻¹ of N in the form of Ca (NO₃)₂ and 40 kg ha⁻¹ of K₂O using KCl, between 50 to 60 d after sowing.

The water was applied by drip irrigation at a daily depth that varied according to the plants' needs depending on climatic conditions, monitored by a weather station located near the experimental area. The crop coefficients (K_c values) used to calculate crop evapotranspiration (ET_c) were those obtained by Freitas and Bezerra (2004) in Canindé, CE, Brazil, for Crimson Sweet watermelon, whose corresponding values were 0.46 to 0.70 at the vegetative stage, 0.89 to 1.22 at the flowering stage, and 1.14 to 0.74 at the fruiting stage.

Phytosanitary treatments were carried out with agrochemicals registered by the Brazilian Ministry of Agriculture, Livestock and Supply, and weeds were removed manually.

Phenotypes of seed size in watermelon populations

During the months of April to August 2018, seeds from all the populations were germinated in expanded polystyrene trays containing commercial substrate for vegetables. At 12 d after sowing, the seedlings were transplanted into the field at the Bebedouro Experimental Station in accordance with a completely randomized design, with each plant considered an experimental unit and with a varied number for each population, in soil previously prepared with gray plastic mulching sheets, with 1 m spacing between plants and 2.5 m between rows. After transplanting, the seedlings were covered with an agrotexile blanket until the flowering period, at which point it was removed for pollination. The seeds of each population came from a single fruit.

At 70 d after planting, fruits were harvested. The seeds were subsequently extracted, washed and left to dry in the shade. Eight, 10, 7, 271, 95 and 94 plants from the P₁, P₂, F₁, F₂, BC₁ and BC₂ populations, respectively, were evaluated.

To investigate seed size inheritance, seed length was measured, as this characteristic is highly correlated with width and weight (Hawkins and Dane, 2001). The lengths of 10 seeds were measured with a digital caliper (Stainless Hardened: JL-YB5474) following the method used by Nascimento *et al.* (2018), with seeds randomly chosen from one fruit of each plant.

The mean and variance [P₁ ($\sigma_{f(P_1)}^2 = \sigma_{m(P_1)}^2 = \sigma_{P_1}^2$), P₂ ($\sigma_{f(P_2)}^2 = \sigma_{m(P_2)}^2 = \sigma_{P_2}^2$), F₁ ($\sigma_{f(F_1)}^2 = \sigma_{m(F_1)}^2 = \sigma_{F_1}^2$); F₂ ($\sigma_{f(F_2)}^2 = \sigma_{g(F_2)}^2 + \sigma_{m(F_2)}^2$), BC₁ ($\sigma_{f(BC_1)}^2 = \sigma_{g(BC_1)}^2 + \sigma_{m(BC_1)}^2$) and BC₂ ($\sigma_{f(BC_2)}^2 = \sigma_{g(BC_2)}^2 + \sigma_{m(BC_2)}^2$)] of the populations were then estimated.

The mean degree of dominance (MDD) of the characteristic was estimated based on the variances and in accordance with the model $MDD = 2 \bar{F}_1 - (\bar{P}_1 + \bar{P}_2)/\bar{P}_1 - \bar{P}_2$. Broad-sense (h_a^2) (1) and narrow-sense (h_r^2) (2) heritability were estimated as follows:

$$h_a^2 = \frac{\sigma_{g(F_2)}^2}{\sigma_{f(F_2)}^2} = \frac{\sigma_{f(F_2)}^2 - (\sigma_{P_1}^2 + 2\sigma_{F_1}^2 + \sigma_{P_2}^2)/4}{\sigma_{f(F_2)}^2} \quad (1)$$

$$h_r^2 = \frac{\sigma_a^2}{\sigma_{f(F_2)}^2} = \frac{2\sigma_{f(F_2)}^2 - [\sigma_{f(BC_1)}^2 + \sigma_{f(BC_2)}^2]}{\sigma_{f(F_2)}^2} \quad (2)$$

Additionally, the values of k1 (based on variance) and k2 (based on the mean) were estimated, where $k = 0$ indicates the absence of dominance ($k=0$ =additive), $0 \leq k \leq 1$ indicates incomplete dominance, $k > 1$ indicates over dominance, and $k=1$ indicates complete dominance.

Additionally, the minimum number of genes (MNG) controlling the characteristic was estimated by the formula $n = (\bar{P}_1 + \bar{P}_2)^2 / 8[\sigma_{g(F_2)}^2]$. In terms of computer resources, the “segregating and non-segregating generations” procedure of the GENES software program was used (Cruz, 2016).

Afterward, the characteristic mean was subjected to the chi-square (X^2) test at a 0.05 significance level for verification of the mendelian pattern within the F_2 population according to the following model (3):

$$\chi^2 = \sum_{i=1}^n \left[\frac{(Obs_i - Esp_i)^2}{Esp_i} \right] \quad (3)$$

where X^2 is the chi-square value calculated and Obs_i and Esp_i are the observed and expected frequencies of the i -th phenotypic class ($i = 1, 2, \dots, n$), respectively (Schuster and Cruz, 2013).

RESULTS AND DISCUSSION

The mean, variance and standard deviation of the seed length of the parents of the segregating populations (Tab. 1) were different, which indicated the existence of genetic variability between them.

Table 1. Number of plants, means, variance and standard deviations of seed size, as evaluated for six watermelon populations (*Citrullus lanatus* var. *lanatus*).

Generation (seed size) ¹	Number of individuals	Mean	Variance	Standard deviation
P ₁ (small)	8	5.90	0.0286	0.17
P ₂ (largest)	10	11.63	0.0934	0.31
F ₁ (medium)	7	6.89	0.0214	0.15
F ₂ (¼ small, ½ medium and ¼ largest)	271	7.71	5.4044	2.32
BC ₁ (½ small and ½ medium)	95	6.24	0.2474	0.50
BC ₂ (½ medium and ½ largest)	94	9.68	5.4711	2.34

¹ Seed size in populations.

The results are in line with the assumption of Cruz *et al.* (2014), who stated that better precision in genetic analyses requires parents whose studied characteristics contrast. According to those authors, genetic parameter estimates allow the identification of the nature of the action of the genes involved in characteristic control. Thus, estimates can be used as an advantageous auxiliary tool in breeding program management.

The variance expressed by the parents and the hybrids showed low values, while the F₂ and RC₂ populations presented high values. Similar results were reported by Souza *et al.* (2006), who, when studying seed size inheritance in watermelon generations, observed high values of variance in the F₂ and BC₂ populations. The high values in both studies possibly occurred because gene segregation within these populations is higher than in the other populations.

Although watermelon is an allogamous species, the low values of variance in the P₁, P₂ and F₁ populations of the present study are due to environmental factors, since the parents used to develop the segregating populations were generated by seven cycles of self-fertilization. Thus, each of the plants of these generations is considered to correspond to the same genotype. However, expecting higher values of the variance of F₂ populations and backcross generations due to the interaction of genetic and environmental factors influences the phenotypic expression of traits (Cruz *et al.*, 2014).

The F₂ population presented the fourth lowest mean. According to Zewdie and Bosland (2000), depending on the interaction of a trait, the mean for this population may increase or decrease, indicating that parents are contributing alleles to increase or decrease the trait under study. Based on this information, the mean value of the F₂ population may have occurred

because most seeds were classified as medium and short, which together represent approximately 75% of the total amount.

According to the genetic parameter estimates for the F₂ population (Tab. 2), in comparison with the environmental variance, the genetic variance had the greatest influence. Nascimento (2017) found similar results for watermelon seed size, with a higher genetic variance than environmental variance. These results show that the environment has little influence on the phenotypic expression of the evaluated characteristic and that the phenotype observed was mostly the result of genetic influence.

Table 2. Genetic parameters related to seed size in a watermelon F₂ population obtained by crossing contrasting varieties.

Parameter	Value
Phenotypic variance (σ_p^2)	5.404
Environmental variance (σ_e^2)	0.041
Genotypic variance (σ_g^2)	5.363
Additive variance (σ_{ad}^2)	5.090
Variance of dominance deviation (σ_d^2)	0.273
Broad-sense heritability (h_a^2)	99.237
Narrow-sense heritability (h_f^2)	98.189
Mean degree of dominance (MDD)	0.327
K values	k1 (0.33) and k2 (0.66)
Minimum number of genes controlling the characteristic (MNG)	1.82
Maximum value	12.8
Minimum value	4.2

The estimates of the phenotypic, genotypic, additive and environmental variances showed that the contribution of the additive-effect genes to the dominance genes in terms of seed size was significant (Tab. 2), indicating additive allelic interaction. Corroborating the results of the present study, Tanaka *et al.* (1995), when examining the inheritance of fruit shape and seed size in two contrasting watermelon populations, also observed that seed size was governed predominantly by additive allelic interaction, with the small-seed phenotype being an easily improved characteristic.

However, Adjoumani *et al.* (2016), when studying the genetics of seed traits resulting from the intraspecific crossbreeding of distinct watermelon varieties, observed that the effects of genes for the characteristic in question showed dominant allelic interactions. The differing results reported by the above authors can be attributed to the genetic differences of the populations evaluated in each study.

It is worth emphasizing that the predominance of additive allelic interaction facilitates the selection stage, since superior individuals also show superior descent (Cruz *et al.*, 2014). Regarding dominant allelic interactions, breeding programs can benefit from developing hybrids.

Within the F₂ population, one superior individual was observed for the smallest seed size (4.2 mm) (Tab. 2). This individual can be further investigated and give rise to a small-seed open-pollinated cultivar and can also produce a small-seed-size variety, which can be used for hybrid combinations that may result in reduced-seed-size genotypes. It is worth noting that small seed size is dominant over large seed size, and medium size is dominant over small and large sizes (Guner and Wehner, 2004).

The use of transgressive genotypes to develop hybrids with reduced seed size can be very promising, as producers may be able to acquire hybrid seeds of relatively large size, making planting easier, whereas the fruits of these hybrids will have relatively small seeds, thus making watermelon consumption more agreeable and providing a market advantage over the other genotypes.

In the present study, individuals from the F₂ population presented the following approximate phenotypic percentages of seed size: small (25%), medium (50%) and large (25%). Previous inheritance studies on watermelon seed traits such as weight showed that monogenic segregation at a 3:1 ratio did not occur as expected but was close enough to suggest that the smallest seed mass was monogenic and dominant over the largest seed mass (Weetman, 1937).

Similar results were observed in the present study for seed size, where the smallest size, although not in accordance with the expected 3:1 ratio, was quite close to a ratio of 1:2:1, thus suggesting that the characteristic is controlled by two genes.

In addition, the seed size characteristic showed high broad-sense and narrow-sense heritability (Tab. 2). Similar results were observed by Amburani (2018), who, when studying the heritability of morphoagronomic traits in 30 watermelon genotypes, obtained heritability estimates of more than 80% for seven of the 13 traits evaluated, including 100-seed weight and number of seeds per fruit. According to Cruz *et al.* (2014), narrow-sense heritability represents the variance of the additive type fixed in the populations as a result of the advancement of generations. Based on this information, the correlation between phenotypic values and genetic values is considered high for the characteristic in question, and it is possible to select individuals with relatively small seed sizes based on their phenotype.

To infer the mean degree of dominance and determine the best strategy to improve various characteristics, the values of k_1 (0.33) and k_2 (0.66) (Tab. 2) were calculated and showed mutual agreement, indicating the existence of incomplete dominance for seed size in the studied populations. Adjoumani *et al.* (2016), when studying the genetics of seed traits resulting from the intraspecific crossing of distinct watermelon varieties, obtained the same results. In both studies, these results reinforce the efficiency of genotype selection of individuals of interest based on the phenotype.

The minimum value for the number of genes involved in the control of the studied characteristic was 1.82 (Tab. 2). This information is important for estimating the type of inheritance that controls a given characteristic (Lobo *et al.*, 2005). In the present study, seed size is controlled by two genes, indicating that simpler breeding methods, such as mass selection, can be used to obtain small-seed-size cultivars.

When we performed the chi-square test on the populations (Tab. 3), there were no significant differences between populations for observed and expected frequencies associated with phenotypic segregation of the BC_1 population, in which all individuals should present the same seed size. However, it was observed that the seeds of the individuals were divided into small and medium sizes, confirming the hypothesis by which two genes control the characteristic under incomplete dominance.

Table 3. Chi-square test results of the watermelon populations to investigate the segregation of two loci based on the 1:2:1 ratio.

Generation	Observed			Expected			X^2	DL	P
	Small	Medium	Large	Small	Medium	Large			
P_1	8	0	0	8.0	0	0	-	-	-
P_2	0	0	10	0	0	10.0	-	-	-
F_1	0	7	0	0	7.0	0	-	-	-
F_2	64	130	49	67.75	135.5	67.75	5.62 ^{ns}	2	0.05
BC_1	45	50	0	47.5	47.5	0	0.263 ^{ns}	1	0.05

^{ns} not significant at the 0.05 significance level (P) by the chi-square test (X^2); DF: degrees of freedom.

Similarly, Poole *et al.* (1941), cited by Prothro *et al.* (2012) studied seed length inheritance by evaluating crosses between plants that produce small and large, small and medium, and medium and large seeds and proposed that this characteristic is controlled by two genes because medium-size seeds were dominant over small- and large-size seeds. Those authors proposed that the recessive *l* and *s* genes afforded large- and small-seed phenotypes, respectively. In addition, among the evaluated genotypes, the *s* gene was epistatic for *l*, and

LL SS was assigned to medium-size seeds; *ll SS*, large-size seeds; and *LL ss* or *ll ss*, small-size seeds.

On the other hand, Tanaka *et al.* (1995) proposed that, in a Sweet Princess watermelon population, tiny seed size was monogenic, controlled by the *Ti* gene and dominant over medium seed size; thus, the tiny seed size characteristic was possibly governed by simple inheritance.

The differences observed in the results of the present study possibly show that, for the characteristic in question and depending on the cultivars involved in the crosses, inheritance may behave differently with respect to the populations under study, so it is not possible to extrapolate these results to other populations. Furthermore, the method adopted here for seed size measurement was quantitative, expressed in millimeters, while the study by Polle *et al.* (1941) categorized seed size into classes; that is, they performed a qualitative assessment, disregarding minimal size differences and limiting the population.

The results obtained by means of the predicted gain from selection (Tab. 4) reinforce that, for the improvement of the characteristic under study, the use of simpler breeding methods can be effective, since the gain from the selection was 59% for relatively large seeds and a decrease of just under 30% for relatively small ones. In other words, it is possible to select and recombine individuals that produce large seeds and small seeds, seeking to develop hybrid combinations that produce seeds of intermediate size; this is in addition to the possibility of exploiting these individuals to develop new cultivars for different purposes.

Table 4. Predicted gain from selection for seed size in a watermelon F₂ population obtained by crossing contrasting varieties.

Parameters	Selection	
	Highest values	Lowest values
MSI	12.577	5.285
S	4.870	-2.422
GS (%)	59.516	-29.605
PM 1° S	12.294	5.425

MSI: mean of selected individuals; S: selection differential; GS (%): gain from selection (in percentage), and PM1°S; predicted mean for a selection cycle.

CONCLUSIONS

In the present work, inheritance of the smallest seed size trait in the studied populations is controlled by two genes with incomplete dominance;

Depending on the population under study, the inheritance behavior for the seed size characteristic may behave differently.

Selection of superior individuals for the smallest seed size within an F₂ population can occur based on phenotype;

Increased numbers of individuals can be exploited within breeding programs to develop lines or hybrids.

ACKNOWLEDGMENTS

To the Coordination for the Improvement of Higher Education Personnel (CAPES) for the granting. The author is grateful to Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB - Bahia Research Foundation) for the doctoral scholarship granted; to Embrapa Semiárido (a unit of the Brazilian Agricultural Research Corporation) for the partnership and the infrastructure made available; to the Graduate Program in Plant Genetic Resources (PGGRGV) of the State University of Feira de Santana for support in the courses; and to the Embrapa Semiárido trainees and the Biological Sciences undergraduates at the University of Pernambuco (UPE), Karina Branco, Jackson, Ingrid Caroline, Débora Eduarda and André Granja for their help in the experiment evaluations.

BIBLIOGRAPHIC REFERENCES

Adjoumani, K., S.B. Bony, G.K. Koffi, L.C. Kouonon, F.K. Brou, and R. Sié. 2016. Genetic evaluation of seed traits from intraspecific crossing of genetically distinct watermelon varieties. Afr. Crop Sci. J. 24(2), 143-154. Doi: <https://doi.org/10.4314/acsj.v24i2.3>

Alves, A.F., I.R. Nascimento, C.A. Ticona-Benavente, M.V. Faria, R.A. Sarmento, A.R. Figueira, and W.R. Maluf. 2014. Herança da resistência do acesso de melancia PI 595201 a isolado de PRSV-W do Estado do Tocantins. Bragantia 73(2), 138-142. Doi: <https://doi.org/10.1590/brag.2014.022>

Amburani, A. 2018. Studies on genetic variability and genetic advance for yield parameters in watermelon (*Citrullus lanatus* Thumb.). Asian J. Hortic. 13 (2), 39-44. Doi: <https://doi.org/10.15740/HAS/TAJH/13.2/39-44>

Cruz, C.D. 2016. Genes Software – extended and integrated with the R, Matlab and Selegen. Acta Sci.-Agron. 38 (4), 547-552. Doi: <https://doi.org/10.4025/actasciagron.v38i3.32629>

Cruz C.D., P.C.S. Carneiro, and A.J. Regazzi. 2014. Modelos biométricos aplicados ao melhoramento genético. 4th ed. Universidade Federal de Viçosa, Viçosa, Brazil.

FAO. 2018. FAOSTAT. Production crops. In: <http://www.fao.org/faostat/es/#data/QC>; consulted: June, 2019.

Farias, R.M., C.F. Barreto, R.R. Zandoná, J.P. Rosado, and C.R. Martins. 2014. Comportamento do consumidor de frutas na região da fronteira Oeste do Rio Grande do Sul com Argentina e Uruguai. Rev. Bras. Frutic. 36(4), 872-883. Doi: <https://doi.org/10.1590/0100-2945-417/13>

Freitas, A.A. and F.M.L. Bezerra. 2004. Coeficientes de cultivo para melancia nas suas fases fenológicas. Rev. Ciênc. Agron. 35(2), 319-325.

Guner, N. and T.C. Wehner. 2004. The genes of watermelon. HortScience 39(6), 1175-1182. Doi: <https://doi.org/10.21273/HORTSCI.39.6.1175>

Gusmini, G., T.C. Wehner, and R.L. Jarret. 2004. Inheritance of Egusi seed type in watermelon. J. Hered. 95(3), 268-270. Doi: <https://doi.org/10.1093/jhered/esh031>

Hawkins, L.K. and F. Dane. 2001. Molecular markers associated with morphological traits in watermelon. HortScience 36(7), 1318-1322. Doi: <https://doi.org/10.21273/HORTSCI.36.7.1318>

Kumar, R. and T.C. Wehner. 2013. Quantitative analysis of generations for inheritance of fruit yield in watermelon. Hortscience 48(7), 844-847. Doi: <https://doi.org/10.21273/HORTSCI.48.7.844>

Li, N., J. Shang, J. Wang, D. Zhou, N. Li, and S. Ma. 2018. Fine mapping and discovery of candidate genes for seed size in watermelon by genome survey sequencing. *Sci. Rep.* 8(17843), 1-11. Doi: <https://doi.org/10.1038/s41598-018-36104-w>

Lobo, V.L.S., Giordano, L.B. and C.A. Lopes. 2005. Herança da resistência à mancha bacteriana em tomate. *Fitopatol. Bras.* 30(4), 343-349. Doi: <https://doi.org/10.1590/S0100-41582005000400002>

Mendes, A.M.S., C.M.B. Faria, and D.J. Silva. 2010. Adubação. In: Dias, R.C.S., G.M. Resende, and N.D. Costa (eds.). Sistema de produção de melancia. *Electronic version*. In: Embrapa Semiárido, <https://sistemasdeproducao.cnptia.embrapa.br/FontesHTML/Melancia/SistemaProducaoMelancia/adubacao.htm>; consulted: april 2019.

Nascimento, T.L. 2017. Divergência genética, capacidade de combinação e heterose em melancia. PhD thesis. Universidade Federal Rural de Pernambuco, Recife, Brazil.

Nascimento, T.L., F.F. Souza, R.C.S. Dias, and E.F. Silva. 2018. Agronomic characterization and heterosis in watermelon genotypes. *Pesqui. Agropecu. Trop.* 48(2), 170-177. Doi: <https://doi.org/10.1590/1983-40632018v48i1779>

Poole, C.F., P.C. Grimball, and D.R. Porter. 1941. Inheritance of seed characters in watermelon. *J. Agric. Res.* 63(3), 433-456.

Prothro, J., K. Sandlin, and H. Adel-Haleem. 2012. Main and epistatic quantitative trait loci associated with seed size in watermelon. *J. Am. Soc. Hortic. Sci.* 137(6), 452-457. Doi: <https://doi.org/10.21273/JASHS.137.6.452>

Schuster, I. and C.D. Cruz. 2013. Estatística genômica aplicada a populações derivadas de cruzamentos controlados. 2nd ed. Universidade Federal de Viçosa, Viçosa, Brazil.

Souza, F.F., E.B.A. Souza, A.C. Silva, L.R.S. Neves, R.C.S. Dias, and M.A. Queiróz. 2006. Estudo de herança do tamanho da semente em melancia. pp. 1613-1616. In: *Proc.* 46

Congresso Brasileiro de Olericultura. Associação Brasileira de Horticultura (ABH), Goiânia, Brazil.

Tanaka, T., S. Wimol, and T. Mizutani. 1995. Inheritance of fruit shape and seed size of watermelon. J. Jpn. Soc. Hortic. Sci. 65 (3), 543-548. Doi: <https://doi.org/10.2503/jjshs.64.543>

Weetman, L.M. 1937. Inheritance and correlation of shape, size, and color in the watermelon, *Citrullus vulgaris* Schrad. Res. Bull: Iowa Agriculture and Home Economics Experiment Station 20(228), 222-256.

Yong, J.K., Y. TaeJin, P.Y. Hoon, L.Y. Jik, K.S. Cheol, K.Y. Kwon, and C.J. Lai. 2009. Development of near isogenic lines with various seed sizes and study on seed size-related characteristics in watermelon. Korean J. Breed. Sci. 41(4), 403-411.

Zewdie, Y. and P. Bosland. 2000. Capsaicinoid inheritance in an interspecific hybridization of *Capsicum annum* 9 *C. chinense*. J. Am. Soc. Hortic. Sci. 125 (4), 448-453. Doi: <https://doi.org/10.21273/JASHS.125.4.448>