

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/353646601>

Large-scale degradation of the Tocantins–Araguaia River basin

Article in *Environmental Management* · August 2021

CITATIONS

0

READS

764

57 authors, including:



Fernando Pelicice

Universidade Federal de Tocantins

106 PUBLICATIONS 5,226 CITATIONS

[SEE PROFILE](#)



Angelo A Agostinho

Universidade Estadual de Maringá

325 PUBLICATIONS 15,327 CITATIONS

[SEE PROFILE](#)



Alberto Akama

Museu Paraense Emilio Goeldi - MPEG

37 PUBLICATIONS 471 CITATIONS

[SEE PROFILE](#)



José D Andrade Filho

Fundação Oswaldo Cruz

131 PUBLICATIONS 2,012 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Taxonomic and bioecological studies of sand flies (Diptera: Psychodidae) collected at Rebenta Bombas cave, Lassance, Minas Gerais, Brazil [View project](#)



Photoperiod Differences in Sand Fly (Diptera: Psychodidae) Species Richness and Abundance in Caves in Minas Gerais State, Brazil [View project](#)



Large-scale Degradation of the Tocantins-Araguaia River Basin

Fernando Mayer Pelicice¹ · Angelo Antonio Agostinho² · Alberto Akama³ · José Dilermando Andrade Filho⁴ · Valter M. Azevedo-Santos⁵ · Marcus Vinicius Moreira Barbosa⁶ · Luis Mauricio Bini⁷ · Marcelo Fulgêncio Guedes Brito⁸ · Carlos Roberto dos Anjos Candeiro⁹ · Érica Pellegrini Caramaschi¹⁰ · Priscilla Carvalho¹¹ · Rodrigo Assis de Carvalho¹² · Leandro Castello¹³ · Davi Borges das Chagas¹⁴ · Carine Cavalcante Chamon¹⁵ · Guarino Rinaldi Colli¹⁶ · Vanessa Salette Daga¹⁷ · Murilo Sversut Dias¹⁸ · José Alexandre Felizola Diniz Filho⁷ · Philip Fearnside¹⁹ · Wagner de Melo Ferreira¹ · Diego Azevedo Zoccal Garcia²⁰ · Tiago Kutter Krolow²¹ · Rodrigo Ferreira Kruger²² · Edgardo Manuel Latrubesse²³ · Dilermando Pereira Lima Junior²⁴ · Solange de Fátima Lolis¹ · Fabyano Alvares Cardoso Lopes²⁵ · Rafael Dias Loyola²⁶ · André Lincoln Barroso Magalhães²⁷ · Adriana Malvasio²⁸ · Paulo De Marco Jr.⁷ · Pedro Ribeiro Martins²⁹ · Rosana Mazzoni³⁰ · João Carlos Nabout¹² · Mário Luis Orsi²⁰ · Andre Andrian Padial³¹ · Hasley Rodrigo Pereira³² · Thiago Nilton Alves Pereira¹⁵ · Phamela Bernardes Perônico¹² · Miguel Petrere Jr.³³ · Renato Torres Pinheiro³⁴ · Etiene Fabbrin Pires³⁵ · Paulo Santos Pompeu³⁶ · Thiago Costa Gonçalves Portelinha³⁷ · Edson Eyji Sano³⁸ · Vagner Leonardo Macedo dos Santos³⁹ · Paloma Helena Fernandes Shimabukuro⁴ · Idelina Gomes da Silva⁴⁰ · Lucas Barbosa e Souza⁴¹ · Francisco Leonardo Tejerina-Garro^{42,43} · Mariana Pires de Campos Telles^{44,45} · Fabrício Barreto Teresa¹² · Sidinei Magela Thomaz⁴⁶ · Livia Helena Tonella⁴⁷ · Ludgero Cardoso Galli Vieira⁴⁸ · Jean Ricardo Simões Vitule⁴⁹ · Jansen Zuanon⁵⁰

Received: 21 June 2021 / Accepted: 21 July 2021 / Published online: 2 August 2021

© The Author(s), under exclusive licence to Springer Science+Business Media, LLC, part of Springer Nature 2021

Abstract

The Tocantins-Araguaia Basin is one of the largest river systems in South America, located entirely within Brazilian territory. In the last decades, capital-concentrating activities such as agribusiness, mining, and hydropower promoted extensive changes in land cover, hydrology, and environmental conditions. These changes are jeopardizing the basin's biodiversity and ecosystem services. Threats are escalating as poor environmental policies continue to be formulated, such as environmentally unsustainable hydropower plants, large-scale agriculture for commodity production, and aquaculture with non-native fish. If the current model persists, it will deepen the environmental crisis in the basin, compromising broad conservation goals and social development in the long term. Better policies will require thought and planning to minimize growing threats and ensure the basin's sustainability for future generations.

Keywords Biodiversity · Conservation · Policy · South America · Sustainability.

Introduction

Among the largest river systems in South America, the Tocantins-Araguaia basin stands out due to its large drainage area (767,000 km²) and spectacular biodiversity (Ribeiro et al. 1995; Lucinda et al. 2007). Located entirely

within Brazil, this basin drains a vast area of the Cerrado savanna and rainforest ecosystems. Two main rivers form the basin: the Tocantins, characterized by a unique ichthyofauna with several exclusive species, and the Araguaia, with one of the largest and most biodiverse floodplains in the world (Latrubesse et al. 2019). The Araguaia River is also one of the few large free-flowing rivers in South America, which harbors essential areas for biodiversity conservation (Latrubesse et al. 2019; Martins et al. 2021). The unique biodiversity of the Tocantins-Araguaia Basin, however, contrasts with a long history of policies and initiatives that have induced extensive environmental

✉ Fernando Mayer Pelicice
fmpelicice@gmail.com

Extended author information available on the last page of the article.

degradation. Over the past 40 years, the expansion of dams, croplands, irrigation, mining, and aquaculture induced extensive changes to land cover, hydrology, and environmental conditions, jeopardizing biodiversity, ecosystems, and associated services (Coe et al. 2017; Strassburg et al. 2017; Latrubesse et al. 2019).

Large-scale Degradation

Currently, seven large hydroelectric dams regulate the main channel of the Tocantins River, with many other dams blocking the course of tributaries (Winemiller et al. 2016; Akama 2017). There are plans to build new dams (ca. 90), mainly in the Araguaia basin (Latrubesse et al. 2019), in addition to waterways and a large-scale water diversion system between the Tocantins and São Francisco basins (Daga et al. 2020). The rapid expansion of agribusiness for soy and other commodities has caused the loss of ~50% of the Cerrado (Scaramuzza et al. 2017), and the activity has advanced progressively over remnant natural areas (Trigueiro et al. 2020). In 2019, pastures and monocultures covered more than 42% of the basin (Fig. 1). The Tocantins-Araguaia basin is currently the most targeted area for expanding agricultural activities, as stated in the Presidential Decree 8447 of 2015, which created the MATOPIBA Federal Plan for Development of the Brazilian Cerrado in the states of Maranhão, Tocantins, Piauí, and Bahia. In the lower portion of the basin, cattle ranching has led to the clearing of extensive rainforest areas (Nepstad et al. 2014). Broad changes in land use have eliminated riparian forests (Swanson and Bohlman 2021) and altered hydrological dynamics (Coe et al. 2009). Moreover, agriculture expansion has dramatically increased the use of pesticides (Martinelli et al. 2010), which invariably end up in aquatic ecosystems. The increasing demand for irrigation has affected regional water supplies (Morais et al. 2014), leading to growing water conflicts. Other activities generally overshadow mining, but its impacts are severe. The basin hosts the two largest iron ore deposits globally, the Carajás mine in Pará State (with 17 billion tons) and the Serra do Carmo iron deposit in Tocantins State (with 159 billion tons). Future exploitation of these deposits is a significant threat, as shown by the recent environmental disasters in southeastern Brazil (Salvador et al. 2020). Terrestrial and aquatic biodiversity has responded negatively to these impacts, as evidenced by major losses in fish diversity (e.g., Araújo et al. 2013; Perônico et al. 2020; Pereira et al. 2021; Santana et al. 2021). Fish constitute a megadiverse and endemic group in the Tocantins-Araguaia basin with over 700 species (Dagosta and de Pinna 2019). According to ICMBio (2018), this

basin has 51 threatened fish species, most of them endemic.

New Threats

Threats are escalating as public policies continue to prioritize maximizing economic growth at the cost of environmental sustainability (Pelicice et al. 2017). The recent expansion of aquaculture is exemplary. Historically, aquaculture in Brazil has been based on low standards of socio-environmental sustainability (Agostinho et al. 2007; Lima Junior et al. 2018), involving non-native organisms subjected to poor confinement conditions and inadequate management practices. Government policies have encouraged the expansion of aquaculture in the Tocantins-Araguaia basin, particularly in hydroelectric reservoirs. Many aquaculture parks have been planned (Lima et al. 2018) and, in 2021, cage aquaculture was authorized in the area impounded by the Peixe Angical Dam (<https://www.gov.br/pt-br/noticias/agricultura-e-pecuaria/2021/04/liberada-aquicultura-em-hidreletrica-no-rio-tocantins>). The authorization is valid for 35 years, and it is the first to be issued in the country after updates in legislation (Federal Decree 10.576 of 2020) that reassigned responsibilities and facilitated the authorization process. This grant was issued shortly after the permission for the farming of the Nile tilapia (*Oreochromis niloticus*), an invasive species, in aquaculture cages in Tocantins State (<https://seagro.to.gov.br/noticia/2018/12/6/criacao-de-tilapia-em-tanque-rede-e-liberada-no-tocantins/>). At the same time, State Decree 337 of 2019 allowed farming non-native fishes (i.e., grass carp *Ctenopharyngodon idella* and Nile tilapia) in the Araguaia basin and other drainage basins in Mato Grosso State. In Tocantins State, a bill is moving towards a vote that would allow the cultivation of the striped catfish (*Pangasianodon hypophthalmus*) (<https://al.to.leg.br/noticia/gabinete/jorge-frederico/9889/peixe-panga-jorge-frederico-a-presenta-projeto-de-lei-que-autoriza-o-cultivo-no-tocantins>), an Asian fish with high invasive potential that is already cultivated in other Brazilian states (Garcia et al. 2018). These initiatives must accelerate the expansion of aquaculture with non-native fish in the Tocantins-Araguaia Basin.

This episode illustrates how government policies, guided by short-term economic profits, ignore scientific advice and the costs of environmental degradation. Non-native organisms are incipient in aquatic ecosystems in this basin (Doria et al. 2021; Santana et al. 2021), and aquaculture expansion will change this scenario, triggering species invasions into other Amazonian drainage basins. The literature on socio-environmental impacts resulting from aquaculture is vast (Agostinho et al. 2007; Diana 2009; Vitule et al. 2009). It includes the unavoidable escapes and invasions (e.g., Azevedo-Santos et al. 2011; Forneck et al. 2021), and direct




Fig. 1 Land use and land cover (LULC) map of the Tocantins-Araguaia River Basin, Brazil. The map shows the scenario in 2019, indicating the area covered by natural vegetation (forest and non-

forests), water and agro-pastoral activities (pastures and crops), and the location of small (PCH) and large (UHE) hydroelectric dams. Data source: MapBiomas (2021)

and indirect negative impacts caused by the Nile tilapia (compiled by Canonico et al. 2005; Pelicice et al. 2014), striped catfish (compiled by Garcia et al. 2018), and many

possible synergistic effects between non-native species and other anthropogenic stressors (e.g., Bezerra et al. 2019). The potential for local biodiversity to boost fish production is

enormous. Supported by science, many native fish species in the basin could provide a basis for local initiatives and markets (Pelicice et al. 2014). There is a research facility concerned with aquaculture development in the basin (Embrapa Pesca e Aquicultura, Palmas, Tocantins State), which could provide adequate expertise and innovation – but it has focused extensively on using non-native fishes.

The Need for Better Policies

For decades, Brazil has fostered policies that pursue development “at any cost”. Previous setbacks have brought severe consequences for environmental policies (De Sousa et al. 2011; Fearnside 2016; Latrubesse et al. 2017; Dobrovolski et al. 2018), and current trends have had immediate impacts on the conservation of the Amazon (Ferrante and Fearnside 2019; Pelicice and Castello 2021). This approach found fertile ground in the current political scenario, characterized by scientific denialism and widespread disregard for the environment (Thomaz et al. 2020). The Tocantins-Araguaia basin proved highly vulnerable to these trends and setbacks since it has been historically targeted for economic development (e.g., Fearnside 2001), and has experienced a rapid expansion of human activities over the last ten years. The persistence of this scenario will deepen the environmental crisis in the basin, compromising broad conservation goals (e.g., UNESCO biosphere reserves for biodiversity hotspots), national policies (e.g., the National Strategy for Exotic Species and the National Action Plan for the Conservation of Endangered Species), and international agreements (e.g., the Convention on Biological Diversity and the United Nations sustainable development goals). The costs arising from environmental degradation are enormous and irreparable, considering the remarkable biodiversity and endemism that characterize the basin. Cultural risks are no less considerable (Doria et al. 2017), threatening ancestral knowledge and the ways of life of several native and traditional groups that live in the basin (Lopes et al. 2021), in addition to archeological and paleontological heritage (Lopes et al. 2019). Prevailing policies fail to recognize that environmental degradation generates socioeconomic impacts, threatening the persistence of human activities in the long term (Burger et al. 2012). The impacts and costs include over-exploitation of hydroelectric resources, loss of freshwater, pollution, deforestation, soil degradation, and losses caused by invasive species. In a situation of high environmental degradation (e.g., river regulation, deforestation, biological invasion), restoration efforts are complex, costly, and sometimes impracticable, especially in aquatic environments, where ecological impacts are less readily perceptible (Vitule 2009). Climate change must complicate this scenario (Colli et al. 2020), with the risk of affecting current policies, intensifying environmental degradation, and compromising conservation efforts.

We recommend greater caution, responsibility, and planning when dealing with environmental issues in the Tocantins-Araguaia basin. Any development in the basin must account for its high regional biodiversity value to establish balanced policies combining economic development, environmental preservation, the best available scientific knowledge, and the interests of the different social groups involved. Protected areas are limited in number and area, especially in the Tocantins River (Azevedo-Santos et al. 2019), so development policies must incorporate some basic tenets (Azevedo-Santos et al. 2021). For example, maintenance of the connectivity and natural flow regimes of remaining free-flowing rivers (especially in the Araguaia Basin), more-stringent control of irrigation projects, climatic monitoring, preservation of riparian forests, control of deforestation in both the savanna and rainforests (especially in the headwaters), revoking the authorization of aquaculture with non-native species, and improving the inspection and control of these activities. Still necessary to achieve these goals are integrative and multidisciplinary studies to explicitly evaluate the impacts of environmental and anthropic drivers on distinct components of biodiversity and at different spatial scales, especially because basic and important knowledge gaps persist (Carvalho and Tejerina-Garro 2019; Machado et al. 2019; Colli et al. 2020). The economic importance of this basin is beyond question (i.e., production of hydroelectricity and commodities), but it does not justify the ongoing environmental degradation. The human-dominated scenario in the Tocantins-Araguaia Basin provides the ideal candidate to marry the already existing economic agenda with conservation of remaining ecosystems in ways that can serve as a model for replication elsewhere. It would be feasible through the involvement and engagement of a wide base of stakeholders, including local and indigenous peoples, farmers and ranchers, the hydro-power industry, academia, the governmental and non-governmental sectors, and maybe the international community. At this moment, economic development and biodiversity preservation require much better planning to minimize escalating conflicts and ensure the sustainability of resources for coming generations.

Acknowledgements The Programa de Pós-Graduação em Biodiversidade, Ecologia e Conservação (PPGBec) at the Universidade Federal do Tocantins (UFT) supported this initiative. Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) provided research grants.

Author Contributions FMP conceived the idea and wrote the first draft. All authors contributed with writing, development, and revision.

Funding Some authors have been supported by different agencies, especially Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (Capes).

Compliance with Ethical Standards

Conflict of interest The authors declare no competing interests.

Consent to participate All authors consent with the content of the paper.

Consent for publication All authors consent with the publication of the paper.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

References

- Agostinho AA, Gomes LC, Pelicice FM (2007) Ecologia e manejo de recursos pesqueiros em reservatórios do Brasil. Eduem, Maringá, p 501
- Akama A (2017) Impacts of the hydroelectric power generation over the fish fauna of the Tocantins river, Brazil: Marabá Dam, the final blow. *Oecologia Aust* 21(3):222–231. <https://doi.org/10.4257/oeco.2017.2103.01>
- Araújo ES, Marques EE, Freitas IS, Neuberger AL, Fernandes R, Pelicice FM (2013) Changes in distance decay relationships after river regulation: similarity among fish assemblages in a large Amazonian river. *Ecol Freshw Fish* 22:543–552
- Azevedo-Santos VM, Rigolin-Sá O, Pelicice FM (2011) Growing, losing or introducing? Cage aquaculture as a vector for the introduction of non-native fish in Furnas Reservoir, Minas Gerais, Brazil. *Neotrop Ichthyol* 9:915–919
- Azevedo-Santos VM, Frederico RG, Fagundes CK, Pompeu PS, Pelicice FM, Padial AA, Nogueira MG, Fearnside PM, Lima LB, Daga VS, Oliveira FJM, Vitule JRS, Callisto M, Agostinho AA, Esteves FA, Lima-Junior DP, Magalhães ALB, Sabino J, Mormul RP, Grasel D, Zuanon J, Vilella FS, Henry R (2019) Protected areas: a focus on Brazilian freshwater biodiversity. *Divers Distrib* 25:442–448. <https://doi.org/10.1111/ddi.12871>
- Azevedo-Santos VM, Rodrigues-Filho JL, Fearnside PM, Lovejoy TE, Brito MFG (2021) Conservation of Brazilian freshwater biodiversity: Thinking about the next 10 years and beyond. *Biodivers Conserv* 30:235–241. <https://doi.org/10.1007/s10531-020-02076-5>
- Bezerra LAV, Freitas MO, Daga VS, Occhi TVT, Faria L, Costa APL, Padial AA, Prodócimo V, Vitule JRS (2019) A network meta-analysis of threats to South American fish biodiversity. *Fish Fish* 20:620–639. <https://doi.org/10.1111/faf.12365>
- Burger JR, Allen CD, Brown JH, Burnside WR, Davidson AD, Fristoe TS, Hamilton MJ, Mercado-Silva N, Nekola JC, Okie JG, Zuo W (2012) The macroecology of sustainability. *PLoS Biol* 10(6): e1001345. <https://doi.org/10.1371/journal.pbio.1001345>
- Canonico GC, Arthington A, McCrary JK, Thieme ML (2005) The effects of introduced tilapias on native biodiversity. *Aquat Conserv* 15:463–483. <https://doi.org/10.1002/aqc.699>
- Carolina RdC, Doria E, Agudelo A, Akama B, Barros M, Bonfim L, Carneiro SR, Briglia-Ferreira L, Nobre CCA, Bonilla-Castillo P, Charvet DTB, dosSantos CHP, da Silva CR, Garcia-Dávila HDB, dos AF, Duponchelle A, Encalada I, Fernandes AC, Florentino PCP, Guarido TL, de Oliveira G, Jimenez-Segura OM, Lasso-Alcalá MR, Macean EE, Marques RNG, Mendes-Júnior G, Miranda-Chumacero JLS, Nunes TVT, Occhi LS, Pereira W, Castro-Pulido L, Soares RGC, Sousa G, Torrente-Vilara PA, Van DJ, Zuanon JR Simões, Vitule (2021) The Silent Threat of Non-native Fish in the Amazon: ANNF Database and Review. *Front in Eco and Evol* 9. <https://doi.org/10.3389/fevo.2021.646702>
- Carvalho RA, Tejerina-Garro FL (2019) Spatial and temporal trends in freshwater fish research: the case of three neotropical river basins from Goiás State, Central Brazil. *Acta Limnol Bras* 31:e25. <https://doi.org/10.1590/S2179-975X5418>
- Coe MT, Brando PM, Deegan LA, Macedo MN, Neill C, Silvério DV (2017) The forests of the Amazon and Cerrado moderate regional climate and are the key to the future. *Trop Conserv Sci* 10:1–6. <https://doi.org/10.1177/1940082917720671>
- Coe MT, Costa MH, Soares-Filho BS (2009) The influence of historical and potential future deforestation on the stream flow of the Amazon River-Land surface processes and atmospheric feedbacks. *J Hydrol* 369:165–174. <https://doi.org/10.1016/j.jhydrol.2009.02.043>
- Colli GR, Vieira CR, Dianese JC (2020) Biodiversity and conservation of the Cerrado: recent advances and old challenges. *Biodivers Conserv* 29:1465–1475. <https://doi.org/10.1007/s10531-020-01967-x>
- Daga VS, Azevedo-Santos VM, Pelicice FM, Fearnside PM, Perbiche-Neves G, Paschoal LRP, Cavallari DC, Erickson J, Ruocco AMC, Oliveira I, Padial AA, Vitule JRS (2020) Water diversion in Brazil threatens biodiversity. *Ambio* 49:165–172. <https://doi.org/10.1007/s13280-019-01189-8>
- Dagosta FCP, de Pinna M (2019) The fishes of the Amazon: distribution and biogeographical patterns, with a comprehensive list of species. *Bull Am Mus Nat Hist* 431:1–163. <https://doi.org/10.1206/0003-0090.431>
- De Sousa PT, Piedade MTF, Candotti E (2011) Ecological oversight: Brazil's forest code puts wetlands at risk. *Nature* 478:458. <https://doi.org/10.1038/478458b>
- Diana JS (2009) Aquaculture production and biodiversity conservation. *BioScience* 59:27–38. <https://doi.org/10.1525/bio.2009.59.1.7>
- Dobrovolski R, Loyola R, Rattis L, Gouveia SF, Cardoso D, Santos-Silva R, Gonçalves-Souza D, Bini LM, Diniz-Filho JAF (2018) Science and democracy must orientate Brazil's path to sustainability. *Perspect Ecol Conserv* 16:121–124. <https://doi.org/10.1016/j.pecon.2018.06.005>
- Doria CRC, Athayde S, Marques EE, Lima MAL, Dutka-Gianelli J, Ruffino ML, Kaplan D, Freitas CEC, IsaacVN (2017) The invisibility of fisheries in the process of hydropower development across the Amazon. *Ambio* 47(4):453–465. <https://doi.org/10.1007/s13280-017-0994-7>
- Fearnside PM (2001) Environmental impacts of Brazil's Tucuruí Dam: Unlearned lessons for hydroelectric development in Amazonia. *Environ Manag* 27(3):377–396. <https://doi.org/10.1007/s002670010156>
- Fearnside PM (2016) Brazilian politics threaten environmental policies. *Science* 353:746–748. <https://doi.org/10.1126/science.aag0254>
- Ferrante L, Fearnside FM (2019) Brazil's new president and 'ruralists' threaten Amazonia's environment, traditional peoples and the global climate. *Environ Conserv* 46:261–663. <https://doi.org/10.1017/S0376892919000213>
- Forneck SC, Dutra FM, Camargo MP, Vitule JRS, Cunico AM (2021) Aquaculture facilities drive the introduction and establishment of non-native *Oreochromis niloticus* populations in Neotropical streams. *Hydrobiologia* 848:1955–1966. <https://doi.org/10.1007/s10750-020-04430-8>
- Garcia DAZ, Magalhães ALB, Vitule JRS, Casimiro ACR, Lima-Junior DP, Cunico AM, Brito MFG, Petreire-Junior M, Agostinho AA, Orsi ML (2018) The same old mistakes in aquaculture: the newlyavailable striped catfish *Pangasianodon hypophthalmus* is on its way to putting Brazilian freshwater ecosystems at risk. *Biodivers Conserv* 27:3545–3558. <https://doi.org/10.1007/s10531-018-1603-1>
- ICMBio (Instituto Chico Mendes de Conservação da Biodiversidade) (2018) Livro Vermelho da Fauna Brasileira Ameaçada de Extinção: Volume VI - Peixes. ICMBio, Brasília, DF, Brazil
- Latrubesse E, Arima E, Dunne T, Park E, Baker V, Horta F, Wight C, Wittmann F, Zuanon J, Baker P, Ribas C, Norgaard R, Filizola N,

- Ansar A, Flyvbjerg B, Stevaux J (2017) Damming the rivers of the Amazon basin. *Nature* 546:363–369. <https://doi.org/10.1038/nature22333>
- Latrubesse EM, Arima E, Ferreira ME, Nogueira SH, Wittmann F, Dias MS, Dagosta FCP, Bayer M (2019) Fostering water resource governance and conservation in the Brazilian Cerrado biome. *Conserv Sci Pract* 1:e77. <https://doi.org/10.1111/csp2.77>
- Lima LB, Oliveira FJ, Giacomini HC, Lima Jr DP (2018) Expansion of aquaculture parks and the increasing risk of non-native species invasions in Brazil. *Rev Aquacult* 10:111–122. <https://doi.org/10.1111/raq.12150>
- Lima Jr DP, Magalhães ALB, Pelicice FM, Vitule JRS, Azevedo-Santos VM, Orsi ML, Simberloff D, Agostinho AA (2018) Aquaculture expansion in Brazilian freshwaters against the Aichi biodiversity targets. *Ambio* 47:427–440. <https://doi.org/10.1007/s13280-017-1001-z>
- Lopes GR, Lima MGB, Reis TNP (2021) Maldevelopment revisited: inclusiveness and social impacts of soy bean expansion over Brazil's Cerrado in Matopiba. *World Dev* 139:105316. <https://doi.org/10.1016/j.worlddev.2020.105316>
- Lopes RF, Candeiro CRA, de Valais S (2019) Geoconservation of the paleontological heritage of the geosite of dinosaur footprints (sauropods) in the locality of São Domingos, municipality of Itaguatins, state of Tocantins, Brazil. *Environ Earth Sci* 78:707. <https://doi.org/10.1007/s12665-019-8722-1>
- Lucinda PHF, Freitas IS, Soares AB, Marques EE, Agostinho CS, Oliveira RJ (2007) Fish, Lajeado Reservoir, rio Tocantins drainage, State of Tocantins, Brazil. *Check List* 3:70–83. <https://doi.org/10.15560/3.2.70>
- Machado KB, Targueta CP, Antunes AM, Soares TN, Telles MPC, Logares R, Vieira LCG, Huszar VLM, Nabout JC (2019) Diversity patterns of planktonic microeukaryote communities in tropical floodplain lakes based on 18S rDNA gene sequences. *J Plankton Res* 41(3):241–256. <https://doi.org/10.1093/plankt/fbz019>
- MapBiomas, 2021. Collection 5 of Brazilian Land Cover & Use Map Series. <https://mapbiomas.org/download> (Accessed March 2021).
- Martinelli LA, Naylor R, Vitousek PM, Moutinho P (2010) Agriculture in Brazil: impacts, costs, and opportunities for a sustainable future. *Curr Opin Environ Sustain* 2:431–438. <https://doi.org/10.1016/j.cosust.2010.09.008>
- Martins PR, Sano EE, Martins ES, Vieira LCG, Salemi LF, Vasconcelos V, Couto Júnior AF (2021) Terrain units, land use and land cover, and gross primary productivity of the largest fluvial basin in the Brazilian Amazonia/Cerrado ecotone: the Araguaia River basin. *Appl Geogr* 127:102379. <https://doi.org/10.1016/j.apgeog.2020.102379>
- Morais PB, Noleto Jr S, Martins ICM (2014) Análise de sustentabilidade do projeto hidroagrícola Javaés/LAGOÁ, no estado do Tocantins. *Cad de Ciência Tecnologia* 34(1):83–111. <https://doi.org/10.35977/0104-1096.cct2017.v34.26297>
- Nepstad D, Mcgrath D, Stickler C, Alencar A, Azevedo A, Swette B, Bezerra T, DiGiano M, Shimada J, Motta RS, Armijo E, Castello L, Brando P, Hansen MC, MsGrath-Horn M, Carvalho O, Hess L (2014) Slowing Amazon deforestation through public policy and interventions in beef and soy supply chains. *Science* 344:1118–1123. <https://doi.org/10.1126/science.1248525>
- Pelicice FM, Castello L (2021). A political tsunami hits Amazon conservation. *Aquatic Conserv Mar Freshw Ecosyst*. 31:1221–1229. <https://doi.org/10.1002/aqc.3565>
- Pelicice FM, Azevedo-Santos VM, Vitule JRS, Orsi ML, Lima Jr DP, Magalhães ALB, Pompeu PS, Petreire Jr M, Agostinho AA (2017) Neotropical freshwater fishes imperiled by unsustainable policies. *Fish Fish* 18:1119–1133. <https://doi.org/10.1111/faf.12228>
- Pelicice FM, Vitule JRS, Lima Jr DP, Orsi ML, Agostinho AA (2014) A serious new threat to Brazilian freshwater ecosystems: the naturalization of nonnative fish by decree. *Conserv Lett* 7:55–60. <https://doi.org/10.1111/conl.12029>
- Pereira HR, Gomes LR, Soares PT, Martins PR, Pelicice FM, Teresa FB, Vieira LCG (2021) Long-term responses of fish diversity to river regulation: a multi-metric approach. *Environ Biol Fish* 104:71–84. <https://doi.org/10.1007/s10641-021-01056-0>
- Perônico PB, Agostinho CS, Fernandes R, Pelicice FM (2020) Community reassembly after river regulation: rapid loss of fish diversity and the emergence of a new state. *Hydrobiologia* 847(2):519–533. <https://doi.org/10.1007/s10750-019-04117-9>
- Ribeiro MCLB, Petreire Jr M, Juras AA (1995) Ecological integrity and fisheries ecology of the Araguaia—Tocantins River basin, Brazil. *Regul Rivers: Res Manag* 11(3–4):325–350. <https://doi.org/10.1002/rrr.3450110308>
- Salvador GN, Leal CG, Brejão GL, Pessali TC, Alves CBM, Rosa GR, Ligeiro R, Montag LFA (2020) Mining activity in Brazil and negligence in action. *Perspect Ecol Conserv* 18:139–144. <https://doi.org/10.1016/j.pecon.2020.05.003>
- Santana MLC, Carvalho FR, Teresa FB (2021) Broad and fine-scale threats on threatened Brazilian freshwater fish: variability across hydrographic regions and taxonomic groups. *Biota Neotropica* 21(2):e20200980. <https://doi.org/10.1590/1676-0611-BN-2020-0980>
- Scaramuzza CAM, Sano EE, Adami M, Bolfe EL, Coutinho AC (2017) Land-use and Land-cover mapping of the Brazilian Cerrado based mainly on Landsat-8 satellite images. *Rev Bras Cartografia* 69(6):1041–1051. 2017
- Strassburg BBN, Brooks T, Feltran-Barbieri R, Iribarrem A, Crouzeilles R, Loyola R, Latawiec AE, Oliveira FJB, Scaramuzza CAD, Scarano FR, Soares B, Balmford A (2017) Moment of truth for the Cerrado hotspot. *Nat Ecol Evol* 1:3. 10.1038/s41559-017-0099
- Swanson AC, Bohlman S (2021) Cumulative impacts of land cover change and dams on the land–water interface of the Tocantins River. *Front Environ Sci* 9:662904. <https://doi.org/10.3389/fenvs.2021.662904>
- Thomaz SM, Gomes Barbosa L, de Souza Duarte MC, Panosso R (2020) The future of nature conservation in Brazil. *Inland Waters* 10:295–303. <https://doi.org/10.1080/20442041.2020.1750255>
- Trigueiro WR, Nabout JC, Tessarolo G (2020) Uncovering the spatial variability of recent deforestation drivers in the Brazilian Cerrado. *J Environ Manag* 275:111243. <https://doi.org/10.1016/j.jenvman.2020.111243>
- Vitule JRS (2009) Introdução de peixes em ecossistemas continentais brasileiros: revisão, comentários e sugestões de ações contra o inimigo quase invisível. *Neotrop Biol Conserv* 4:111–122. <https://doi.org/10.4013/5123>
- Vitule JRS, Freire CA, Simberloff D (2009) Introduction of non-native freshwater fish can certainly be bad. *Fish Fish* 10:98–108. <https://doi.org/10.1111/j.1467-2979.2008.00312.x>
- Winemiller KO, McIntyre PB, Castello L, Fluet-Chouinard E, Giarrizzo T, Nam S, Baird IG, Darwall W, Lujan NK, Harrison I, Stiassny MLJ, Silvano RAM, Fitzgerald DB, Pelicice FM, Agostinho AA, Gomes LC, Albert JS, Baran E, Petreire Jr M, Zarfl C, Mulligan M, Sullivan JP, Arantes CC, Sousa LM, Koning AA, Hoenighaus DJ, Sabaj M, Lundberg JG, Armbruster J, Thieme ML, Petry P, Zuanon J, Torrente Vilara G, Snoeks J, Ou C, Rainboth W, Pavanelli CS, Akama A, van Soesbergen A, Sáenz L (2016) Balancing hydropower and biodiversity in the Amazon, Congo, and Mekong. *Science* 351:128–129. <https://doi.org/10.1126/science.aac7082>

Affiliations

Fernando Mayer Pelicice ¹ · Angelo Antonio Agostinho² · Alberto Akama³ · José Dilermando Andrade Filho⁴ · Valter M. Azevedo-Santos⁵ · Marcus Vinicius Moreira Barbosa⁶ · Luis Mauricio Bini⁷ · Marcelo Fulgêncio Guedes Brito⁸ · Carlos Roberto dos Anjos Candeiro⁹ · Érica Pellegrini Caramaschi¹⁰ · Priscilla Carvalho¹¹ · Rodrigo Assis de Carvalho¹² · Leandro Castello¹³ · Davi Borges das Chagas¹⁴ · Carine Cavalcante Chamon¹⁵ · Guarino Rinaldi Colli¹⁶ · Vanessa Salette Daga¹⁷ · Murilo Sversut Dias¹⁸ · José Alexandre Felizola Diniz Filho⁷ · Philip Fearnside¹⁹ · Wagner de Melo Ferreira¹ · Diego Azevedo Zoccal Garcia²⁰ · Tiago Kutter Krolow²¹ · Rodrigo Ferreira Kruger²² · Edgardo Manuel Latrubesse²³ · Dilermando Pereira Lima Junior²⁴ · Solange de Fátima Lolis¹ · Fabyano Alvares Cardoso Lopes²⁵ · Rafael Dias Loyola²⁶ · André Lincoln Barroso Magalhães²⁷ · Adriana Malvasio²⁸ · Paulo De Marco Jr.⁷ · Pedro Ribeiro Martins²⁹ · Rosana Mazzoni³⁰ · João Carlos Nabout¹² · Mário Luis Orsi²⁰ · Andre Andrian Padial³¹ · Hasley Rodrigo Pereira³² · Thiago Nilton Alves Pereira¹⁵ · Phamela Bernardes Perônico¹² · Miguel Petrere Jr.³³ · Renato Torres Pinheiro³⁴ · Etiene Fabbrin Pires³⁵ · Paulo Santos Pompeu³⁶ · Thiago Costa Gonçalves Portelinha³⁷ · Edson Eyji Sano³⁸ · Vagner Leonardo Macedo dos Santos³⁹ · Paloma Helena Fernandes Shimabukuro⁴ · Idelina Gomes da Silva⁴⁰ · Lucas Barbosa e Souza⁴¹ · Francisco Leonardo Tejerina-Garro^{42,43} · Mariana Pires de Campos Telles^{44,45} · Fabrício Barreto Teresa¹² · Sidinei Magela Thomaz⁴⁶ · Livia Helena Tonella⁴⁷ · Ludgero Cardoso Galli Vieira⁴⁸ · Jean Ricardo Simões Vitule⁴⁹ · Jansen Zuanon⁵⁰

¹ Núcleo de Estudos Ambientais, Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil

² Programa de Pós Graduação em Ecologia de Ambientes Aquáticos Continentais (PEA), Universidade Estadual de Maringá (UEM), Maringá, Brazil

³ Museu Paraense Emílio Goeldi, Belém, Brazil

⁴ Grupo de Estudos em Leishmanioses, Instituto René Rachou, Fundação Oswaldo Cruz (FIOCRUZ), Belo Horizonte, Brazil

⁵ Universidade Estadual Paulista “Júlio de Mesquita Filho”, Botucatu, Brazil

⁶ Museu de Zoologia e Taxidermia José Hidasi, Universidade Estadual do Tocantins (Unitins), Porto Nacional, Brazil

⁷ Departamento de Ecologia, Universidade Federal de Goiás (UFG), Goiânia, Brazil

⁸ Laboratório de Ictiologia, Departamento de Biologia, Universidade Federal de Sergipe (UFS), São Cristóvão, Brazil

⁹ Laboratório de Paleontologia e Evolução, Curso de Geologia, Universidade Federal de Goiás (UFG), Aparecida de Goiânia, Brazil

¹⁰ Departamento de Ecologia, Universidade Federal do Rio de Janeiro (UFRJ), Rio de Janeiro, Brazil

¹¹ Universidade Federal de Goiás (UFG), Goiânia, Brazil

¹² Laboratório de Biogeografia e Ecologia Aquática, Universidade Estadual de Goiás, Anápolis, Brazil

¹³ Department of Fish and Wildlife Conservation, Virginia Polytechnic Institute and State University, Virginia, USA

¹⁴ Herbário do Tocantins, Núcleo de Estudos Ambientais, Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil

¹⁵ Laboratório de Ictiologia Sistemática, Núcleo de Estudos Ambientais, Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil

¹⁶ Departamento de Zoologia, Universidade de Brasília, Brasília-DF 70910-900, Brazil

¹⁷ Researcher, Tampa, FL, USA

¹⁸ Departamento de Ecologia, Universidade de Brasília (UnB), Brasília-DF, Brazil

¹⁹ Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil

²⁰ Laboratório de Ecologia de Peixes e Invasões Biológicas, Universidade Estadual de Londrina (UEL), Londrina, Brazil

²¹ Laboratório de Entomologia, Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil

²² Universidade Federal de Pelotas (UFPeL), Pelotas, Brazil

²³ Environmental Sciences Program-CIAMB, Universidade Federal de Goiás (UFG), Goiânia, Brazil

²⁴ Universidade Federal de Mato Grosso (UFMT), Campus Universitário do Araguaia, Pontal do Araguaia, Brazil

²⁵ Laboratório de Microbiologia Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil

²⁶ Fundação Brasileira para o Desenvolvimento Sustentável (FBDS) & Universidade Federal de Goiás (UFG), Goiânia, Brazil

²⁷ Researcher, Belo Horizonte, Brazil

²⁸ Laboratório de Ecologia e Zoologia (LABECZ), Universidade Federal do Tocantins (UFT), Palmas, Brazil

²⁹ Researcher, Brasília, Brazil

³⁰ Laboratório de Ecologia de Peixes, Departamento de Ecologia, Universidade do Estado do Rio de Janeiro (UERJ), Rio de Janeiro, Brazil

- 31 Laboratório de Análise e Síntese em Biodiversidade, Departamento de Botânica, Universidade Federal do Paraná (UFPR), Curitiba, Brazil
- 32 Secretaria de Estado da Educação de Goiás-(Seduc-GO), Luziânia, Goiás, Brazil
- 33 UNISANTA, PPGEOMAR, Santos, Brazil
- 34 Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil
- 35 Laboratório de Paleobiologia, Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil
- 36 Departamento de Ecologia e Conservação, Universidade Federal de Lavras (UFLA), Lavras, Brazil
- 37 Laboratório de Caracterização de Impactos Ambientais (LCIA), Universidade Federal do Tocantins (UFT), Palmas, Brazil
- 38 Embrapa Cerrado, Planaltina, Brazil
- 39 Instituto de Biodiversidade e Sustentabilidade, Universidade Federal do Rio de Janeiro, Macaé, Brazil
- 40 Programa de Pós-Graduação em Ecologia, Universidade Federal do Pará (UFPA), Belém, Brazil
- 41 Laboratório de Análises Geoambientais (LGA), Universidade Federal do Tocantins (UFT), Porto Nacional, Brazil
- 42 Centro de Biologia Aquática, Pontifícia Universidade Católica de Goiás, Goiânia, Brazil
- 43 Laboratório de Biodiversidade, Universidade Evangélica de Goiás, Anápolis, Brazil
- 44 Escola de Ciências Agrárias e Biológicas, Pontifícia Universidade Católica (PUC), Goiânia, Brazil
- 45 Laboratório de Genética & Biodiversidade - ICB/UFG, Goiânia, Brazil
- 46 Núcleo de Pesquisas em Limnologia, Ictiologia e Aquicultura (Nupelia), Universidade Estadual de Maringá (UEM), Maringá, Brazil
- 47 Departamento de Direito, Universidade Federal do Tocantins (UFT), Palmas, Brazil
- 48 Núcleo de Estudos e Pesquisas Ambientais e Limnológicas (Nepal), Universidade de Brasília (UnB), Planaltina-DF, Brazil
- 49 Laboratório de Ecologia e Conservação (LEC), Universidade Federal do Paraná (UFPR), Curitiba, Brazil
- 50 Coordenação de Biodiversidade, Instituto Nacional de Pesquisas da Amazônia (INPA), Manaus, Brazil