











Rainfall variability affects the chemical composition, gas production and degradability of cacti

Variabilidad de la precipitación afecta la composición química, producción de gas y degradabilidad de los cactus

Daniel Ribeiro Menezes¹ ; Juliana Muniz dos Santos¹ ; Cintia Raquel Nunes de Oliveira² ; Polyana Deyse Rodrigues Marcelino² ; Anna Paula da Mata Araújo Pinheiro¹ ; Dalinne Tamara Queiroz de Carvalho² ; Júlio Cesar Silva Nascimento¹ ; Glayciane Costa Gois^{1*} ; Tadeu Vinhas Voltolini³ ; Mario Adriano Ávila Queiroz¹ .

¹Universidade Federal do Vale do São Francisco, Campus de Ciências Agrárias, Petrolina/PE, Brazil.

²Universidade Federal da Bahia, Escola de Medicina Veterinária e Zootecnia, Salvador/BA, Brazil.

³Empresa Brasileira de Pesquisa Agropecuária, Embrapa Semiárido, Petrolina/PE, Brazil.

*Correspondence: glayciane_gois@yahoo.com.br

Recepción: diciembre 2021 | Aprobación: febrero 2022 | Publicación: febrero 2022

ABSTRACT

Objective. The objective was to evaluate the effect of rainfall variability on the chemical composition, *in vitro* degradability and gas production of cacti in Semiarid region. **Materials and methods.** A completely randomized experimental design in a factorial arrangement of 2 rainfall periods (high and low rainfall) x 5 cactus species, with 5 replications was adopted. Cacti evaluated were *Pilosocereus gounellei* (Weber ex K. Schum). Bly ex Rowl, *Cereus jamacaru* DC., *Opuntia ficus indica* Mill, *Nopalea cochenillifera* Salm. Dyck and *Opuntia stricta* (Haw.) Haw, all fresh. **Results.** There was an interaction between rainfall variation and cactus species for dry matter, crude protein, neutral detergent fiber and acid detergent fiber ($p < 0.05$), *in vitro* gas production ($p < 0.05$) and for *in vitro* dry matter degradability ($p < 0.05$). Among cacti, *Pilosocereus gounellei* presented a lower proportion of organic matter and an increase in the content of mineral matter in relation to the other cacti ($p < 0.05$). **Conclusions.** The chemical composition, the gas production coefficients and the *in vitro* degradability were influenced by the cacti species and by the rainfall variation. Cacti species increase the dry matter, crude protein and neutral detergent fiber contents during the low rainfall. Varieties of cactus pear (*Opuntia* and *Nopalea*) stood out among the cacti tested in relation to gas production and *in vitro* degradability.

Keywords: *Cactaceae*; crassulacean acid metabolism; dryland; dry matter; food shortage; neutral detergent fiber (Source: CAB).

RESUMEN

Objetivo. El objetivo fue evaluar el efecto de la variabilidad de las lluvias sobre la composición química, la degradabilidad *in vitro* y la producción de gas de cactus en la región Semiárida. **Materiales y métodos.** Este fue un diseño experimental completamente al azar en un arreglo factorial de 2 períodos de lluvia (lluvia alta y baja) x 5 especies de cactus, con 5 repeticiones. Los cactus evaluados fueron *Pilosocereus gounellei* (Weber ex K. Schum). Bly ex Rowl, *Cereus jamacaru* DC., *Opuntia ficus indica* Mill, *Nopalea cochenillifera* Salm. Dyck y *Opuntia stricta* (Haw.) Haw, todas frescas. **Resultados.** Hubo una interacción entre la variación de la lluvia y las especies de cactus para materia seca, proteína cruda, fibra detergente neutra y fibra detergente ácido ($p < 0.05$), producción de gas *in vitro* ($p < 0.05$) y para la degradabilidad *in vitro* de materia seca ($p < 0.05$). Entre los cactus, *Pilosocereus gounellei* presentó una menor proporción de materia orgánica y un aumento en el contenido de materia mineral en relación a los demás cactus ($p < 0.05$). **Conclusiones.** La composición química, los coeficientes de producción de gas y la degradabilidad *in vitro* fueron influenciados por las especies de cactus y por la variación de las precipitaciones. Las especies de cactus aumentan el contenido de materia seca, proteína cruda y fibra de detergente neutro durante la escasez de precipitaciones. Las variedades de nopal (*Opuntia* y *Nopalea*) se destacaron entre los cactus ensayados en relación a la producción de gas y la degradabilidad *in vitro*.

Palabras clave: *Cactaceae*; escasez de alimentos; fibra detergente neutra; materia seca; metabolismo del ácido crasuláceo; regiones de tierras secas (Fuente CAB).

Como citar (Vancouver).

Menezes DR, dos Santos JM, Nunes de Oliveira CR, Marcelino PDR, Pinheiro APMA, de Carvalho DTQ et al. Rainfall variability affects the chemical composition, gas production and degradability of cacti. Rev Colombiana Cienc Anim. Recia. 2022; 14(1):e895. <https://doi.org/10.24188/recia.v14.n1.2022.895>

Publicado por la Revista Colombiana de Ciencia Animal - RECIA, bajo los términos de la licencia 

ISSN: 2027-4297

INTRODUCTION

Rainfall is one of the most important drivers of ecosystem dynamics and productivity. In this context, arid and semiarid regions in the world are characterized by a variation in rainfall pulses, which are concentrated in only a few months of the year (1).

Considering the local soil and climate conditions, it can be said that the forage production in dry land regions is challenging (2). The lack of rain limits the growth of native and cultivated species, such as grasses and legumes of great water demand, causing a decrease in the biomass and the carrying capacity of pastures (3). Thus, cacti have been frequently used as one of the main forage supports for ruminants during the period of food shortage (4).

Native cacti from arid and semiarid regions such as Xique-xique (*Pilosocereus gounellei* (Weber ex K. Schum). Bly ex Rowl), Mandacaru (*Cereus jamacaru* DC.), cactus pear "Gigante" (*Opuntia ficus indica* Mill), cactus pear "Miúda" (*Nopalea cochenillifera* Salm. Dyck) and cactus pear "Mexican Elephant Ear" (*Opuntia stricta* (Haw.) Haw) have high productive potential in environments with low and erratic rainfall (3, 4). This is due to the characteristics that these cacti have, such as physiological adaptability to soil and climate conditions remaining succulent during dry periods, high digestibility of dry matter and high-water content making them an excellent dietary strategy where this nutrient is limited, in addition to being excellent sources of energy for being rich in non-fiber carbohydrates (2).

Although the Caatinga vegetation in Brazilian semiarid is driven by low rainfall with low productivity and reduced biomass, it shows large spatial and temporal variations (5). Thus, studies on the nutritional value and *in vitro* degradation kinetics of cacti in periods of low and high rainfall, are necessary and, to the best of our knowledge, studies comparing exclusively the cacti found in the semiarid regions are scarce (2), being evaluated as part of the ingredients of the diet, compromising the results of its real nutritional value.

The objective of this study was to evaluate the effect of rainfall variability on the chemical composition, *in vitro* degradability and the gas production of cacti in Semiarid region.

MATERIAL AND METHODS

Experimental site. The experiment was carried out at the premises of the Laboratory of Animal Requirement and Metabolism (LEMA), belonging to the Federal University of the São Francisco Valley (UNIVASF), Petrolina, state of Pernambuco, Brazil (9°24'38" South latitude, 40°29'56" West longitude, 377 m altitude).

The climate is hot semiarid, with rainy season (BSh), concentrated between November and April, with an average annual rainfall of 501.5 mm, irregularly distributed, average annual temperature of 26°C and average annual relative humidity of 60%.

Experimental design, samples and rainfall. This was a completely randomized experimental design in a factorial arrangement of 2 rainfall periods x 5 cactus species, with 5 replications. Samples were collected at different sites in the Caatinga experimental area, belonging to the Brazilian Agricultural Research Corporation (Embrapa Semiarid), in Petrolina, state of Pernambuco, Brazil. The collection area is maintained without irrigation and the soil is classified as plinic yellow Argisol (6).

The cactus species collected were *Pilosocereus gounellei* (Weber ex K. Schum). Bly ex Rowl (Common name: Xique-xique), *Cereus jamacaru* DC. (Common name: Mandacaru), *Opuntia ficus indica* Mill (Common name: cactus pear "Gigante"), *Nopalea cochenillifera* Salm. Dyck (Common name: Cactus pear "Miúda") and *Opuntia stricta* (Haw.) Haw (Common name: cactus pear "Mexican Elephant Ear"), all fresh. All cacti had 3 years of establishment in the field.

Collections were carried out in two periods: period with low rainfall (LR), between August and November, with total and average rainfall of 50.6 mm and 12.6 mm/month, respectively; and period with high rainfall (HR), between January and April, with total and average rainfall of 112.0 mm and 28.0 mm/month (Figure 1), respectively. Samplings were carried out in the last month of each experimental period, with analysis of 5 plants of each species of cactus.

Sample preparation. Considering animal consumption, spines of *P. gounellei* (Weber ex K. Schum). Bly ex Rowl and *C. jamacaru* DC. were removed by burning with a flamethrower. For the varieties of cactus pear (*Opuntia* and *Nopalea*), the secondary and tertiary cladodes of the plants were used (2). Samples were ground in a stationary shredder (PP-35, Pinheiro, Itapira - SP) to an average particle size of approximately 5.0 cm. After this procedure, samples were pre-dried in a forced ventilation oven at 55°C for 72-h and ground into 1 mm and 3 mm particles in a knife mill (Wiley, Marconi, MA 580, Piracicaba, Brazil) to determine the chemical composition, gas production and the *in vitro* degradability test.




Figure 1. Monthly agrometeorological data for periods of high rainfall (a) and low rainfall (b) during the experimental period (Temp – Temperature; RH – Relative Humidity; Prec – Precipitation; ETo – Reference evapotranspiration).

Chemical analysis. Analyses were performed using the methods described by AOAC (7) to determine the content of dry matter (DM; method 967.03), mineral matter (MM; method 942.05), crude protein (CP; method 981.10) and acid detergent fiber (ADF; method 973.18). The neutral detergent fiber (NDF) content was determined using the methodology described by Van Soest et al (8). Hemicellulose (HEM) was determined by the difference between NDF and ADF.

In vitro gas production. *In vitro* gas production was conducted according technique adapted by Menezes et al (9). The pressure (P; in psi) was measured by a portable pressure transducer (GE Druck Series DPI 705) and readings were taken at 2, 4, 6, 8, 9, 11, 12, 14, 17, 20, 24, 28, 34, 48, 72, 96 and 120-h after incubation. Pressure data were converted to gas volume (1 psi = 4.859 mL gas).

Cumulative gas production data were analyzed by the Gompertz two-compartmental model (10):

$$V(t) = Vf1/[1 + e^{(2 - 4m1(L - T))}] + Vf2/[1 + e^{(2 - 4m2(L - T))}]$$

where: $V(t)$ = total maximum volume of gas produced; $Vf1$ = maximum volume of gas for the fast digesting fraction (non-fiber carbohydrates; NFC); $Vf2$ = maximum volume of gas for the slow digesting fraction (fibrous carbohydrates; FC); $m1$ = specific growth rate for the rapid degradation fraction; $m2$ = specific growth rate for the slow degradation fraction; L = duration of initial digestion events (latency phase), common to both phases, and; T = fermentation time.

In vitro dry matter degradability. The *in vitro* dry matter degradability (IVDMD) was estimated from the insertion of nylon bags (20mg/cm² weight and 50 microns of porosity) containing 600 mg sample in flasks with 60 mL buffer solution (combination of solutions A + B with pH 6.8) and 15 mL inoculum collected from 2 fistulated Santa Inês sheep, kept on a diet consisting of elephant grass (*Pennisetum purpureum*), concentrate based on corn and soybean, in addition to mineral salt (Ovinofós, Tortura, Porto Alegre, Brazil). The rumen content was filtered through gauze, constantly injecting CO₂ to maintain the anaerobic environment and stored at 39°C. Samples were incubated at times 0, 2, 6, 12, 24, 48, 96 and 120-h. After *in vitro* fermentation, bags were washed and dried in an oven at 105°C for 4-h, and weighed. Samples at time 0 were just washed with distilled water at 39°C for 5 min, and then dried and weighed (11).

To determine potential degradability (PD), the Ørskov and McDonald (12) model was used:

$$PD = a + b (1 - e^{-ct})$$

where: a = water-soluble fraction; b = water-insoluble fraction, but potentially degradable; c = degradation rate of the fraction “ b ”; and t = incubation time in hours. The letter “ e ” is the natural log of “ $-ct$ ”.

The effective degradability (ED) was calculated using the formula:

$$ED = a + (b \times c) / (c + k)$$

where: k is the rate of passage. The gas production rate obtained by the semi-automated gas production technique ($m1+m2$) was used to estimate the rate of passage (k) used in the degradability test.

Statistical analysis. Data were initially analyzed using the UNIVARIATE procedure to check the normal distribution of the data. After observing normal distribution, analysis of variance was applied, using the PROC ANOVA command from the Statistical Analysis System (SAS, version 9.1). Parameters of the gas production and degradability model were estimated by the nonlinear regression procedure (NLIN) of SAS (version 9.1). The following statistical model was adopted:

$$Y_{ijk} = \mu + \alpha_i + \beta_j + \alpha\beta_{ij} + e_{ijk}$$

where Y is the observed value of the variable *ijk* that refers to the *k*-th repetition of the combination of the *i*-th level of factor A with the *j*-th level of factor B; μ is the mean of all experimental units of the variable; α_i is the cactus species effect on the observed value Y_{ij} ; β_j is the rainfall periods effects on the observed value Y_{ijk} ; $\alpha\beta_{ij}$ is the interaction effect between cactus species and rainfall periods; and e_{ijk} is the error associated with Y_{ijk} 's observation. Tukey's test was applied to compare the means, with probability values below 5% ($p < 0.05$) being considered significant.

RESULTS

There was an interaction effect of rainfall variation and cactus species for DM, CP, NDF and ADF ($p < 0.05$; Table 1). Isolated effect was observed for the OM content, in which *P. gounellei* presented the lowest levels for this variable, in relation to the other studied species ($p < 0.05$; Table 1). There was an isolated effect for the MM content, in which *P. gounellei* showed the highest levels for this variable, in relation to the other studied cacti ($p < 0.05$; Table 1).

Table 1. Chemical composition of cactus species collected in high rainfall (HR) and low rainfall (LR) periods in the Brazilian semiarid (n=5).

Period	Cactus species					SEM	p-value		
	<i>Pilosocereus gounellei</i>	<i>Cereus jamacaru</i>	<i>Opuntia ficus indica</i> Mill	<i>Nopalea cochenillifera</i> Salm. Dyck	<i>Opuntia stricta</i> Haw		C	R	CxR
<i>Dry matter (g/kg NM)</i>									
HR	62.5Bc	144.4Ba	114.4Bb	144.7Ba	144.7Aa	0.74	*	*	*
LR	190.6Ab	238.3Aa	157.5Ac	214.7Aab	147.2Ac		*	*	*
<i>Organic matter (g/kg DM)</i>									
HR	725.4Ac	839.3Ab	840.4Ab	862.4Aab	887.5Aa	0.74	*	ns	ns
LR	783.2Ac	859.3Aab	857.5Aab	857.8Aab	841.4Ab		*	ns	ns
<i>Mineral matter (g/kg DM)</i>									
HR	274.6Aa	160.7Ab	159.6Ab	137.6Ac	112.5Ad	0.65	*	ns	ns
LR	216.8Aa	140.7Ab	142.5Ab	142.2Ab	158.6Ab		*	ns	ns
<i>Crude protein (% DM)</i>									
HR	29.1Ab	34.4Ba	37.9Ba	30.8Ab	23.6Bb	0.23	*	*	*
LR	39.1Ac	53.6Ab	71.4Aa	36.8Ac	56.3Ab		*	*	*
<i>Neutral Detergent Fiber (% DM)</i>									
HR	309.2Ab	284.1Bc	281.0Bc	382.6Ba	279.4Bc	1.97	*	*	*
LR	316.7Ad	404.6Ac	462.5Ab	496.6Aa	491.0Aa		*	*	*
<i>Acid Detergent Fiber (g/kg DM)</i>									
HR	146.8Bbc	238.3Aa	171.7Ab	137.8Ac	124.5Ac	1.35	*	*	*
LR	201.8Aa	121.0Bc	157.3Ab	126.3Ac	120.7Ac		*	*	*

DM: Dry matter; C: Cactus species effect; R: Rainfall periods effects; C x P: interaction effect between cactus species and rainfall periods; SEM: Standard error of the mean; Capital letters in the column indicate the statistical effect of rainfall periods on cactus species; Lower case letters on the line indicate statistical effect on cactus species within rainfall periods; Significant at the 5% level by the Tukey test; * $p < 0.05$; ns: Not significant.

There was an interaction effect of rainfall variation and cactus species for *in vitro* gas production ($p < 0.05$; Table 2). The period of low rainfall provided a greater cumulative production of gases, in which *O. stricta* Haw showed a higher production of gases from non-fiber carbohydrates ($Vf1 = 122.16$ mL/g DM) among the other evaluated cacti.

The reduction in gas production from non-fiber carbohydrates during the period of high rainfall was more evident for *P. gounellei*, which showed a reduction of 36.86 mL/g DM during the transition between the low and high periods of rainfall (Table 2).

With respect to the rainfall periods and the studied cacti, *P. gounellei* presented the lowest values for gas production from the fiber fraction ($Vf2$; 15.31 mL/g DM in high rainfall and 18.32 mL/g DM in low rainfall) and the highest values for the rates of gas production from the fiber fraction ($m1$; 0.09 mL/g DM/h in high rainfall and 0.19 mL/g DM in low rainfall) ($p < 0.05$; Table 2).

The varieties *N. cochenillifera* and *O. stricta* Haw in the period of high rainfall showed the highest gas production rates of the non-fiber fraction (m2), with average value of 0.015 mL/g DM/h, respectively, for the two cacti ($p < 0.05$; Table 2).

Table 2. *In vitro* gas production from cactus species collected in high rainfall (HR) and low rainfall (LR) periods in the Brazilian semiarid (n=5).

Period	Cactus species					SEM	p-value		
	<i>Pilosocereus gounellei</i>	<i>Cereus jamacaru</i>	<i>Opuntia ficus indica</i> Mill	<i>Nopalea cochenillifera</i> Salm. Dyck	<i>Opuntia stricta</i> Haw		C	R	CxR
<i>Vf1 (mL/g DM)</i>									
HR	57.35Bb	85.38Aa	86.52Ba	83.00Ba	92.95Ba	1.89	*	*	*
LR	94.21Ab	87.40Ab	94.21Ab	95.97Ab	122.16Aa		*	*	*
<i>Vf2 (mL/g DM)</i>									
HR	15.31Ad	19.36Bc	73.74Aab	68.17Ab	93.59Aa	5.87	*	*	*
LR	18.32Ad	29.49Ac	61.43Ba	64.42Aa	41.21Bb		*	*	*
<i>m1 (mL/g DM/h)</i>									
HR	0.09Ba	0.03Bc	0.04Bc	0.06Ab	0.07Bb	0.01	*	*	*
LR	0.19Aa	0.05Ac	0.06Ac	0.06Ac	0.10Ab		*	*	*
<i>m2 (mL/g DM/h)</i>									
HR	0.011Bc	0.010Ac	0.013Ab	0.015Aa	0.015Ba	0.001	*	*	*
LR	0.021Aa	0.011Ac	0.015Ab	0.015Ab	0.025Aa		*	*	*
<i>Vt (mL/g DM)</i>									
HR	72.66Bd	104.74Ac	160.26Ab	151.17Ab	186.54Aa	7.42	*	*	*
LR	112.52Ac	116.89Ab	155.64Aa	160.38Aa	163.38Ba		*	*	*
<i>M (mL/g DM/h)</i>									
HR	0.09Ba	0.04Bb	0.06Bab	0.08Aa	0.09Ba	0.01	*	*	*
LR	0.21Aa	0.06Ac	0.07Ac	0.08Ac	0.12Ab		*	*	*
<i>L (hours)</i>									
HR	13.02Ab	21.53Aa	12.97Ab	7.36Bc	9.41Ab	0.72	*	*	*
LR	9.13Bb	18.74Ba	8.84Bb	8.95Ab	9.99Ab		*	*	*

Vf1: Maximum gas production potential of non-fibrous carbohydrates; Vf2: Maximum gas production potential of fibrous carbohydrates; m1: rate of non-fibrous carbohydrate gas production; m2: rate of production of fibrous carbohydrate gases; Vt: maximum gas production potential of total carbohydrates; M: total gas production rate; L: duration of initial digestion events (latency phase), common to both phases; DM: Dry matter; C: Cactus species effect; R: Rainfall periods effects; C x P: interaction effect between cactus species and rainfall periods; SEM: Standard error of the mean; Capital letters in the column indicate the statistical effect of rainfall periods on cactus species; Lower case letters on the line indicate statistical effect on cactus species within rainfall periods; Significant at the 5% level by the Tukey test; * $p < 0.05$; ns: Not significant.

O. stricta Haw in the period of high rainfall showed a higher value for the potential of total gas production (186.54 mL/g DM) in relation to the other cacti. *P. gounellei*, on the other hand, showed a reduction in the total gas production with increasing rainfall (72.66 mL/g DM; Table 2).

The latency phase (L) during the high rainfall period was reduced in *N. cochenillifera*. Regarding the period of low rainfall and cacti, *C. jamacaru* had a longer colonization time ($p < 0.05$; Table 2).

There was an interaction effect of rainfall variation and cactus species for *in vitro* dry matter degradability ($P < 0.05$; Table 3). *O. ficus indica* Mill and *O. stricta* Haw reduced the water-soluble fraction (a) and the degradation rate of fraction B (C; %/hour) during the transition from high to low rainfall ($p < 0.05$; Table 3).

P. gounellei and *N. cochenillifera* reduced the fraction insoluble in water, but potentially degradable (b) in relation to the other cacti, during the period of high rainfall ($p < 0.05$; Table 3).

Table 3. *In vitro* degradability from cactus species collected in high rainfall (HR) and low rainfall (LR) periods in the Brazilian semiarid (n=5).

Period	Cactus species					SEM	p-value		
	<i>Pilosocereus gounellei</i>	<i>Cereus jamacaru</i>	<i>Opuntia ficus indica</i> Mill	<i>Nopalea cochenillifera</i> Salm. Dyck	<i>Opuntia stricta</i> Haw		C	R	CxR
			A (%)						
HR	28.14Bab	21.72Bc	28.98Aab	27.88Bb	37.22Aa	0.91	*	*	*
LR	42.61Aa	34.67Aab	24.59Ab	42.29Aa	23.42Bb		*	*	*
			B (%)						
HR	58.67Aab	53.18Bb	67.31Aa	70.49Aa	54.05Bb	1.26	*	*	*
LR	35.16Bb	60.85Aa	71.26Aa	48.43Bb	69.01Aa		*	*	*
			C (%/hora)						
HR	0.017Bc	0.023Bb	0.043Aa	0.023Bb	0.041Aa	0.03	*	*	*
LR	0.066Ab	0.736Aa	0.020Bc	0.755Aa	0.033Ac		*	*	*
			PD (%)						
HR	86.81Abc	74.90Bc	96.29Aab	98.37Aa	91.28Ab	0.86	*	*	*
LR	77.76Bb	95.52Aa	95.85Aa	90.73Ba	92.44Aa		*	*	*
			ED (%)						
HR	37.44Bc	39.72Bb	58.24Aa	43.71Bab	54.10Aa	2.08	*	*	*
LR	51.04Ab	90.64Aa	39.69Bc	85.97Aa	37.67Bc		*	*	*

A: Water-soluble fraction; B: fraction insoluble in water, but potentially degradable; C: degradation rate of fraction "B"; PD: potential degradability; ED: effective degradability; C: Cactus species effect; R: Rainfall periods effects; C x P: interaction effect between cactus species and rainfall periods; SEM: Standard error of the mean; Capital letters in the column indicate the statistical effect of rainfall periods on cactus species; Lower case letters on the line indicate statistical effect on cactus species within rainfall periods; Significant at the 5% level by the Tukey test; * p<0.05; ns: Not significant.

DISCUSSION

Cacti are of great importance for dry land regions mainly in periods of low or zero rainfall, becoming a source of water and one of the main forages to feed herds, as they adapt to the soil and climate conditions of the region, have the capacity to store water in its structure and be persistent to drought (13). This adaptation is because these plants are classified as plants with the Crassulacean Acid Metabolism (MAC), a mechanism that gives them high efficiency in the use of water (14). This efficiency was verified in the present study, where, with the exception of *Opuntia stricta* Haw, which remained stable in the two periods evaluated, the other studied cacti increased their moisture content with the highest water supply received.

Protein and fiber contents are directly related to climatic factors, season, water availability, plant age, part of the plant collected and handling of the plant during sample preparation (2). The high rainfall, in addition to providing a reduction in the content of DM in the cacti, also promoted a reduction in the content of CP and NDF. According to Matias et al (15), cacti have low content of DM, CP and NDF and when supplied as exclusive food to ruminants, they can cause metabolic disturbances in animals, with a consequent drop in performance, and it is recommended to combine them with other rich foods in highly effective fiber and protein, so they can meet the nutritional needs of species and animal categories (16).

Since, for the growth and development of ruminal microorganisms responsible for the degradation of nutrients from the fiber fraction of the forage, the diet selected by the animal should contain values equal to or greater than 7 g/kg DM crude protein (17). Thus, only *O. ficus indica* Mill during the period of low rainfall, had a CP content considered sufficient to meet the minimum requirements for the development of rumen microorganisms (Table 1). Diets with low nitrogen availability can reduce fiber digestion and thus reduce intake due to the slow passage of food through the rumen (18).

Regarding the ADF content, *P. gounellei*, among the other cacti, presented higher contents of this nutrient during the period of low rainfall. The opposite was found for *C. jamacaru*, whose ADF content was higher in the period of high rainfall, compared to the other cacti. ADF values obtained for these cacti are lower than those presented by Carvalho et al (4), who reported values of 28.10% DM and 32.47% DM of ADF for *P. gounellei* and *C. jamacaru*, respectively. These variations in fiber concentrations of the studied species affirm the active participation of the semiarid climate in the caatinga of northeastern Brazil, offering unique features to the different species of cacti in the region (19).

Despite the higher ADF content for *P. gounellei* and *C. jamacaru*, all cacti studied showed low contents of this component and this is explained by the amount of carbohydrates, especially non-fiber carbohydrates (sugar, starch, organic acids

and fructose) present in these plants, which are an important source of energy for ruminants that use them for the development of the microorganisms that colonize the rumen (20).

The increase in the MM content for *P. gounellei* in relation to the other cacti can be explained by the reduction in organic matter, since these variables are inversely proportional. According to Furtado et al (21), cacti usually have a high content of mineral matter in their composition, with *P. gounellei* standing out among cacti because they have higher levels of MM, a fact that corroborates our findings.

Gas production is an important indicator of the degradability and energy density of food, varying due to the concentration of fibers, proteins and the content of tannins (22). The non-fiber fraction of carbohydrates in this cactacea is probably more readily available for ruminal fermentation during the low rainfall period due to a possible adaptation strategy of this cactus for the next rainy season, which will use their reserves to increase its production capacity (23).

The results observed for gas production of *P. gounellei* are possibly associated with the fiber content of carbohydrates of this cactus, which is related to the degradation rate and colonization time by ruminal microorganisms (24).

It is known that *in vitro* gas production results almost entirely from the carbohydrates present in the incubated material and indicates degradation by ruminal microorganisms (25). Thus, the higher gas production rates of the non-fibrous fraction (m2) obtained by the varieties *N. cochenillifera* and *O. stricta* Haw in the period of high rainfall are correlated with the low content of acid detergent fiber that these cacti presented in relation to others during the analyzed period, justifying the observed results.

The reduction in total gas production with the increase in precipitation presented by *P. gounellei* can be explained by its anatomical and physiological characteristics, whose stems and branches are divided into vascular cylinder and central stem, with the highest percentages of soluble carbohydrates in the central stem (26). Thus, probably, during a period of low rainfall there was a thickening of this structure and accumulation of soluble carbohydrates to help plant metabolism during the period of water scarcity.

The short colonization time by ruminal microorganisms (latency phase) observed for *N. cochenillifera* during the period of high rainfall is due to its physical and chemical characteristics, as the soluble fraction is an energetic substrate for microorganisms and has rapid fermentation, which facilitates the processes of adhesion and colonization in the substrate. This can increase the fermentation of fibrous carbohydrates and shorten the latency period (27). Regarding the period of little rain and cacti, *C. jamaçaru* had a longer colonization time, which is not good, as the microorganisms will adhere to the particles more slowly. The higher latency phase, the slower the fiber degradation (28).

Some cacti, in the presence of rain, increase their metabolizable nutritional fractions for growth and multiplication of cladodes and, thus, increase the area of the cladodes, mainly to increase photosynthetic capacity (29). This was observed in the present study by the reduction of the water-insoluble but potentially degradable fraction (b) presented by the cacti *P. gounellei* and *N. cochenillifera*.

With the exception of *C. jamaçaru*, during the high rainfall period and *P. gounellei*, in the low rainfall period, the studied cacti obtained a potential degradability of over 80%, regardless of the rainfall periods analyzed, suggesting a high proportion of digestible and fermentable nutrients in these cacti.

The transition from the low-rainfall to the high-rainfall period resulted in a decline in the effective degradability of all studied cacti. Only *O. ficus indica* Mill and *O. stricta* Haw during the period of high rainfall presented degradability above 50%, which is satisfactory. This degradability is a reflection of the high rate of degradation of fraction B, which in these cacti was higher than the others during the period of high rainfall, making higher percentages of plants more readily available for ruminal fermentation (30).

The chemical composition, the coefficients of gas production and the degradability of the cacti were influenced by the species of cacti and the rainfall variation. Varieties of cactus pear (*Opuntia* and *Nopalea*) stood out among the cacti tested in relation to gas production and *in vitro* degradability.

Acknowledgments

We are grateful to the Brazilian Agricultural Research Corporation (Embrapa Semi-arid) for providing the area for sample collection and laboratory structure for chemical analysis.

Competing interests

The authors declare that they have no competing interests.

REFERENCES

1. Salimon C, Anderson L. How strong is the relationship between rainfall variability and Caatinga productivity? A case study under a changing climate. *An Acad Bras Ci.* 2019; 90:2121–2127. <http://dx.doi.org/10.1590/0001-3765201720170143>.
2. Magalhães ALR, Teodoro AL, Gois GC, Campos FS, Souza JSR, Andrade AP, et al. Chemical and mineral composition, kinetics of degradation and *in vitro* gas production of native cactus. *J Agric Stud.* 2019; 7:119–137. <https://doi.org/10.5296/jas.v7i4>.
3. Edvan RL, Mota RRM, Dias-Silva TP, Nascimento RR, Sousa SV, Silva AL, et al. Resilience of cactus pear genotypes in a tropical semi-arid region subject to climatic cultivation restriction. *Scient Rep.* 2020; 10:1–10. <https://doi.org/10.1038/s41598-020-66972-0>.
4. Carvalho CBM, Edvan RL, Carvalho MLAM, Reis ALA, Nascimento RR. Uso de cactáceas na alimentação animal e seu armazenamento após colheita. *Arch Zootec.* 2018; 67:440-446. <https://doi.org/10.21071/az.v67i259.3803>.
5. Barbosa HA, Kumar TVL, Paredes F, Elliott S, Ayuga JG. Assessment of Caatinga response to drought using Meteosat-SEVIRI normalized difference vegetation index (2008–2016). *ISPRS J Phot Rem Sen.* 2019; 148:235-252. <https://doi.org/10.1016/j.isprsjprs.2018.12.014>.
6. Santos HG, Jacomine PKT, Anjos LHC, et al. Sistema brasileiro de classificação de solos. 5th ed. Brasília, DF: Embrapa; 2018.
7. Aoac. Association of Official Analytical Chemists. Official methods of analysis. 20th ed. Washington, DC: Latimer Jr, G.W; 2016.
8. Van Soest PJ, Robertson JB, Lewis BA. Methods for dietary fiber, neutral detergent fiber, and non-starch polysaccharides in relation to animal nutrition. *J Dairy Sci.* 1991; 74:3583–3597. [https://doi.org/10.3168/jds.S0022-0302\(91\)78551-2](https://doi.org/10.3168/jds.S0022-0302(91)78551-2).
9. Menezes DR, Costa RG, Araújo GGL, Pereira LGR, Nunes ACB, Henrique LT, et al. Cinética ruminal de dietas contendo farelo de mamona destoxificado. *Arq Bras Med Vet Zootec.* 2015; 67:636-641. <https://doi.org/10.1590/1678-7040>.
10. Schofield P, Pitt RE, Pell AN. Kinetics of fiber digestion from *in vitro* gas production. *J Anim Sci.* 1994; 72:2980–2991. <http://dx.doi.org/10.2527/1994.72112980x>.
11. Tilley JMA, Terry RA. A two-stage technique for the *in vitro* digestion of forage crops. *J Brit Grass Soc.* 1963; 18:104–111. <http://dx.doi.org/10.1111/j.1365-2494.1963.tb00335.x>.
12. Ørskov ER, McDonald I. The estimation of protein degradability in the rumen from incubation measurements weighted according to rate of passage. *J Agric Sci.* 1979; 92:499-503. <https://doi.org/10.1017/S0021859600063048>.
13. Alves FAL, Andrade AP, Bruno RLA, Silva MG, Souza MFV, Santos DC. Seasonal variability of phenolic compounds and antioxidant activity in prickly pear cladodes of *Opuntia* and *Nopalea* genes. *Food Sci Techn.* 2017; 37:536-543. <http://dx.doi.org/10.1590/1678-457X.19316>.
14. Davis SC, Simpson J, Gil-Veja KC, Niechayev NA, van Tongerlo E, Castano NH, et al. Undervalued potential of crassulacean acid metabolism for current and future agricultural production. *J Exp Bot.* 2019; 70:6521–6537. <https://doi.org/10.1093/jxb/erz223>.
15. Matias AGS, Araújo GGL, Campos FS, Moraes SA, Gois, GC, Silva TS, et al. Fermentation profile and nutritional quality of silages composed of cactus pear and maniçoba for goat feeding. *J Agric Sci.* 2020; 158:304-312. <https://doi.org/10.1017/S0021859620000581>.
16. Gouws CA, Georgousopoulou EN, Mellor DD, McKune A, Naumovski N. Effects of the consumption of prickly pear cacti (*Opuntia* spp.) and its products on blood glucose levels and insulin: A systematic review. *Med.* 2019; 55:1-18. <https://dx.doi.org/10.3390/medicina55050138>.
17. Pereira DS, Lana RP, Carmo DL, Costa YKS. Chemical composition and fermentative losses of mixed sugarcane and pigeon pea silage. *Acta Scient. Anim Sci.* 2019; 41:e43709. <https://dx.doi.org/10.4025/actascianimsci.v41i1.43709>.

18. Hristov AN, Bannink A, Crompton LA, Huhtanen P, Kreuzer M, McGee M, et al. Invited review: Nitrogen in ruminant nutrition: A review of measurement techniques. *J Dairy Sci.* 2019; 102:5811–5852. <https://doi.org/10.3168/jds.2018-15829>.
19. Albuquerque AS, Freire FJ, Barbosa MD, Marangon LC, Feliciano ALP. Efficiency of biological utilization of micronutrients by forests species in hypoxerophytic Caatinga. *Flor Amb.* 2018; 25:e20170925. <https://doi.org/10.1590/2179-8087.092517>.
20. Carvalho CBM, Edvan RL, Nascimento KS, Nascimento RR, Bezerra LR, Jácome DLS, et al. Methods of storing cactus pear genotypes for animal feeding *Afr J Range For Sci.* 2020; 37:173-179. <https://doi.org/10.2989/10220119.2020.1734084>.
21. Furtado RN, Carneiro MSS, Pereira ES, Moreira Filho EC, Magalhães JA, Oliveira SMP. Intake, milk yield, and physiological parameters of lactating cows fed on diets containing different quantities of xique xique (*Pilosocereus gounellei*). *Semina: Ci Agr.* 2016; 37:483-494. <https://doi.org/10.5433/1679-0359.2016v37n1p483>.
22. Cordova-Torres AV, Mendoza-Mendoza JC, Bernal-Santos G, Gasca TG, Kawas J, Costa RG, et al. Nutritional composition, *in vitro* degradability and gas production of *Opuntia ficus indica* and four other wild cacti species. *Life Sci J.* 2015; 12:42–54. <https://doi.org/10.7537/marslsj1202s15.07>.
23. Vazquez-Mendoza P, Miranda-Romero LA, Aranda-Osorio G, Burgueno-Ferreira JA, Salem AZM. Evaluation of eleven Mexican cultivars of prickly pear cactus trees for possibly utilization as animal fed: *In vitro* gas production. *Agrof Syst.* 2017; 91:749–756. <http://dx.doi.org/10.1007/s10457-016-9947-6>.
24. Ribeiro IA, Voltolini TV, Simões WL, Ferreira MAJF, Menezes DR, Gois GC. Morphological responses, fruit yield, nutritive value and *in vitro* gas production of forage watermelon genotypes on semi-arid condition. *Biol Rhyt Res.* 2019; 50:1–9. <http://dx.doi.org/10.1080/09291016.2019.1629218>.
25. Pinto CS, Magalhães ALR, Teodoro AL, Gois GC, Vêras RML, Campos FS, et al. Potential alternative feed sources for ruminant feeding from the biodiesel production chain by-products. *South Afr J Anim Sci.* 2020; 50:69–77. <http://dx.doi.org/10.4314/sajas.v50i1.8>.
26. Silva JGM, Silva DS, Pereira WE, Diniz MCNM, Silva GJAM, Medeiros MR. Características morfológicas e produção do xique-xique cultivado em diferentes densidades. *Rev Cent.* 2011; 2:08–17.
27. Pinho RMA, Santos EM, Oliveira JS, Carvalho GGP, Silva TC, Macêdo AJS, et al. Does the level of forage neutral detergent fiber affect the ruminal fermentation, digestibility and feeding behavior of goats fed cactus pear? *Anim Sci J.* 2018; 89:1424-1431. <http://dx.doi.org/10.1111/asj.13043>.
28. Yansari AT. Ruminal kinetics of nutrients degradation, hydration, and functional specific gravity of three types of beet pulp. *Iranian J Appl Anim Sci.* 2017; 7:17-26. http://ijas.iaurasht.ac.ir/article_528790.html
29. Alves FAL, Andrade AP, Bruno RLA, Santos DC. Study of the variability, correlation and importance of chemical and nutritional characteristics in cactus pear (*Opuntia* and *Nopalea*). *Afr J Agric Res.* 2016; 11:2882-2892. <https://doi.org/10.5897/AJAR2016.11025>
30. Doorenbos J, Martín-Tereso J, Dijkstra J, Van Laar H. Effect of different levels of rapidly degradable carbohydrates calculated by a simple rumen model on performance of lactating dairy cows. *J Dairy Sci.* 2017; 100:5422-5433. <https://doi.org/10.3168/jds.2016-12278>

Copyright of Revista Colombiana de Ciencia Animal - RECIA is the property of Revista Colombiana de Ciencia Animal - RECIA and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.