

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/372717977>

Carcass yield, non-carcass components and economic viability of using licuri oil in the diet for Santa Inês ewe's

Article in *Revista Colombiana de Ciencias Pecuarias* · July 2023

DOI: 10.17533/udea.rccp.v36n4a4

CITATIONS

0

READS

25

14 authors, including:



Glayciane Costa Gois
Universidade Federal do Maranhão

169 PUBLICATIONS 563 CITATIONS

[SEE PROFILE](#)



Araújo Gherman Garcia Leal de
Brazilian Agricultural Research Corporation (EMBRAPA)

277 PUBLICATIONS 2,068 CITATIONS

[SEE PROFILE](#)



Maria Helena Tavares De Matos
Universidade Federal do Vale do São Francisco (UNIVASF)

70 PUBLICATIONS 840 CITATIONS















[SEE PROFILE](#)



Carcass yield, non-carcass components, and economic viability of licuri oil addition to the diet of Santa Ines ewes

Rendimiento de la canal, componentes ajenos a la canal y viabilidad económica de la adición de aceite de licuri en la dieta de ovejas Santa Inés

Rendimento de carcaça, componentes não carcaça e viabilidade econômica do uso do óleo de licuri na dieta de ovelhas Santa Inês

Cynthia P Lima-Cavalcanti¹ ; Taís J Silva-Macedo² ; Vanessa Peripolli^{3*} ; Glayciane Costa-Gois¹ ; Vanúzia Gonçalves-Menezes² ; Alane P Oliveira-Monte¹ ; Dielen J Menezes-Silva¹ ; Paulo V Nunes-Nascimento³ ; Elves Oliveira-Silva¹ ; Gherman G Leal-Araújo⁴ ; Rafael T Souza-Rodrigues¹ ; Áurea Wischral² ; Maria H Tavares-Matos¹ ; Mário A Ávila-Queiroz^{1*} .

¹Universidade Federal do Vale do São Francisco, Petrolina, PE, Brazil.

²Universidade Federal Rural de Pernambuco, Recife, PE, Brazil.

³Universidade Estadual do Sudoeste da Bahia, Itapetinga, BA, Brazil.

⁴Embrapa Semiárido, Petrolina, PE, Brazil.

To cite this article:

Lima-Cavalcanti CP, Silva-Macedo TJ, Costa-Gois G, Gonçalves-Menezes V, Oliveira-Monte AP, Domingos-Silva A, Menezes-Silva DJ, Nunes-Nascimento PV, Oliveira-Silva E, Leal-Araújo GG, Souza-Rodrigues RT, Wischral A, Tavares-Matos MH, Ávila-Queiroz MA. Carcass yield, non-carcass components, and economic viability of licuri oil addition to the diet of Santa Ines ewes. *Rev Colomb Cienc Pecu* 2023; 36(4):196–209. <https://doi.org/10.17533/udea.rccp.v36n4a4>

Abstract

Background: The production of biofuels has caused an increase in the prices of agricultural commodities. Thus, the ecological footprint, social inclusion and profitability of production systems have encouraged the use of agro-industrial products as an alternative in ruminant feeds. **Objective:** To evaluate carcass yields, non-carcass components, and the economic viability of including licuri oil in diet of Santa Ines ewes. **Methods:** A total of 32 Santa Ines ewes (multiparous, non-lactating, 2–4 years old, and 36.7±0.87 kg of body weight—BW) were allotted to a randomized block design with four treatments (diets containing 0, 2, 4 or 5% licuri oil) with eight replicates per treatment, and confined for 77 days. **Results:** The incremental inclusion of licuri oil promoted a quadratic response on slaughter BW, carcass and true yields, chest widths, heart fat, pancreas, omental fat, and ribeye

Received: February 16, 2022. Accepted: March 18, 2023

*Corresponding author. Petrolina, 56300-990, PE, Brazil. Telephone: +55 8721014861. E-mail: mario.queiroz@univasf.edu.br



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License, which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

© 2023 Universidad de Antioquia. Publicado por Universidad de Antioquia, Colombia.

eISSN: 2256-2958

Rev Colomb Cienc Pecu 2023; 36(4, Oct-Dec):196–209

<https://doi.org/10.17533/udea.rccp.v36n4a4>

area ($p < 0.05$). Carcass, heart and loin weight, shank yield, fat thickness and loin fat were reduced ($p < 0.05$); while cooling losses, left half-carcass weight, saw blade yield, and proportion of meat in the loin increased ($p < 0.05$) with increasing dietary levels of licuri oil. The control diet resulted in the best gross revenue (USD\$609.39); however, there was a loss (USD\$50.96) regarding economic performance indicators. **Conclusion:** The use of up to 2% licuri oil in the diet increased carcass yield of discard ewes.

Keywords: *carcass weight; carcass yield; cooling losses; drylands; economic performance; energy; energy sources; ewe meat; ewes; fat; feed alternatives; licuri; loin; small ruminants; vegetable oil.*

Resumen

Antecedentes: La producción de biocombustibles ha provocado un aumento en los precios de los productos agrícolas básicos. Por lo tanto, la huella ecológica, la inclusión social y la rentabilidad de los sistemas productivos han incentivado la búsqueda de productos agroindustriales como alternativa para la alimentación de rumiantes. **Objetivo:** Evaluar el rendimiento en canal, los componentes ajenos a la canal, y la viabilidad económica de incluir aceite de licuri en dietas para ovejas de raza Santa Ines. **Métodos:** Un total de 32 ovejas Santa Ines (múltiparas, no lactantes, de 2 a 4 años y de $36,7 \pm 0,87$ kg de peso corporal—PC) fueron asignadas a un diseño de bloques al azar con cuatro tratamientos (dieta basal adicionada con 0, 2, 4 o 5% de aceite de licuri) con ocho repeticiones por tratamiento, y confinadas por 77 días. **Resultados:** Los niveles incrementales de aceite de licuri tuvieron un efecto cuadrático sobre el PC al sacrificio, el rendimiento real y de la canal, el ancho del pecho, la grasa del corazón, el páncreas, la grasa del epiplón y el área del ojo del lomo ($p < 0,05$). Se redujo ($p < 0,05$) el peso de la canal, el corazón y el lomo, el rendimiento del jarrete, el grosor de la grasa y la grasa del lomo; y aumentaron las pérdidas por enfriamiento, el peso de media canal restante, el rendimiento de la hoja de sierra y la proporción de carne en el lomo ($p < 0,05$) con niveles incrementales de aceite de licuri. La dieta control resultó en el mejor ingreso bruto (USD\$609,39); sin embargo, hubo una pérdida (USD\$50,96) en los indicadores de desempeño económico. **Conclusión:** El uso de hasta un 2% de aceite de licuri en la dieta incrementa el rendimiento en canal de las ovejas de descarte.

Palabras clave: *aceite vegetal; alimentos alternativos; carne de oveja; energía; fuentes de energía; grasa; licuri; lomo; ovejas; pequeños rumiantes; pérdidas por enfriamiento; peso de la canal; rendimiento económico; rendimiento en canal; tierras secas.*

Resumo

Antecedentes: A produção de biocombustíveis tem causado um aumento nos preços das commodities agrícolas. Assim, a pegada ecológica, a inclusão social e a rentabilidade do sistema de produção têm incentivado o uso de produtos agroindustriais como alternativa alimentar para ruminantes. **Objetivo:** Avaliar os rendimentos de carcaça, componentes não carcaça e a viabilidade econômica da inclusão do óleo de licuri em dietas para ovelhas Santa Inês. **Métodos:** Um total de 32 ovelhas Santa Inês (múltiparas, não lactantes, 2-4 anos e $36,7 \pm 0,87$ kg de peso corporal—PC) foram distribuídas em delineamento em blocos casualizados com 4 tratamentos (dietas contendo óleo de licuri a 0, 2, 4 e 5% de matéria seca) com 8 repetições por tratamento e confinadas por 77 dias. **Resultados:** A inclusão de óleo de licuri promoveu efeito quadrático no PC de abate, rendimentos de carcaça e verdadeiros, largura de peito, gordura do coração, pâncreas, gordura omental e área de olho de lombo ($p < 0,05$). Os pesos de carcaça, coração e lombo, rendimento de pernil, espessura de gordura e gordura do lombo foram reduzidos ($p < 0,05$), e as perdas por resfriamento, peso de meia carcaça esquerda, rendimento de lâmina de serra e proporção de carne no lombo aumentaram ($p < 0,05$) com um aumento nos níveis de óleo de licuri. A dieta controle resultou na melhor receita bruta (USD\$609,39); no entanto, houve uma perda (USD\$50,96) em relação aos indicadores de desempenho econômico. **Conclusão:** O uso de até 2% de óleo de licuri na dieta proporcionou aumento nos rendimentos de carcaça de ovelhas de descarte.

Palavras-chave: *alternativas de alimentação; carne de ovelha; energia; fontes de energia; gordura; licuri; lombo; óleo vegetal; ovelhas; pequenos ruminantes; perdas por resfriamento; performance econômica; peso da carcaça; rendimento de carcaça; terras secas.*

Introduction

The rapid growth of the bio biofuel industry has caused an increase in the prices of agricultural commodities, including corn and soybean. As a result, goat and sheep farmers are looking for alternative ingredients to include in animal diets (Makkar, 2018) with the aim of reducing costs without affecting productivity. In this context, the ecological footprint, social inclusion, and farm profitability are encouraging the use of agro-industrial products as alternatives for ruminants (Pulina *et al.*, 2017). Inclusion of vegetable oils as alternative ingredients in sheep diets is a promising approach for feedlot systems, as it increases the energy density of the diet, influences fat deposition, modifies the fatty-acid profile (FA) of meat (Parente *et al.*, 2018; Bagaldo *et al.*, 2019; Vargas *et al.*, 2020), decreases the short-chain fatty-acid content (SCFA), and increases the levels of 18:1 t11 and 18:2 c9t11 acids (Palmquist and Jenkins 2017).

Brazil has a many palm species with potential to produce vegetable oil. Among these is the licuri palm (*Syagrus coronata* (Martius) Beccari) of the Arecaceae family, native to the Brazilian semiarid regions of Caatinga and covering the eastern and central areas of northern Minas Gerais to southern Pernambuco (Araújo *et al.*, 2019). The licuri palm produces about four bunches a year, with an average of 1,357 small coconuts (licuri). The almond in these coconuts is rich in oils (54% of the almond weight; Rodrigues *et al.*, 2020) widely used in the food, cosmetic, pharmaceutical, and biodiesel industries (Araújo *et al.*, 2019; Santos *et al.*, 2019; Daza *et al.*, 2020). However, the use of licuri oil as an additive for animal feed is still little explored (Silva *et al.*, 2011; Lima *et al.*, 2015; Oliveira *et al.*, 2016).

About 84.9% of the fatty-acid content in licuri oil is saturated, containing 36% lauric acid (Araújo *et al.*, 2019), which may be important in tropical regions, since oils with high unsaturated acid levels are subject to oxidation. Although saturated fatty acids have hypercholesterolemic properties, most of them are highly oxidized after absorption. This can contribute to decrease the accumulation

of body fat in small ruminants, which may be of interest to consumers (Parente *et al.*, 2020). To the best of our knowledge, no scientific study has been conducted on the inclusion of licuri oil in diets for ewes and its effects on the carcass traits of confined animals.

The economic analysis of the investments made in a feedlot is an important assessment tool that validates the viability of a project by describing all the details of the production chain and seeking the necessary changes for greater productive efficiency (Santos *et al.*, 2017). Studies evaluating the economic viability of licuri oil as an ingredient in the diet of confined ewes are scarce in the literature. Thus, we hypothesized that ewes fed diets containing licuri oil would have carcass traits similar to those fed the control diet, but that licuri oil would reduce the deposition of fat in tissues due to the greater oxidation of lauric acid. Additionally, licuri oil could reduce production costs and improve the economic response in confined discard ewes.

Therefore, the objective of the present study was to evaluate carcass yield, non-carcass components, and the economic viability of including licuri oil in diets for Santa Ines ewes.

Materials and Methods

Ethical considerations

This study was approved and certified by the Ethics Committee on Human and Animal Studies of UNIVASF (Protocol N° 0002/24102017).

Location

The experiment was conducted at Universidade Federal do Vale do São Francisco (UNIVASF), Petrolina, PE, Brazil. The predominant climate of the region, according to Köppen and Geiger (1928), is semi-arid BSh with summer rains. The rainy season starts in November and ends in April, with an annual rainfall of 376 mm. During the experimental period, the maximum and minimum temperatures were 33.8 and 24.6 °C, respectively, with relative humidity of 50.5 and 73.6%, respectively.

Animals, treatments, and experimental diets

Thirty-two Santa Ines discard ewes (multiparous, non-lactating, 2–4 years old and 36.7 ± 0.87 kg of body weight—BW) were distributed in a randomized block design with four treatments (diets) and eight replicates per treatment. The confinement lasted 77 days preceded by 15 days of adaptation. The ewes were housed in individual pens (2.42 m²) and provided with feeders and drinking fountains.

The treatments consisted of increasing levels of licuri oil in the diet (0, 2, 4 and 5%, based on dry matter). The experimental diets (Table 1) were formulated to be isoproteic, in a roughage/concentrate ratio of 50:50, based on dry matter, and balanced to allow an average daily gain of 40 g/day, following NRC (2007) recommendations. Diet samples were collected to determine the fatty-acids profile by gas chromatography (Table 1) following the methodology of Visentainer (2012). The chemical composition of the ingredients of the experimental diets are shown in Table 2.

Table 1. Ingredients and chemical composition of experimental diets.

Ingredient (% dry matter)	Inclusion level of licuri oil (% DM)			
	0	2	4	5
Ground corn	34.3	32.3	29.8	28.5
Soybean meal	13.8	13.8	14.3	14.6
Licuri oil	0	2.0	4.0	5.0
Mineral mixture ¹	1.6	1.6	1.6	1.6
Dicalcium phosphate	0.3	0.3	0.3	0.3
Elephant grass	50.0	50.0	50.0	50.0
Chemical composition (% DM)				
Dry matter ²	54.9	55.1	55.3	56.1
Ash	8.6	8.6	8.7	8.7
Crude protein	11.6	11.5	11.5	11.6
Ether extract	2.0	3.9	5.8	6.8
Neutral detergent fiber	47.2	46.6	45.9	45.5
Metabolizable energy – SRNS (Mcal/kg)	2.5	2.6	2.7	2.7
Fatty acids (%)				
Caproic (C6:0)	0.04	0.04	0.034	0.04
Capric (C10:0)	0.04	0.17	0.29	0.36
Lauric (C12:0)	1.57	2.41	3.24	3.67
Miristic (C14:0)	0.49	0.78	1.06	1.21
Palmitic (C16:0)	19.54	19.46	19.40	19.36
Palmitoleic (C16:1)	0.14	0.14	0.14	0.13
Heptadecanoic (C17:0)	0.14	0.14	0.14	0.13
Stearic (C18:0)	3.45	3.49	3.54	3.57
Oleic (C18:1n9c/t)	13.51	13.29	13.03	12.89
Linoleic (C18:2n6c)	36.13	35.02	33.88	33.31
α -Linolenic (C18:3)	18.49	18.46	18.47	18.48
Archaic (C20:0)	0.14	0.14	0.13	0.13
Eicosanoic/gadoleic (C20:1)	0.06	0.06	0.06	0.05
Dihomo- γ -linolenic (C20:3)	0.07	0.07	0.07	0.06

¹Composition= 7.5% P; 19% Ca; 1% Mg; 7% S; 14.3% Na; 21.8% Cl; 500 ppm Fe; 300 ppm Cu; 4,600 ppm Zn; 1,100 ppm; Mn; 80 ppm I; 405 ppm Co; 30 ppm Se. ²% of natural matter.

Table 2. Chemical composition of diet ingredients.

(% dry matter)	Elephant grass	Ground corn	Soybean meal	Licuri oil
Dry matter ¹	26.2	87.6	88.6	100
Mineral matter	7.9	2.0	6.5	-
Crude protein	6.9	8.9	48.8	-
Ether extract	2.9	4.5	1.9	100
Neutral detergent fiber	70.9	13.0	14.8	-
Acid detergent fiber	41.9	7.7	8.7	-
Total carbohydrates	83.0	72.0	42.8	-
Non-fiber carbohydrates	17.4	77.8	27.6	-
Total digestible nutrients	57.0	86.0	81.2	-

¹% of natural matter.

Feeding was carried out twice a day (9 am and 4 pm) as complete ration, allowing for refusals of 10% of offered feed (DM basis). Water was provided *ad libitum*. Refused feed was collected and weighed daily to adjust the feed offered and calculate dry matter intake (DMI). During the experimental period, the ewes showed average dry matter intake, ether extract and average daily gain (in g.day⁻¹) of 1,609, 44, and 91.3 (0% licuri oil); 1,759, 64, and 103.9 (2% licuri oil); 1,356, 63, and 63.9 (4% licuri oil); and 1,307, 79, and 48.6 (5% licuri oil), respectively.

Samples of ingredients, feed supplied, and leftovers were collected weekly and stored at -20 °C for further analysis. Subsequently, the samples were pre-dried in a forced ventilation oven at 55 °C for 72 hours and ground to 1-mm particle size (Wiley mill, Marconi, MA-580, Piracicaba, Brazil). Chemical analyses were performed using the procedures described by AOAC (2016) for dry matter (DM; method 967.03), mineral matter (MM; method 942.05), crude protein (CP; method 981.10), and ether extract (EE; method 920.29). Neutral detergent fiber was determined according to Mertens (2002).

Slaughter and carcass yield

After 77 days in feedlot the ewes were individually weighed to obtain BW after solid fasting of 16 hours, according to animal welfare standards. After this period, the animals were weighed to determine body weight at slaughter

(BWS). The animals were first desensitized by cerebral concussion and immediately slaughtered by bleeding through the carotid arteries and jugular veins, according to the current rules of the Regulation for the Inspection of Industrial Sanitation of Products of Animal Origin (Brazil 2017).

After bleeding the animals were skinned. The head was separated from the carcass and the limbs separated by section of the carpometacarpal joints (forelimbs) and the tarsometatarsal joints (hind limbs). Evisceration was performed through an opening along the abdominal midline to remove the non-carcass components: chips (reproductive and respiratory tracts + bladder), gastrointestinal tract, liver, spleen, heart, heart fat, omental fat, pancreas and skin, which were washed and weighed separately for later calculation of the total weight of the organs, in order to determine their yield in relation to the weight of the empty body and the body weight at slaughter (Silva-Sobrinho 2001).

The carcasses were identified and weighed with the kidneys and renal fat to obtain the hot carcass weight (HCW). The carcasses were transferred to a cold chamber where they were hung on appropriate hooks and kept refrigerated at 5 °C for 24 hours. After this period, carcasses were weighed to obtain the cold carcass weight (CCW). Hot carcass yield (HCY; $HCY = (HCW/BWS) \times 100$), cold carcass yield (CCY; $CCY =$

($CCW/BWS \times 100$) and determination of cooling losses (CL; $CL = [(HCW - CCW)/HCW] \times 100$) were calculated according to Silva Sobrinho *et al.* (2005). The real yield (RY; $RY = HCW/Empty\ BW$) was obtained according to Sañudo and Sierra (1986). The kidneys and kidney fat were removed from the chilled, weighed carcasses, and the weights obtained were subtracted from the HCW and CCW.

Morphometric evaluation of carcasses

After the refrigeration period, measurements were made on the carcasses according to the methodology of Cezar and Souza (2007), with the following measurements (in cm) being determined: external carcass length (ECL); internal carcass length (ICL); first chest width (FCW); second chest width (SCW); croup width (CW); croup perimeter (CP); leg length (LL); leg perimeter (LP) and depth of the chest (DC).

Evaluation of commercial cuts and ribeye area

The carcasses were divided longitudinally and the left half-carcass (LHC) was weighed and divided into six commercial cuts: neck, shoulder, ribs, loin, leg and saw. The cuts were weighed separately and then the yields of each cut calculated in relation to the weight of the left half-carcass (Silva Sobrinho 2001). After obtaining the commercial cuts, the loins were identified, vacuum packed and frozen at $-18\text{ }^{\circ}\text{C}$ for further dissection. Between the 12th and the 13rd ribs, a cross-section was made to measure the loin eye area (LEA) of the *Longissimus dorsi* muscle, tracing its outline on transparent slides for later measurement with the AutoCAD[®] software (Autodesk, Inc, Brazil). The thickness of the subcutaneous fat was measured with a digital caliper (150 mm, MTX-316119, Franca, Brazil).

The loins were thawed under refrigeration ($8\text{ }^{\circ}\text{C}$; 12 hours) and dissected manually. Tissues were separated into total fat (subcutaneous and intermuscular combined), muscle (total muscle dissected, after complete removal of all adhered subcutaneous and intermuscular fat) and bones (dissected after complete removal of all muscle

and adhered subcutaneous and intermuscular fat), which were weighed individually, to be expressed as a percentage of the weight of the loin, as described by Purchas *et al.* (1991) and Hopkins *et al.* (1997).

Economic analysis

The economic analysis was performed to verify the viability of using licuri oil in diets. The cost of feed per kg of roughage (irrigated and fertilized elephant grass) was 0.06, corn was 0.16, soybean meal 0.24; mineral mixture of 0.52, licuri oil of 0.84, and bicalcium phosphate USD\$ 0.64 (cost of ingredients in the São Francisco Valley region in May 2018, expressed in USD\$ kg^{-1} DM; dollar quotation on 12/22/2022; 1 BRL = 0.1928 USD, Central Bank of Brazil).

The purchase price of each ewe was USD\$44.28. The capital invested in the construction of the containment was estimated at USD\$3,052.56. The opportunity cost was calculated as 6% of the capital invested, divided by 360 days and multiplied by the 77 days of confinement, which was later divided by four experimental treatments (diets). The annual depreciation was calculated using the straight-line method (constant quotas), using the amount spent on the facilities and subtracting the residual value (10%), which was divided by the 15 years of useful life. Finally, it was adjusted to 77 days and divided by the four experimental treatments (diets). The cost of labor was calculated based on the minimum wage in force in May 2018 (USD\$168.96), plus 37% social charges. For this, the labor apportioned to the work performed for 0300 h daily was adopted, corresponding to the time needed to feed the ewe and to clean the facilities.

To determine the cost of production, the Total Operating Cost of Production methodology, described by Matsunaga *et al.* (1976), which defines and takes into account all costs related to the production of a good, product or service. The economic indicators analyzed were: Gross Income (GI, USD\$/year) = total production in kg BW \times market selling price; Effective Operating

Cost (EOC, USD\$/year) = costs of operations (maintenance of facilities and equipment) + costs of contracted labor + costs of inputs (food, medications, energy); Total operating cost (TOC, USD\$/year) = EOC + depreciation; Total cost (TC, USD\$/year) = TOC + return on average capital invested in animals and improvements; Gross margin (GM, USD\$/year) = GI – EOC; Net margin of activity (NM, USD\$/year) = GI – TOC; Result (profit or loss) = GI – TC.

Economic performance indicators were assessed by profitability (P; USD\$/year) and rentability (capital invested, %). Profitability was obtained by the following equation: $P = (GI - TC * 100) / GI$. To calculate gross income (GI), gross income from the sale of shredded meat, income from the sale of skin and manure were considered. The sales values per kg of shredded meat, per piece of raw leather and per kg of manure were USD\$3.36, USD\$0.88, and USD\$0.02, respectively. Values were based on prices current in the São Francisco Valley region of northeastern Brazil in 2018.

Statistical analysis

The data were submitted to the Shapiro–Wilk test to verify the normal distribution of residues and homogeneity of variances, using the univariate procedure. Analysis of variance was performed using the GLM procedure using orthogonal polynomials. Finally, with a significant effect at 5% probability, regression equations were estimated using the REG procedure. The SAS statistical package, version 9.1 (SAS Institute, Cary, NC, USA; 2003) was used for all tests.

Results

The inclusion of licuri oil in the diets of sheep promoted a quadratic effect for BWS ($p=0.03$), HCY ($p=0.014$), CCY ($p=0.013$), and RY ($p=0.007$). The HCW ($p<0.001$) and CCW ($p<0.001$) were reduced and CL increased ($p<0.001$) according to the increase in licuri oil levels in the diets (Table 3). The increase in licuri oil levels provided a

quadratic effect for first chest width ($p<0.001$) and second chest width ($p=0.03$) measured in the chest of the carcass. There were no effects of the levels of licuri oil on ILC, ECL, LL, CW, CP, LP and DC ($p>0.05$; Table 3).

A reduction in heart weight ($p=0.004$) was observed as the levels of licuri oil increased. A quadratic effect was seen on the weight of heart fat ($p=0.001$), pancreas ($p<0.001$) and omental fat ($p<0.001$), increasing from 0 to 2% inclusion of licuri oil (Table 4). No effect of licuri oil levels was observed on the weight of spleen, liver, kidneys, trimmings, rumen + reticulum + omasum + abomasum, small intestine, large intestine, legs and skin ($p>0.05$; Table 4).

The addition of licuri oil to the diets promoted a linear increase of LHC weight ($p<0.001$) and saw yield ($p<0.001$), and a reduction of leg yield ($p<0.001$). There was no effect of licuri oil on rib, loin, shoulder, and neck yields ($p>0.05$; Table 5).

A quadratic effect ($p=0.02$) of licuri oil on LEA was observed, for levels 0 and 2% of licuri oil. Fat thickness ($p=0.02$) and loin weight ($p<0.001$) decreased linearly according to the inclusion level of licuri oil. Regarding tissue composition of the loin, licuri oil resulted in linear increase in the proportion of meat ($p<0.001$) and reduction in fat proportion ($p<0.001$). There was no effect of licuri oil on the proportion of bone ($p>0.05$; Table 4).

The selling price per kg of carcass during 2018 in the different regions of Brazil ranged between 2.13 and 3.19 for lambs/ram and between 1.59 and USD\$2.13 for discard ewes. The economic analysis was performed considering the purchase cost of the animal. The treatment that resulted in the best gross revenue was the one with 0% inclusion of licuri oil in the diet, making a value of USD\$663.35, however, considering the economic performance indicators, the result was negative with a loss of approximately USD\$55.48 (Table 6).

Table 3. Weight (kg), yield (%), and morphometry (cm) of carcasses of Santa Ines ewes fed diets containing increasing levels of licuri oil.

Variable	Licuri oil level				SEM	p-value	
	0%	2%	4%	5%		L	Q
Initial body weight (kg)	34.85	37.52	36.54	34.53	0.90	0.80	0.27
Slaughter body weight (kg)	41.88	45.52	40.46	38.35	0.94	0.06	0.03 ¹
Hot carcass weight (kg)	22.81	22.25	19.77	17.73	0.55	<0.001 ²	0.15
Cold carcass weight (kg)	22.18	21.99	19.15	17.14	0.54	<0.001 ³	0.14
Hot carcass yield (%)	52.25	52.04	48.80	43.87	0.82	<0.001	0.014 ⁴
Cold carcass yield (%)	50.78	50.58	47.25	42.40	0.81	<0.001	0.013 ⁵
Real yield (%)	59.94	59.98	55.76	48.85	1.08	<0.001	0.007 ⁶
Cooling losses (%)	2.78	2.80	3.17	3.35	0.06	<0.001 ⁷	0.07
Internal carcass length (cm)	69.9	68.3	68.8	68.7	0.54	0.54	0.48
External carcass length (cm)	73.4	73.9	73.6	70.9	0.71	0.31	0.28
Leg length (cm)	35.9	34.4	35.9	36.8	0.48	0.41	0.14
Croup width (cm)	25.9	26.4	26.1	24.9	0.26	0.23	0.09
First chest width (cm)	25.7	27.6	25.6	23.7	0.31	0.005	<0.001 ⁸
Second chest width (cm)	19.6	19.8	18.4	17.7	0.22	<0.001	0.03 ⁹
Croup perimeter (cm)	68.1	68.2	67.2	65.9	0.49	0.12	0.35
Leg perimeter (cm)	38.8	38.5	36.3	38.2	0.55	0.34	0.58
Thorax perimeter (cm)	30.8	28.9	30.2	29.6	0.53	0.53	0.85

SEM= standard error of the mean; L= Linear Effect; Q= Quadratic Effect; Significance was determined at the 5% probability level. Equations: ¹y= 42.10+2.81x-0.74x², R²= 0.57; ²y= 0.23-0.99x, R²= 0.51; ³y= 22.73-0.99x, R²= 0.42; ⁴y= 52.12+1.39x-0.59x², R²= 0.56; ⁵y= 50.67+1.36x-0.59x², R²= 0.57; ⁶y= 59.75+2.17x-0.85x², R²= 0.58; ⁷y= 0.27+0.12x, R²= 0.48; ⁸y= 25.72+1.76x-0.43x², R²= 0.63; ⁹y= 19.60+0.33x-0.15x², R²= 0.49.

Table 4. Weight (kg), yield (%), and morphometry (cm) of carcasses of Santa Ines ewes fed diets containing increasing levels of licuri oil.

Variable (g)	Licuri oil level				SEM	p-value	
	0%	2%	4%	5%		L	Q
Spleen	96.0	89.0	86.5	86.0	3.86	0.35	0.76
Heart	229.9	214.2	196.1	180.7	6.48	0.004 ¹	0.72
Heart fat	132.0	172.0	118.7	105.9	5.86	0.002	0.001 ²
Liver	602.6	615.5	614.4	649.7	24.47	0.56	0.83
Pancreas	36.5	56.5	34.0	33.1	2.22	0.007	<0.001 ³
Kidneys	111.7	108.0	99.5	105.9	4.73	0.29	0.51
Cutting	0.9	0.7	0.8	0.8	0.03	0.12	0.20
Rumen+Reticle+Omasum+Abomasum	1.5	1.6	1.6	1.5	0.04	0.75	0.46
Small intestine	0.5	0.5	0.5	0.5	0.02	0.99	0.39
Large intestine	0.7	0.6	0.6	5.8	0.04	0.27	0.82
Omental fat	3.1	3.5	2.4	1.7	0.01	<0.001	<0.001 ⁴
Feet	0.9	0.9	0.9	0.9	0.01	0.57	0.19
Skin	2.6	2.6	2.6	2.7	0.06	0.68	0.63

SEM = standard error of the mean; L= Linear Effect; Q= Quadratic Effect; Significance was determined at the 5% probability level. Equations: ¹y= 0.24-0.14x, R²= 0.59; ²y= 0.13+0.28x-0.70x², R²= 0.49; ³y= 0.38+0.12x-0.28x², R²= 0.58; ⁴y= 1.08+0.14x-0.52x², R²= 0.44.

Table 5. Left half-carcass weight, yields from commercial cuts, loin eye area (LEA), fat thickness of the LEA, and tissue composition of the loin of Santa Ines ewes fed increasing levels of licuri oil.

Variable (g)	Licuri oil level				SEM	p-value	
	0	2	4	5		L	Q
Left half-carcass (kg)	10.6	10.1	9.2	8.5	0.22	<0.001 ¹	0.42
Rib (%)	26.4	26.8	26.8	26.1	0.27	0.84	0.30
Loin (%)	9.1	8.5	8.9	8.6	0.13	0.33	0.41
Saw (%)	5.9	5.4	5.0	4.3	0.14	<0.001 ²	0.43
Shoulder (%)	16.9	17.1	17.5	17.9	0.22	0.11	0.61
Leg (%)	31.3	32.3	32.8	33.3	0.19	<0.001 ³	0.68
Neck (%)	10.2	9.9	8.8	9.8	0.24	0.18	0.39
LEA (cm ²)	12.5	12.9	10.7	10.8	0.22	<0.001	0.02 ⁴
Fat thickness (mm)	2.4	2.5	1.9	1.6	0.15	0.02 ⁵	0.25
Loin (kg)	0.94	0.88	0.81	0.68	0.29	<0.001 ⁶	0.27
Tissue composition (%)							
Meat	50.5	53.5	55.6	58.7	0.79	<0.001 ⁷	0.64
Fat	27.5	26.4	20.0	17.4	1.10	<0.001 ⁸	0.20
Bone	22.0	20.1	24.4	23.9	0.63	0.06	0.18

SEM= standard error of the mean; L= Linear Effect; Q= Quadratic Effect. Significance was determined at the 5% probability level. Equations: ¹y= 10,747-409.05x, R²= 0.43; ²y= 6.02-0.30x, R²= 0.55; ³y= 31.41+0.38x, R²= 0.51; ⁴y= 12.63+0.16x-0.12x², R²= 0.55; ⁵y= 2.58-0.17x, R²= 0.56; ⁶y= 950.7-38.08x, R²= 0.58; ⁷y= 50.39+1.52x, R²= 0.44; ⁸y= 28.65-22.11x, R²= 0.44.

Discussion

Carcass yield parameters are directly affected by BWS (Costa *et al.*, 2019a) and these were strongly influenced by the dietary levels of licuri oil, showing progressive decreases that caused a reduction in the measurements made in the thorax of the discard ewes. The change in carcass yield in relation to the increasing levels of licuri oil may have been influenced by the reduced deposition of adipose tissue and the increase in CL in animals supplemented with 4 and 5% licuri oil, which is directly related to lower DM intake.

In addition to DMI, one of the factors that may have contributed to a reduction of carcass yield with high levels of licuri oil could be the weight of the liver. Although there was no effect of treatment on this organ, at the highest level of licuri oil (5%), the liver weight stood out in relation to the other treatments (649.7 g). This result can be explained by the high metabolic rate of the organ, and this activity is intensified by the increase in the EE level in the diets (Bauchart *et al.*, 1996), a result seen

in this study. The liver is more developed to meet the demand for nutrient metabolism, since it is the central organ that gathers the absorbed nutrients and those from circulation in general, moderating and distributing them to the peripheral tissues and acting as the main site for the regulation of nutrient supply (Meikle *et al.*, 2018).

Once ingested, medium-chain fatty acids in licuri oil are absorbed by the abomasum, reach the liver through the hepatic portal system, and behave in a similar way to glucose (Daza *et al.*, 2020). This readily available energy source results in greater development of organs involved in energy balance. Thus, the increase in pancreas weight by the addition of up to 2% licuri oil to the diet may be related to changes in lipolytic activities, which require more of this gland to guarantee the best use of fatty acids, causing its hypertrophy (Delgado-Puga *et al.*, 2020). Thus, the lower values of pancreas weight due to the greater inclusions levels of licuri oil may reflect possible changes in the influx of nutrients in the duodenum and in the secretion of lipases (Xu *et al.*, 2009) caused by the low DMI.

Table 6. Operating costs, gross revenue result indicators, and economic performance indicators for the acquisition and termination of ewes, receiving a 50% concentrate diet and different levels of licuri oil.

Specification	Unit	Licuri oil level (%)			
		0	2	4	5
<i>Effective operating cost (EOC)</i>					
Purchase of animals		385.57	385.57	385.57	385.57
Labor		60.63	60.63	60.63	60.63
Food		251.30	257.14	214.05	204.22
Total EOC		697.50	703.34	660.25	650.42
Depreciation		10.66	10.66	10.66	10.66
Total operating cost		708.10	714.00	670.91	661.06
Remuneration of invested capital	USD\$	10.66	10.66	10.66	10.66
Total cost		718.82	724.66	681.57	671.72
Sale of shredded meat		649.77	644.38	561.16	502.17
Sale of skin		7.71	7.71	7.71	7.71
Sale of manure		5.87	3.73	4.95	3.29
Gross Revenue		663.35	655.82	573.83	513.17
<i>Result indicators</i>					
Gross Margin		(-34.16)	(-47.52)	(-86.43)	(-137.25)
Net Margin	USD\$	(-44.82)	(-58.18)	(-97.09)	(-147.91)
Result		(-55.48)	(-68.84)	(-107.75)	(-158.57)
<i>Economic performance indicators</i>					
Profitability	%	(-1.61)	(-2.02)	(-3.62)	(-5.96)
Rentability		(-0.32)	(-0.40)	(-0.62)	(-0.92)

Values expressed in US Dollars (USD\$); Dollar quote on 12/22/2022; 1 BRL = 0.1928 USD\$ (Central Bank of Brazil).

Such an event may also have caused a reduction in heart weight, which suffered a decrease of 49.12 g (21.37%); that is, losses of 9.82 g for each 1% inclusion of licuri oil in the diet. These losses may be related to a low availability and absorption of nutrients as a result of the damage caused to the metabolism and rumen fermentation.

The inclusion of vegetable oils in ruminant diets modifies the production of volatile fatty acids in the rumen, increasing the proportion of propionate and reducing acetate. Propionate is a gluconeogenic precursor and therefore can increase glucose supply for lipogenesis, resulting in greater fat deposition (Park *et al.*, 2018). This factor, combined with the fatty-acid composition of licuri oil, the intake of DM and the inherent characteristics of the animal such as early growth, lower adult weight and greater accumulation of fat

due to sex hormones, makes females more efficient, whereas fat metabolism in males (Argente-Arizón *et al.*, 2016) explains the greater deposition of omental and heart fat with dietary inclusion of 2% licuri oil.

Leg was the cut that showed the highest yield with the rise in licuri oil level in the diets. The highest percentage of carcass muscle mass is found in this cut, with a higher edible yield and, in addition to predicting the total content of the tissues, it is considered the most noble cut in sheep (Araújo *et al.*, 2017). Saw yield and loin weight were reduced by 0.33 and 5.53%, respectively, in their proportion and weight for each 1% licuri oil added to the diet. These reductions are related to the lower HCW and CCW observed with the increase in licuri oil in diets, a reduction in fat thickness and the later maturity of cuts.

The LEA is a measure that reflects the amount of muscle in the carcass, being positively correlated with muscularity. Comparing inclusion levels of 0 and 5% licuri oil, we observed a reduction of 10 g (36.84%) in the proportion of fat in the total composition of the loin, which could explain the increase in the proportion of meat in this cut.

The EOC, describing the amount of resources used to cover the costs of confinement (Costa *et al.*, 2019b), underwent changes due to the levels of licuri oil in the diet, reflecting the fluctuations in the DM intake according to the level of inclusion. The DM intake of the animals associated with decrease in the costs of concentrates in diets containing licuri oil was the determining factor in EOC. Similarly, it affected the total operating cost, which, in addition to expenses, also includes the depreciation of improvements, machinery and equipment.

Considering the results and the economic performance indicators, the addition of licuri oil to the diet was unsatisfactory. In the short and medium term, the gross and net margins were negative, revealing that the income obtained from the activity did not even cover the actual operating costs and an interruption of the production process would be indicated. In the long run, the results (profit or loss) were also negative, meaning that return on the capital invested in the activity would be lower than the opportunity interest rate, and therefore, neither treatment was economically viable.

Within the system, the effective cost of animal feeding is mainly responsible for compromising the remuneration generated by the sale of products (Ferreira *et al.*, 2017), even though it has worked with cut meat as a strategy to add value to the meat discard sheep. Thus, in order to achieve a return on the capital invested in the activity of at least US\$ 19.28 above the opportunity interest rate adopted, it would be necessary to reduce feed cost with the inclusions of 2, 4, or 5% licuri oil. In these conditions, if profitability and rentability of each diet were compared, the system including 5% licuri oil in the feed would have better profitability and rentability.

Under the experimental conditions, the inclusion of up to 2% licuri oil in the diet provided an increase in carcass yield of discard ewes. However, the acquisition or purchase of this category of animal for feedlots was not profitable, nor was the inclusion of higher levels of licuri oil in the diet.

Declarations

Acknowledgements

To the Research Support Foundation of Pernambuco (FACEPE) and the National Council for Scientific and Technological Development (CNPq) for the scholarship granted.

Funding

The Research Support Foundation of Pernambuco (FACEPE; process: APQ-0895-5.05/14) provided financial support for this project.

Conflicts of interest

The authors declare they have no conflicts of interest with regard to the work presented in this report.

Author contributions

Conception and design: MAAQ, AW, MHTM, GGLA, and RTSR. Acquisition of data: CPLC, TJSM, VGM, APOM, ADS, DJMS, PVNN, and EOS. Analysis and interpretation of data: CPLC, MAAQ, and RTSR. Drafting the article: CPLC, GCG, and MAAQ. Critical review of important intellectual content: GCG and MAAQ. All authors gave a final approval of the version to be published.

References

AOAC. Official methods of analysis. Association of Analytical Chemists International. 20th ed. Washington (DC): G.W. Latimer Jr; 2016.

- Araújo PHM, Maia AS, Cordeiro AMTM, Gondim AD, Santos NA. Catalytic deoxygenation of the oil and biodiesel of Licuri (*Syagrus coronata*) to obtain n-alkanes with chains in the range of biojet fuels. ACS Omega 2019; 4(14):15849–15855. <https://doi.org/10.1021/acsomega.9b01737>
- Araújo M, Fontes Cezar M, Gomes MG, Souza WH, Morais J, Torres AS, Henriques G, Campos R. Commercial cuts and carcass characteristics of sheep and goat supplemented with multinutritional blocks. Rev MVZ 2017; 22(3):6180–6190. <https://dx.doi.org/10.21897/rmvz.1123>
- Bagaldo AR, Miranda GS, Soares Júnior MSF, Araújo FL, Matoso RVM, Chizzotti ML, Bezerra LR, Oliveira RL. Effect of Licuri cake supplementation on performance, digestibility, ingestive behavior, carcass traits and meat quality of grazing lambs. Small Rum Res 2019; 177(1):18–24. <https://doi.org/10.1016/j.smallrumres.2019.05.020>
- Bauchart D, Gruffat D, Durand D. Lipid absorption and hepatic metabolism in ruminants. Proc Nut Soc 1996; 55(1B):39–47. <https://doi.org/10.1079/PNS19960010>
- Brazil. Decreto Nº 9.013, de 29 de março de 2017. Regulamenta a Lei Nº 1.283, de 18 de dezembro de 1950, e a Lei Nº 7.889, de 23 de novembro de 1989, que dispõem sobre a inspeção industrial e sanitária de produtos de origem animal. Diário Oficial União. 2017. <https://www.gov.br/agricultura/pt-br/assuntos/aquicultura-e-pesca/legislacao/legislacao-geral-da-pesca/decreto-no-9-013-de-29-03-2017.pdf/view>
- Costa CA, Carvalho FFR, Guim A, Andrade GP, Cardoso DB, Maciel MV, Silva GG, Nascimento AGO. Carcass characteristics of lambs fed diets with increasing levels of crude glycerin. Asian-Austral J Anim Sci 2019a; 32(12):1882–1888. <https://doi.org/10.5713/ajas.18.0825>
- Costa JB, Oliveira RL, Silva TM, Ribeiro OL, Ribeiro RDX, Pinto LFB, Nascimento TVC. Economic analysis of the finishing of lambs under confinement conditions using licuri cake (*Syagrus coronata* Mart. Becc.). Rev Bras Saúde Prod Anim 2019b; 20(e0252019):01–15. <http://dx.doi.org/10.1590/S1519-9940200252019>
- Daza A, Souza JG, Monerrat JPIS, Ribeiro CVDM. Performance of growing lambs supplemented with ground licuri (*Syagrus coronata*). Asian-Austral J Anim Sci 2020; 33(01):01–23. <https://doi.org/10.5713/ajas.20.0199>
- Hopkins DL, Fogarty NM, Menzies DJ. Differences in composition, muscularity, muscle:bone ratio and cut dimensions between six lamb genotypes. Meat Sci 1997; 45(4):439–450. [https://doi.org/10.1016/S0309-1740\(96\)00134-9](https://doi.org/10.1016/S0309-1740(96)00134-9)
- Köppen W, Geiger R. Climate der Erde. Gotha: Verlag Justus Perthes 1928; Wall-map 150cmx200cm.
- Lima LS, Oliveira RL, Garcez Neto AF, Bagaldo AR, Abreu CL, Silva TM, Carvalho ST, Bezerra LR. Licuri oil supplements for lactating cows on pasture. Can J Anim Sci 2015; 95(1):617–624. <https://doi.org/10.4141/CJAS-2014-165>
- Makkar HPS. Review: Feed demand landscape and implications of food-not feed strategy for food security and climate change. Anim 2018; 12(8):1744–1754. <https://doi.org/10.1017/S175173111700324X>
- Matsunaga M, Bemelmans PF, Toledo PEN, Dullely RD, Okawa H, Pedroso IA. Metodologia de custo de produção utilizada pelo IEA. Agricultura em São Paulo – SP 1976; 23(1):123–139. <http://www.iea.sp.gov.br/out/verTexto.php?codTexto=11566>
- Meikle A, Brun V, Carriquiry M, Soca P, Sosa C, Adrien ML, Chilibruste P, Abecia JA. Influences of nutrition and metabolism on reproduction of the female ruminant. Anim Rep 2018; 15(1):899–911. <http://dx.doi.org/10.21451/1984-3143-AR2018-0017>
- Mertens DR. Gravimetric determination of amylase-treated neutral detergent fiber in feeds with refluxing in beaker or crucibles: collaborative study. J AOAC Int 2002; 85:1217–1240. <https://pubmed.ncbi.nlm.nih.gov/12477183/>

NRC. National Research Council. Nutrient requirements of small ruminants: sheep, goats, cervids, and new world camelids. Washington (DC): The National Academy Press. 2007.

Oliveira RL, Neto SG, Lima FHS, Medeiros AN, Bezerra LR, Pereira ES, Bagaldo AR, Pellegrini CB, Correia BR. Composition and fatty acid profile of milk from cows supplemented with pressed oilseed cake. *J Anim Sci* 2016; 87(10):1225–1232. <http://dx.doi.org/10.1111/asj.12571>

Palmquist DL, Jenkins TC. A 100-Year Review: Fat feeding of dairy cows. *J Dairy Sci* 2017; 100(12):10061–10077. <https://doi.org/10.3168/jds.2017-12924>

Parente MOM, Rocha KS, Bessa RJB, Parente HN, Zanine AM, Machado NAF, Lourenço Júnior JB, Bezerra LR, Landim AV, Alves SP. Effects of the dietary inclusion of babassu oil or buriti oil on lamb performance, meat quality and fatty acid composition. *Meat Sci* 2020; 160:e107971. <https://doi.org/10.1016/j.meatsci.2019.107971>

Parente MOM, Susin I, Nolli CP, Ferreira EM, Gentil RS, Polizel DM, Vaz Pires A, Alves SA, Bessa RJB. Effects of supplementation with vegetable oils, including castor oil, on milk production of ewes and on growth of their lambs. *J Anim Sci* 2018; 96(1):354–363. <https://doi.org/10.1093/jas/skx015>

Park SJ, Beak SH, Jung DJS, Kim SY, Jeong IH, Piao MY, Kang HJ, Fassah DM, Na SW, Yoo SP, Baik M. Genetic, management, and nutritional factors affecting intramuscular fat deposition in beef cattle — A review. *Asian-Austral J Anim Sci* 2018; 31(7):1043–1061. <https://doi.org/10.5713/ajas.18.0310>

Pulina G, Francesconi AHD, Stefanon B, Sevi A, Calamari L, Lacetera N, Dell’Orto V, Pilla F, Marsan PA, Mele M, Rossi F, Bertoni G, Crovetto GM, Ronchi, B. Sustainable ruminant production to help feed the planet. *Italian J Anim Sci* 2017; 16(1):140–171. <https://doi.org/10.1080/1828051X.2016.1260500>

Purchas RW, Davies AS, Abdullah AY. An objective measure of muscularity: changes with animal growth and differences between genetic lines of southdown sheep. *Meat Sci* 1991; 30(1):81–94. [https://doi.org/10.1016/0309-1740\(91\)90037-Q](https://doi.org/10.1016/0309-1740(91)90037-Q)

Rodrigues LKO, Mendes JUL, Jesus DS. Physicochemical analysis of *in - natura* and transesterified licuri (*Syagrus Coronata*) oil. *Res Soc Dev* 2020; 9(8):e554985793. <http://dx.doi.org/10.33448/rsd-v9i8.5793>

SAS. Statistical Analysis System. SAS user’s guide. Version 9.1, for Windows. Cary, NC, USA: SAS Institute Inc., 2003.

Santos HO, Howell S, Earnest CP, Teixeira FJ. Coconut oil intake and its effects on the cardiometabolic profile—A structured literature review. *Prog Cardio Dis* 2019; 62(5):436–443. <https://doi.org/10.1016/j.pcad.2019.11.001>

Santos PB, Santana Júnior HA, Araújo MJ, Oliveira AP, Freitas TB, Viana PT. Production and economic viability of feedlot beef cattle categories. *Acta Scient J Anim Sci* 2017; 39(2):195–199. <https://doi.org/10.4025/actascianimsci.v39i2.33452>

Sañudo C, Sierra I. Calidad de la canal en la especie ovina. *Ovino* 1986; 1:127–153.

Silva TM, Oliveira RL, Barbosa LP, Garcez Neto AF, Bagaldo AR, Pazzanese D. Preliminary study on meat quality of goats fed levels of licury oil in the diet. *Asian-Austral J Anim Sci* 2011; 24(8):1112 – 1119. <https://doi.org/10.5713/ajas.2011.11053>

Silva Sobrinho AG, Purchas RW, Kadim IT, Yamamoto SM. Características de qualidade da carne de ovinos de diferentes genótipos e idades ao abate. *Rev Bras Zootec* 2005; 34(3):1070–1078. <http://dx.doi.org/10.1590/S1516-35982005000300040>

Silva Sobrinho AG. Criação de ovinos. 3rd ed. Jaboticabal: Funep. 2001.

Vargas JE, Andrés S, López-Ferreras L, Snelling TJ, Yáñez-Ruíz DR, García-Estrada C, López S. Dietary supplemental plant oils reduce methanogenesis from anaerobic microbial fermentation in the rumen. *Scient Rep* 2020; 10(e1613):1–9. <https://doi.org/10.1038/s41598-020-58401-z>

Visentainer JV. Aspectos analíticos da resposta do detector de ionização em chama para ésteres de ácidos graxos em biodiesel e alimentos. *Quím Nova* 2012; 35(2):274–279. <https://doi.org/10.1590/S0100-40422012000200008>

Xu M, Du S, Wang J, Yu ZP, Harmon DL, Yao JH. Influence of rumen escape starch on pancreatic exocrine secretion of goats. *J Anim Phys Anim Nut* 2009. 93(1):122–129. <https://doi.org/10.1111/j.1439-0396.2007>