



UNIVERSIDAD DISTRITAL  
FRANCISCO JOSÉ DE CALDAS



## Callogenesis and Morphohistological Characterization of *Hancornia speciosa* Gomes

### Calogénesis y caracterización morfohistológica de *Hancornia speciosa* Gomes

Ana da Silva Ledo<sup>1</sup> , Caroline de Araújo Machado<sup>2</sup> , Annie Carolina Araújo de Oliveira<sup>2</sup> ,  
Maria de Fátima Arrigoni-Blank<sup>2</sup> , Evaristo Mauro de Castro<sup>3</sup> , Ana Veruska Cruz da Silva<sup>1</sup> 

Ledo, A. da S., Machado, C. de A., de Oliveira, A. C. A., Arrigoni-Blank, M. de F., de Castro, E. M., & Cruz da Silva, A. V. (2023). Callogenesis and morphohistological characterization of *Hancornia speciosa* Gomes. *Colombia Forestal*, 26(2), 5-14.

**Received:** July 18, 2022

**Approved:** April 10, 2023

#### Resumen

*Hancornia speciosa* Gomes belongs to the family Apocynaceae and is distributed across different regions of Brazil. The objective of this study was to evaluate callus induction and histodifferentiation in five mangaba accessions under different *in vitro* culture conditions. Five accessions from the Active Germplasm Bank of Mangaba (Embrapa, Brazil) were evaluated. *In vitro*-germinated plant seedlings were used for the excision of different explants (internode and node segments, and foliar section). These segments were inoculated in a culture medium containing different concentrations of 6-Benzylaminopurine (BA) and 2,4-Dichlorophenoxyacetic acid (2,4-D). There was a significant effect of accessions, treatments, and time on the callus mass (g). There was no callus induction in the absence of regulators. After 60 days of *in vitro* culture, all treatments exhibited a linear positive cellular growth. The highest callus mass was observed in the BI accession, in the presence of 22.62  $\mu\text{M}$  2,4-D and 11.10  $\mu\text{M}$  BA.

**Keywords:** *in vitro* tissue culture, cellular dedifferentiation, mangaba.

#### Abstract


*Hancornia speciosa* Gomes pertenece a la familia Apocynaceae, distribuida en diferentes regiones de Brasil. El objetivo de este trabajo fue evaluar la calogénesis y la histodiferenciación en cinco accesiones de mangaba bajo diferentes condiciones de cultivo *in vitro*. Se evaluaron cinco accesiones del Banco de Germoplasma Activo de Mangaba (Embrapa, Brasil). Las plántulas germinadas *in vitro* se utilizaron para la escisión de explantes (segmentos nodales e internodales y secciones foliares). Estos segmentos fueron inoculados en un medio de cultivo que contenía diferentes concentraciones de 6-benzilaminopurina (BA) y ácido 2,4-diclorofenoxiacético (2,4-D). Hubo un efecto significativo de las accesiones, los tratamientos y el tiempo del cultivo en la masa de los callos. No hubo inducción de callos en ausencia de reguladores. A los 60 días de cultivo *in vitro*, los otros tratamientos presentaron un crecimiento celular lineal positivo. La mayor masa de callo se observó en la accesión BI, en presencia de 22,62  $\mu\text{M}$  2,4-D y 11,10  $\mu\text{M}$  BA.

**Palabras clave:** cultivo de tejidos *in vitro*, dediferenciación celular, mangaba.

1 Embrapa Tabuleiros Costeiros. Aracaju, Sergipe, Brazil.

2 Universidade Federal de Sergipe, Programa de Pós-graduação em Agricultura e Biodiversidade. São Cristóvão, Sergipe, Brazil.

3 Universidade Federal de Lavras, Departamento de Botânica Aplicada, Lavras. Minas Gerais, Brazil.

 Corresponding author.

## INTRODUCTION

Native to Brazil, *Hancornia speciosa* Gomes is a fruitful species of the dicotyledonous group, which belongs to the Apocynaceae family (Silva *et al.*, 2017). The mangaba fruit exhibits great market potential in the tropical fruit sector, due to its wide variety of products and byproducts. It also has nutritional and pharmaceutical traits, such as a high vitamin C concentration (Santos *et al.*, 2017). Moreover, studies show that mangaba has a high antioxidant activity (Rufino *et al.*, 2009; Rufino *et al.*, 2010), potentially reducing the incidence of cancer, as well as of cardiovascular and cerebrovascular diseases.

Plant tissue culture provides an alternative to the use of entire specimens to produce bioactive secondary metabolites, thus allowing for active environmental control, regardless of seasonal and geographical conditions (Cardoso *et al.*, 2019). Alternatives to using plant material from natural resources are important, and the application of plant tissue culture techniques offers opportunities for a sustainable access to natural products (Wawrosch & Zotchev, 2021). One of the possibilities for producing valuable metabolites *in vitro* is the plant tissue callus induction method (Grąbkowska *et al.*, 2016, Cardoso *et al.*, 2019).

Plant growth regulators play a vital role in plant growth and development. Thus, 2,4-D (2,4-Dichlorophenoxyacetic acid), NAA (Naphthalene acetic acid), and other auxins are widely used for callus induction (Akram & Aftab, 2016). Studies have recently reported the *in vitro* formation of calluses in *H. speciosa*, concluding that NAA and BA regulators are superior for callogenesis induction when compared to others such as TDZ, IBA, and IAA (Prudente *et al.*, 2016; Oliveira *et al.*, 2016). However, information on the callogenesis potential of different mangaba accessions is scarce. Understanding the growth dynamics of plant cells is crucial in order to manipulate all related parameters, so that the potential of plants can be fully utilized (Puad & Abdullah, 2018).

The main objective of this research was to evaluate different combinations of 2,4-dichlorophenoxyacetic acid and benzylaminopurine for the induction of calluses in five mangaba accessions, establishing growth curves and performing histological analyzes.

## MATERIAL AND METHODS

### Location and plant material

Accessions from the Active Germplasm Bank of Mangaba, located in Itaporanga d'Ajuda, SE, Brazil (11°06'40"N and 37°11'15"E), were selected, given this bank's ability to provide both mature fruits and information on the location of origin (Accessions: AB-Água Boa – Pará/0°46'31"N; 48°31'05"E; CA-Costa Azul – Bahia/11°33'32"N; 47°47'06"E; BI-Barra do Itariri – Bahia/11°48'39"N; 37°36'40"E; PT-Indiaroba – Sergipe/11°31'10"N; 37°30'47"E; TC-Terra Caída – Sergipe/11°46'31"N; 37°30'47"E). After harvesting the mature fruits of the selected accessions, they were pulped to obtain the seeds, which remained at room temperature for 24 hours. In a flow chamber, the seeds were submitted to asepsis with immersion in 70% alcohol for 60 seconds and in 2-2.5% sodium hypochlorite for 15 minutes, as well as to triple washing in sterile water. The sterile seeds were inoculated in an MS culture medium (Murashige & Skoog, 1962), with 30 g.L<sup>-1</sup> sucrose and 3.5 g.L<sup>-1</sup> Phytigel® (Ref. P8169, Sigma). The pH was adjusted to 5.8, and the medium was then autoclaved at 121 °C for 15 minutes. Explants obtained from *in vitro* aseptic seedlings were used to induce callogenesis.

The experiments were carried out at the Plant Tissue Culture Laboratory of Embrapa Tabuleiros Costeiros, in Aracaju, SE, Brazil, and at the Electronic Microscopy and Ultrastructural Analysis Laboratory of the Federal University of Lavras, MG, Brazil.

## Induction of callogenesis

After 45 days of seed inoculation, internode and node segments (1 cm long) and foliar sections of the central leaf zone (0.5 cm<sup>2</sup>) were excised from the *in vitro* seedlings. For callus induction, the internode and nodal segments were inoculated in a longitudinal position, and the foliar section with abaxial side in contact with the MS culture medium. The medium was supplemented with 30 g.L<sup>-1</sup> sucrose and 3.5 g.L<sup>-1</sup> Phytigel® (Ref. P8169, Sigma) containing respectively different combinations of 2,4-dichlorophenoxyacetic acid (2,4-D-Ref. D7299, Sigma) and benzylaminopurine (BA-Ref. B3408, Sigma). The pH of the media was adjusted to 5.8 before autoclaving at 121 °C for 15 minutes.

The explants were weighed at 10, 20, 30, 40, and 60 days of *in vitro* culture, so the callus mass could be measured. The cultures were maintained in a growth room at 25±2 °C, in the absence of light to avoid phenolic exudation (Amente & Chimdessa, 2021) throughout the experimental period.

The experiment was completely randomized; it employed a 5 (accessions) × 3 (explants) × 5 (growth regulator treatments) × 5 (days) factorial design. Each treatment had four replicates, each containing one explant. The data were subjected to an analysis of variance, and a Scott–Knott test at 5% probability was performed on the means of the qualitative treatments, which were also subjected to a regression analysis.

For the statistical analyses, due to the absence of callus formation in T0 and their high oxidation in T4 treatments, only three combinations of growth regulators were considered for callus mass (g) evaluation. The SISVAR statistical program (Ferreira, 2019) was used.

## Plotting the callus growth curves

To determine the callus growth curves originating from internode, node, and leaf explants, explants were grown in a culture medium able to provide maximum callus production.

The calluses were weighed on a precision scale, on the first day of inoculation (time 0) and then in 10-day intervals for 60 days, with three replicates per evaluation time. Each replicate consisted of a Petri dish with four explants. The growth curve was plotted from the average of the three measurements of fresh mass for each evaluation time (0, 10, 20, 30, 40, and 60 days).

## Histological studies

This study followed the methodology of Alves *et al.* (2008). For the histological test, callus samples from different accessions and types of explants (node, internode, and leaf segments) were used. The samples were kept in Karnovsky's modified solution (1965). Dehydration was performed using graded ethyl alcohol series. Then, the callus was infiltrated in pure resin and embedded in Leica® Histo-resin. The slides were analyzed under an optical microscope (Zeiss/Jenamed2, Germany), in which photomicrographs (AxioCam software, Zeiss, Germany) of the prepared slides were obtained.

The analysis of the callus samples under a scanning electron microscope also employed the methodology developed by Alves *et al.* (2008). To fix the samples, a glutaraldehyde (25%) (Ref. 104239, Merck), paraformaldehyde (16%) (Ref. P6148, Sigma-Aldrich), sodium cacodylate buffer (0.2 M, pH 7.2) (Ref. C0250, Sigma), CaCl<sub>2</sub> (0.1 M) (Ref. C1016, Sigma-Aldrich), and distilled water (Karnovsky, 1965 modified) solution was used. For the assays, the samples were washed in 0.05 M cacodylate buffer and post fixed in 1% osmium tetroxide (Ref.251755, Sigma-Aldrich®). Afterwards, they were progressively dehydrated in acetone subjected to the critical point of CO<sub>2</sub> desiccation in a BAL-TEC equipment, CPD-030 (Bal-Tec, Los Angeles, California, USA), fixed to a metallic support and covered with metallic gold (10 nm) (01O1003.01.AA, Synth<sup>©</sup>) while using a BAL-TEC apparatus, SCD-050 (Bal-Tec, Los Angeles, California, USA). The sample was prepared, observed, and electromicrographed in a LEO EVO 40

XVP scanning electron microscope (Carl Zeiss AG, Germany).

## RESULTS

### *In vitro* callus induction

Regarding callus mass, a significant difference in interaction between the growth regulator treatment and the accession was observed. There was no significant effect of the explant type on callus mass. Among the accessions studied, no statistical difference was identified.

Callus fresh mass had a linear and positive behavior with respect to all factors combined. There was a greater mass increase over time in the 45.25

$\mu\text{M}$  2,4-D and 11.10  $\mu\text{M}$  BA (T3) combination for all accessions (Table 1). The highest callus mass was observed in the BI accession at 60 days, in the presence of 22.62  $\mu\text{M}$  2,4-D and 11.10  $\mu\text{M}$  BA. Under the same conditions, BI had a callus mass of approximately 1.4 g, in comparison with 0.1 g in the AB accession. In all treatments, a friable callus of light-yellow color was observed at all evaluation times.

### Callus growth curve

To plot the callus growth curve from different explants (leaf, node, and internode segments) evaluated for up to 60 days of *in vitro* culture, samples induced in the presence of 45.25  $\mu\text{M}$  2,4-D + 22.20  $\mu\text{M}$  BA were used, as well as accessions with the greatest

**Table 1.** Effect of growth regulators on the formation of fresh callus mass (g) in mangaba accessions at 10, 20, 30, 40, and 60 days of *in vitro* culture. (T1) 4.52  $\mu\text{M}$  2,4-D + 2.22  $\mu\text{M}$  BA; (T2) 22.62  $\mu\text{M}$  2,4-D + 11.10  $\mu\text{M}$  BA; (T3) 45.25  $\mu\text{M}$  2,4-D + 22.20  $\mu\text{M}$  BA. Accessions: Água Boa (AB); Barra do Itariri (BI); Costa Azul (CA); Pontal (PT); Terra Caída (TC).

Days	10	20	30	40	60	Regression equation
AB						
T1	0.0158b	0.0742b	0.1350b	0.2375a	0.2983a	$\hat{y}^{**} = 0.0059x - 0.0369$ ; $R^2 = 0.9638$
T2	0.0192a	0.0383a	0.0925a	0.1283a	0.1583a	$\hat{y}^{**} = 0.003x - 0.0077$ ; $R^2 = 0.9453$
T3	0.0442d	0.1108d	0.2400c	0.4425b	0.8450a	$\hat{y}^{**} = 0.0165x - 0.1915$ ; $R^2 = 0.9696$
BI						
T1	0.0300c	0.0567c	0.1558b	0.2383b	0.4317a	$\hat{y} = 0.0083x - 0.0843$ ; $R^2 = 0.9801$
T2	0.0200c	0.0575c	0.1767b	0.3208b	0.6250a	$\hat{y}^* = 0.0126x - 0.1618$ ; $R^2 = 0.9703$
T3	0.0292d	0.1283d	0.4225c	0.7858b	1.2950a	$\hat{y}^* = 0.0267x - 0.3223$ ; $R^2 = 0.9822$
CA						
T1	0.0250a	0.0500a	0.0816a	0.1475a	0.2125a	$\hat{y}^{**} = 0.0039x - 0.0224$ ; $R^2 = 0.9803$
T2	0.0233b	0.0491b	0.0850b	0.1425b	0.3116a	$\hat{y}^{**} = 0.0058x - 0.0635$ ; $R^2 = 0.9450$
T3	0.0408d	0.1266d	0.3041c	0.5333b	0.8350a	$\hat{y}^{**} = 0.0166x - 0.1644$ ; $R^2 = 0.9874$
PT						
T1	0.0150b	0.0491b	0.0800b	0.1150b	0.2783a	$\hat{y}^{**} = 0.0052x - 0.0576$ ; $R^2 = 0.9384$
T2	0.0250c	0.0433c	0.1266b	0.1983b	0.4941a	$\hat{y}^{**} = 0.0065x - 0.0487$ ; $R^2 = 0.9223$
T3	0.0108b	0.0450b	0.2083a	0.2116a	0.3183a	$\hat{y}^{**} = 0.0099x - 0.1426$ ; $R^2 = 0.9480$
TC						
T1	0.0252a	0.0400a	0.0625a	0.0733a	0.1258a	$\hat{y}^{**} = 0.002x - 0.0002$ ; $R^2 = 0.9861$
T2	0.0208b	0.0308b	0.0683b	0.1433a	0.2066a	$\hat{y}^{**} = 0.004x - 0.0351$ ; $R^2 = 0.9577$
T3	0.0358d	0.0891d	0.2716c	0.4125b	0.7541a	$\hat{y}^{**} = 0.004x - 0.0351$ ; $R^2 = 0.9577$

\* Means followed by the same lowercase letters did not differ from each other in the Scott-Knott test ( $p < 0.05$ ). \* Significance:  $p < 0.05$ . CV(%) 90.58


contribution to the accumulation of callus fresh mass (Table 1) as absolute values. Different stages of growth in induced callogenesis were evaluated in different explants during the 60 days of culture (Figure 1).

There is a difference in the callus mass between the evaluated explant types. In all the studied accessions, the node explant had a greater callus mass formation. In the AB, BI, CA, PT, and TC accessions, phases I (10 to 20 days), II (20 to 40

days), and III (40 to 60 days) were reached within the same experimental periods. The stationary and decline or deceleration phases were not observed until 60 days had passed.

## Histological analysis

The callus observed in all treatments were friable and yellowish and had increasing mass increments (g). The following morphological structures




**Figure 1.** Callus growth in mangaba explants (Leaf ◆, Node ■, and Internode ▲) of AB, BI, CA, PT, and TC accessions in an MS medium with 45.25  $\mu\text{M}$  2,4-D + 22.20  $\mu\text{M}$  BA at 10, 20, 30, 40, and 60 days of culture. I - Lag phase, II - Exponential phase and III - Linear phase. Accessions: Água Boa (AB); Barra do Itariri (BI); Costa Azul (CA); Pontal (PT); Terra Caída (TC).



were observed in BI accession, which are also identified in [Figure 2](#): round agglomerated cells (rac) – rounded cells, dispersed or associated with agglomerated cells; agglomerated elongated cells (aec) – elongated cells, dispersed or associated with cellular agglomerations; small agglomerated

cells (sac) – formed by small and isodiametric cells; small scattered (sc) cells – comprised of small, dispersed cells; and vascular bundle (vb) – an anatomical structure of the stem. Cells in de-differentiation were observed in all the studied accessions.



**Figure 2.** Anatomical structures in different mangaba in vitro cultures after 30 days in an MS medium supplemented with 45.25  $\mu\text{M}$  2,4-D + 22.20  $\mu\text{M}$  BA. (A) foliar AB; (B) internode AB; (C) node AB; (D) foliar BI; (E) internode BI; (F) node BI; (G) foliar CA; (H) internode CA; (I) node CA; (J) foliar PT; (K) internode PT; (L) node PT; (M) foliar TC; (N) internode TC; (O) node TC. Bar 100  $\mu\text{m}$ . Type: (rac) round agglomerated cells, rounded cells dispersed or associated with cellular clusters; (aec) agglomerated elongated cells, elongated cells dispersed or associated with cell agglomerates; (sac) small, agglomerated cells, small and isodiametric cells, forming agglomerates; (sc) small scattered, formed by small and dispersed cells; (vb) vascular bundle, structure of the vascular bundle of the stem. Accessions: Água Boa (AB); Barra do Itariri (BI); Costa Azul (CA); Pontal (PT); Terra Caída (TC).

In the leaf segment, rac and aec cells were observed (Figure 2D). The callus from the node segment (Figure 2E) had sac cells and a vb anatomical structure, with the beginning of dedifferentiated cell formation. This type of cell was also observed in all studied accessions of *H. speciosa*. Regarding the formation of callus from internode segments (Figure 2E, N), there was an initial development of dedifferentiated cells, and it was possible to identify the other anatomical structures present in the plant stem area. The callus exhibited a similar pigmentation and consistency across accessions, despite the fact that different treatments modified the yellow coloration intensity.

## DISCUSSION

### *In vitro* callus induction

*In vitro* callus induction depends on several biotic and abiotic factors such as culture medium composition, plant growth regulators, sucrose compositions, pH, and incubation conditions. These factors may influence the induction potential and morphology of callogenesis (Osman *et al.*, 2016). Callus induction from different explants of *Mandevilla guanabara* (Apocynaceae) (apical meristems, leaf, node segments, internode segments, roots, and epicotyl and hypocotyl) was performed by Cordeiro *et al.* (2014), who evaluated cell development in different explants and culture medium conditions. Their study is a good reference for the selection of the explants and growth regulators able to induce higher callus growth rates.

Mustafa *et al.* (2011) reported that friable callus is mainly used for cells under suspension and for the study of metabolites, given that, in a culture medium, the cells rapidly divide and disperse. Auxins are capable of initiating cell division and controlling the processes of cell growth and elongation (Perrot-Rechenmann, 2010). The combination of endogenous phytohormones and 7.74  $\mu\text{M}$

2,4-D in a WPM culture medium stimulated the cell division of *H. speciosa* parenchyma regions, forming callus from internode segments (Prudente *et al.*, 2016). Callus induced from the leaf, young stem, and shoot tip of *Melientha suavis* Pierre were observed by Premjet *et al.* (2020) in an MS basal medium supplemented with various formulations of 2,4-D and 6-BA. Another promising auxin in the induction of callus in species of the family Apocynaceae is NAA. Studies with the apical parts of *Rauwolfia serpentina* L. in MS media enriched with NAA resulted in excellent callus development (Varnika *et al.*, 2020). The balance between auxin and cytokinin fosters a greater callus mass in different segments of the plant. The color and texture of the callus varied depending on the type of explants, as well as on the concentrations of growth regulators used in both media (Das *et al.*, 2018). Therefore, more studies are needed to define the growth regulator.

### Callus growth curve

Callus growth curves have five phases: lag (I), exponential (II), linear (III), stationary (IV), and deceleration (V) (Bathia *et al.*, 2015). In the AB, BI, CA, PT, and TC accessions, phases I (10 to 20 days), II (20 to 40 days), and III (40 to 60 days) were reached within the same experimental periods. The stationary and decline or deceleration phases were not observed until 60 days had gone by. Thus, determining the *in vitro* callus growth curve is important for understanding the time of maximum mass increment (in days).

In studies with *Aspidosperma cylindrocarpon* (Apocynaceae), cell suspension cultures showed a four-day growth lag phase for 15 days, and, after 20 days, the cells showed a stationary growth (Cornelio, 2018). The growth acceleration of *Bletilla striata* callus started after seven days, gradually decreasing thereafter until day 24 of the cultivation period (Pan *et al.*, 2020). This result is an important contribution to *in vitro* studies on

mangaba cell suspensions, and it indicates the high induction and proliferation capacity of *in vitro* callus mass over long periods of time.

### Histological analysis

All studied cell types are more likely to form embryogenic callus in *Byrsonima verbascifolia* (Malpighiaceae) (Chiavegatto *et al.*, 2015). Oliveira *et al.* (2016) also observed the formation of friable calluses on the basis of node segments with variation in callus pigmentation as a function of the growth regulator concentration. *H. speciosa* callus cultivated in WPM media in the presence of BA and 2,4-D were friable and easy to dissociate, and they initially exhibited a white coloration (Prudente *et al.*, 2016).

In histological studies on callus of *Asteracantha longifolia* Nees, Kumar and Nandi (2015) observed that the meristematic regions were formed by small cells with densely agglomerated nuclei. These aec-like cell structures were observed in this research when inducing callus in the node segment of the PT accession (Figures B and C). Bhagya and Chandrashekar (2013) detected the initial formation of callus from the exchange region of *C. peltata* stems, with disorganized cells arranged around a prominent nucleus. The results obtained in this work for *H. speciosa* are important due to the lack of studies on callogenesis and histology in different accessions.

### CONCLUSIONS

After 60 days of *in vitro* culture, all treatments exhibited a linear positive cellular growth. The BI accession showed the maximum increase in callus mass in the presence of 22.62  $\mu\text{M}$  2,4-D and 11.10  $\mu\text{M}$  BA. The callus mass from the foliar section and the node and internode segments in the MS medium with 45.25  $\mu\text{M}$  2,4-D + 5.0  $\text{mg}\cdot\text{L}^{-1}$  BA induced morphological alterations in the AB, BI, CA, TC, and PT accessions.

### ACKNOWLEDGEMENTS

The authors would like to thank the CAPES (Coordination of Superior-Level Staff Improvement, Brazil), the CNPq (National Council for Scientific and Technological Development, Brazil), and Embrapa (Brazilian Agricultural Research Corporation) for their financial support and scholarship.

### CONFLICT OF INTEREST

The authors declare no conflict of interest.

### REFERENCES

- Akram, M., & Aftab, F. (2016). Establishment of embryogenic cultures and efficient plant regeneration system from explants of forced softwood shoots of teak (*Tectona grandis* L.). *Horticultural Plant Journal*, 2(5), 293-300.  
<https://doi.org/10.1016/j.hpj.2017.01.008>
- Alves, E., Leite, B., Marucci, R. C., Pascholati, S. F., Lopes, J. R. S., & Andersen, P. C. (2008). Retention sites for *Xylella fastidiosa* in four sharpshooter vectors (*Hemiptera cicadellidae*) analyzed by scanning electron microscopy. *Current Microbiology*, 56(5), 531-8.  
<https://doi.org/10.1007/s00284-008-9119-7>
- Amente, G., & Chindessa, E. (2021). Control of browning in plant tissue culture: A review. *Journal of Scientific Agriculture*, 5, 67-71.  
<https://doi.org/10.25081/jsa.2021.v5.7266>
- Bhagya, N., & Chandrashekar, K. R. (2013). Effect of growth regulators on callus induction from *Cyclea peltata* (Lam.) Hook. f. Thoms. *Asian Journal Pharmaceutica and Clinical Research*, 6(4), 85-88.  
<https://innovareacademics.in/journals/index.php/ajpcr/article/view/497>
- Bathia, S. (2015). Plant tissue culture. In S. Bathia, T. Bera, & K. Sharma (Eds.), *Modern Applications of Plant Biotechnology in Pharmaceutical Sciences* (pp. 31-107). Academic Press.  
<https://doi.org/10.1016/C2014-0-02123-5>



- Cardoso, J. C., Oliveira, M. E. B. S., & Cardoso. F. de C. I.** (2019). Advances and challenges on the in vitro production of secondary metabolites from medicinal plants. *Horticultura Brasileira*, 37(2), 124-132. <https://doi.org/10.1590/S0102-053620190201>
- Cordeiro, S. Z., Simas, N., Henriques, A. B., & Sato, A.** (2014). Micropropagation and callogenesis in *Mandevilla guanabara* (Apocynaceae), an endemic plant from Brazil. *Crop Breeding and Applied Biotechnology*, 14, 108-115. <https://doi.org/10.1590/1984-70332014v14n2a19>
- Cornelio, M. N.** (2018). *Chemical analysis and establishment of in vitro culture of Aspidosperma cylindrocarpon* Muell. Arg. and *Aspidosperma polyneuron* Muell. Arg. USP [Doctoral thesis, Universidade de São Paulo]. <https://doi.org/10.11606/T.46.2018.tde-31072018-151847>
- Chiavegatto, R. B., Castro, A. H. F., Marçal, M. G., & Pádua, M. S.** (2015). Cell Viability, mitotic index and callus morphology of *Byrsonima verbascifolia* (Malpighiaceae). *Tropical Plant Biology*, 8, 87-97. <https://doi.org/10.1007/s12042-015-9150-3>
- Das, P., Tanti, B., & S. K. Borthakur, S. K.** (2018). In vitro callus induction and indirect organogenesis of *Brucea mollis* Wall. ex Kurz – A potential medicinal plant of Northeast India. *South African Journal of Botany*, 119, 203-211. <https://doi.org/10.1016/j.sajb.2018.09.012>
- Ferreira, D. F.** (2019). A computer analysis system to fixed effects split plot type designs: Sisvar. *Brazilian Journal of Biometrics*, 37(4), 529-535. <https://doi.org/10.28951/rbb.v37i4.450>
- Grąbkowska, R., Matkowski, A., Grzegorzczuk-Karolak, I., & Wysokińska, H.** (2016). Callus cultures of *Harpagophytum procumbens* (Burch.) DC. Ex Meisn.; production of secondary metabolites and antioxidant activity. *South African Journal of Botany*, 103, 41-48. <https://doi.org/10.1016/j.sajb.2015.08.012>
- Karnovsky, M. J.** (1965). A formaldehyde-glutaraldehyde fixative of high osmolality for use in electron microscopy. *Journal of Cell, Biology*, 27(2), 137-138. <http://www.jstor.org/stable/1604673>
- Kumar, M. S., & Nandi, S. C.** (2015). High frequency plant regeneration with histological analysis of organogenic callus from internode explants of *Asteracantha longifolia* Nee. *Journal of Genetic Engineering and Biotechnology*, 76(1), 31-37. <https://doi.org/10.1016/j.jgeb.2014.12.002>
- Murashige, T., & Skoog, F.** (1962). A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*, 15(3), 473-497. <https://doi.org/10.1111/j.1399-3054.1962.tb08052.x>
- Mustafa, N. R., Winter, W., Iren, F., & Verpoorte, R.** (2011). Initiation, growth and cryopreservation of plant cell suspension cultures. *Nature Protocols*, 6(6), 715-742. <https://doi.org/10.1038/nprot.2010.144>
- Oliveira, K. S., Freire, F. A. M., & Aloufa, M. A. I.** (2016). Efeito de 6-benzilaminopurina e ácido naftalenoacético sobre propagação in vitro de *Hancornia speciosa* Gomes. *Revista Floresta*, 46, 335-342. <https://doi.org/10.5380/RF.V46I3.43993>
- Osman, N. I., Sidik, N. J., & Awal, A.** (2016). Effects of variations in culture media and hormonal treatments upon callus induction potential in endosperm explant of *Barringtonia racemosa* L. *Asian Pacific Journal Tropical Biomedicine*, 6, 143-147. <https://doi.org/10.1016/j.apjtb.2015.10.007>
- Pan, Y., Li, L., Xiao, S., Chen, Z., Sarsaiya, S., & Zhang, S.** (2020). Callus growth kinetics and accumulation of secondary metabolites of *Bletilla striata* Rchb.f. using a callus suspension culture. *PLoS ONE*, 15(2), e0220084. <https://doi.org/10.1371/journal.pone.0220084>
- Perrot-Rechenmann, C.** (2010). Cellular responses to auxin: Division versus expansion. *Cold Spring Harbor Perspectives in Biology*, 2(5), 1-15. <https://doi.org/10.1101/cshperspect.a001446>
- Premjet, D., Obeng, A. K., & Premjet, S.** (2020). Establishment of callus culture of *Melientha suavis* Pierre. *Chilean Journal of Agricultural Research*, 8(3), 459-465. <https://doi.org/10.4067/S0718-58392020000300459>
- Prudente, D. O., Paiva, R., Nery, F. C., Máximo, F. C., Flausino, W. P., & Coutinho L. C.** (2016).

Organogênese indireta *in vitro* de *Hancornia speciosa* Gomes. *Bioscience Journal*, 32(3), 721-729.

<https://pdfs.semanticscholar.org/dfc1/129120d-469d290e00e42163936e0e3acf977.pdf>

**Puad N. I. M. & Abdullah, T. A.** (2018). Monitoring the growth of plant cells in suspension culture. In A. Amid, S. Sulaiman, D. N. Jimat, N. Fadhillah M. Azmin (Eds.), *Multifaceted Protocol in Biotechnology* (pp. 203-214). Springer.

[https://doi.org/10.1007/978-981-13-2257-0\\_17](https://doi.org/10.1007/978-981-13-2257-0_17)

**Rufino, M. S. M., Alves, R. E., de Brito, E. S., Pérez-Jiménez, J., Saura-Calixto, F., & Mancini-Filho, J.** (2010). Bioactive compounds and antioxidant capacities of 18 non-traditional tropical fruits from Brazil. *Food Chemistry*, 121, 996-1002.

<https://doi.org/10.1016/j.foodchem.2010.01.037>

**Rufino, M. S. M., Fernandes, F. A. N., Alves, R. E. & de Brito, E. S.** (2009). Free radical-scavenging behaviour of some north-east Brazilian fruits in a DPPH system. *Food Chemistry*, 114, 693-695.

<https://doi.org/10.1016/j.foodchem.2008.09.098>

**Santos, P. S., Freitas, L. dos S., Santana, J. G. S., Muniz, E. N. & Rabbani, A. R. C.** (2017). Genetic diversity and the quality of mangabeira tree fruits (*Hancornia speciosa* Gomes – Apocynaceae), a native species from Brazil. *Scientia Horticulturae*, 226, 372-378.

<https://doi.org/10.1016/j.scienta.2017.09.008>

**Silva, A. V. C., Soares, N. R. A., Ledo, A. da S. & Costa, T. S.** (2017). Uses and technological prospects for the mangaba, a native fruit of Brazil. *African Journal of Biotechnology*, 16(7), 302-311.

<http://dx.doi.org/10.5897/AJB2016.15786>

**Varnika, S., Varnika V., Sharma, R., Singh, A., Shalini, S. & Sharma, N.** (2020). Micropropagation and screening of phytochemicals present among in vitro raised and wild plants of *Rauvolfia serpentina*. *Walailak Journal Sciences & Technology*, 17(11), 1177-1193.

<http://dx.doi.org/10.48048/wjst.2020.6492>

**Wawrosch, C. & Zotchev, S. B.** (2021). Production of bioactive plant secondary metabolites through in vitro technologies-status and outlook. *Applied Microbiology and Biotechnology*, 105(18), 6649-6668.

<http://dx.doi.org/10.1007/s00253-021-11539-w>

