BEEF QUALITY FROM ANIMALS RAISED IN THREE DIFFERENT GRAZING SYSTEMS

Suelen Cristina S. Carvalho^{1*}, Vanessa C. Francisco², Althieres J. Furtado³, Sophia A. M.

Chamilete¹, Kalisa F. Silva¹, Paulo H. M. Rodrigues³, Renata T. Nassu²

¹Faculdade de Ciências Farmacêuticas (FCFAr)/UNESP, Araraquara-São Paulo, Brazil ²Embrapa Pecuária Sudeste, São Carlos, SP, Brazil.

³ Faculdade de Medicina Veterinária e Zootecnia (FMVZ)/USP, Pirassununga, SP, Brazil.
*Corresponding author email: renata.nassu@embrapa.br

I. INTRODUCTION

The increase in greenhouse gas emissions (GHG) is considered one of the leading causes of global warming, and enteric methane (CH₄) from beef cattle represents an important emission source in the agricultural sector [1]. In the pigeon pea (*Cajanus cajan* cv. BRS Mandarin) and *Urochloa spp*. consortium, the legume acts as a protein bank and green manure, reducing GHG emissions [2]. Intensified systems and nutritional strategies improve production efficiency and can influence meat quality. The present study aimed to evaluate the quality of aged beef produced in three pasture systems: degraded, recovered and consortium with pigeon pea.

II. MATERIALS AND METHODS

Twenty-seven *Nellore* steers, initially weighing 221±7 kg and 15-16 months of age, were distributed into three treatments with three replications: degraded pasture (*Urochloa spp.*), recovered pasture (*Urochloa spp.* fertilized with 200 kg of N/ha⁻¹) and consortium pasture with pigeon pea (*Cajanus cajan* cv. BRS Mandarin) and *Urochloa spp.*, for two years. The animals were slaughtered in a commercial slaughterhouse following standards and guidelines for the industrial and sanitary inspection of products of animal origin. After 24 hours in a cold chamber, samples of the *Longissimus thoracis* muscle (between the 12th and 13th rib) from the left half carcass of each animal were removed and taken to the Meat Analysis Laboratory of Embrapa Southeastern Livestock, São Carlos, São Paulo, Brazil. 2.5 cm thick steaks were aged for 14 days at 0-2 °C, and the following analyses were performed at 0, 7, and 14 days: pH; instrumental color measurement, using a portable Hunter Lab Miniscan XE colorimeter to measure variations within the CIE system color space (L*, a*, b*) [3]; water holding capacity (WHC) [4], cooking loss (CL) and shear force (WBSF) [5]. The obtained data were analyzed by analysis of variance (ANOVA) using PROC MIXED in SAS software, and mean comparisons were performed using Tukey's test (5%).

III. RESULTS AND DISCUSSION

The results obtained for the meat quality traits are presented in Table 1. Based on the analyzed data, the grazing systems had no effect (p>0.05). On the other side, there was a significant effect (p<0.01) of aging times on all the meat quality traits, except for a^{*}.

Effects		Variables						
Grazing systems	Aging Time	Color		pН	CL (%)	WBSF (kgf)		
	(days)	a*	b*					
Degraded		13.32	10.80	6.05	24.23	8.35		
Recovered		13.74	10.46	6.05	25.13	7.59		
Pigeon pea+Urochloa spp.		14.09	11.32	5.97	23.01	7.71		
	0	13.81	10.61 ^B	5.96 ^B	26.77 ^A	9.34 ^A		
	7	13.66	11.26 ^A	6.04 ^A	23.00 ^B	8.25 ^B		
	14	13.68	10.70 ^B	6.07 ^A	22.60 ^B	6.06 ^C		
Average		13.71	10.86	6.02	24.12	7.89		
SEM		0.20	0.25	0.04	0.71	0.40		

Table 1 - Quality traits of beef produced in different pasture systems.

Statistical Probabilities (P value)									
Treatment	0.2469	0.3420	0.6702	0.2546	0.6693				
Aging time	0.5608	<.0001	<.0001	0.0003	<.0001				
Treat.*AgingTime	0.3285	0.6256	0.8625	0.7061	0.9371				

Lowercase letters differ among grazing systems (P≤0.05). Uppercase letters differ (P≤0.05) among aging times. CL = cooking loss; WBSF = Warner Bratzler Shear Force

A significant interaction was observed in L* (P=0.0484) and WHC (P=0.0241) of the meat produced in the pigeon pea + *Urochloa spp.* treatment at the 7-day aging time (Figure 1).



Figure 1 - Effect of the interaction between grazing systems and aging times for luminosity (L*) (A) and water holding capacity (WHC) (B) of the produced beef. Lowercase letters differ among grazing systems (P \leq 0.05). Uppercase letters differ (P \leq 0.05) among aging times.

For L*, it is observed a significant difference (p<0.05) in zero and seven days, and not at 14 days for the grazing systems (Figure 1A). For WHC (Figure 1B), the behavior between aging times and grazing systems was different: meat of animals from the recovered system showed a lower value on day zero and higher at day 7, indicating that depending on aging time, there was a difference between WHC (%) values in the grazing systems.

IV. CONCLUSION

The pigeon pea+*Urochloa spp* consortium system, in addition to being a strategy to mitigate GHG emissions, generally did not affect beef quality, except for luminosity and water holding capacity.

ACKNOWLEDGEMENTS

The authors acknowledge the FAPESP (2017/20084-5), coordinated by the Faculty of Veterinary Medicine and Animal Science (FMVZ/USP—Pirassununga/SP, Brazil), with collaboration from the Institute of Animal Science (IZ—Nova Odessa/SP, Brazil) and Embrapa Southeastern Livestock (São Carlos/SP, Brazil); CAPES—funding code 001 and CNPq.

REFERENCES

- Ouatahar, L.; Bannink, A.; Lanigan, G.; & Amon, B. (2021) Modelling the effect of feeding management on greenhouse gas and nitrogen emissions in cattle farming systems. Science of the Total Environment, 776, 145932.
- da Silva, A.; de Sousa Santos, F. L.; de Mattos Barretto, V. C.; de Freitas, R. J. & Kluthcouski, J. (2018) Recuperação de pastagem degradada pelo consórcio de milho, Urochloa brizantha cv. Marandu e guandu. Revista de Agricultura Neotropical, 5: 39-47.
- 3. AMSA. (2012) American Meat Science Association meat color measurement guidelines. Am. Meat Sci.
- 4. Hamm, R. (1986) Functional properties of the myofibrillar system and their measurement. In: Bechtel PJ. (Ed.). Muscle as food. Orlando: Academic Press. 135-199.
- 5. Wheeler, T. L.; Shackelford, S. D.; & Koohmaraie, M. (1997) Standardizing collection and interpretation of Warner-Bratzler shear force and sensory tenderness data. In Proc. recip. meat conf 50: 68-77.