

Natural Sciences

Original article

Hemato-immunological responses of juvenile tambaqui (*Colossoma macropomum*) after commercial clay dietary supplementation

Respuestas hematoimmunológicas de juveniles de tambaqui (*Colossoma macropomum*) después de la suplementación dietética con arcilla comercial

Yuri Ian Carvalho Furtado^{1,*}, Fernanda Layza Souza de Souza², Mosa Valdina Ferreira Moreira¹, Amanda Mendes Pacheco², Clara Brito Salomão², Eliane Tie Oba Yoshioka^{1,3}

¹ Programa de Pós-Graduação em Biodiversidade Tropical, Universidade Federal do Amapá – UNIFAP, Brasil

² Universidade do Estado do Amapá – UEAP, Colegiado de Engenharia de Pesca, Brasil

³ Empresa Brasileira de Pesquisa Agropecuária, Embrapa Amapá, Brasil

Abstract

The present study evaluated the physiological and biochemical responses of tambaqui (*Colossoma macropomum*) to different dietary supplementation levels of bentonite. We evaluated four treatments with three replicates (10 fish *per* tank). The sodium bentonite adsorbent (Buntech) was incorporated into the commercial feed at three levels in addition to the control group (0%, 0.5%, 1.0%, and 2%). After 10 and 20 days of feeding, we collected blood samples from five fish in each replicate for analysis. We observed the changes in hematocrit, hemoglobin, erythrocytes, and hematimetric indices of the tambaqui. Their glucose concentration, total proteins, total cholesterol, triglycerides, and albumin differed significantly due to the bentonite inclusion. We also recorded reductions in *C. macropomum* defense cells due to physiological damage caused by the adsorbent agent, resulting in thrombocytopenia, leukopenia, lymphocytopenia, and monocytopenia. Such reductions indicated a reaction and migration to inflammatory foci; the addition of this product did not reflect any improvement in the hematological-biochemical profile of this fish species. Our results suggest that sodium bentonite dietary supplementation has deleterious effects on *C. macropomum* and is not indicated as a food additive.

Keywords: *Colossoma macropomum*; Hematology; Immunostimulant; Freshwater fish.

Resumen

El presente estudio evaluó las respuestas fisiológicas y bioquímicas del tambaqui (*Colossoma macropomum*) a diferentes niveles de suplementación dietética con bentonita. Se evaluaron cuatro tratamientos con tres réplicas (10 peces por tanque). El adsorbente de bentonita sódica (Buntech) se incorporó al alimento comercial en tres niveles además del grupo control (0 %, 0,5 %, 1,0 % y 2 %). A los 10 y los 20 días de alimentación, se recolectaron muestras de sangre de cinco peces de cada réplica para su análisis. Se observaron cambios en el hematocrito, la hemoglobina, los eritrocitos y los índices hematimétricos del tambaqui. La concentración de glucosa, las proteínas totales, el colesterol total, los triglicéridos y la albúmina difirió significativamente debido a la inclusión de la bentonita. Hubo reducciones en el número de las células de defensa de *C. macropomum* debido al daño fisiológico causado por el agente adsorbente, lo que resultó en trombocitopenia, leucopenia, linfocitopenia y monocitopenia. Dichas reducciones sugerían una reacción y la migración a focos inflamatorios; la adición del producto no reflejó una mejoría en el perfil hematológico-bioquímico de esta especie de pez. Nuestros resultados indican que la suplementación dietética con bentonita sódica tiene efectos nocivos sobre *C. macropomum*, por lo cual no está indicada como aditivo alimentario.

Palabras clave: *Colossoma macropomum*; Hematología; Inmunoestimulante; Peces de agua dulce.

Citación: Furtado, YIC, *et al.*
Hemato-immunological responses of juvenile tambaqui (*Colossoma macropomum*) after commercial clay dietary supplementation. Revista de la Academia Colombiana de Ciencias Exactas, Físicas y Naturales. 49(190):90-101, enero-marzo de 2025. doi: <https://doi.org/10.18257/raccefyn.3077>

Editor: Martha Patricia Ramírez Pinilla

*Correspondencia:

Yuri Ian Carvalho Furtado;
yuri_furtado@hotmail.com

Recibido: September 21, 2024

Aceptado: February 18, 2025

Publicado en línea: March 12, 2025



This is an open access article distributed under the terms of the Creative Commons Attribution License.

Introduction

Colossoma macropomum, commonly known as tambaqui, belongs to the Serrasalminae subfamily and is native to the Amazon and Orinoco rivers and their tributaries (Goulding & Carvalho, 1982; Hilsdorf *et al.*, 2022; Mirande, 2010). The tambaqui has several desirable traits for aquaculture, including hardiness, ease of adaptation to captive management, excellent meat, and high commercial value (Garcez *et al.*, 2021). The global demand for high-quality and economically sustainable aquaculture feeds is increasing, driven by the rapid development of the aquaculture sector (Prabhu *et al.*, 2019). More than 70% of Brazil's fish production comes from intensive farming (Instituto Brasileiro de Geografia e Estatística - IBGE, 2014).

Although there is much uncertainty about the complex interplay between immune function and resistance to disease in fish, it is widely recognized that adequate nutrition plays a crucial role in strengthening the immune system (Hadiuzzaman *et al.*, 2022; Ucar *et al.*, 2019; Martin and Król, 2017). In fish, the immune system is composed of innate and adaptive components, including leukocytes such as the lymphocytes, neutrophils, and macrophages, which act against pathogens (Ranzani-Paiva *et al.*, 2013).

The innate immune response, acting as the first line of defense, is mediated by physical and chemical barriers, in addition to cells such as neutrophils and macrophages (Uribe *et al.*, 2011), whereas the adaptive response, which is more specific, involves lymphocytes that recognize specific antigens and have a complex interaction between the two systems to protect efficiently the organism (Secombes *et al.*, 2012). Efforts to modulate the immune response through dietary supplements have increased significantly (Ribeiro *et al.*, 2016; Dias *et al.*, 2019; Hoshino *et al.*, 2020).

Bentonite, a type of clay with sorption/absorption properties that also acts as a mycotoxin adsorbent, has the potential to modulate the immune response in fish through different mechanisms. One of the hypotheses proposes that, by adsorbing toxins and other harmful compounds in the digestive tract, bentonite reduces the burden on the immune system and allows it to function more efficiently, for which there have been studies to demonstrate its potential as immunomodulatory agent in aquaculture (Kiron, 2012; Fazio, 2019). Another proposal is that it interacts directly with immune cells, influencing the production of immunoglobulins, kidney function, and liver enzymes, and improving immunological and antioxidant parameters. Besides, bentonite may affect the intestinal microbiota, which plays a key role in modulating the immune response (El-Dahhar *et al.*, 2024; Abdel-Rahim *et al.*, 2023).

Previous studies have demonstrated the positive effects of bentonite in the diet of hybrid groupers (*Epinephelus fuscoguttatus* x *Epinephelus lanceolatus*) (Arshad *et al.*, 2021), the European seabass (*Dicentrarchus labrax*), the rainbow trout (*Oncorhynchus mykiss*) (Ucar *et al.*, 2019), the gilthead seabream (*Sparus aurata*) (Kanyilmaz & Tekelioğlu, 2016), and the Nile tilapia (*Oreochromis niloticus*) (Farrag *et al.*, 2009), leading to improvements in innate immunity, growth performance, and feed intake, as well as reducing the toxic effects of lead oxide. However, the beneficial effects of bentonite as a feed additive for captive-raised tambaqui have not yet been explored. In this context, we studied the potential use of natural bentonite clay on the hematological and biochemical responses of *C. macropomum*.

Materials and methods

The present study was authorized by the Embrapa Amapá Ethics Committee for the Use of Animals in Experiments (CEUA) (Number 019-CEUA/CPAFAP) and registered in the National System for Management of Genetic Heritage and Associated Traditional Knowledge (SisGen) under the identification number A0D6DC0.

Experimental design and diets

The experiments were conducted at the Embrapa Amapá, Aquaculture and Fisheries Laboratory, Macapá, Amapá State, Brazil. Tambaqui (*C. macropomum*) specimens (n = 120)

with an initial mean weight of 125.82 ± 21.67 g were randomly assigned to 12 experimental tanks (100 L polypropylene water containers). The sodium bentonite adsorbent (Buntech Tecnologia em Insumos, Indaiatuba, SP, Brazil) was incorporated into the commercial feed at three levels. The experimental groups were run in triplicate as follows: a) control (0%, with no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%), and d) 20 g of bentonite/kg of feed (2%). We collected 100 g of feed from each to analyze the centesimal composition of the experimental diets in triplicate. **Table 1** shows the chemical composition of the experimental diets.

Fish were fed with commercial rations (grain size 3.00 mm) containing 32% crude protein (Acqua Line, Rações Supra, Alisul Alimentos S.A., São Leopoldo, RS, Brazil) four times a day (at 08:00, 11:00, 14:00, and 17:00) during the acclimatization and all the experiment periods. The feed offered per day was equivalent to 5% of the total biomass of each experimental tank. During the experiment period, oxygen levels, temperature, and pH of the water were monitored using a multiparametric probe (Horiba, model AK88) and we obtained the following means and standard deviations: dissolved oxygen, 6.01 ± 0.38 mg/L; temperature, $29.47 \pm 0.12^\circ\text{C}$, and pH, 5.12 ± 0.68 , showing that water quality in experimental tanks was adequate for tambaqui (Aride *et al.*, 2007).

Growth parameters

After the feeding period (10 and 20 days), five tambaqui specimens from each replicate ($n=120$) were captured for blood collection, and then the total length and total weight of the specimens were registered.

Hematological analysis

We collected a blood sample (1,0 mL) per specimen by puncturing the caudal vein with a 3 mL syringe containing anticoagulant (EDTA 5%). We determined the following hematological parameters: hematocrit (Ht), i.e., the percentage of erythrocytes in the blood obtained through centrifugation of capillary tubes in a microhematocrit centrifuge for 5 minutes (Micro Spin, model CE120, Hangzhou, China) and read the results using a reading card (Goldenfarb *et al.*, 1971); we determined the hemoglobin concentration (Hb) using the cyanomethemoglobin method expressing the values in g dL^{-1} (Collier, 1944); finally, red blood cell count (RBC) was done by diluting blood samples in a formol-citrate solution and counting in erythrocytes $\times 10^6 \mu\text{L}^{-1}$ in a Neubauer chamber under an optical microscope (Boeco, model BOE-01, Hamburg, Germany). We calculated the following hematimetric indices based on these results: mean corpuscular volume (MCV, in fL), mean corpuscular hemoglobin (MCH, in g dL^{-1}), and mean corpuscular hemoglobin concentration (MCHC, in g dL^{-1}) (Ranzani-Paiva *et al.*, 2013). The leukocyte respiratory activity (respiratory burst) was determined as described by Sahoo *et al.* (2005) and Biller-Takahashi *et al.* (2013) with absorbance readings at

Table 1. Proximate chemical composition (%) of the experimental diets: a) Control (0%, with no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%); d) 20 g of bentonite/kg of feed (2%)

Parameters	0%	0.5%	1%	2%
Dry matter (%)	90.78 ± 0.11^a	91.11 ± 0.01^a	92.08 ± 1.62^a	90.09 ± 1.32^a
Crude protein (%)	32.67 ± 0.17^a	31.83 ± 0.78^a	32.50 ± 0.85^a	31.62 ± 1.09^a
Ether extract (%)	5.31 ± 0.33^a	4.64 ± 0.54^a	4.27 ± 0.19^b	4.45 ± 0.23^a
Ash (%)	11.93 ± 0.03^a	13.64 ± 1.6^a	12.62 ± 0.18^a	12.37 ± 1.64^a
Calcium (%)	1.62 ± 0.37^a	0.99 ± 0.12^b	1.02 ± 0.15^b	1.18 ± 0.10^a
Phosphorous (%)	1.61 ± 0.11^a	1.60 ± 0.10^a	1.50 ± 0.04^a	1.70 ± 0.14^a

Data expressed as mean $\pm$ standard deviation. Letters indicate significant differences ($p < 0.05$).

540 nm in a spectrophotometer (Biospectro, SP-220, Curitiba, Paraná, Brazil). Blood smears were prepared in duplicate and stained with the May-Grünwald-Giemsa-Wright staining to obtain total leukocytes, total thrombocytes, and differential leukocytes counts by the indirect method (Ishikawa *et al.*, 2008). After centrifugation of the blood (at 75 G, for 10 minutes) (Centrifuge 5424, Eppendorf, Hamburg, Germany), the plasma obtained was used to determine the total proteins, albumin, glucose, total cholesterol, and triglyceride concentrations (Labtest Diagnóstica S.A., Lagoa Santa, Minas Gerais, Brazil). The samples were read using a spectrophotometer (Biospectro SP-220, Curitiba, Paraná, Brazil) at specific wavelengths for each metabolite.

Statistical analysis

The data were subjected to normality and homoscedasticity tests using the Shapiro-Wilk and Levene methods, respectively, and, when necessary, they were transformed (total erythrocyte count). We used one-way and two-way variance analyses (ANOVA) and Tukey's *a posteriori* tests to compare the means. Diet and time were used as the main factors. Differences were considered significant at 5% probability (Zar, 2010). Tests were performed using the statistical software Sigmaplot 12.0.

Results and discussion

Adsorbents in aquaculture are increasingly used to prevent and mitigate animal health issues (Palm *et al.*, 2022; Ucar *et al.*, 2019). Prevention can be achieved by enhancing immunity and suppressing pathogens (El-Dahhar *et al.*, 2024). Table 2 shows the growth performance and survival rates of animals fed with different concentrations of the adsorbent for 20 days. Bentonite in the diet of *C. macropomum* did not significantly increase the total length or weight gain. Growth is a multifactorial process, and bentonite, although important, does not by itself explain growth performance. However, the species showed adequate development at this stage compared to the control group, and no mortality was recorded.

Hematological parameters provide valuable information about the health status of fish and are fundamental indicators for assessing animal well-being and optimal farming conditions (Ranzani-Paiva *et al.*, 2013; Fazio, 2019; Hoshino *et al.*, 2020). The innate immune system of fish is the primary line of defense against a wide range of pathogens, and it has a more significant role in fish than mammals (Saurabh & Sahoo, 2008). In our study, hematocrit (Ht) in animals fed with 1% and 2% bentonite for 10 days significantly increased compared to the 0% and 0.5% groups (Table 3), but these values remained within the expected range for the species and were consistent with specimens' health (Aride *et al.*, 2017; Tavares-Dias *et al.*, 2009).

Tambaqui receiving a diet including 0.5% bentonite adsorbent for 10 days showed a decrease in hemoglobin (Hb) concentration compared to the other groups. However, after 20 days of 0.5%, 1%, and 2% bentonite feeding, Hb and red blood cell count (RBC)

Table 2. Initial and final weight of juvenile tambaqui (*Colossoma macropomum*) fed with the following experimental diets: Control (0%, no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%); d) 20 g of bentonite/kg of feed (2%)

Parameters	0%	0.5%	1%	2%
Initial weight (g)	128.23±21.71 ^a	123.40±21.22 ^a	123.06±22.87 ^a	128.60± 21.31 ^a
Final weight (g)	247.66±18.64 ^a	234.66±26.36 ^a	255.50± 44.23 ^a	263.07± 32.96 ^a
Initial length (cm)	15.39±1.21 ^a	15.69 ± 0.94 ^a	15.76 ± 1.08 ^a	15.70 ± 0.96 ^a
Final length (cm)	19.11 ± 1.15 ^a	19.08 ± 0.85 ^a	19.40± 1.53 ^a	19.98 ± 1.14 ^a
Fish survival (%)	100	100	100	100

Data expressed as mean±standard deviation. Letters indicate significant differences (p<0.05).

Table 3. Hematological variables and hematimetric indices (mean $\pm$ standard deviation) of *Colossoma macropomum* fed with these experimental diets: Control (0%, no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%); d) 20 g of bentonite/kg of feed (2%) for 10 and 20 days, respectively

Parameters	Factors			
	0%	0.5%	1%	2%
10 days				
Ht (%)	27.00 $\pm$ 2.61 ^b	26.76 $\pm$ 3.58 ^b	28.70 $\pm$ 2.96 ^a	30.40 $\pm$ 3.85 ^a
Hb (g dL ⁻¹)	7.60 $\pm$ 0.95 ^a	6.88 $\pm$ 1.37 ^b	8.06 $\pm$ 0.76 ^a	7.52 $\pm$ 0.91 ^a
RBC (x 10 ⁶ μ L ⁻¹)	1.57 $\pm$ 0.30 ^a	1.47 $\pm$ 0.22 ^a	1.65 $\pm$ 0.40 ^a	1.34 $\pm$ 0.18 ^a
VCM (fL)	185.65 $\pm$ 45.82 ^a	178.30 $\pm$ 26.62 ^b	184.21 $\pm$ 46.88 ^b	220.35 $\pm$ 31.01 ^a
HCM (g dL ⁻¹)	46.92 $\pm$ 14.95 ^a	47.63 $\pm$ 11.69 ^a	52.35 $\pm$ 16.09 ^a	56.88 $\pm$ 10.47 ^a
CHCM (g dL ⁻¹)	28.18 $\pm$ 2.97 ^a	25.85 $\pm$ 4.12 ^a	28.07 $\pm$ 1.34 ^a	25.10 $\pm$ 3.47 ^a
Parameters	Factors			
	0%	0.5%	1%	2%
20 days				
Ht (%)	27.39 $\pm$ 3.13 ^a	27.50 $\pm$ 2.46 ^a	27.85 $\pm$ 1.02 ^a	28.66 $\pm$ 1.15 ^a
Hb (g dL ⁻¹)	8.16 $\pm$ 0.98 ^a	6.61 $\pm$ 0.97 ^b	6.87 $\pm$ 0.82 ^b	7.10 $\pm$ 0.60 ^b
RBC (x 10 ⁶ μ L ⁻¹)	1.46 $\pm$ 0.34 ^a	1.08 $\pm$ 0.20 ^b	1.01 $\pm$ 0.12 ^{b*}	1.11 $\pm$ 0.17 ^{b*}
VCM (fL)	177.27 $\pm$ 21.93 ^b	249.65 $\pm$ 40.35 ^{a*}	276.30 $\pm$ 28.37 ^{a*}	268.86 $\pm$ 39.74 ^{a*}
HCM (g dL ⁻¹)	56.03 $\pm$ 11.18 ^a	58.15 $\pm$ 12.99 ^a	64.37 $\pm$ 13.76 ^a	64.05 $\pm$ 11.15 ^a
CHCM (g dL ⁻¹)	30.70 $\pm$ 4.89 ^{a*}	23.97 $\pm$ 2.00 ^b	24.42 $\pm$ 3.25 ^{b*}	24.69 $\pm$ 2.15 ^b

Letters indicate significant differences in two-way ANOVA followed by post-hoc Tukey ($p < 0.05$); *: significant difference ($p < 0.05$) when comparing 10- and 20-day-period groups using the same treatment and for the same parameter. Ht: Hematocrit; Hb: Hemoglobin concentration; RBC: Red blood cells count; MCV: Mean corpuscular volume; MCH: Mean corpuscular hemoglobin; MCHC: Mean corpuscular hemoglobin concentration

reduced significantly compared to the control group, which indicates an interaction between the diet and the feeding period, particularly in the 1% group, and suggests normochromic anemia due to the extended period of adsorbent consumption (Ranzani-Paiva *et al.*, 2013; Kanyilmaz & Tekelioğlu, 2016), despite that, according to Tavares-Dias *et al.* (2009), Hb concentration of *C. macropomum* ranged from 6.3 to 13.7 g dL⁻¹, showing similar values to those reported in the present study. This was also the case in the study by Costa *et al.* (2022) for tambaqui diets containing silage made from fish and vegetable residues. Hb, Ht, and RBC are the main hematological parameters reflecting the initial response of the erythrocytic series, which indicates blood oxygen transport capacity and its subsequent utilization by the body. The RBC, Hb, and Ht values found in the tambaqui in the present study were similar to those described by Tavares-Dias *et al.* (2009). However, RBC reduced after 20 days of feeding, particularly in the groups receiving bentonite, suggesting decreased oxygen transport capacity possibly leading to viral, bacterial, and parasitic infections, exposure to toxins, nutritional deficiencies, and blood loss (Currie *et al.*, 2022; Witeska *et al.*, 2015).

The mean corpuscular volume (MCV) was higher at 10 days in the 0% and 2% groups compared to the 0.5% and 1% groups (Table 3). At 20 days, MCV was similar in the groups with bentonite supplementation and statistically higher than the control group (0%). Besides the additive, there was a strong positive interaction with time, with an MCV significant increase in the 0.5%, 1%, and 2% groups. The mean RBC, MCV, and MCHC values in fish fed with bentonite indicated morphological adjustments where RBC reduction was compensated by the increase in the volume of this cell, which resulted in

more space for hemoglobin. Consequently, it did not harm the exchange of respiratory gases, a basic function of erythrocytes in the blood (Tavares-Dias & Moraes, 2010; Ranzani-Paiva *et al.*, 2013). However, erythrocyte indices (MCV, MCH, MCHC) help in the differential diagnosis of anemia and indicated here a hypochromic macrocytic anemic process (Ranzani-Paiva *et al.*, 2013). This also suggests cellular impairment as a result of RBC reduction after 20 days with bentonite in *C. macropomum* diets. No differences in MCH were observed ($p > 0.05$) among the groups or between different feeding times. After 20 days of testing, the groups with bentonite inclusion in the diet showed the lowest MCHC values ($p < 0.05$), resulting from the lower Hb concentration in these groups. This supports the findings for rainbow trout (*O. mykiss*) using bentonite and copper (Cu) in experimental diets (Ucar *et al.*, 2019).

The addition of 1% and 2% bentonite in diets for tambaqui significantly increased the respiratory burst activity of leukocytes after 20 days (Table 4), which has been used as a marker of innate immunity in fish (Biller-Takahashi *et al.*, 2013) and might be a mechanism for restoring homeostasis. During burst, pathogens are destroyed through phagocytosis, increasing intracellular oxygen consumption by leukocytes and producing reactive oxygen species (ROS), which result in free radicals and cellular metabolism byproducts (Biller & Takahashi, 2018; Biller-Takahashi *et al.*, 2015; Dong *et al.*, 2017). These byproducts play a crucial role in destroying invading agents. Thus, burst can act as defense against pathogens, but it also has a reactive effect on the animal's immune system, triggered in tambaqui by the higher bentonite levels in the diet.

Additionally, *C. macropomum* fed for 20 days with diets containing different levels of adsorbents showed a decrease in the total number of leukocytes, thrombocytes, lymphocytes, monocytes, and neutrophils in circulation, alongside an increase in respiratory burst activity (Table 4). These leukocyte count alterations may be related to stress or physiological damage caused by the use of clay supplementation in the diets compared to the control. It is important to note that leukocytes are essential for evaluating the fish immune system (Witeska *et al.*, 2023; Fazio, 2019; Tavares-Dias *et al.*, 2007), as they work together to maintain tissue integrity against infectious agents and are responsible for the fish immune balance migrating through the bloodstream to sites of injury or tissue infection during adverse inflammatory processes (Ranzani-Paiva *et al.*, 2013; Kiron, 2012; Fazio, 2019).

The stress response in fish can be assessed through indicators such as serum or plasma glucose and cortisol and lysozyme activity, which change in response to stressful stimuli (Urbinati *et al.*, 2020). Our results showed a significant decrease in plasma glucose levels in the fish after 20 days of cultivation compared to 10 days (Table 5). Such reduction is

Table 4. Total thrombocytes and total and differential leukocytes count in juvenile tambaqui (*Colossoma macropomum*) fed with experimental diets: Control (0%, no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%); d) 20 g of bentonite/kg of feed (2%) after 20 days

Parameters	0%	0.5%	1%	2%
Burst (OD)	0.25±0.03 ^b	0.26±0.03 ^b	0.29±0.04 ^a	0.29±0.03 ^a
Thrombocytes (x10 ³ µL)	28.55±12.80 ^a	19.02±19.02 ^b	20.76±6.56 ^b	20.61±5.91 ^b
Leukocytes (x10 ³ µL)	143.43±41.55 ^a	108.14±24.14 ^b	99.35±14.11 ^b	112.95±23.30 ^b
Lymphocytes (x10 ³ µL)	63.68±21.02 ^a	51.66±14.69 ^a	53.64±22.10 ^a	55.77±13.34 ^a
Monocytes (x10 ³ µL)	55.47±16.23 ^a	39.73±8.96 ^b	34.40±9.31 ^b	42.215±9.90 ^b
Neutrophils (x10 ³ µL)	20.73±7.74 ^a	13.57±4.99 ^b	13.44±3.11 ^b	13.25±6.12 ^b
Eosinophils (x10 ³ µL)	0.53±1.17 ^a	0.07±0.01 ^a	0.17±0.25 ^a	0.21±0.32 ^a
LG-PAS (x10 ³ µL)	3.05±1.81 ^a	3.13±1.41 ^a	2.35±1.57 ^a	1.69±1.32 ^a

Data expressed as mean±standard deviation. Letters mean significant differences ($p < 0.05$). LG-PAS: Leukocyte granular-PAS positive

Table 5. Plasma metabolite profile (n=15) of *Colossoma macropomum* fed with experimental diets: Control (0%, no bentonite); b) 5 g of bentonite/kg of feed (0.5%); c) 10 g of bentonite/kg of feed (1%); d) 20 g of bentonite/kg of feed (2%) for 10 and 20 days

Parameters	Factors			
	0%	0.5%	1%	2%
10 days				
Glucose (mg dL ⁻¹)	71.48±15.48 ^a	82.55±14.53 ^a	77.25±17.63 ^a	78.00±19.76 ^a
Total protein (g dL ⁻¹)	3.10±0.37 ^a	3.26±0.21 ^a	3.29±0.23 ^a	3.24±0.26 ^a
Total Cholesterol (mg dL ⁻¹)	96.63±9.73 ^b	118.93±23.26 ^a	88.60±14.28 ^b	78.31±16.26 ^b
Triglycerides (mg dL ⁻¹)	147.52±24.59 ^a	151.57±16.36 ^a	136.63±22.05 ^{ab}	126.01±15.85 ^b
Albumin (g dL ⁻¹)	0.82±0.07 ^{cd}	0.44±0.13 ^b	1.04±0.11 ^a	0.93±0.10 ^{ad}
Globulin (g dL ⁻¹)	2.16 ± 0.36 ^a	2.38 ± 0.34 ^a	2.18 ± 0.34 ^a	2.30 ± 0.22 ^a
Parameters	Factors			
	0%	0.5%	1%	2%
20 days				
Glucose (mg dL ⁻¹)	44.15 ± 11.34 ^{a*}	56.34 ± 7.57 ^{a*}	62.27 ± 7.80 ^{a*}	63.13 ± 10.88 ^{a*}
Total protein (g dL ⁻¹)	3.69 ± 0.30 ^{a*}	3.27 ± 0.25 ^b	3.40 ± 0.29 ^b	3.55 ± 0.41 ^{a*}
Total Cholesterol (mg dL ⁻¹)	85.65 ± 7.21 ^b	116.39 ± 18.06 ^a	128.13 ± 17.00 ^{a*}	100.29 ± 15.07 ^{b*}
Triglycerides (mg dL ⁻¹)	132.18 ± 19.81 ^b	158.88 ± 22.61 ^a	176.29 ± 17.64 ^{a*}	163.71 ± 32.68 ^{a*}
Albumin (g dL ⁻¹)	0.66 ± 0.06 ^{a*}	0.96 ± 0.20 ^{d*}	0.49 ± 0.08 ^{b*}	1.51 ± 0.18 ^{a*}
Globulin (g dL ⁻¹)	3.03 ± 0.30 ^{a*}	2.06 ± 0.61 ^{b*}	2.91 ± 0.33 ^{a*}	2.08 ± 0.40 ^b

Data expressed as mean±standard deviation. Letters mean significant differences in the ANOVA two-way test followed by post-hoc Tukey ($p<0.05$). *: significant difference ($p<0.05$) when comparing 10- and 20-day groups with the same treatment and parameter

often observed with immunostimulatory diets due to the reduction in stress effects (Hoshino *et al.*, 2017; Dias *et al.*, 2019). A likely explanation is that the diet may have stimulated insulin activity, which serves as a secondary stress (Brandão *et al.*, 2006) indicator and results in reduced glucose levels. Furthermore, it may be related to energy metabolism regulation under the influence of clay (Schell *et al.*, 1993; Shannon *et al.*, 2017), which is known for its adsorbent and stabilizing properties (Fazio, 2019; Kiron, 2012).

Total protein concentration in fish reflects aspects of their physiology, including nutritional status, general health, stress level, humoral defense capacity, and well-being (Abdel-Rahim *et al.*, 2023). After 20 days of feeding, the 0.5% and 1% bentonite groups presented lower total protein levels compared to the other treatments. According to Chagas *et al.* (2007), tambaqui specimens are considered well-nourished when plasma total protein levels are above 2.0 g dL⁻¹, which aligns with the present study and the findings of Costa *et al.* (2022) and Silva *et al.* (2020).

Interactions between diet and time were observed with 1% and 2% bentonite, resulting in a significant increase in plasma lipid profile levels ($p<0.05$), which could be a crucial benefit in membrane structure; besides, lipids are precursors to all steroid hormones. Such interactions may be triggered by liver and kidney dysfunction, leading to elevated cholesterol levels in the bloodstream (Öner *et al.*, 2008) due to the stress induced by bentonite in their diet. Another possible explanation for the increase in total cholesterol and triglycerides levels is related to the increase in feed viscosity, endogenous feed losses, and fermentation in fish intestines, among other causes (Rosas *et al.*, 2008). Jiang *et al.* (2010) presented similar results with the inclusion of clays in piglets' diets with and without the mycotoxin zearalenone, which significantly increased plasma cholesterol levels.

Stress in fish triggers an increase in energy demand and mobilization of plasma proteins used to meet energy needs, maintain osmotic balance, and improve the immune response, besides reflecting liver and kidney health (Javed & Usmani, 2015; Costa *et al.*, 2019). At 10 days, there was a variation in albumin levels between groups, with the 1% group presenting the highest value and the 0.5% group the lowest. At 20 days, the variation was even more evident. The 2% group had the highest albumin value, while the 1% group had the lowest. Regarding globulin, significant differences were observed only at 20 days. The 0% and 1% groups exhibited the highest globulin values, while the 0.5% and 2% groups had the lowest. Albumin and globulin are important proteins in the blood of fish: the first contributes to osmotic pressure and the transport of substances (Mobarhan, 1988), while globulin is related to the immune response (De Souza *et al.*, 2020). The relationship between albumin and globulin can be a health indicator. In our study, there appeared to be a change in this relationship over time, especially in the groups treated with bentonite, suggesting a possible negative influence of bentonite on the fish protein metabolism. The albumin increase in the 1% (10 days) and 2% (20 days) groups can be attributed to the synthesis of proteins needed to meet the high energy demand since the synthesis occurs in the liver, besides indicating an increase in the tambaqui immune response (Morante *et al.*, 2021).

Conclusion

Our study is the first to provide hemato-immunological analyses of tambaqui fed with sodium bentonite-supplemented diets. The findings highlight that the adsorbent supplementation did not improve the hematological-biochemical profile of this fish species. Physiological damage caused by the adsorbent, such as thrombocytopenia, leukopenia, lymphocytopenia, and monocytopenia, pointed to the response and migration of the cells to inflammatory sites. In conclusion, our results indicate that dietary supplementation with sodium bentonite had deleterious effects on the organism of the *C. macropomum* evaluated and, therefore, it is not recommended as a food additive for this fish species.

Acknowledgments

The authors thank Empresa Brasileira de Pesquisa Agropecuária (Embrapa Amapá) and the Amapá State Research Support Foundation (FAPEAP) for their financial support and the scholarship granted to F.L.S. Souza. We also thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for Y.I.C. Furtado's scholarship (grant 88887.636974/2021-00).

Author contributions

Conceptualization, E.T.O.Y. and Y.I.C.F.; methodology, E.T.O.Y., Y.I.C.F., F.L.S.S., M.V.F.M., A.M.P., C.B.S.; software Y.I.C.F.; validation Y.I.C.F. and E.T.O.Y.; investigation Y.I.C.F., F.L.S.S. and E.T.O.Y.; resources Y.I.C.F., F.L.S.S., M.V.F.M., A.M.P., C.B.S., and E.T.O.Y.; original draft preparation Y.I.C.F. and E.T.O.Y.; manuscript review and editing Y.I.C.F., F.L.S.S., M.V.F.M., A.M.P., C.B.S., and E.T.O.Y. All authors read and agreed on the final version of the manuscript.

Conflicts of interest

The authors declare no conflict of interest.

Funding

This research was funded by Embrapa, UNIFAP, and FAPEAP.

References

- Abdel-Rahim, M. M., Elhetawy, A. I., Refaey, W. M., Lotfy, A. M., El-Zaeem, S. Y., Elebiary, E. H., El-Dahhar, A. A. (2023). Potential benefits of natural bentonite as a water clarifier

- on ammonia removal, performance, immunity, antioxidant, and histology of *Dicentrarchus labrax*. *The Egyptian Journal of Aquatic Research*, 49(2), 253-260. <https://doi.org/10.1016/j.ejar.2023.04.002>
- Aride, P. H. R., Oliveira, A. M., Batista, R. B., Ferreira, M. S., Pantoja-Lima, J., Ladislau, D. S., Castro, P. D. S., Oliveira, A. T. (2017). Changes on physiological parameters of tambaqui (*Colossoma macropomum*) fed with diets supplemented with Amazonian fruit Camu camu (*Myrciaria dubia*). *Brazilian Journal of Biology*, 78, 360-367. <https://doi.org/10.1590/1519-6984.169442>
- Aride, P. H. R., Roubach, R., Val, A. L. (2007). Tolerance response of tambaqui *Colossoma macropomum* (Cuvier) to water pH. *Aquaculture Research*, 38(6), 588-594. <https://doi.org/10.1111/j.1365-2109.2007.01693.x>
- Arshad, S. E., Jeffrey, F. N. M., Amin, Z., Shapawi, R. (2021). Effects of sodium bentonite clay as a feed additive on the growth and haematology parameters of hybrid grouper, *Epinephelus fuscoguttatus* x *Epinephelus lanceolatus*. *Songklanakarin Journal of Science and Technology*, 43(2), 505-511. <https://sjst.psu.ac.th/journal/43-2/27.pdf>
- Biller, J. D., & Takahashi, L. S. (2018). Oxidative stress and fish immune system: phagocytosis and leukocyte respiratory burst activity. *Anais da Academia Brasileira de Ciências*. 90, 3403-3414. <https://doi.org/10.1590/0001-3765201820170730>
- Biller-Takahashi, J. D., Takahashi, L. S., Mingatto, F. E., Urbinati, E. C. (2015). The immune system is limited by oxidative stress: dietary selenium promotes optimal antioxidative status and greatest immune defense in pacu *Piaractus mesopotamicus*. *Fish & Shellfish Immunology*, 47(1), 360-367. <https://doi.org/10.1016/j.fsi.2015.09.022>
- Biller-Takahashi, J. D., Takahashi, L. S., Saita, M. V., Gimbo, R. Y., Urbinati, E. C. (2013). Leukocytes respiratory burst activity as indicator of innate immunity of pacu *Piaractus mesopotamicus*. *Brazilian Journal of Biology*. 73(2) 425-429. <https://doi.org/10.1590/S1519-69842013000200026>
- Brandão, F. R., Gomes, L. D. C., Chagas, E. C. (2006). Respostas de estresse em pirarucu (*Arapaima gigas*) durante práticas de rotina em piscicultura. *Acta Amazonica*, 36, 349-356, 2006. <https://doi.org/10.1590/S0044-59672006000300010>
- Instituto Brasileiro de Geografia e Estatística (IBGE). August 10, 2023. Produção da aquicultura, por tipo de produto. 2014. <http://www.sidra.ibge.gov.br/bda/tabela/listabl.asp?c=3940andz=tando=21>
- Centeno, L., Silva-Acuña, R., Barrios, R., Salazar Lugo, R., Matute, C., L Pérez, J. (2007). Características hematológicas de la cachama (*Colossoma macropomum*) en tres etapas de crecimiento cultivadas en el estado Delta Amacuro, Venezuela. *Zootecnia tropical*, 25(4), 237-243. <https://ve.scielo.org/pdf/zt/v25n4/art01.pdf>
- Chagas, E.C., Gomes, L. D.C., Martins Júnior, H., Roubach, R. (2007). Produtividade de tambaqui criado em tanque-rede com diferentes taxas de alimentação. *Ciência Rural*, 37, 1109-1115. <https://doi.org/10.1590/S0103-84782007000400031>
- Costa, O. T. F., Dias, L. C., Malmann, C. S. Y., de Lima Ferreira, C. A., do Carmo, I. B., Wischneski, A. G., Sousa, R. L., Caverro, B. A. S., Lameiras, J. L. V., Dos-Santos, M. C. (2019). The effects of stocking density on the hematology, plasma protein profile and immunoglobulin production of juvenile tambaqui (*Colossoma macropomum*) farmed in Brazil. *Aquaculture*, 499, 260-268. <https://doi.org/10.1016/j.aquaculture.2018.09.040>
- Costa, M. N. F., Furtado, Y. I. C., Monteiro, C. C., Brasiense, A. R. P., Yoshioka, E. T. (2022). Physiological responses of tambaqui (*Colossoma macropomum*) fed diets supplemented with silage from fish and vegetables residues. *Brazilian Journal of Biology*, 84, e255493. <https://doi.org/10.1590/1519-6984.255493>
- Currie, A. R., Cockerill, D., Diez-Padrisa, M., Haining, H., Henriquez, F. L., Quinn, B. (2022) Anemia in salmon aquaculture: Scotland as a case study. *Aquaculture*. 546, 737313. <https://doi.org/10.1016/j.aquaculture.2021.737313>
- De Souza, R. C., Baldisserotto, B., Melo, J. F. B., da Costa, M. M., de Souza, E. M., Copatti, C. E. (2020). Dietary *Aloysia triphylla* essential oil on growth performance and biochemical and haematological variables in Nile tilapia. *Aquaculture*, 519, 734913. <https://doi.org/10.1016/j.aquaculture.2019.734913>
- Dias, M. K. R., Yoshioka, E. T. O., Rodriguez, A. F. R., Ribeiro, R. A., Faria, F. S. E. D. V., Ozório, R. O. A., Tavares-Dias, M. (2019). Growth, physiological and immune responses of *Arapaima gigas* (Arapaimidae) to *Aeromonas hydrophila* challenge and handling stress following feeding with immunostimulant supplemented diets. *Fish & shellfish immunology*, 84, 843-847. <https://doi.org/10.1016/j.fsi.2018.10.045>

- Dong, M., Liang, Y., Ramalingam, R., Tang, S. W., Shen, W., Ye, R., Gopalakrishnan, S., Au, D. W. T., Lam, Y. W. (2017). Proteomic characterization of the interactions between fish serum proteins and waterborne bacteria reveals the suppression of anti-oxidative defense as a serum-mediated antimicrobial mechanism. *Fish & shellfish immunology*, 62, 96-106. <https://doi.org/10.1016/j.fsi.2017.01.013>
- El-Dahhar, A. A., Elhetawy, A. I., Refaey, W. M., El-Zaeem, S. Y., Elebiary, E. H., Lotfy, A. M., Abdel-Rahim, M. M. (2024). Bentonite-supplemented diets improved fish performance ammonia excretion haemato-biochemical analyses immunity antioxidants and histological characteristics of European seabass *Dicentrarchus labrax*. *Scientific Reports*, 14(1), 13868. <https://doi.org/10.1038/s41598-024-63936-6>
- Farrag, F. H., Khalil, F. F., Mehrim, A. I. (2009). Reduction of lead oxide toxicity by using bentonite in mono-sex nile tilapia *Oreochromis niloticus* diets. *Abbassa Int. J. Aqua*, 429-451. <https://kenanaonline.com/files/0027/27570/05%20Pb%20and%20Bentonite%202009.pdf>
- Fazio, F. (2019). Fish hematology analysis as an important tool of aquaculture: a review. *Aquaculture*, 500, 237-242. <https://doi.org/10.1016/j.aquaculture.2018.10.030>
- Garcez, J. R., da Nóbrega, V. S. L., Torres, T. P., Signor, A. A. (2021). Cultivo de tambaqui (*Colossoma macropomum*) em tanques-rede: Aspectos técnicos. *Research, Society and Development*, 10(8), e45810817560-e45810817560. <https://doi.org/10.33448/rsd-v10i8.17560>
- Goulding, M. & Carvalho, M. L. (1982). Life history and management of the tambaqui (*Colossoma macropomum*, Characidae): an important Amazonian food fish. *Revista Brasileira de Zoologia*, 1, 107-133. <https://doi.org/10.1590/S0101-81751982000200001>
- Goldenfarb, P. B., Bowyer, F. P., Hall, E., Brosious, E. (1971). Reproducibility in the hematology laboratory: the microhematocrit determination. *American Journal of Clinical Pathology*, 56(1), 35-39. <https://doi.org/10.1093/ajcp/56.1.35>
- Hadiuzzaman, M., Moniruzzaman, M., Shahjahan, M., Bai, S. C., Min, T., & Hossain, Z. (2022). β -Glucan: Mode of action and its uses in fish immunomodulation. *Frontiers in Marine Science*. 9, 905986. <https://doi.org/10.3389/fmars.2022.905986>
- Hilsdorf, A. W. S., Hallerman, E., Valladão, G. M. R., Zaminhan-Hassemer, M., Hashimoto, D. T., Dairiki, J. K., Takahashi, L. S., Albergaria, F. C., Gomes, M. E. S., Venturieri, R. L. L., Moreira, R. G., Cyrino, J. E. P. (2022). The farming and husbandry of *Colossoma macropomum*: From Amazonian waters to sustainable production. *Reviews in Aquaculture*, 14(2), 993-1027. <https://doi.org/10.1111/raq.12638>
- Hoshino, M. D. F. G., Marinho, R. D. G. B., Pereira, D. F., Yoshioka, E. T. O., Tavares-Dias, M., Ozorio, R. O. D. A., Rodriguez, A. F. R., Ribeiro, R. A., Faria, F. S. E. D. V. D. (2017). Hematological and biochemical responses of pirarucu (*Arapaima gigas*, Arapaimidae) fed with diets containing a glucomannan product derived from yeast and algae. *Acta Amazonica*, 47(2), 87-94. <https://doi.org/10.1590/1809-4392201700781>
- Hoshino, M. D. F. G., da Silva Ramos, R., Lopes, J. R. T., Ribeiro, R. A., Rodriguez, A. F. R., da Silva, T. C., Faria, F. S. E. D. V. D., Tavares-Dias, M., Ozorio, R. O. D. A., Yoshioka, E. T. O. (2020). Innate immune response of pirarucu improved with yeast-supplemented diets. *Aquaculture Reports*, 18, 100421. <https://doi.org/10.1016/j.aqrep.2020.100421>
- Ishikawa, N. M., Ranzani-Paiva, M. J. T., Lombardi, J. V. (2008). Metodologia para quantificação de leucócitos totais em peixe, *Oreochromis niloticus*. *Archives of Veterinary Science*, 13, 54-63. <https://doi.org/10.5380/avs.v13i1.11560>
- Javed, M. & Usmani, N. (2015). Stress response of biomolecules (carbohydrate, protein and lipid profiles) in fish *Channa punctatus* inhabiting river polluted by Thermal Power Plant effluent. *Saudi Journal of Biological Sciences*, 22(2), 237-242. <https://doi.org/10.1016/j.sjbs.2014.09.021>
- Jiang, S., Yang, Z., Yang, W., Gao, J., Liu, F., Chen, C. C., Chi, F. (2010). Physiopathological effects of zearalenone in post-weaning female piglets with or without montmorillonite clay adsorbent. *Livestock Science*, 131(1), 130-136. <https://doi.org/10.1016/j.livsci.2010.02.022>
- Kanyılmaz, M. & Tekelioğlu, N. (2016). Effects of dietary zeolite levels on some blood parameters of gilthead seabream (*Sparus aurata*) juveniles. *Journal of Aquaculture Engineering and Fisheries Research*, 2(3), 119-127. <https://doi.org/10.3153/JAEFR16014>
- Kiron, V. (2012). Fish immune system and its nutritional modulation for preventive health care. *Animal feed science and technology*, 173(1-2), 111-133. <https://doi.org/10.1016/j.anifeeds.2011.12.015>

- Martin, S. A. M. & Król, E.** (2017). Nutrigenomics and immune function in fish: new insights from omics technologies. *Developmental & Comparative Immunology*, 75, 86-98. <https://doi.org/10.1016/j.dci.2017.02.024>
- Mirande, J. M.** (2010). Phylogeny of the family Characidae (Teleostei: Characiformes): from characters to taxonomy. *Neotropical Ichthyology*, 8, 385-568. <https://doi.org/10.1590/S1679-62252010000300001>
- Mobarhan, S.** (1988). The role of albumin in nutritional support. *Journal of the American College of Nutrition*, 7(6), 445-452. <https://doi.org/10.1080/07315724.1988.10720260>
- Morante, V. H. P., Copatti, C. E., Souza, A. R. L., Costa, M. M., Braga, L. G. T., Souza, A. M., Melo, F. V. S. T., Camargo, A. C. S., Melo, J. F. B.** (2021). Assessment the crude grape extract as feed additive for tambaqui (*Colossoma macropomum*), an omnivorous fish. *Aquaculture*, 544(2021), 737068. <https://doi.org/10.1016/j.aquaculture.2021.737068>
- Öner, M., Atli, G., Canli, M.** (2008). Changes in serum biochemical parameters of freshwater fish *Oreochromis niloticus* following prolonged metal (Ag, Cd, Cr, Cu, Zn) exposures. *Environmental Toxicology and Chemistry: An International Journal*, 27(2), 360-366. <https://doi.org/10.1897/07-281R.1>
- Ranzani-Paiva, M. J. T. R., Pádua, S. B., Tavares-Dias, M., Egami, M. I.** (2013). *Métodos para análise hematológica em peixes*. Editora da Universidade Estadual de Maringá-EDUEM.
- Ribeiro, S. C., Malheiros, D. F., Guiloizki, I. C., Majolo, C., Chaves, F. C. M., Chagas, E. C., Assis, H. C. S., Tavares-Dias, M., Yoshioka, E. T. O.** (2018). Antioxidants effects and resistance against pathogens of *Colossoma macropomum* (Serassalmidae) fed *Mentha piperita* essential oil. *Aquaculture*, 490, 29-34. <https://doi.org/10.1016/j.aquaculture.2018.02.024>
- Rosas, C., Tut, J., Baeza, J., Sánchez, A., Sosa, V., Pascual, C., Arena, A., Domingues, P., Cuzon, G.** (2008). Effect of type of binder on growth, digestibility, and energetic balance of *Octopus maya*. *Aquaculture*, 275(1-4), 291-297. <https://doi.org/10.1016/j.aquaculture.2008.01.015>
- Sahoo, P. K., Kumari, J., Mishra, B. K.** (2005). Non-specific immune responses in juveniles of Indian major carps. *Journal of Applied Ichthyology*, 21, 151-155. <https://doi.org/10.1111/j.1439-0426.2004.00606.x>
- Secombes, C. J. & Wang, T.** (2012). *The innate and adaptive immune system of fish*. In *Infectious disease in aquaculture* (pp. 3-68). Woodhead Publishing. <https://doi.org/10.1533/9780857095732.1.3>
- Shannon, T. A., Ledoux, D. R., Rottinghaus, G. E., Shaw, D. P., Daković, A., Marković, M.** (2017). The efficacy of raw and concentrated bentonite clay in reducing the toxic effects of aflatoxin in broiler chicks. *Poultry science*, 96(6), 1651-1658.
- Palm, H. W., Berchtold, E., Gille, B., Knaus, U., Wenzel, L. C., Baßmann, B.** (2022). Growth and welfare of African catfish (*Clarias gariepinus* Burchell, 1822) under dietary supplementation with mixed-layer clay mineral montmorillonite-illite/muscovite in commercial aquaculture. *Aquaculture journal*, 2(3), 227-245. <https://doi.org/10.3390/aquacj2030013>
- Prabhu, P. A. J., Fountoulaki, E., Maas, R., Heinsbroek, L. T. N., Eding, E. H., Kaushik, S. J., Schrama, J. W.** (2019). Dietary ingredient composition alters faecal characteristics and waste production in common carp reared in recirculation system. *Aquaculture*, 512, 734357. <https://doi.org/10.1016/j.aquaculture.2019.734357>
- Saurabh, S. & Sahoo, P. K.** (2008) Lysozyme: an important defence molecule of fish innate immune system. *Aquaculture research*, 39(3), 223-239. <https://doi.org/10.1111/j.1365-2109.2007.01883.x>
- Schell, T. C., Lindemann, M. D., Kornegay, E. T., Blodgett, D. J., Doerr, J. A.** (1993). Effectiveness of different types of clay for reducing the detrimental effects of aflatoxin-contaminated diets on performance and serum profiles of weanling pigs. *Journal of Animal Science*, 71(5), 1226-1231. <https://doi.org/10.2527/1993.7151226x>
- Silva, A. F., Copatti, C. E., de Oliveira, E. P., Bonfá, H. C., de Melo, F. V. S. T., de Silva Camargo, A. C., Melo, J. F. B.** (2020). Effects of whole banana meal inclusion as replacement for corn meal on digestibility, growth performance, haematological and biochemical variables in practical diets for tambaqui juveniles (*Colossoma macropomum*). *Aquaculture Reports*, 17, 100307. <https://doi.org/10.1016/j.aqrep.2020.100307>
- Tavares-Dias, M. & Moraes, F. R.** (2004) *Hematologia de peixes teleósteos*. Ribeirão Preto.
- Tavares-Dias, M. & Sandrim, E. F. S.** (1998). Características hematológicas de teleósteos brasileiros. I. Série vermelha e dosagens de cortisol e glicose do plasma sanguíneo de espécimes de *Colossoma macropomum* em condições de cultivo. *Acta Scientiarum*, 20(2), 157-160. <https://doi.org/10.4025/actasciobiolsci.v20i0.4466>

- Tavares-Dias, M., Barcellos, J. F. M., Marcon, J. L., Menezes, G. C., Ono, E. A., Affonso, E. G.** (2007). Hematological and biochemical parameters for the pirarucu *Arapaima gigas* Schinz, 1822 (Osteoglossiformes, Arapaimatidae) in net cage culture. *Electronic Journal of Ichthyology*, 2, 61-68. <https://citeseerx.ist.psu.edu/document?repid=rep1&type=pdf&doi=57cb2fe77124e64b8f401aac1168d7d2ac5a7af>
- Tavares-Dias, M., Ishikawa, M. M., Martins, M. L., Satake, F., Hisano, H., de Pádua, S. B., Jerônimo, G. T., Sá, A. R. S.** (2009). *Hematologia: ferramenta para o monitoramento do estado de saúde de peixes em cultivo* (p. 43-80). In: Saran Net, A.; Mariano, W. Dos S.; Sória, SFP (Org.). Tópicos especiais em saúde e criação animal. São Carlos, SP: Pedro & João Editores.
- Tavares-Dias, M. & Moraes, F. R.** (2010). Biochemical parameters for *Piaractus mesopotamicus*, *Colossoma macropomum* (Characidae) and hybrid tambacu (*P. mesopotamicus* X *C. macropomum*). *Ciência Animal Brasileira*, 11(2), 361-368.
- Ucar, A., Atamanalp, M., Kocaman, E. M., Alak, G., Topal, A., Fakioglu, O., Parlak, V., Yanik, T.** (2019). Effects of dietary bentonite on improvements in hematology and enzyme in copper exposed rainbow trout (*Oncorhynchus mykiss*). *Journal of Applied and Physical Sciences*, 5(2), 67-72. https://www.tafpublications.com/gip_content/paper/Japs-5.2.4.pdf
- Urbinati, E.C., Zanuzzo, F., Biller, J.D.** (2020). Stress and immune system in fish. In: Baldisserotto, B.; Urbinati, E.C.; Cyrino, J.E.P (Ed.). *Biology and Physiology of Freshwater Neotropical Fish* (p.93-114). Academic Press.
- Uribe, C., Folch, H., Enríquez, R., Moran, G. J. V. M.** (2011). Innate and adaptive immunity in teleost fish: a review. *Veterinárni medicína*, 56(10), 486-503. <https://vetmed.agriculturejournals.cz/pdfs/vet/2011/10/02.pdf>
- Witeska, M.** (2015). Anemia in teleost fishes. *Bulletin of the European Association of Fish Pathologists*, 35, 4.
- Witeska, M., Kondera, E., Bojarski, B.** (2023). Hematological and hematopoietic analysis in fish toxicology—a review. *Animals*, 13(16), 2625. <https://doi.org/10.3390/ani13162625>
- Zar, J. H.** (2010). *Biostatistical analysis. 5th Edition*. Prentice Hall.