

ISSN 1678-3921

Journal homepage: [www.embrapa.br/pab](http://www.embrapa.br/pab)

For manuscript submission and journal contents,  
access: [www.scielo.br/pab](http://www.scielo.br/pab)

## Diversification of oil-bearing biomass sources for decarbonization and sustainability

**Abstract** – The objective of this review was to analyze the diversification of oil-bearing biomass sources as a strategic pathway to support decarbonization and sustainability, with emphasis on bioenergy in Brazil. The review identifies opportunities, challenges, and conditions to expand the use of diverse oily crops. The work is based on scientific publications, technical reports, and institutional documents addressing biomass and bioenergy production. It highlights the potential of oily crops, such as *macaúba* and tropicalized canola, to expand biomass supply for biofuels and bioproducts. In addition, it examines integrated agricultural systems that combine biomass production with environmental restoration. Such an approach shows that diversifying oil-bearing crops can reduce pressure on traditional feedstocks such as soybean and oil palm, which face sustainability concerns abroad and can be highly affected by climate change. Technological, regulatory, and financial challenges that must be overcome for large-scale adoption of new crops were identified. The review also presents lessons from past failures, showing that success depends on long-term research, breeding, and risk management. Diversifying oil-bearing biomass sources strengthens climate mitigation, rural development, and bioeconomy strategies, positioning Brazil as a leader in the production of sustainable biomass.

**Index terms:** *Acrocomia* spp., energy transition, integrated crop-livestock-forestry systems, low-carbon bioeconomy, regenerative agriculture, tropical canola.

### Diversificação das fontes de biomassa oleaginosa para descarbonização e sustentabilidade

**Resumo** – O objetivo desta revisão foi analisar a diversificação de fontes de biomassa oleaginosa como um caminho estratégico para apoiar a descarbonização e a sustentabilidade, com ênfase na bioenergia no Brasil. O artigo de revisão identifica oportunidades, desafios e condições para a ampliação do uso de diversas culturas oleaginosas. O trabalho baseia-se em publicações científicas, relatórios técnicos e documentos institucionais que tratam da produção de biomassa e bioenergia. Destaca o potencial de culturas oleaginosas, como a macaúba e a canola tropicalizada, para ampliar a oferta de biomassa destinada à produção de biocombustíveis e bioproductos. Além disso, também examina sistemas agrícolas integrados que combinam a produção de biomassa com a restauração ambiental. Essa abordagem mostra que a diversificação de culturas oleaginosas pode reduzir a pressão sobre matérias-primas tradicionais, como a soja e a palma de óleo, que enfrentam questões de sustentabilidade no exterior e podem ser fortemente afetadas

Mauricio Antonio Lopes (✉)   
Embrapa Agroenergia, Brasília, DF, Brazil.  
E-mail: [mauricio.lopes@embrapa.br](mailto:mauricio.lopes@embrapa.br)

Bruno Galveas Laviola   
Embrapa Agroenergia, Brasília, DF, Brazil.  
E-mail: [bruno.laviola@embrapa.br](mailto:bruno.laviola@embrapa.br)

Simone Palma Favaro   
Embrapa Agroenergia, Brasília, DF, Brazil.  
E-mail: [simone.favaro@embrapa.br](mailto:simone.favaro@embrapa.br)

✉ Corresponding author

Received  
May 09, 2025

Accepted  
July 09, 2025

**How to cite**  
LOPES, M.A.; LAVIOLA, B.G.; FAVARO, S.P.  
Diversification of oil-bearing biomass sources  
for decarbonization and sustainability. **Pesquisa  
Agropecuária Brasileira**, v.61, e04127, 2026.  
DOI: <https://doi.org/10.1590/S1678-3921.pab2026.v61.04127>.

por mudanças climáticas. Foram identificados desafios tecnológicos, regulatórios e de mercado que precisam ser superados para a adoção em larga escala de novas culturas. A revisão também apresenta lições aprendidas com fracassos anteriores, evidenciando que o sucesso depende de pesquisa em longo prazo, melhoramento genético e gestão de riscos. Diversificar as fontes de biomassa oleaginosa fortalece as estratégias de mitigação climática, o desenvolvimento rural e a bioeconomia, posicionando o Brasil como líder na produção de biomassas sustentáveis.

**Termos para indexação:** *Acrocomia* spp., transição energética, sistemas de integração lavoura-pecuária-floresta, bioeconomia de baixo carbono, agricultura regenerativa, canola tropical.

## Introduction

The urgency of transitioning to a low-carbon economy has elevated biomass use to a pivotal position in global strategies for climate change mitigation and sustainable development. Society's growing awareness of planetary boundaries and the finitude of natural resources demand a profound reconfiguration of development models traditionally reliant on non-renewable inputs (Sikdar, 2003). International frameworks, such as the Paris Agreement and the United Nations' 2030 Agenda for Sustainable Development, underscore the necessity of reconciling human systems with the environment, highlighting the critical role of renewable resources, such as biomass, in achieving targets for global emission reduction (United Nations, 2015; IPCC, 2021, 2022).

Among the many biomass types with potential to drive decarbonization and sustainability, oil-bearing biomass occupies a particularly strategic position due to its market versatility, high energy density, and compatibility with existing processing and distribution infrastructure. Unlike lignocellulosic or starchy feedstocks, vegetable oils generate liquid biofuels – such as biodiesel, renewable diesel, and sustainable aviation fuels (SAFs) – that are critical for decarbonizing hard-to-electrify sectors, including aviation, heavy-duty transport, and many industrial processes (IEA, 2023; Gilio et al., 2024; Bardon & Massol, 2025).

Additionally, oil-rich species yield valuable co-products such as high-protein meals and biochemicals, reinforcing their relevance within integrated food-energy-feed systems (Alves & Lopes,

2021; Khan, 2023; Motoike & Hilger, 2024; United States, 2025). Several of these species show a relatively high yield per hectare, potential for cultivation in degraded or low-competition areas, and adaptability to crop rotation and agroforestry schemes, further enhancing their systemic contribution to sustainable intensification and landscape regeneration (Colombo et al., 2018; Cortez & Rosillo-Calle, 2023; Zhang et al., 2023). For tropical countries such as Brazil, the strategic development of diversified oil-bearing biomass sources represents a dual opportunity to reduce carbon emissions, while also strengthening rural economies through inclusive and innovation-driven value chains (Alves & Lopes, 2021).

In Brazil, despite the strategic importance of oil-bearing biomass, current vegetable oil production and use remain heavily concentrated in a few traditional oilseed feedstocks, mainly soybean [*Glycine max* (L.) Merr.], accounting for 96% of the 13.26 million tonnes of vegetable oil production, followed by cottonseed and peanut (*Arachis hypogaea* L.), each contributing to 3.1 and 0.6% of the total amount, respectively (USDA-FAS, 2025b). An alternative is the worldwide-used oil palm (*Elaeis guineensis* Jacq.) that accounts for 37% of the output as palm oil or palm kernel oil (USDA-FAS, 2025a). However, in Brazil, this feedstock is produced on a smaller land area, covering 220 thousand hectares, which only render 600 thousand tonnes per year, making the country the tenth producer globally, with a value that is very far from the 46 million tonnes of Indonesia, the main producer (USDA-FAS, 2025b).

This overreliance on a narrow range of oilseed crops not only constrains the diversification potential of the Brazilian bioeconomy but also amplifies supply-chain vulnerabilities, intensifies land-use conflicts, and raises significant environmental and socio-economic trade-offs (Koh & Ghazoul, 2008; Goldemberg et al., 2014; Valin et al., 2015; Carlson & Garrett, 2018; Abreu et al., 2022; IEA, 2022a). Furthermore, global food and agricultural systems, which are among the largest users of soil, water, and biodiversity resources, continue to operate under carbon-intensive paradigms (Rockström et al., 2009; Herrero et al., 2021). The accelerating climate crisis, coupled with unsustainable land management practices and escalating resource consumption, challenges these systems to innovate toward models rooted in circularity, eco-efficiency, and

resilience (IIASA, 2018; Ellen MacArthur Foundation, 2019; IPCC, 2022).

Emerging concepts – such as regenerative agriculture, circular economy, and bioeconomy – designed to mimic nature's closed-loop efficiencies have gained traction as viable models for fostering sustainability (D'Amato et al., 2017; Ellen MacArthur Foundation, 2017). In this context, the diversification of oil-bearing biomass sources presents a critical opportunity to mitigate greenhouse gas emissions, promote biodiversity conservation, enhance resource-use efficiency, and strengthen the food-energy-environment nexus (Jeswani et al., 2020; Lima & Palme, 2022; Subbarao et al., 2023; Millinger et al., 2025).

Tropical countries, particularly Brazil, hold a comparative advantage due to their year-round biomass productivity, biodiversity richness, and established agricultural research and development capacity (Alves & Lopes, 2021; Cortez & Rosillo-Calle, 2023; Souza et al., 2023). Advances in production systems – as integrated crop-livestock-forestry (ICLF) models – and the development or adaptation of underutilized oil-rich crops further strengthen Brazil's position to lead the transition toward a diversified, sustainable, and inclusive bioeconomy. Diversified sources that show possibilities of expansion include: *macaúba* (*Acrocomia* spp.), tropical canola (*Brassica napus* L.), sunflower (*Helianthus annuus* L.), Abyssinian mustard (*Brassica carinata* A.Braun), and oil palm, among others (Henderson et al., 1995; Tomm et al., 2011; Bungenstab & Almeida, 2014; César et al., 2015; Pereira et al., 2018; Motoike & Hilger, 2024).

The targeted transformation will require more than technological advances, also depending on enabling policy frameworks, assessing integrated sustainability, de-risking investment environments, and land-use strategies that align biomass production with ecological integrity and social inclusion (Rathore et al., 2013; Pereira et al., 2017a; Denny, 2020). The resilience and success of future oil-bearing biomass systems will hinge upon balancing productivity with ecosystem health, climate mitigation, and equity, particularly in vulnerable tropical landscapes (Lopes, 2022; Vandenberghe et al., 2022).

The objective of this review was to analyze the diversification of oil-bearing biomass sources as a

strategic pathway to support decarbonization and sustainability, with emphasis on bioenergy in Brazil.

## Materials and Methods

This review critically examines the strategic pathways for diversifying oil-bearing biomass sources to foster global decarbonization and sustainability. For this, it explores the role of alternative feedstocks in reducing emissions across the energy, industrial, and transportation sectors; discusses the integration of biomass production into sustainable land-use systems, including ICLF models; and assesses the regulatory frameworks, investment mechanisms, and technological innovations necessary to enable scalable and sustainable biomass expansion.

This paper adopts a narrative and integrative approach, combining scientific literature, policy documents, and technical reports to provide a comprehensive overview of strategies for diversifying oil-bearing biomass sources in support of decarbonization and sustainability. The literature search was conducted using academic databases such as Scopus (Elsevier, 2025b), Web of Science (Clarivate, 2025), and ScienceDirect (Elsevier, 2025a), as well as institutional sources from the International Energy Agency – IEA (2025b), Food and Agriculture Organization – FAOSTAT (FAO, 2025), International Air Transport Association – IATA (2024b), Brazilian Agricultural Research Corporation – Embrapa (2025), and Brazil's Ministry of Mines and Energy / Energy Research Office – MME/EPE (2025).

Inclusion criteria prioritized peer-reviewed articles and institutional reports published within the last ten years (2014–2024), with emphasis on studies addressing oilseed and other oil-bearing crops for bioenergy use, particularly in tropical or subtropical contexts. Classical or foundational references outside this time window were also retained when relevant to conceptual framing (Rockström et al., 2009; Ellen MacArthur Foundation, 2017). Studies focusing on starch- or lignocellulose-based biomass were excluded unless they contributed comparative insights relevant to vegetable oil production.

This methodological approach allowed the authors to synthesize multidisciplinary evidence and to identify key opportunities, bottlenecks, and governance issues associated with the expansion of sustainable oilseed

biomass systems in Brazil and comparable tropical regions.

### **Biomass role in multisectoral decarbonization**

Biomass is increasingly recognized as a cornerstone of global decarbonization strategies, providing a renewable and adaptable resource to curb greenhouse gas emissions and reduce environmental pollutants across multiple sectors (Zhou et al., 2022; Leslie-Bole et al., 2025; Millinger et al., 2025; United States, 2025). In the energy sector, expanding the portfolio of biomass feedstocks facilitates the shift away from fossil-based electricity and heat generation, significantly reducing emissions of CO<sub>2</sub>, NO<sub>x</sub>, and particulate matter (Mantulet et al., 2020; Lodato et al., 2022). Technological advances such as biomass gasification and anaerobic digestion further enhance the sustainability of bioenergy systems by maximizing energy recovery and mitigating methane emissions from organic waste streams (Mantulet et al., 2020; Alves & Lopes, 2021; Styles et al., 2022; Subbarao et al., 2023).

The World Bioenergy Association (2023) provided a study of the evolution and diversification of biomass-based electricity generation over two decades, indicating the growing contribution of biomass to the global energy matrix, with the dominant generation, in 2020, of solid biomass (69%), followed by municipal and industrial waste (17%), biogas (13%), and liquid biofuels (1.5%). This distribution highlights both the progress made and the continued concentration in a limited set of feedstocks, reinforcing the central argument of this article regarding the need for a broader diversification to enhance sustainability, resilience, and energy security. In this context, it is important to discuss global trends and the strategic role that tropical countries, such as Brazil, can play in expanding the portfolio of biomass sources and systems.

In the industrial sector, the integration of biomass-derived input into high-temperature processes, such as cement and steel production, shows substantial potential for carbon abatement. Recent studies have estimated that substituting even a modest portion of fossil fuels with biomass residues could drive significant reductions in industrial CO<sub>2</sub> emissions by mid-century (Kusuma et al., 2022). Simultaneously, the development of bio-based chemicals and advanced biomaterials opens new avenues for displacing

carbon-intensive inputs, reinforcing circular economy strategies (Alves & Lopes, 2021; Carmona-Martínez et al., 2024).

Oil-bearing biomass is especially important in sectors such as aviation and long-haul transportation, where electrification is not yet viable at scale and drop-in liquid fuels are required. Moreover, the production of advanced biofuels – including cellulosic ethanol, renewable diesel, and SAFs – is crucial for reducing the carbon footprint of these industries (Bardon & Massol, 2025).

In particular, the aviation sector is experiencing a rapidly growing demand for SAFs, driven by international climate commitments, carbon reduction mandates, and the adoption of net-zero roadmaps by airlines and global aviation alliances (IEA, 2025a). According to IATA (2024a), SAF could contribute up to 65% of the sector's required carbon mitigation by 2050. However, SAF production must increase from an estimated 8 million tonnes in 2025 to approximately 23 million tonnes by 2030, and continue increasing toward 2050 under net-zero aligned scenarios. (ATAG, 2024; IATA, 2024b). This exponential growth trajectory underscores the significant role SAFs are expected to play in decarbonizing aviation. However, current production remains below 1% of total jet fuel use, highlighting the urgent need for feedstock diversification and investment in scalable and sustainable supply chains (Pereira et al., 2017b; IEA, 2023; Gilio et al., 2024; Watson et al., 2024). Therefore, expanding the portfolio of biomass sources, particularly from underutilized tropical species, is essential to meet this demand, while ensuring climate and land-use goals are not compromised.

On a global scale, strategic initiatives are accelerating efforts to diversify biomass supplies for the aviation sector. The European Union's "Fit for 55" package is a set of proposals aimed at reducing greenhouse gas emissions by at least 55% by 2030, compared to the levels in 1990 (European Commission, 2023). It promotes the sustainable mobilization of agricultural residues, forestry biomass, and novel energy crops to meet ambitious decarbonization targets. The European Court of Auditors and the European Council detail how the package aims to increase renewable energy's share to at least 42.5% by 2030, with sector-specific measures targeting transport, buildings, and industry to support this transition (Council of the European

Union, 2023). In the United States, the Department of Energy prioritizes research into nonfood bioenergy crops such as switchgrass (*Panicum virgatum* L.) and *Miscanthus* spp., focusing on increasing biomass yields without exacerbating land-use pressures (IEA, 2024). Similarly, India's National Bioenergy Programme fosters the use of nonedible oilseeds and agricultural residues to enhance rural energy security and support low-carbon transitions (Sinha, 2023; Mabee, 2024).

These examples are indicative that, in addition to scaling up SAF production, diversifying feedstocks is also essential to meet future demands and ensure the sustainability of the aviation sector. The Bioenergy Strategic Plan of IEA (2022b) underscores the role of modern bioenergy systems as a critical component of clean energy portfolios, while promoting the diversification of biomass processing and waste use, such as gasification, supporting the European Union's commitment to achieve net-zero emissions by 2050.

### **Oil-bearing biomass: strategic role, opportunities, and diversification**

Plant oils play a crucial role in global agriculture, with oil crops accounting for around 12 % of total agricultural production—positioning them among the top four global crop categories after cereals, sugar crops, and vegetables. Vegetable oils are also a major food commodity, an important source of dietary energy, and economically significant for producers (Mannucci et al., 2023; FAO, 2025). Crops cultivated for their oil richness include oil palm, canola, sunflower, and soybean; the latter is primarily used as a protein source, which serves diverse functions in human diets and in industrial applications as cooking oil, shortenings, and biofuels (Ahmad et al., 2021).

There has been an increase in the cultivation of oily crops due to the growing demand for vegetable oils and biofuels, attributed to population growth, changes in dietary habits, and rising global affluence (Gladek et al., 2017; USDA-ERS, 2024; OECD-FAO, 2025). The global production and consumption of oil-bearing crops have shown a significant upward trend, particularly in emerging economies in South America and Southeast Asia, which benefit from favorable agroclimatic conditions and investments in agricultural technology (Ahmad et al., 2021; Khan, 2023).

As the demand for renewable energy continues to rise, the strategic role of oil-bearing biomass is expected to expand. Research and development efforts are focusing on source diversification, improving the efficiency of biomass conversion technologies and optimizing biomass production systems to enhance yield, while minimizing environmental impacts (United States, 2025). According to National Oilseed Processors Association (NOPA, 2024), by fostering innovations in agricultural practices and processing technologies, oil-bearing biomass can significantly contribute to a sustainable energy future and facilitate multisectoral decarbonization.

The use of oil-bearing biomass offers several environmental benefits, such as reduced lifecycle carbon emissions compared with those of fossil fuels, as well as enhanced carbon sequestration through sustainable agricultural practices (Zhou et al., 2022; Leslie-Bole et al., 2025). However, this potential is accompanied by controversies surrounding land competition, food security, and the environmental impacts of cultivation, raising questions about sustainability and agricultural practices in the context of a rising global demand for vegetable oils and biofuels (Zhang et al., 2023; United States, 2025). In this scenario, innovations in conversion technologies – such as pyrolysis, hydrotreated esters and fatty acids, and transesterification – aim to optimize biomass conversion and energy yields, while minimizing emissions, which highlights the significance of research and development in this field (Masi et al., 2021).

In response to these challenges and opportunities, the strategic role of oil-bearing biomass is expected to expand as previously noted, with ongoing research focusing on improving biomass conversion efficiency and integrating oil-bearing crops into sustainable agricultural practices (NOPA, 2024; United States, 2025). This evolution reflects a broader trend toward a circular economy that prioritizes renewable resources, biodiversity enhancement, and waste reduction, essential for achieving multisectoral decarbonization and climate mitigation goals (El-Araby, 2024; Leslie-Bole et al., 2025). As governments and stakeholders develop supportive policies and financial incentives to promote the adoption of biofuels and sustainable practices, oil-bearing biomass stands at the forefront of the global transition toward a more

sustainable and resilient energy future (Masi et al., 2021; NOPA, 2024).

To illustrate the transition in Brazil, the country's engagement with biofuels began in the 1970s with the implementation of an ethanol program primarily based on sugarcane, aimed at enhancing energetic security (Carvalho, 2007). Over the decades, this initiative positioned Brazil as a global leader in biofuel production, with the development of a mature and sustainable production chain that encompasses a variety of renewable energy sources. The introduction of biodiesel in the 2000s further expanded the country's renewable energy portfolio with the use of vegetable oils and animal fats to diversify biofuel offerings (Hofsetz & Silva, 2012; Bergmann et al., 2013; USDA-ERS, 2024; Branco et al., 2025).

Although soybean is the traditionally used source due to its productivity and economic viability, it has shown low fuel productivity per hectare compared with other oily crops (Bergmann et al., 2013; Huang et al., 2016; Osman et al., 2024), in addition to being subjected to risks related to seasonal availability, price volatility, and regional concentration. Therefore, as the biodiesel industry matures, Brazil has increasingly sought alternative oily crops to supplement soybean use. This diversification is driven by an expanding domestic demand for biodiesel, spurred by regulatory changes that have increased blending mandates, which reached the historical blend of 15% in 2025 (Brasil, 2024).

In recent years, several initiatives have emerged to diversify the vegetable feedstock base, aiming to reduce environmental and socioeconomic vulnerabilities, while enabling the expansion of a low-carbon bioeconomy (Bergmann et al., 2013). Some of the species used as an alternative to soybean are presented below.

### Oil palm

Oil palm is a high-yielding perennial oily crop, originally from West Africa and widely used for biodiesel production in Southeast Asia, especially in Indonesia and Malaysia (Bergmann et al., 2013; Murphy et al., 2021). Due to the restricted environmental conditions to its optimal growth, oil palm is cultivated mostly in the humid zones of the Legal Amazon (Almeida et al., 2020; USDA-FAS, 2022a). In Brazil, cultivation is concentrated in the state of Pará, representing 85%

of total production, and in northern Bahia (USDA-FAS, 2025b), where the crop benefits from favorable conditions. However, the risk of deforestation and land tenure conflicts has triggered environmental concerns. In response, Brazil has implemented the agroecological zoning for oil palm, identifying suitable expansion areas with a low environmental risk (Brasil, 2010). Despite this, commercial expansion has been slow and cautious (Carvalho et al., 2021).

Oil palm renders two kinds of oil from its fruits, each with distinct characteristics. The most abundant, known as palm oil, comes from the mesocarp, and the other, an oilseed, comes from the kernel, accounting for approximately 90% and 10% of the extracted oil, respectively (Bergman et al., 2013; FAO, 2018). With yields ranging from 3.0 to 8.0 tonnes of oil per hectare, a value about ten times that of the oil yield of soybean, oil palm is considered one of the most productive oil crops globally (Bergmann et al., 2013). Under Brazilian current conditions and agricultural practices, the average yield of palm oil is 3 tonnes per hectare (USDA-FAS, 2025b).

Despite palm oil's high productivity and versatility for food and oleochemical uses, it remains marginal in Brazil's biodiesel matrix. In 2024, soybean oil supplied approximately 73–74% of biodiesel feedstock, while beef tallow contributed around 6–8%, and palm oil – along with other vegetable oils – accounted for only about 1–4% (USDA-FAS, 2024; Pinheiro, 2025).

The expansion of oil palm in Brazil is limited by some technical and agronomic challenges (César et al., 2013; Englund et al., 2015). First, palm oil is highly susceptible to quality degradation due to the rapid hydrolysis of triacylglycerides, requiring processing within 24–48 hours post-harvest, and manual harvesting also imposes high labor costs (Ali et al., 2014). Another critical obstacle has been the spread of fatal yellowing, a disease of unknown etiology that causes necrosis and fruit loss, responsible for decimating large plantation areas in Northern Brazil (Silva et al., 2025). This has driven breeding efforts to produce varieties resistant to fatal yellowing, i.e., an interspecific hybrid developed from oil palm and the native *Elaeis oleifera* (Kunth) Cortés. Although disease-resistant, these hybrids have lower oil yields than traditional African varieties and require manual pollination (Freitas et al., 2021; Anhô et al., 2022).

Overall, oil palm remains a promising candidate for large-scale biodiesel and bio-based product

supply chains in Brazil. Its sustainable expansion will depend on overcoming phytosanitary and logistical constraints, improving mechanization and scaling up cultivars resistant to fatal yellowing in order to enhance productivity and resilience.

### ***Macaúba***

*Macaúba*, a palm with a long history of use by pre-Columbian peoples, is rapidly emerging as one of Brazil's most promising biomass resources for sustainable development (Scariot et al., 1995; Evaristo et al., 2016; Vargas-Carpintero et al., 2021; Duque et al., 2025). Widely distributed across the country and adaptable to various soils and climates, *macaúba*, like oil palm, produces high-value oils from both its mesocarp and kernel, which are rich in fatty acids, with applications across the food, cosmetic, pharmaceutical, chemical, and energy sectors (Ciconini et al., 2013; Del Río et al., 2016). These renewable and cost-effective oils enhance the value of a wide array of industrial products (Favaro & Rocha, 2022).

In addition to its oil, *macaúba* provides abundant biomass for energy and other byproducts, making it a multipurpose species ideal for agroforestry systems, integrated crop-livestock-forest systems, and the regeneration of degraded lands (Evaristo et al., 2016; Favaro & Rocha, 2022; Biesdorf et al., 2024). Its resilience to environmental stressors and genetic diversity further strengthen its potential as a climate-smart crop (Del Río et al., 2016; Colombo et al., 2018). As research advances in ecophysiology, reproductive biology, and pest management, *macaúba* is increasingly seen not just as a plant species, but as a strategic ally in creating a more sustainable and resilient agricultural model. By valuing and managing *macaúba* appropriately, Brazil can enhance biodiversity, improve food security, and support the socio-economic well-being of rural communities, while conserving natural resources (Henderson et al., 1995; Lorenzi et al., 1996; Cortez & Rosillo-Calle, 2023; Motoike & Hilger, 2024; Benatti et al., 2025).

Although early studies on the species were sporadic and focused mainly on basic botanical and ethnobotanical aspects, the 2000s marked a turning point, characterized by a sharp rise in publications driven by the growing interest in bioenergy, processing technologies, and potential for sustainable value chains. This surge in scientific output reflects

a broader recognition of the versatility of *macaúba* as a source of vegetable oil and bio-based products (Vargas-Carpintero et al., 2021; Ampese et al., 2023; Duque et al., 2025; Sorita et al., 2025), underscoring its emerging relevance within global discussions on bioeconomy and renewable resources.

In summary, *macaúba* stands out as a promising option for diversifying Brazil's oil-bearing biomass portfolio, combining economic, ecological, and social benefits. Its strategic role in the Brazilian bioeconomy is expected to grow, provided that supportive institutional frameworks, technological innovations, and sustainable agricultural practices continue to advance.

### **Canola**

Canola has also been gaining momentum in Brazil as a versatile biomass feedstock with multiple applications in vegetable oil production, animal feed, and biofuels. Originally introduced in the Southern region of Brazil, particularly in the state of Rio Grande do Sul, canola cultivation has expanded rapidly due to its potential for agricultural diversification, energy independence, and sustainable production systems (Araújo et al., 2021).

In recent years, there has been a notable growth in canola acreage and production. By 2024, Brazilian canola production reached approximately 226,000 tonnes, cultivated across 151,000 hectares, reflecting a growing recognition of its agronomic and economic benefits (MNAgro, 2025). This growth is supported by research and development initiatives focused on developing tropical-adapted hybrids that can thrive beyond traditional temperate zones, targeting regions such as the Cerrado biome and semi-arid areas of the Northeastern region (Araújo et al., 2021). This "tropicalization" strategy is crucial for unlocking canola's broader potential in Brazil's diverse agricultural landscapes.

Economically, canola holds a significant promise. It is widely used to produce edible oil, due to its favorable nutritional profile, and biodiesel, due to its desirable oil properties such as low viscosity and high oxidative stability (Oliveira et al., 2024b; Pimentel et al., 2024). Additionally, canola meals serve as a high-protein component in animal feed, enhancing the crop's value as a multipurpose agricultural commodity (Oliveira et al., 2024b; Pimentel et al., 2024).

From an environmental perspective, canola offers important advantages, fitting well into crop rotation systems, where it contributes to soil health and reduces pressure on large-scale systems such as soybean. Studies suggest that canola could occupy up to 20% of the area traditionally used for soybean, promoting land-use efficiency and crop diversity (Proforest, 2024). Furthermore, the use of genetically modified herbicide-tolerant varieties has been associated with reductions in fuel use and greenhouse gas emissions due to decreased tillage and herbicide applications (Brookes & Barfoot, 2018).

Despite the benefits of the species, challenges persist. The Brazilian canola sector still lags major global producers such as Canada and the European Union, and there are agronomic hurdles related to pest and disease management, climate resilience, and market competitiveness (Canola Council of Canada, 2025; MNAgro, 2025). Additional constraints include regulatory complexities, particularly regarding environmental compliance in sensitive areas such as the Cerrado biome (Pimentel et al., 2024).

Looking ahead, initiatives such as the BRS Canola project, a collaboration between Embrapa and private-sector partners, aim to accelerate the development of high-performing, disease-resistant, and herbicide-tolerant hybrids suited to Brazil's Midwestern and Southern regions (Araújo et al., 2021). According to the same authors, the integration of canola as a winter crop alongside summer staples, as soybean and corn (*Zea mays* L.), offers additional opportunities to enhance land use, improve farm income, and support the expansion of sustainable bioenergy systems.

While Brazil's canola production remains modest on the global stage, its strategic development offers clear opportunities for diversifying the country's oilseed biomass base, enhancing energy security and advancing environmental sustainability.

### **Incorporation of species into diversified agricultural mixed systems**

Integrated crop-livestock (ICL) and ICLF systems have fostered the integration of diversified crops, such as tropical-adapted canola, into regenerative agricultural models, enhancing ecosystem services such as soil health, biodiversity conservation, and water regulation (Telhado & Capdeville, 2021; Cortez & Rosillo-Calle, 2023). Evidence from over

65 peer-reviewed publications shows that adopting ICLF systems significantly benefits soil quality and ecosystem services in Brazil (Oliveira et al., 2024a).

Among the species described previously, *macaúba* is a suitable option for mixed systems. In the context of energy crops, the strategic incorporation of species such as *macaúba* and tropical-adapted canola into diversified agricultural systems offers substantial promises.

In addition to tropical canola and *macaúba*, other crops such as castor bean (*Ricinus communis* L.), crambe (*Crambe abyssinica* Hochst. ex R.E.Fr.), sunflower, Abyssinian mustard, safflower (*Carthamus tinctorius* L.), mustard (*Brassica* spp.), pongam oiltree [*Pongamia pinnata* (L.) Pierre], and *pequi* (*Caryocar brasiliense* Cambess.) may also contribute to Brazil's biomass diversification efforts.

### **Castor bean**

Castor bean is well adapted to Brazil's diverse climates, especially in semi-arid regions such as the Northeast (Ribeiro et al., 2014). Its cultivation is expanding due to the development of more productive and disease-resistant cultivars (Fontgalland, 2021). Castor oil has seen growing industrial demand—especially in pharmaceuticals, cosmetics, and lubricants—and offers a renewable, bio-based alternative to petroleum-derived chemicals (Mubofu, 2016). Moreover, castor bean is considered an excellent option for crop rotation in Brazil, particularly as a second crop following soybean or corn (Witt et al., 2023).

### **Crambe**

Crambe is recognized for its high oil content of 35–60% and short growth cycle, making it a promising feedstock for biodiesel production in Brazil (Bassegio et al., 2016). The crop is tolerant to drought and temperature variations, has a low production cost, and can be grown as a second crop after soybean or corn, using the same equipment; in addition, its cultivation supports crop rotation and soil cover, contributing to sustainable agricultural practices (Queiroz et al., 2021). Crambe oil is rich in erucic acid, valuable for various industrial applications, including lubricants, plastics, and insulating fluids (Bassegio et al., 2016).

## Sunflower

Sunflower is highly adaptable to Brazil's diverse climates, from the South to the North of the country, and is often cultivated as a second crop after soybean or corn, especially in the Cerrado biome and in states like Goiás and Mato Grosso (Castro & Leite, 2018). Although Brazil imports a significant portion of its sunflower oil, domestic production is on the rise—with planted area more than doubling since 2022 and overall production growing by 167%, alongside productivity gains averaging 1.6 t/ha (USDA-FAS, 2025a). Sunflower enhances crop rotation systems, improves ecological balance, and optimizes the use of land and resources, also supporting integrated agricultural systems when its meal is used as animal feed (Iowa State University, 2021). The crop is noted for its drought tolerance and ability to thrive under various climatic conditions, making it a valuable option for diversifying oilseed production (Castro et al., 2014; Jovic et al., 2015; Martínez-Force et al., 2015).

## Abyssinian mustard

Abyssinian mustard is recognized as a non-food oilseed, with a high potential for advanced biofuels such as low-carbon diesel and SAFs, suitable oil for drop-in renewable fuels, and meal that can be used for animal feed (Seepaul et al., 2021). Abyssinian mustard produces a high biomass and can be integrated into existing cropping systems as a cover crop, supporting soil health and crop diversity (Abraha et al., 2024). In addition, its cultivation can improve nutrient cycling and reduce greenhouse gas emissions, making it a scalable and sustainable option (Malins, 2022).

## Safflower

Safflower is recognized for its resilience to drought, high temperatures, saline soil, and low humidity (Thoday-Kennedy et al., 2023). It is suitable for cultivation in various Brazilian regions, as the semi-arid, and during off-season periods due to its deep taproot system and ability to extract water and nutrients from deeper soil layers (Daniel et al., 2024). Research in the states of Mato Grosso and São Paulo showed a significant genetic diversity among safflower genotypes, supporting ongoing breeding programs to improve yield, drought tolerance, and

adaptability to Brazilian climates (Lira et al., 2021). Safflower seeds can contain up to 50% oil and are rich in linoleic and oleic acids, making them valuable for food, pharmaceutical, and industrial applications (Deliorman Orhan et al., 2022). The oil of the species is used in the production of biofuels, paints, varnishes, and cosmetics, for example (Muscalu et al., 2023). Research on safflower as a feedstock for biodiesel and bioethanol production has shown the potential of the species as a sustainable bioenergy crop in Brazil (Lira et al., 2021; Vieira et al., 2021).

## Mustard

Mustard (*Brassica juncea*) is being investigated as a viable alternative feedstock source for biodiesel in Brazil, owing to its favorable agronomic attributes and oil profile (Jham et al., 2009). The crop, which can be grown in rotation with soybean and corn, is adaptable to Brazilian conditions, showing potential as an alternative oilseed crop for energy use despite its still limited commercial cultivation (Madalena et al., 2017).

## Pongam oiltree

Pongam oiltree is a leguminous tree that grows well on degraded and marginal lands, offering potential for land rehabilitation and sustainable biomass production (Hasnah et al., 2020). Due to its resilience to drought and poor soils, the species is a candidate for expanding oilseed production in less fertile areas. As a nitrogen-fixing species, it can also improve soil fertility and support agroforestry systems. The seeds of pongam oiltree are rich in oil, being suitable for biodiesel and other industrial uses.

## Pequi

*Pequi* is a native fruit tree of the Brazilian Cerrado, with significant cultural, culinary, and economic importance, being traditionally used in food, cosmetics, and potentially as industrial fat (Guedes et al., 2017). Its fruit has a high oil content of up to 66% dry weight in the mesocarp, and its productivity can reach up to 2.4 tonnes of oil per hectare with improved agronomic practices (Guedes et al., 2017). Most *pequi* production is extractive, supporting thousands of smallholder families, while contributing to the conservation of the

Cerrado biome (Pinto et al., 2019). However, there are efforts to domesticate and cultivate the species, which could increase yields and the support to sustainable rural livelihoods (Guedes et al., 2017).

Although less visible, the aforementioned crops – crambe, Abyssinian mustard, safflower, mustard, pongam oiltree, and *pequi* – offer diverse agronomic and environmental benefits that can support Brazil's efforts to diversify its biomass base. Their further development and adoption can enhance sustainability, rural livelihoods, and resilience in the face of climate and market challenges.

Regional initiatives across the transition zones between Brazil's Cerrado and Amazon biomes are showing how the integration of diversified biomass sources into local bioenergy value chains can simultaneously drive emission reductions, stimulate socioeconomic development, and foster landscape restoration (Azevedo et al., 2025).

Brazil's longstanding leadership in sugarcane ethanol production offers critical technological, institutional, and policy lessons for scaling the next generation of sustainable biomass systems, including those based on oil-bearing and oilseed species (Silva et al., 2021; Cortez & Rosillo-Calle, 2023). However, the scaling of energy crops also presents risks that must be proactively managed. The unchecked expansion of monocultures – even of native species such as *macaúba* – could threaten biodiversity, reduce landscape resilience, and exacerbate water resource pressures. Therefore, sustainable biomass strategies must prioritize integrated land-use planning, polyculture multifunctional models, and landscape restoration approaches that harmonize biomass expansion with ecosystem integrity and long-term sustainability goals (Telhado & Capdeville, 2021; Lopes, 2022).

### **Sustainable production systems for oil-bearing biomass supply**

The sustainable expansion of biomass supply must be anchored in innovative production systems that simultaneously enhance productivity, restore degraded ecosystems, and foster socioeconomic inclusion. Systemic approaches such as ICL and ICLF and agroforestry systems have emerged as strategic pathways to reconcile biomass production with environmental stewardship, biodiversity conservation,

and climate resilience (Telhado & Capdeville, 2021; Rodrigues et al., 2023; Mattos et al., 2024).

Successful integration of energy crops into sustainable production systems requires strategic selection of species and genotypes suited to local soils and climates, application of ecologically resilient management practices such as crop rotations and soil conservation, and adaptive land-use strategies (Jungers et al., 2023; Dida, 2024; von Cossel et al., 2025). Ecologically, diversified systems contribute to restoring degraded ecosystems, enhancing landscape connectivity, regulating microclimates, and supporting critical ecosystem services such as pollination and water regulation (Carvalho et al., 2024).

Socioeconomically, biomass diversification stimulates rural development by creating new bio-based value chains, attracting investments in biorefineries and decentralized energy systems, while also promoting social inclusion, particularly for small and medium-sized farmers (Sakai et al., 2020; Azevedo et al., 2025). Pilot initiatives in Brazilian states such as Minas Gerais and Mato Grosso have shown that integrating energy crops into sustainable farming models can generate employment, elevate rural incomes, and reinforce the nexus between food, energy, and environmental security (Rodrigues et al., 2023). To maximize the intended impact, it is essential that smallholders and traditional communities are integrated into biomass markets through inclusive financing instruments, technical support programs, and equitable value-chain governance structures. Therefore, sustainable biomass systems must move beyond mere yield maximization, also embracing holistic models that harmonize agronomic productivity, ecological health, and social equity, ensuring that biomass expansion serves as a catalyst for regenerative rural landscapes and climate-smart development.

Brazil's experience in crop tropicalization and in the development of integrated farming systems offers valuable lessons to guide biomass diversification in other tropical regions. For instance, the development of tropical canola hybrids adapted to the Cerrado, combined with the use of bioinputs to reduce chemical fertilizer dependency, represents meaningful progress in sustainable land use (Araújo et al., 2021; MNAgro, 2025). These outcomes reinforce the role of agricultural research as a driver of innovation in the tropical

bioeconomy and as a foundation for an effective public policy (Martha Júnior & Lopes, 2023).

In addition to agronomic and environmental benefits, research and development efforts on alternative biomass in Brazil have also created opportunities for industrial innovation and rural development (Rosillo-Calle, 2000). The integration of plant breeding, agroecological practices, and environmental profile assessment has enabled the design of low-carbon and socially inclusive biomass value chains (OECD, 2015). Such progress is strategic not only for meeting international sustainability standards but also for consolidating a national bioeconomy model anchored in science, technology, and innovation, capable of addressing climate change and energy security challenges (Scheiterle & Birner, 2020; Martha Júnior & Lopes, 2023).

### **Lessons learned from past experiences in oil crop diversification in Brazil**

As Brazil advances to diversify and sophisticate its biomass production capacity, it is important to learn from unsuccessful experiences with oil-bearing crops in the past, among which physic nut (*Jatropha curcas* L.) is the most significant case.

The plant is drought-resistant and can grow under a wide range of climatic and soil conditions, including poor soils, which makes it attractive for cultivation in marginal lands (Durães et al., 2012; van Loo & Jongschaap, 2014; Mazumdar et al., 2018). The species is native to tropical regions of the Americas, including Mexico and Central America, and is characterized as a deciduous shrub or small tree, whose height is typically from 2.0 to 6.0 m, leaves are heart-shaped and lobed, and fruits are green to brown, containing two to three black seeds rich in oil (Durães et al., 2012; van Loo & Jongschaap, 2014; Mazumdar et al., 2018).

As an oil crop, physic nut seeds contain approximately 27–40% oil, which can be processed into high-quality biodiesel compatible with standard diesel engines or used in thermal energy storage and as a heat transfer fluid (Durães et al., 2012; Mazumdar et al., 2018). The crop has been explored for biofuel production due to its resilience and oil yield potential, although commercial success has been limited by agronomic challenges (van Loo & Jongschaap, 2014; Mazumdar et al., 2018). The seeds and other parts of the plant, for example, contain toxic compounds,

limiting their use as food when the proper processing or selection of nontoxic varieties is not carried out (Durães et al., 2012; Mazumdar et al., 2018).

Despite its potential, physic nut remains largely a quasi-undomesticated species in Brazil. This means there is limited scientific knowledge and few established agronomic practices tailored to maximize the productivity of the species under Brazilian conditions. The lack of well-developed cultivation protocols, including planting density, nutrient management, and intercropping systems, hampers consistent yields and the adoption of the crop by farmers (Durães et al., 2012; Silva et al., 2024).

Although physic nut (*Jatropha curcas*) is frequently cited as tolerant of poor soils, empirical studies demonstrate that attaining high fruit and oil yields depends on fertile soils with corrected acidity and adequate fertilization, particularly with phosphorus (P) and potassium (K). Consequently, nutrient deficiencies prevalent in many Brazilian soils restrict physic nut productivity. Moreover, intercropping with cover crops or grains may exacerbate nutrient competition and further diminish yields (Souza et al., 2011; Cañadas-López et al., 2020; Cristiano & Kurihara, 2024). Under a low phosphate availability, physic nut plants are highly dependent on arbuscular mycorrhizal fungi, i.e., in soils with poor mycorrhizal populations or low phosphorus, plant growth and productivity are constrained unless a proper inoculation or fertilization is applied (Balota et al., 2011; Cristiano & Kurihara, 2024).

In summary, the main causes for the unsuccessful cultivation of physic nut in Brazil include: its quasi-undomesticated status, with insufficient agronomic knowledge; nutrient limitations, mainly of phosphorus and potassium; high mycorrhizal dependency; herbicide sensitivity; pest management challenges; and lack of consistent yield data to support its large-scale adoption. Addressing these issues through targeted research and improved cultivation practices is essential to evaluate physic nut's potential as a biofuel crop in Brazil (Balota et al., 2011; Durães et al., 2012; Cristiano & Kurihara, 2024).

There are other cases of an unsuccessful or limited development of oilseed species both in Brazil and globally, often due to agronomic, environmental, and economic challenges such as those faced by physic nut.

Another case is that of castor bean, whose development in Brazil is still slow despite its potential

as a biodiesel feedstock and its adaptability to poor soils. Challenges include limited mechanization, susceptibility to diseases such as gray mold, and the use of genotypes not fully adapted to Brazilian conditions, which restrict its expansion when compared with that of major crops as soybean and corn (Witt et al., 2023).

Another example is that of cottonseed, whose production in Brazil remains a minor part of the oilseed sector, accounting for about 3.4% of the total oilseed volume (USDA-FAS, 2022b). Similarly, sunflower cultivation is still small and vulnerable to climatic stresses such as drought and heat waves, which have caused significant yield and quality losses, making oil extraction economically unfeasible in some scenarios (USDA-FAS, 2025a).

These cases illustrate that the unsuccessful or limited development of oil-bearing crops often stems from a combination of biological constraints, inadequate breeding and agronomic practices, environmental stress, disease and pest pressures, and economic factors such as market demand and infrastructure. The physic nut case, specifically, teaches us that the successful introduction of new oily crops in Brazil requires a comprehensive approach: domestication and breeding, soil and nutrient management, understanding of ecological interactions, integrated pest and weed control, and long-term commitment to research and farmer support. These lessons can help avoid pitfalls and accelerate the development of other promising but less-known species for biomass diversification in the country.

### **Bases for biomass diversification: regulatory frameworks, economic instruments, and scaling up**

The successful diversification of biomass sources requires not only technological and agronomic innovations but also the establishment of robust regulatory frameworks, financial incentives, and risk mitigation mechanisms that support scalability and sustainability. Public policies must deliver consistent and long-term signals that encourage investment in diversified bioenergy systems, while ensuring environmental integrity and social inclusion (Grangeira et al., 2022; Souza et al., 2023).

Brazil stands out as a dynamic example in which biomass diversification aligns with climate ambition and bioeconomic development (Alves & Lopes, 2021; Bhandari & Sessa, 2020). RenovaBio, officially

known as the national biofuel policy of Brazil, was established by Law N° 13.576/2017 (Brazil, 2017) to promote the predictable expansion of biofuels in the country's energy matrix. It aims to support Brazil's commitments under the Paris Agreement by fostering the decarbonization of the transport sector, while introducing annual decarbonization targets for fossil fuel distributors, which must be met through the acquisition of decarbonization credits issued by certified biofuel producers based on their energy efficiency and life-cycle greenhouse gas emissions. RenovaBio creates market signals that reward a more sustainable and efficient biofuel production, encourage investment in low-carbon technologies, and integrate environmental performance into the energy sector. It is regarded as one of the most advanced global frameworks for promoting sustainable biofuels through scientifically grounded and market-based mechanisms. The program has been instrumental in advancing carbon-intensity-based certification frameworks, encouraging the production of low-carbon biofuels and promoting the adoption of new feedstocks beyond traditional crops such as sugarcane and soybean (Goldemberg et al., 2018; Salina et al., 2019; Denny, 2020; Martinelli et al., 2022; Ribeiro & Cunha, 2022; Müller-Langers, 2023; Sá et al., 2023).

Certification schemes and comprehensive life-cycle assessment methodologies (Osman et al., 2021; Rossi et al., 2024) are indispensable to ensure that biomass expansion does not trigger adverse impacts such as deforestation, biodiversity loss, or social displacement (Ogwu et al., 2025). Examples include voluntary standards and sector-specific certifications, as the roundtable on sustainable biomaterials (RSB, 2025) and Bonsucro (2025) for sugarcane, respectively, which provide transparency and traceability across biomass supply chains. The integration of life-cycle assessment frameworks into national and international policies enables a precise accounting for greenhouse gas emissions, water use, land-use changes, and overall environmental impacts (Parada et al., 2022).

In this scenario, developing risk mitigation tools is essential to unlock private capital for sustainable biomass initiatives. Instruments such as climate risk insurance, blended finance mechanisms, and public-private de-risking facilities are increasingly deployed to navigate infrastructural and environmental uncertainties, particularly in tropical regions (Ogwu et al., 2025). Furthermore, innovative financial

solutions – including ecosystem service payments, carbon credits from agroforestry bioenergy systems, and green bonds linked to biomass value chains – are providing new revenue streams and enhancing project bankability (Henry et al., 2023).

Looking forward, the investment outlook for the tropical bioeconomy remains highly promising. Tropical nations, specifically Brazil, show comparative advantages such as year-round biomass productivity and an immense biodiversity, which can be leveraged to competitively produce low-carbon fuels, biomaterials, and certified carbon credits (Souza et al., 2023). However, achieving transformative impacts will depend on establishing enabling ecosystems that integrate smallholders into biomass markets, invest in logistics and biorefinery infrastructure, and ensure equitable benefit distribution (Hiloidhari et al., 2023).

Ultimately, the future of biomass diversification will rely on building integrated ecosystems in which regulatory certainty, financial incentives, certification standards, and risk-sharing mechanisms converge synergistically. Such comprehensive approaches are essential for unlocking the full potential of tropical bioeconomy to drive energy transitions, climate mitigation, and inclusive rural development (Iriarte et al., 2021; Alves & Lopes, 2021; Lopes, 2022; Hiloidhari et al., 2023).

### Concluding Remarks

This review shows that diversifying oil-bearing biomass sources is not only feasible but imperative for advancing decarbonization and sustainability in Brazil and other tropical regions. The presented findings clearly confirm that expanding the portfolio of oily crops – such as *macaúba* and tropical-adapted canola – offers substantial opportunities to enhance biomass supply for bioenergy, bio-based products, and regenerative agricultural systems, while simultaneously promoting ecosystem restoration and rural development.

Brazil's distinctive combination of ecological conditions, agricultural expertise, and bioeconomy policy frameworks places the country in a strong position to lead this transition. However, several unresolved challenges remain, mainly regarding institutional, regulatory, and financial barriers that hinder the scaling of diversified plant oil-based systems

beyond pilot initiatives. Unlocking the full potential of these crops will require integrated approaches that align agricultural innovation with climate policies, inclusive business models, and multifunctional land-use strategies.

At the same time, it is essential to recognize that the global demand for plant oils is growing rapidly, driven by population growth, dietary shifts, industrial applications, and the accelerating push for biofuels. This trend places increasing pressure on already dominant systems, particularly soybean and oil palm, which face significant sustainability concerns related to deforestation, biodiversity loss, greenhouse gas emissions, and social conflicts. The overreliance on these crops risks exacerbating land-use tensions and undermining global climate and conservation goals. Therefore, diversifying the oily crop portfolio through the development of underutilized species and regenerative production models is not merely an economic strategy, but a necessary response to mitigate systemic risks and ensure that the expansion of biomass supply does not further compromise environmental integrity or social equity.

Moreover, the lessons learned from past failures with species such as physic nut highlight the need for long-term investments in domestication, breeding, agronomic management, and risk assessment. Without such foundational work, even promising species risk repeating the shortcomings of earlier efforts.

Future research must deepen the understanding of how diversified systems can contribute simultaneously to energy transitions, biodiversity conservation, and food security. This includes the development of advanced metrics for ecological and social co-benefits, new models for integrating smallholders into value chains, and financing mechanisms that reward landscape-level sustainability outcomes.

In sum, plant oil diversification represents not only a pathway for biomass expansion but a catalyst for systemic transformation, offering Brazil and similar regions an opportunity to reconcile agricultural development with climate mitigation, economic resilience, and social inclusion. Moving forward, success will depend on building robust research, policy, and investment ecosystems capable of sustaining this transformation over the long term.

## References

- ABRAHA, R.; YGZAW, W.; SHAIBU, A.; BADJI, A.; EDEMA, R.; EKWAMU, A.; CLEREK, M. Ethiopian mustard emerging as non-food feedstock and environmentally sustainable crop: current status, constraints, and future prospects. **African Journal of Rural Development**, v.9, p.37-52, 2024.
- ABREU, M.; SILVA, L.; RIBEIRO, B.; FERREIRA, A.; ALVES, L.; PAIXÃO, S.M.; GOUVEIA, L.; MOURA, P.; CARVALHEIRO, F.; DUARTE, L.C.; FERNANDO, A.L.; REIS, A.; GÍRIO, F. Low indirect land use change (ILUC) energy crops to bioenergy and biofuels - a review. **Energies**, v.15, art.4348, 2022. DOI: <https://doi.org/10.3390/en15124348>.
- AHMAD, M.; WARAICH, E.A.; SKALICKY, M.; HUSSAIN, S.; ZULFIQAR, U.; ANJUM, M.Z.; ZULFIQAR, U.; ANJUM, M.Z.; HABIB UR RAHMAN, M.; BRESTIC, M.; RATNASEKERA, D.; LAMILLA-TAMAYO, L.; AL-ASHKAR, I.; EL SABAGH, A. Adaptation strategies to improve the resistance of oilseed crops to heat stress under a changing climate: an overview. **Frontiers in Plant Science**, v.12, art.767150, 2021. DOI: <https://doi.org/10.3389/fpls.2021.767150>.
- ALI, F.S.; SHAMSUDIN, R.; YUNUS, R. The effect of storage time of chopped oil palm fruit bunches on the palm oil quality. **Agriculture and Agricultural Science Procedia**, v.2, p.165-172, 2014. DOI: <https://doi.org/10.1016/j.aaspro.2014.11.024>.
- ALMEIDA, A.S.; VIEIRA, I.C.G.; FERRAZ, S.F. Long-term assessment of oil palm expansion and landscape change in the eastern Brazilian Amazon. **Land Use Policy**, v.90, art.104321, 2020.
- ALVES, A.A.; LOPES, M.A. O setor sucroenergético na emergente bioeconomia. **Revista Opiniões**, v.18, p.58-59, 2021. OpAA69. Available at: <https://sucroenergetico.revistaopinioes.com.br/pt-br/revista/detalhes/16-o-setor-sucroenergetico-na-emergente-bioeconomia/>. Accessed on: July 1 2025.
- AMPESE, L.C.; BULLER, L.S.; MONROY, Y.M.; GARCIA, M.P.; RAMOS-RODRIGUEZ, A.R.; FORSTER-CARNEIRO, T. Macaúba's world scenario: a bibliometric analysis. **Biomass Conversion and Biorefinery**, v.13, p.3329-3347, 2023. DOI: <https://doi.org/10.1007/s13399-021-01376-2>.
- ANHÊ, B.B.; SANTOS, A.V.F. dos; SILVA, T.A.F. da; CARVALHO, L.L.B. de; FARIAS, P.R.S. Spatial-temporal distribution of fatal yellowing in different oil palm genetic materials in eastern amazon. **Information Processing in Agriculture**, v.9, p.365-377, 2022. DOI: <https://doi.org/10.1016/j.inpa.2021.08.002>.
- ARAÚJO, L.N. de; ROSADO, T.B.; RODRIGUES, E.V.; SANTOS, A. dos; LAVIOLA, B.G. Tropicalization of canola (*Brassica napus* L.): commercial hybrids show potential for cultivation in the Brazilian Cerrado. **Revista de la Facultad de Ciencias Agrarias UNCuyo**, v.53, p.20-26, 2021. DOI: <https://doi.org/10.48162/rev.39.035>.
- ATAG. Air Transport Action Group. **Sustainable aviation fuel**. 2024. Available at: <https://atag.org/industry-topics/sustainable-aviation-fuel>. Accessed on: Sept. 5 2025.
- AZEVEDO, G.; ALBA, J.G.; SILVA, T. de C.P. e. Restoring Brazil's rainforest can power an environmental and economic transformation. **IDB Invest blog**, May 31 2025. Available at: <https://idbinvest.org/en/blog/development-impact/restoring-rainforest-can-drive-powerful-environmental-and-economic>. Accessed on: Feb. 13 2026.
- BALOTA, E.L.; MACHINESKI, O.; TRUBER, P.V.; SCHERER, A.; SOUZA, F.S. de. Physic nut plants present high mycorrhizal dependency under conditions of low phosphate availability. **Brazilian Journal of Plant Physiology**, v.23, p.33-44, 2011. DOI: <https://doi.org/10.1590/S1677-04202011000100006>.
- BARDON, P.; MASSOL, O. Decarbonizing aviation with sustainable aviation fuels: myths and realities of the roadmaps to net zero by 2050. **Renewable and Sustainable Energy Reviews**, v.211, art.115279, 2025. DOI: <https://doi.org/10.1016/j.rser.2024.115279>.
- BASSEGIO, D.; ZANOTTO, M.D.; SANTOS, R.F.; WERNCKE, I.; DIAS, P.P.; OLIVO, M. Oilseed crop crambe as a source of renewable energy in Brazil. **Renewable and Sustainable Energy Reviews**, v.66, p.311-321, 2016. DOI: <https://doi.org/10.1016/j.rser.2016.08.010>.
- BENATTI, G.S. de S.; BUAINAIN, A.M.; CAVALCANTE FILHO, P.G.; VARGAS-CARPINTERO, R.; ASVELD, L.; OSSEWEIJER, P. Macaw palm (*Acrocomia* spp.): an opportunity for including smallholders in Brazil's biodiesel production. **Cleaner and Circular Bioeconomy**, v.10, art.100134, 2025. DOI: <https://doi.org/10.1016/j.clcb.2025.100134>.
- BERGMANN, J.C.; TUPINAMBÁ, D.D.; COSTA, O.Y.A.; ALMEIDA, J.R.M.; BARRETO, C.C.; QUIRINO, B.F. Biodiesel production in Brazil and alternative biomass feedstocks. **Renewable and Sustainable Energy Reviews**, v.21, p.411-420, 2013. Special Agriculture 4.0. DOI: <https://doi.org/10.1016/j.rser.2012.12.058>.
- BHANDARI, R.; SESSA, V. Energy in agriculture in Brazil. **Revista Ciência Agronômica**, v.51, e20207784, 2020. Special Agriculture 4.0. DOI: <https://doi.org/10.5935/1806-6690.20200098>.
- BIESDORF, E.M.; FÁVARO, S.P.; CONCEIÇÃO, L.D.H.C.S. da; SÁ, S.F. de; PIMENTEL, L.D. Production and quality of leaf biomass from *Acrocomia aculeata* for bioenergy. **Industrial Crops and Products**, v.214, art.118501, 2024. DOI: <https://doi.org/10.1016/j.indcrop.2024.118501>.
- BONSUCRO. Available at: <https://bonsucro.com/>. Accessed on: Aug. 7 2025.
- BRANCO, J.E.H.; PRATES, R.C.; OLIVEIRA, A.L. do N.; CARVALHO, A.K.F. de. Evaluation of the greenhouse gas emissions of the Brazilian biodiesel and the impact of the mandatory blending of the biofuel into commercial diesel fuel. **Chemosphere**, v.376, art.144268, 2025. DOI: <https://doi.org/10.1016/j.chemosphere.2025.144268>.
- BRASIL. **Decreto nº 7.172, de 7 de maio de 2010**. Aprova o zoneamento agroecológico da cultura da palma de óleo e dispõe sobre o estabelecimento pelo Conselho Monetário Nacional de normas referentes às operações de financiamento ao segmento da palma de óleo, nos termos do zoneamento. Available at: <https://>

[www.planalto.gov.br/ccivil\\_03/\\_ato2007-2010/2010/decreto/d7172.htm](http://www.planalto.gov.br/ccivil_03/_ato2007-2010/2010/decreto/d7172.htm)>. Accessed on: Sept. 5 2025.

BRASIL. **Lei nº 14.993, de 8 de outubro de 2024**. Dispõe sobre a promoção da mobilidade sustentável de baixo carbono e a captura e a estocagem geológica de dióxido de carbono; institui o Programa Nacional de Combustível Sustentável de Aviação (ProBioQAV), o Programa Nacional de Diesel Verde (PNDV) e o Programa Nacional de Descarbonização do Produtor e Importador de Gás Natural e de Incentivo ao Biometano; altera as Leis nºs 9.478, de 6 de agosto de 1997, 9.847, de 26 de outubro de 1999, 8.723, de 28 de outubro de 1993, e 13.033, de 24 de setembro de 2014; e revoga dispositivo da Lei nº 10.438, de 26 de abril de 2002. Available at: <<https://www2.camara.leg.br/legin/fed/lei/2024/lei-14993-8-outubro-2024-796443-publicacaooriginal-173317-pl.html>>. Accessed on: Sept. 5 2025.

BRAZIL. **Law No. 13,576 of December 26, 2017**. Provides for the Brazilian Biofuels Policy (RenovaBio). Available at: <<https://www.gov.br/anp/pt-br/assuntos/renovabio/arg/law-13576-2017.pdf>>. Accessed on: Sept. 5 2025.

BROOKES, G.; BARFOOT, P. Environmental impacts of genetically modified (GM) crop use 1996-2016: impacts on pesticide use and carbon emissions. **GM Crops & Food**, v.9, p.109-139, 2018.

BUNGENSTAB, D.J.; ALMEIDA, R.G. de. (Ed.). **Integrated crop-livestock-forestry systems: a Brazilian experience for sustainable farming**. Brasília: Embrapa, 2014.

CAÑADAS-LÓPEZ, Á.; RADE-LOOR, D.; SIEGMUND-SCHULTZE, M.; ZAMBRANO-CEDENO, X.; VARGAS-HERNÁNDEZ, J.; WEHENKEL, C. Pruning and fertilization effects on jatropha yields under smallholder's conditions in a tropical dry forest from Ecuador. **Revista Facultad Nacional de Agronomía Medellín**, v.73, p.9089-9097, 2020.

CANOLA COUNCIL OF CANADA. Plant establishment: environmental effects. In: CANOLA COUNCIL OF CANADA. **Canola encyclopedia**. 2025. Available at: <<https://www.canolacouncil.org/canola-encyclopedia/plant-establishment/environmental-effects/>>. Accessed on: July 1 2025.

CARLSON, K.M.; GARRETT, R.D. Environmental impacts of tropical soybean and palm oil crops. In: OXFORD Research Encyclopedia of Environmental Science. Oxford: Oxford University Press, 2018. DOI: <https://doi.org/10.1093/acrefore/9780199389414.013.234>.

CARMONA-MARTÍNEZ, A.A.; RUEDA, A.; JARAUTA-CÓRDOBA, C.A. Deep decarbonization of the energy intensive manufacturing industry through the bioconversion of its carbon emissions to fuels. **Fuel**, v.371-A, art.131922, 2024. DOI: <https://doi.org/10.1016/j.fuel.2024.131922>.

CARVALHO, C.B. de MACÊDO; MELLO, A.C.L. de; CUNHA, M.V. da; APOLINÁRIO, V.X. de O.; DUBEUX JÚNIOR, J.C.B.; SILVA, V.J. da; MEDEIROS, A.S.; IZIDRO, J.L.P.S.; BRETAS, I.L. Ecosystem services provided by silvopastoral systems: a review. **The Journal of Agricultural Science**, v.162, p.417-432, 2024. DOI: <https://doi.org/10.1017/S0021859624000595>.

CARVALHO, C.M. de; IWAMA, A.Y.; LA ROVERE, E.L. Scenarios for oil palm expansion in degraded and deforested

lands in the Brazilian Amazon to meet biodiesel demand. **Sustainability in Debate**, v.12, p.90-107, 2021. DOI: <https://doi.org/10.18472/SustDeb.v12n2.2021.35201>.

CARVALHO, L.C.C. Ethanol: a Brazilian successful story. In: WROBEL, P. (Ed). **Clean energy: the Brazilian ethanol experience**. London: [s.n.], 2007.

CASTRO, C. de; OLIVEIRA, F.A.; OLIVEIRA JUNIOR, A.; RAMOS, N.P. Nutrition and fertilization of sunflowers in Brazilian cerrado. In: ARRIBAS, J.E. (Ed.). **Sunflowers: growth and development, environmental influences and pests/diseases**. New York: Nova Science Pubs, 2014. p.257-279.

CASTRO, C.; LEITE, R.M.V.B.C. Main aspects of sunflower production in Brazil. **OCL**, v.25, art.D104, 2018. DOI: <https://doi.org/10.1051/ocl/2017056>.

CÉSAR, A. da S.; ALMEIDA, F. de A.; SOUZA, R.P. de; SILVA, G.C.; ATABANI, A.E. The prospects of using *Acrocomia aculeata* (macaúba) a non-edible biodiesel feedstock in Brazil. **Renewable and Sustainable Energy Reviews**, v.49, p.1213-1220, 2015. DOI: <https://doi.org/10.1016/j.rser.2015.04.125>.

CÉSAR, A. da S.; BATALHA, M.O.; ZOPELARI, A.L.M.S. Oil palm biodiesel: Brazil's main challenges. **Energy**, v.60, p.485-491, 2013. DOI: <https://doi.org/10.1016/j.energy.2013.08.014>.

CICONINI, G.; FAVARO, S.P.; ROSCOE, R.; MIRANDA, C.H.B.; TAPETI, C.F.; MIYAHIRA, M.A.M.; BEARARI, L.; GALVANI, F.; BORSATO, A.V.; COLNAGO, L.A.; NAKA, M.H. Biometry and oil contents of *Acrocomia aculeata* fruits from the Cerrados and Pantanal biomes in Mato Grosso do Sul, Brazil. **Industrial Crops and Products**, v.45, p.208-214, 2013. DOI: <https://doi.org/10.1016/j.indcrop.2012.12.008>.

CLARIVATE. **Web of Science**. Platform overview. 2025. Available at: <<https://clarivate.com/webofsciencelgroup/solutions/web-of-science/>>. Accessed on: Sept. 5 2025.

COLOMBO, C.A.; BERTON, L.H.C.; DIAZ, B.G.; FERRARI, R.A. Macauba: a promising tropical palm for the production of vegetable oil. **OCL**, v.25, art.D108, 2018. DOI: <https://doi.org/10.1051/ocl/2017038>.

CORTEZ, L.A.B.; ROSILLO-CALLE, F. **The future role of biofuels in the new energy transition: lessons and perspectives of biofuels in Brazil**. São Paulo: Blucher, 2023.

COUNCIL OF THE EUROPEAN UNION. **Fit for 55**. Brussels: European Council, 2023. Available at: <<https://www.consilium.europa.eu/en/policies/fit-for-55/>>. Accessed on: July 1 2025.

CRISTIANO, M.Ã.; KURIHARA, C.H. Soil chemical properties and production of physic nut intercropped with forage plants and grain crops. **Journal of Agricultural Science**, v.16, p.1-57, 2024.

D'AMATO, D.; DROSTE, N.; ALLEN, B.; KETTUNEN, M.; LÄHTINEN, K.; KORHONEN, J.; LESKINEN, P.; MATTHIES, B.D.; TOPPINEN, A. Green, circular, bio economy: a comparative analysis of sustainability avenues. **Journal of Cleaner Production**, v.168, p.716-734, 2017. DOI: <https://doi.org/10.1016/j.jclepro.2017.09.053>.

DANIEL, C.; SANTOS, R.F.; PINTO, J. de S.; HUBNER, V.; PAZUCH, F.A. Effect of water deficit on safflower cultivation at different phenological stages. **Engenharia Agrícola**, v.44,

- e20230067, 2024. DOI: <https://doi.org/10.1590/1809-4430-Eng-Agric.v44e20230067/2024>.
- Del RÍO, J.C.; EVARISTO, A.B.; MARQUES, G.; MARTÍN-RAMOS, P.; MARTÍN-GIL, J.; GUTIÉRREZ, A. Chemical composition and thermal behavior of the pulp and kernel oils from macauba palm (*Acrocomia aculeata*) fruit. **Industrial Crops and Products**, v.84, p.294-304, 2016. DOI: <https://doi.org/10.1016/j.indcrop.2016.02.018>.
- DELİORMAN ORHAN, D.; PEKACAR, S.; ULUTAŞ, O.K.; ÖZÜPEK, B.; SÜMMEOĞLU, D.; BERKKAN, A. Assessment of commercially safflower oils (*Carthami Oleum raffinatum*) in terms of European Pharmacopoeia criteria and their weight control potentials. **Turkish Journal of Pharmaceutical Sciences**, v.19, p.273-279, 2022. DOI: <https://doi.org/10.4274/tjps.galenos.2021.84484>.
- DENNY, D.M.T. Competitive renewables as the key to energy transition – RenovaBio: the Brazilian biofuel regulation. In: GUIMARÃES, L.N. (Ed.). **The Regulation and Policy of Latin American Energy Transitions**. London: Elsevier, 2020. p.223-242. DOI: <https://doi.org/10.1016/B978-0-12-819521-5.00013-9>.
- DIDA, G. Biotechnology towards energy crops. **CABI Agriculture and Bioscience**, v.5, art.45, 2024. DOI: <https://doi.org/10.1186/s43170-024-00245-y>.
- DUQUE, T.S.; BARROSO, G.M.; BORGES, C.E.; MENDES, D.S.; SILVA, R.S. da; EVARISTO, A.B.; SANTOS, J.B. dos. Current and future development of *Acrocomia aculeata* focused on biofuel potential and climate change challenges. **Scientific Reports**, v.15, art.8120, 2025. DOI: <https://doi.org/10.1038/s41598-025-92681-7>.
- DURÃES, F.O.M.; LAVIOLA, B.G.; ALVES, A.A. Potential and challenges in making physic nut (*Jatropha curcas* L.) a viable biofuel crop: the Brazilian perspective. **CABI Reviews**, v.7, p.1-8, 2012. DOI: <https://doi.org/10.1079/PAVSNNR20116043>.
- EL-ARABY, R. Biofuel production: exploring renewable energy solutions for a greener future. **Biotechnology for Biofuels and Bioproducts**, v.17, art.129, 2024. DOI: <https://doi.org/10.1186/s13068-024-02571-9>.
- ELLEN MACARTHUR FOUNDATION. **Completing the picture: how the circular economy tackles climate change**. 2019. Available at: <https://ellenmacarthurfoundation.org/completing-the-picture>. Accessed on: Sept. 1 2025.
- ELLEN MACARTHUR FOUNDATION. **What is the circular economy?** 2017. Available at: <https://ellenmacarthurfoundation.org>. Accessed on: Sept. 1 2025.
- ELSEVIER. **ScienceDirect**: about. 2025a. Available at: <https://www.sciencedirect.com/>. Accessed on: Sept. 5 2025.
- ELSEVIER. **Scopus**: about: coverage. 2025b. Available at: <https://www.elsevier.com/solutions/scopus>. Accessed on: Sept. 5 2025.
- EMBRAPA. Brazilian Agricultural Research Corporation. **Portal Embrapa: Publications and Libraries**. 2025. Available at: <https://www.embrapa.br/>. Accessed on: Sept. 5 2025.
- ENGLUND, O.; BERNDES, G.; PERSSON, U.M.; SPAROVEK, G. Oil palm for biodiesel in Brazil—risks and opportunities. **Environmental Research Letters**, v.10, art.044002, 2015. DOI: <https://doi.org/10.1088/1748-9326/10/4/044002>.
- EPE. Empresa de Pesquisa Energética. **Brazilian Energy Balance 2025: year 2024**. Rio de Janeiro, 2025. Available at: <https://www.epe.gov.br/pt/publicacoes-dados-abertos/publicacoes/balanco-energetico-nacional-ben>. Accessed on: Sept. 5 2025.
- EUROPEAN COMMISSION. **2030 climate targets**. 2023. Available at: [https://climate.ec.europa.eu/eu-action/climate-strategies-targets/2030-climate-targets\\_en](https://climate.ec.europa.eu/eu-action/climate-strategies-targets/2030-climate-targets_en). Accessed on: Sept. 5 2025.
- EVARISTO, A.B.; MARTINO, D.C.; FERRAREZ, A.H.; DONATO, D.B.; CARNEIRO, A. de C.O.; GROSSI, J.A.S. Potencial energético dos resíduos do fruto da macaúba e sua utilização na produção de carvão vegetal. **Ciência Florestal**, v.26, p.571-577, 2016. DOI: <https://doi.org/10.5902/1980509822757>.
- FAO. Food and Agriculture Organization of the United Nations. **Faostat: oil, palm fruit**. 2018. Available at: <https://web.archive.org/web/20180817161440/http://ref.data.fao.org/dimension-member?entryId=2095d9b1-bd16-41ac-a315-2673fb88549a>. Accessed on: Sept. 5 2025.
- FAO. Food and Agriculture Organization of the United Nations. **Faostat Statistical Database**. 2025. Available at: <https://www.fao.org/faostat/>. Accessed on: Sept. 5 2025.
- FAVARO, S.P.; ROCHA, J.D. **A nova cadeia produtiva da macaúba para bioprodutos e descarbonização**. Brasília: Embrapa Agroenergia, 2022. 31 p. (Embrapa Agroenergia. Documentos, 46). Available at: <https://www.infoteca.cnptia.embrapa.br/infoteca/handle/doc/1149154>. Accessed on: Sept. 1 2025.
- FONTGALLAND, I.L. Castor bean oil and the matrix of energy in Brazil. **e-Acadêmica**, v.2, e012332, 2021. DOI: <https://doi.org/10.52076/eacad-v2i3.32>.
- FREITAS, A.F. de; LOPES, R.; CUNHA, R.N.V. da; PINA, A.J. de A.; RESENDE, M.D.V. de; GOMES JÚNIOR, R.A. Genotypic values and genetic correlations for components and oil content of bunch of hybrids between caiaué and dendê. **Colloquium Agrariae**, v.17, p.79-92, 2021.
- GILIO, L.; BORGES, P.; JANK, M.S.; RIBEIRO, P.H.C. A bioenergia tropical na transição energética global. **Inspere Agro Global**, jan. 2024. Available at: <https://agro.insper.edu.br/storage/papers/January2024/1AGbioenergiaTransicaoEnergeticaGlobal.pdf>. Accessed on: Sept. 1 2025.
- GOLDEMBERG, J.; COELHO, S.T.; GUARDABASSI, P. The sustainability of ethanol production from sugarcane. In: SORENSEN, B. (Ed.). **Renewable Energy**. London: Routledge, 2018. v.3, p.321-345.
- GOLDEMBERG, J.; MELLO, F.F.; CERRI, C.E.; DAVIES, C.A.; CERRI, C.C. Meeting the global demand for biofuels in 2021 through sustainable land use change policy. **Energy Policy**, v.69, p.14-18, 2014. DOI: <https://doi.org/10.1016/j.enpol.2014.02.008>.
- GRANGEIA, C.; SANTOS, L.; LAZARO, L.L.B. The Brazilian biofuel policy (RenovaBio) and its uncertainties: an assessment of technical, socioeconomic, and institutional aspects. **Energy**

- Conversion and Management: X**, v.13, art.100156, 2022. DOI: <https://doi.org/10.1016/j.ecmx.2021.100156>.
- GUEDES, A.M.M.; ANTONIASSI, R.; FARIA-MACHADO, A.F. de. Pequi: a Brazilian fruit with potential uses for the fat industry. **OCL**, v.24, art.D507, 2017. DOI: <https://doi.org/10.1051/ocl/2017040>.
- HASNAH, T.; LEKSONO, B.; SUMEDI, N.; WINDYARINI, E.; ADINUGRAHA, H.A.; BARAL, H.; ARTATI, Y. Pongamia as a potential biofuel crop: oil content of Pongamia pinnata from the best provenance in Java, Indonesia. In: 2020 INTERNATIONAL CONFERENCE AND UTILITY EXHIBITION ON ENERGY, ENVIRONMENT AND CLIMATE CHANGE, 2020, Pattaya. **Proceedings**. [S.l.]: IEEE, 2020. 6p. ICUE 2020. DOI: <https://doi.org/10.1109/ICUE49301.2020.9307094>.
- HENDERSON, A.; GALEANO, G.; BERNAL, R. **Field guide to the palms of the Americas**. Princeton: Princeton University Press, 1995.
- HENRY, G.; TRIGO, E.; JARAMILLO, E.H. de. **New framework for sustainable growth in Latin America**: lessons from the bioeconomy strategies. Montpellier: CIRAD, 2023. Available at: [https://agritrop.cirad.fr/592946/8/ID592946\\_EN.pdf](https://agritrop.cirad.fr/592946/8/ID592946_EN.pdf). Accessed on: Sept. 1 2025.
- HERRERO, M.; THORNTON, P.K.; MASON-D'CROZ, D.; PALMER, J.; BODIRSKY, B.L.; PRADHAN, P.; BARRETT, C.B.; BENTON, T.G.; HALL, A.; PIKAAR, I.; BOGARD, J.R.; BONNETT, G.D.; BRYAN, B.A.; CAMPBELL, B.M.; CHRISTENSEN, S.; CLARK, M.; FANZO, J.; GODDE, C.M.; JARVIS, A.; LOBOGUERRERO, A.M.; MATHYS, A.; MCINTYRE, C.L.; NAYLOR, R.L.; NELSON, R.; OBERSTEINER, M.; PARODI, A.; POPP, A.; RICKETTS, K.; SMITH, P.; VALIN, H.; VERMEULEN, S.J.; VERVOORT, J.; VAN WIJK, M.; VAN ZANTEN, H.H.E.; WEST, P.C.; WOOD, S.A.; ROCKSTRÖM, J. Articulating the effect of food systems innovation on the Sustainable Development Goals. **The Lancet Planetary Health**, v.5, p.e50–e62, 2021. DOI: [https://doi.org/10.1016/S2542-5196\(20\)30277-1](https://doi.org/10.1016/S2542-5196(20)30277-1).
- HILOIDHARI, M.; SHARNO, M.A.; BARUAH, D.C.; BEZBARUAH, A.N. Green and sustainable biomass supply chain for environmental, social and economic benefits. **Biomass and Bioenergy**, v.175, art.106893, 2023. DOI: <https://doi.org/10.1016/j.biombioe.2023.106893>.
- HOFSETZ, K.; SILVA, M. Brazilian sugarcane bagasse: energy and non-energy consumption. **Biomass and Bioenergy**, v.46, p.564–573, 2012. DOI: <https://doi.org/10.1016/j.biombioe.2012.06.038>.
- HUANG, H.; LONG, S.; SINGH, V. Techno-economic analysis of biodiesel and ethanol co-production from lipid-producing sugarcane. **Biofuels, Bioproducts and Biorefining**, v.10, p.299–315, 2016. DOI: <https://doi.org/10.1002/bbb.1640>.
- IATA. International Air Transport Association. **Developing Sustainable Aviation Fuel (SAF)**. 2024a. Available at: <https://www.iata.org/en/programs/sustainability/sustainable-aviation-fuels/>. Accessed on: Sept. 5 2025.
- IATA. International Air Transport Association. **Net zero 2050**: sustainable aviation fuels (SAF): fact sheet. [Montreal], 2025a. Available at: <https://www.iata.org/en/iata-repository/pressroom/fact-sheets/fact-sheet-sustainable-aviation-fuels/>. Accessed on: Feb. 13 2026.
- IATA. International Air Transport Association. **Net Zero by 2050 – IATA's approach / SAF ramp-up materials**. 2024b. Available at: <https://www.iata.org/>. Accessed on: Sept. 5 2025.
- IATA. International Air Transport Association. **SAF production to triple in 2024 but more opportunities for diversification needed**. 2 June 2024c. Available at: <https://www.iata.org/en/pressroom/2024-releases/2024-06-02-03/>. Accessed on: Sept. 5 2025.
- IEA. International Energy Agency. **Aviation**. Paris, 2025a. Available at: <https://www.iea.org/energy-system/transport/aviation>. Accessed on: Feb. 13 2026.
- IEA. International Energy Agency. **Bioenergy**. Paris, 2022a. Available at: <https://www.iea.org/energy-system/renewables/bioenergy>. Accessed on: Feb. 13 2026.
- IEA. International Energy Agency. **Data and statistics**. Paris, 2025b. Available at: <https://www.iea.org/data-and-statistics>. Accessed on: Sept. 5 2025.
- IEA. International Energy Agency. **Is the biofuel industry approaching a feedstock crunch?** Paris, 2022b. Available at: <https://www.iea.org/reports/is-the-biofuel-industry-approaching-a-feedstock-crunch>. Accessed on: Sept. 1 2025.
- IEA. International Energy Agency. **Net Zero Roadmap**: a global pathway to keep the 1.5 °C goal in reach. Paris, 2023. Available at: <https://www.iea.org/reports/net-zero-roadmap-a-global-pathway-to-keep-the-15-c-goal-in-reach>. Accessed on: Feb. 13 2026.
- IEA. International Energy Agency. **Renewables 2023**: analysis and forecast to 2028. Paris, 2024. Available at: [https://iea.blob.core.windows.net/assets/96d66a8b-d502-476b-ba94-54ffda84cf72/Renewables\\_2023.pdf](https://iea.blob.core.windows.net/assets/96d66a8b-d502-476b-ba94-54ffda84cf72/Renewables_2023.pdf). Accessed on: Feb. 13 2026.
- IIASA. International Institute for Applied Systems Analysis. **Transformations to achieve the Sustainable Development Goals**. Laxenburg, 2018. Available at: <https://pure.iiasa.ac.at/id/eprint/15347/>. Accessed on: Feb. 13 2026.
- IOWA STATE UNIVERSITY. Extension and Outreach. **Sunflower**. 2021. Available at: <https://www.extension.iastate.edu/alternativeag/sunflower>. Accessed on: Sept. 5 2025.
- IPCC. Intergovernmental Panel on Climate Change. **Climate Change 2021**: the physical science basis. Contribution of Working Group I to the Sixth Assessment Report. Cambridge: Cambridge University Press, 2021. Edited by V. Masson-Delmotte, P. Zhai, A. Pirani, S.L. Connors, C. Péan, S. Berger, N. Caud, Y. Chen, L. Goldfarb, M.I. Gomis, M. Huang, K. Leitzell, E. Lonnoy, J.B.R. Matthews, T.K. Maycock, T. Waterfield, O. Yelekçi, R. Yu, and B. Zhou. DOI: <https://doi.org/10.1017/9781009157896>.
- IPCC. Intergovernmental Panel on Climate Change. Summary for policymakers. In: IPCC. Intergovernmental Panel on Climate Change. **Climate change and land**: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and greenhouse gas fluxes in terrestrial ecosystems. Cambridge: Cambridge University Press, 2022. Edited by P.R. Shukla, J. Skea, E. Calvo Buendia,

- V. Masson-Delmotte, H.- O. Pörtner, D.C. Roberts, P. Zhai, R. Slade, S. Connors, R. van Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, and J. Malley. DOI: <https://doi.org/10.1017/9781009157988.001>.
- IRIARTE, L.; FRITSCH, U.R.; VAN DAM, J. **Sustainability governance of bioenergy and the broader bioeconomy**. Technical paper prepared for IEA Bioenergy Task 45 and the Global Bioenergy Partnership (GBEP) Task Force on Sustainability. 2021. Available at: <<https://task45.ieabioenergy.com/wp-content/uploads/sites/13/2021/10/IINAS-2021-Sustainability-governance-of-bioenergy-and-bioeconomy-final.pdf>>. Accessed on: Sept. 5 2025.
- JESWANI, H.K.; CHILVERS, A.; AZAPAGIC, A. Environmental sustainability of biofuels: a review. **Proceedings of the Royal Society A**, v.476, art.20200351, 2020. DOI: <https://doi.org/10.1098/rspa.2020.0351>.
- JHAM, G.N.; MOSER, B.R.; SHAH, S.N.; HOLSER, R.A.; DHINGRA, O.D.; VAUGHN, S.F.; BERHOW, M.A.; WINKLER MOSER, J.K.; ISBELL, T.A.; HOLLOWAY, R.K.; WALTER, E.L.; NATALINO, R.; ANDERSON, J.C.; STELLY, D.M. Wild Brazilian mustard (*Brassica juncea* L.) seed oil methyl esters as biodiesel fuel. **Journal of the American Oil Chemists' Society**, v.86, p.917-926, 2009. DOI: <https://doi.org/10.1007/s11746-009-1431-2>.
- JOCIC, S.; MILADINOVIC, D.; KAYA, Y.; FORCE, E.; DUNFORD, N.T.; SALAS, J.J. Breeding and genetics of sunflower. In: MARTÍNEZ-FORCE, E.; DUNFORD, N.T.; SALAS, J.J. (Ed.). **Sunflower: chemistry, production, processing, and utilization**. Urbana: AOCS Press, 2015. p.1-26.
- JUNGERS, J.M.; MITCHELL, P.D.; GAYDON, D.S.; HAUGEN-KOGAN, A.; KOHLER, C.; MCPHERSON, M.; MCKENZIE, D.; SINCLAIR, T.R.; WELTZ, S.E. **Perennial grain resiliency: risk and opportunity in climate-smart cropping systems**. Minnesota: University of Minnesota Sustainable Cropping Systems Lab., 2023.
- KHAN, I.U. Biomass & bioenergy; case studies bioresources, chemical and biological processes, biomass products for sustainable, renewable energy and materials. **Chemical Engineering and Process Techniques**, v.8, art.1080, 2023.
- KOH, L.P.; GHAZOUL, J. Biofuels, biodiversity, and people: understanding the conflicts and finding opportunities. **Biological Conservation**, v.141, p.2450-2460, 2008. DOI: <https://doi.org/10.1016/j.biocon.2008.08.005>.
- KUSUMA, R.T.; HIREMATH, R.B.; RAJESH, P.; KUMAR, B. Sustainable transition towards biomass-based cement industry: a review. **Renewable and Sustainable Energy Reviews**, v.162, art.112427, 2022. DOI: <https://doi.org/10.1016/j.rser.2022.112503>.
- LESLIE-BOLE, Y.; DENVIR, A.; LASHOF, D.; SCAFIDI, A.; RIBEIRO, C.M. **Biomass and land use in a decarbonizing US economy**. Washington: WRI, 2025. (Working Paper). DOI: <https://doi.org/10.46830/wriwp.24.00033>.
- LIMA, M.G.B.; PALME, U. The bioeconomy-biodiversity nexus: enhancing or undermining nature's contributions to people? **Conservation**, v.2, p.7-25, 2022. DOI: <https://doi.org/10.3390/conservation2010002>.
- LIRA, J.P.E.; BARELLI, M.A.A.; SILVA, V.P. da; FELIPIN-AZEVEDO, R.; SANTOS, D.T. dos; GALBIATI, C.; POLETINE, J.P. Safflower genetic diversity based on agronomic characteristics in Mato Grosso state, Brazil, for a crop improvement program. **Genetics and Molecular Research**, v.20, p.1-12, 2021. DOI: <http://dx.doi.org/10.4238/gmr18698>.
- LODATO, C.; HAMELIN, L.; TONINI, D. Towards sustainable methane supply from local bioresources: anaerobic digestion, gasification, and gas upgrading. **Applied Energy**, v.323, art.119641, 2022. DOI: <https://doi.org/10.1016/j.apenergy.2022.119568>.
- LOPES, M.A. **Descarbonização e circularidade: respostas dos sistemas alimentar e agroindustrial aos Objetivos de Desenvolvimento Sustentável**. Brasília: Embrapa Agroenergia, 2022. (Embrapa Agroenergia. Documentos, 47). Available at: <<http://www.infoteca.cnptia.embrapa.br/infoteca/handle/doc/1149409>>. Accessed on: Sept. 1 2025.
- LORENZI, H.; SOUZA, H.M.; TORRES, M.A.V.; BACHER, L.B. **Palmeiras no Brasil: nativas e exóticas**. Nova Odessa: Instituto Plantarum, 1996.
- MABEE, W.E. **Literature review on social and economic sustainability of bioenergy systems**. Paris: IEA Bioenergy, 2024. Available at: <[https://www.ieabioenergy.com/wp-content/uploads/2025/01/Mabee-2024-Literature-review-on-social-and-economic-sustainability\\_final.pdf](https://www.ieabioenergy.com/wp-content/uploads/2025/01/Mabee-2024-Literature-review-on-social-and-economic-sustainability_final.pdf)>. Accessed on: Sept. 1 2025.
- MADALENA, L.C. de S.; CHANG, P.; OLIVEIRA, A.F. de; ROSSETO, R.E.; ROEHRS, S.A.; SANTOS, R.F.; FRIGO, E.P. Uma revisão sobre a mostarda (*Brassica* spp.) e seu possível uso com alternativa para produção de biodiesel. **Acta Iguazu**, v.6, p.23-34, 2017.
- MALINS, C. Multiple and cover cropping in Brazil: status and opportunities for biofuel production. **Cerology**, 2022. Available at: <<https://theicct.org/wp-content/uploads/2022/01/cover-cropping-biofuel-brazil-jan22.pdf>>. Accessed on: Sept. 1 2025.
- MANNUCCI, P.M.; JOLLIET, O.; MEIJAARD, E.; SLAVIN, J.; RASETTI, M.; ALETA, A.; MORENO, Y.; AGOSTONI, C. Sustainable nutrition and the case of vegetable oils to match present and future dietary needs. **Frontiers in Public Health**, v.11, art.1106083, 2023. DOI: <https://doi.org/10.3389/fpubh.2023.1106083>.
- MANTULET, G.; BIDAUD, A.; MIMA, S. The role of biomass gasification and methanisation in the decarbonisation strategies. **Energy**, v.193, art.116737, 2020. DOI: <https://doi.org/10.1016/j.energy.2019.116737>.
- MARTHA JÚNIOR, G.B.; LOPES, M.A. Charting new sustainable agricultural innovation pathways in Brazil. **Scientia Agricola**, v.80, e20230067, 2023. DOI: <https://doi.org/10.1590/1678-992X-2023-0067>.
- MARTINELLI, F.S.; BIBER-FREUDENBERGER, L.; STEIN, G.; BÖRNER, J. Will Brazil's push for low-carbon biofuels contribute to achieving the SDGs? A systematic expert-based assessment. **Cleaner Environmental Systems**, v.5, art.100075, 2022. DOI: <https://doi.org/10.1016/j.cesys.2022.100075>.

- MARTÍNEZ-FORCE, E.; DUNFORD, N.T.; SALAS, J.J. (Ed.). **Sunflower: chemistry, production, processing, and utilization**. Urbana: AOCS Press, 2015.
- MASI, M.; ODDO, E.; RULLI, M.C.; SEABRA, J.E. **Roadmap to 2050: the land-water-energy nexus of biofuels**. New York: Sustainable Development Solutions Network; Fondazione Eni Enrico Mattei, 2021. 164p. Available at: <<https://roadmap2050.report/biofuels/biofuels-technologies/>>. Accessed on: Sept. 1 2025.
- MATTOS, V.Z.A. de; TAVARES, B.G.; BARRETO, R. da C.; GUIMARÃES, G.P.; FREITAS, M.A.V. de. Low-carbon agricultural technologies improve forage and feed production in the Caatinga biome, Brazil: characteristics, comparison, effects of climate change, resilience, local development, and food security. **Scientia Agropecuaria**, v.15, n.4, p.629-639, 2024. DOI: <https://doi.org/10.17268/sci.agropecu.2024.046>.
- MAZUMDAR, P.; SINGH, P.; BABU, S.; SIVA, R.; HARIKRISHNA, J.A. An update on biological advancement of *Jatropha curcas* L.: new insight and challenges. **Renewable and Sustainable Energy Reviews**, v.91, p.903-917, 2018. DOI: <https://doi.org/10.1016/j.rser.2018.04.082>.
- MILLINGER, M.; HEDENUS, F.; ZEYEN, E.; NEUMANN, F.; REICHENBERG, L.; BERNDEN, G. Diversity of biomass usage pathways to achieve emissions targets in the European energy system. **Nature Energy**, v.10, p.226-242, 2025. DOI: <https://doi.org/10.1038/s41560-024-01693-6>.
- MNAGRO. **Canola expansion in Brazil**. 2025. Available at: <<https://www.mnagro.com.br/news/30/1/canola-expansion-in-brazil>>. Accessed on: July 1 2025.
- MOTOIKE, S.Y.; HILGER, T. (Org.). **Macaúba: do plantio à colheita**. São Paulo: Oficina de Textos, 2024.
- MUBOFU, E.B. Castor oil as a potential renewable resource for the production of functional materials. **Sustainable Chemical Processes**, v.4, art.11, 2016. DOI: <https://doi.org/10.1186/s40508-016-0055-8>.
- MÜLLER-LANGER, F. (Ed.). **Assessment of successes and lessons learned for biofuels deployment**: report work package 2: meta-analysis of existing studies. [Paris]: IEA Bioenergy, 2023. (IEA Bioenergy TCP – Intertask Project of Task 39, Task 40 and Task 45). Available at: <<https://www.ieabioenergy.com/wp-content/uploads/2023/08/IEA-Bioenergy-ITP-Assessment-of-successes-and-lessons-learned-for-biofuels-deployment.pdf>>. Accessed on: Feb. 13 2026.
- MURPHY, D.J.; GOGGIN, K.; PATERSON, R.R.M. Oil palm in the 2020s and beyond: challenges and solutions. **CABI Agriculture and Bioscience**, v.2, art.39, 2021. DOI: <https://doi.org/10.1186/s43170-021-00058-3>.
- MUSCALU, A.; TUDORA, C.; SORICĂ, C.; ANGHELACHE, D.; IONESCU, A.; CONSTANTINESCU, M.; FENDRIHAN, S. **Extraction methods and uses of safflower (*Carthamus tinctorius* L., Asteraceae family) oil**. 2023. Available at: <<https://www.cabidigitallibrary.org/doi/pdf/10.5555/20240010956>>. Accessed on: Sept. 5 2025.
- NOPA. National Oilseed Processors Association. **Feedstocks from U.S. Oilseeds Offer Significant Decarbonization Opportunities and Should Not Be Left Behind**. 2024. Available at: <<https://www.nopa.org/nopa-to-usda-feedstocks-from-u-s-oilseeds-offer-significant-decarbonization-opportunities-and-should-not-be-left-behind/>>. Accessed on: July 1 2025.
- OECD. Organisation for Economic Co-operation and Development. **Innovation, agricultural productivity and sustainability in Brazil**. Paris, 2015. (OECD Food and Agricultural Reviews). DOI: <https://doi.org/10.1787/9789264237056-en>.
- OECD-FAO Agricultural Outlook 2025–2034. Paris: OECD; Rome: FAO, 2025. Available at: <[https://www.oecd.org/content/dam/oecd/en/publications/reports/2025/07/oecd-fao-agricultural-outlook-2025-2034\\_3eb15914/601276cd-en.pdf](https://www.oecd.org/content/dam/oecd/en/publications/reports/2025/07/oecd-fao-agricultural-outlook-2025-2034_3eb15914/601276cd-en.pdf)>. Accessed on: Sept. 5 2025.
- OGWU, M.C.; KOSOE, E.A.; OSAWARU, M.E. *Raw materials for sustainable bioeconomy development in the Global South*. In: OGWU, M.C.; IZAH, S.C.; MERRITT, H.; WISE, R.D.; JINTRAWET, A. (Ed.). **Sustainable Bioeconomy Development in the Global South**. Singapore: Springer Nature, 2025. p.37-63. DOI: [https://doi.org/10.1007/978-981-96-0640-5\\_2](https://doi.org/10.1007/978-981-96-0640-5_2).
- OLIVEIRA, D.M.S.; SANTOS, R.S.; CHIZZOTTI, F.H.M.; BRETAS, I.L.; FRANCO, A.L.C.; LIMA, R.P.; FREITAS, D.A.F.; CHERUBIN, M.R.; CERRI, C.E.P. Crop, livestock, and forestry integration to reconcile soil health, food production, and climate change mitigation in the Brazilian Cerrado: a review. **Geoderma Regional**, v.37, art.e00796, 2024a. DOI: <https://doi.org/10.1016/j.geodrs.2024.e00796>.
- OLIVEIRA, M.B. de; POGORZELSKI, E. da S.; PFEIFEMBERG, R.; KNIES, A.E.; OLIVEIRA, Z.B. de; SANTOS, M.S.N. dos; ZABOT, G.L.; TRES, M.V. Analysis of the cultivation of canola hybrids at different sowing dates. **Acta Scientiarum. Agronomy**, v.46, e67694, 2024b. DOI: <https://doi.org/10.4025/actasciagron.v46i1.67694>.
- OSMAN, A.I.; MEHTA, N.; ELGARAHY, A.M.; AL-HINAI, A.; AL-MUHTASEB, A.A.H.; ROONEY, D.W. Conversion of biomass to biofuels and life cycle assessment: a review. **Environmental Chemistry Letters**, v.19, p.4075-4118, 2021. DOI: <https://doi.org/10.1007/s10311-021-01273-0>.
- OSMAN, W.N.A.W.; ROSLI, M.H.; MAZLI, W.N.A.; SAMSURI, S. Comparative review of biodiesel production and purification. **Carbon Capture Science & Technology**, v.13, art.100264, 2024. DOI: <https://doi.org/10.1016/j.ccsst.2024.100264>.
- PARADA, M.P.; VANDERPUTTEN, W.H.; VANDERWIELEN, L.A.M.; OSSEWEIJER, P.; VAN LOOSDRECHT, M.; KAMALI, F.P.; POSADA, J.A. Sustainability tensions and opportunities for aviation biofuel production in Brazil. In: YOUSUF, A.; GONZALEZ-FERNANDEZ, C. (Ed.). **Sustainable Alternatives for Aviation Fuels**. Amsterdam: Elsevier, 2022. p.237-262. DOI: <https://doi.org/10.1016/B978-0-323-85715-4.00007-0>.
- PEREIRA, L.G.; CAVALETT, O.; BONOMI, A. **Comparison of biofuel life cycle assessment tools**. Paris, Feb. 2017a. (IEA. Task 39 – Commercializing Liquid Biofuels). Available at: <<https://task39.ieabioenergy.com/wp-content/uploads/sites/37/2019/04/Task-39-GHS-models-Final-Report-Phase-1.pdf>>. Accessed on: Feb. 13 2026.

- PEREIRA, L.G.; MACLEAN, H.L.; SAVILLE, B.A. Financial analyses of potential biojet fuel production technologies. **Biofuels, Bioproducts and Biorefining**, v.11, p.665-681, 2017b. DOI: <https://doi.org/10.1002/bbb.1775>.
- PEREIRA, M. de A.; COSTA, F.P.; ALMEIDA, R.G. de. Is the “F Word” an option for Brazilian farmers? The place of forestry in future integrated farming systems. **International Journal of Agricultural Management**, v.6, p.134-140, 2018. DOI: <https://doi.org/10.5836/ijam/2017-06-134>.
- PIMENTEL, G.V.; MARTINS, I.A.; SANTIAGO, A.C.; COSTA, N.; SILVA, R.N.; MOREIRA, S.G. Agronomic performance of canola hybrids in Campos das Vertentes-MG. **Revista Ceres**, v.71, e71021, 2024. DOI: <https://doi.org/10.1590/0034-737X2024710021>.
- PINHEIRO, P. Commodities 2025: Brazil's biodiesel production and feedstocks outlook. **S&P Global**, Jan. 2 2025. Available at: <https://www.spglobal.com/energy/en/news-research/latest-news/agriculture/010225-commodities-2025-brazil-boosts-biodiesel-production-eyes-market-growth>. Accessed on: Feb. 13 2026.
- PINTO, L.C.L.; RODRIGUES, I.P.S.; DRUMOND, M.A. Population structure and fruit productivity analyses in support of the use of *Caryocar brasiliense*. **Floresta e Ambiente**, v.26, e20170995, 2019. DOI: <https://doi.org/10.1590/2179-8087.099517>.
- PROFOREST. **Monitoring Protocol defines environmental commitments for livestock farming in the Cerrado**. 2024. Available at: <https://www.proforest.net/news-and-events/news/monitoring-protocol-defines-environmental-commitments-for-livestock-farming-in-the-cerrado/>. Accessed on: July 1 2025.
- QUEIROZ, I.R.; COSTA, A.S.V. da; ALMEIDA, I.C.; BARROS, G.F.; ALVES, W.M.; SOUZA, M.C. de; FRANCO, M.L.; POMPERMAYER, R. de S.; SOUZA, A.O.; FERREIRA, A.C. Biofuels in Brazil: potentialities of *Crambe abyssinica* culture for biodiesel production. **Research, Society and Development**, v.10, e11510514618, 2021. DOI: <https://doi.org/10.33448/rsd-v10i5.14618>.
- RATHORE, D.; PANT, D.; SINGH, A. A comparison of life cycle assessment studies of different biofuels. In: SINGH, A.; PANT, D.; OLSEN, S.I. (Ed.). **Life cycle assessment of renewable energy sources**. London: Springer, 2013. p.269-289. DOI: [https://doi.org/10.1007/978-1-4471-5364-1\\_12](https://doi.org/10.1007/978-1-4471-5364-1_12).
- RIBEIRO, C.H.; CUNHA, M.P. da. The economic and environmental impacts of Brazilian National Biofuel Policy. **Biofuels, Bioproducts and Biorefining**, v.16, p.413-434, 2022. DOI: <https://doi.org/10.1002/bbb.2326>.
- RIBEIRO, P.R.; FERNANDEZ, L.G.; CASTRO, R.D. de; LIGTERINK, W.; HILHORST, H.W.M. Physiological and biochemical responses of *Ricinus communis* seedlings to different temperatures: a metabolomics approach. **BMC Plant Biology**, v.14, art.223, 2014.
- ROCKSTRÖM, J.; STEFFEN, W.; NOONE, K.; PERSSON, Å.; CHAPIN III, F.S.; LAMBIN, E.F.; LENTON, T.M.; SCHEFFER, M.; FOLKE, C.; SCHELLNHUBER, H.J.; NYKVIST, B.; DE WIT, C.A.; HUGHES, T.; VAN DER LEEUW, S.; RODHE, H.; SÖRLIN, S.; SNYDER, P.K.; COSTANZA, R.; SVEDIN, U.; FALKENMARK, M.; KARLBERG, L.; CORELL, R.W.; FABRY, V.J.; HANSEN, J.; WALKER, B.; LIVERMAN, D.; RICHARDSON, K.; CRUTZEN, P.; FOLEY, J.A. A safe operating space for humanity. **Nature**, v.461, p.472-475, 2009. Available at: <https://www.nature.com/articles/461472a>. Accessed on: Feb. 13 2026.
- RODRIGUES, R. de A.R.; FERREIRA, I.G.M.; SILVEIRA, J.G. da; SILVA, J.J.N. da; SANTOS, F.M.; CONCEIÇÃO, M.C.G. da. Crop-livestock-forest integration systems as a sustainable production strategy in Brazil. In: SØNDERGAARD, N.; SÁ, C.D. de; PLATIAU, A.F.B. (Ed.). **Sustainability challenges of Brazilian agriculture**: governance, inclusion, and innovation. Cham: Springer, 2023. p.165-182. DOI: [https://doi.org/10.1007/978-3-031-29853-0\\_9](https://doi.org/10.1007/978-3-031-29853-0_9).
- ROSILLO-CALLE, F. The role of biomass energy in rural development. In: ENCONTRO DE ENERGIA NO MEIO RURAL, 3., 2000, Campinas. **Proceedings**. Campinas: Unicamp, 2000. p.12-15. Available at: [http://www.proceedings.scielo.br/scielo.php?script=sci\\_arttext&pid=MSC000000022000000200011&lng=en&nrm=iso](http://www.proceedings.scielo.br/scielo.php?script=sci_arttext&pid=MSC000000022000000200011&lng=en&nrm=iso). Accessed on: Sept. 1 2025.
- ROSSI, C.; SHEN, L.; JUNGINGER, M.; WICKE, B. Sustainability certification of bio-based products: systematic literature review of socio-economic impacts along the supply chain. **Journal of Cleaner Production**, v.435, art.143079, 2024. DOI: <https://doi.org/10.1016/j.jclepro.2024.143079>.
- RSB. Available at: <https://rsb.org/>. Accessed on: Aug. 7 2025.
- SÁ, C.D. de; SØNDERGAARD, N.; BARIONI, L.G.; CAMARGO, R.C. The Brazilian way of farming: potential and challenges to agricultural decarbonization. In: SØNDERGAARD, N.; SÁ, C.D. de; PLATIAU, A.F.B. (Ed.). **Sustainability challenges of Brazilian agriculture**: governance, inclusion, and innovation. Cham: Springer, 2023. p.145-163.
- SAKAI, P.; AFIONIS, S.; FAVRETTO, N.; STRINGER, L.C.; WARD, C.; SAKAI, M.; WEIRICH NETO, P.H.; ROCHA, C.H.; GOMES, J.A.; SOUZA, N.M. de; AFZAL, N. Understanding the implications of alternative bioenergy crops to support smallholder farmers in Brazil. **Sustainability**, v.12, art.2146, 2020. DOI: <https://doi.org/10.3390/su12052146>.
- SALINA, F.H.; ALMEIDA, I.A. de; BITTENCOURT, F.R. RenovaBio opportunities and biofuels outlook in Brazil. In: SAYIGH, A. (Ed.). **Renewable Energy and Sustainable Buildings: Selected Papers from the World Renewable Energy Congress WREC 2018**. Cham: Springer International Publishing, 2019. p.391-399. DOI: [https://doi.org/10.1007/978-3-030-18488-9\\_30](https://doi.org/10.1007/978-3-030-18488-9_30).
- SCARIOT, A.O.; LLERAS, E.; HAY, J.D. Reproductive biology of the palm *Acrocomia aculeata* in Central Brazil. **Biotropica**, v.27, p.384-392, 1995. DOI: <https://doi.org/10.2307/2388683>.
- SCHEITERLE, L.; BIRNER, R. Considerations on the role of institutions and networks in the bioeconomy: three case studies from Ghana and Brazil. **Bonn**: University of Bonn, 2020. (*ZEF Working Paper Series*, n.196). Available at: <https://hdl.handle.net/10419/246465>. Accessed on: Sept. 1 2025.

- SEEPAL, R.; KUMAR, S.; IBOYI, J.E.; BASHYAL, M.; STANSLY, T.L.; BENNETT, R.; BOOTE, K.J.; MULVANEY, M.J.; SMALL, I.M.; GEORGE, S.; WRIGHT, D.L. *Brassica carinata*: biology and agronomy as a biofuel crop. **GCB Bioenergy**, v.13, p.582-599, 2021. DOI: <https://doi.org/10.1111/gcbb.12804>.
- SIKDAR, S.K. Sustainable development and sustainability metrics. **AIChE Journal**, v.49, p.1928-1932, 2003. DOI: <https://doi.org/10.1002/aic.690490802>.
- SILVA, J.A.N. da; SOUZA, C.M.A. de; ARCOVERDE, S.N.S.; KURIHARA, C.H.; SILVA, C.J. da. Soil chemical properties and production of physic nut intercropped with forage plants and grain crops. **Journal of Agricultural Science**, v.16, p.57-71, 2024. DOI: <https://doi.org/10.5539/jas.v16n7p57>.
- SILVA, J.N. da; FARIAS, P.R.S.; SILVA, A.G. da; TINÔCO, R.S.; SANTOS, A.V.F. dos; CARDOSO, L.A.S.; MONFORT, L.H.F.; OLIVEIRA, F.J. de. Spatial distribution of fatal yellowing cases in oil palm plantations in the Eastern Amazon. **Discover Plants**, v.2, art.158, 2025. DOI: <https://doi.org/10.1007/s44372-025-00251-y>.
- SILVA, S.P. da; COSTA, A.S.V. da; SANTOS, S.L.B. dos; LAIA, M.L. de. A importância da biomassa na matriz energética brasileira. **Pensar Acadêmico**, v.19, p.557-583, 2021.
- SINHA, P. New report estimates district-wise crop residue potential for bioenergy production. **Mongabay India**, July 17 2023. Available at: <https://india.mongabay.com/2023/07/new-report-unveils-district-wise-crop-residue-estimates-for-bioenergy-production/>. Accessed on: Feb. 13 2026.
- SORITA, G.D.; FAVARO, S.P.; GAMBETTA, R.; AMBROSI, A.; DI LUCCIO, M. *Macauba (Acrocomia spp.) fruits*: a comprehensive review of nutritional and phytochemical profiles, health benefits, and sustainable oil production. **Comprehensive Reviews in Food Science and Food Safety**, v.24, e70097, 2025. DOI: <https://doi.org/10.1111/1541-4337.70097>.
- SOUZA, N.R.D. de; PETRIELLI, G.P.; HERNANDES, T.A.D.; LEDUC, S.; Di FULVIO, F.; HENZLER, D. de S.; CHAGAS, M.F.; JUNQUEIRA, T.L.; CAVALETT, O. Bioenergy-livestock integration in Brazil: unraveling potentials for energy production and climate change mitigation. **Journal of Cleaner Production**, v.422, art.138586, 2023. DOI: <https://doi.org/10.1016/j.jclepro.2023.138586>.
- SOUZA, P.T. de; SILVA, E. de B.; GRAZZIOTTI, P.H.; FERNANDES, L.A. NPK fertilization on initial growth of physic nut seedlings in Quartzarenic Neossol. **Revista Brasileira de Ciência do Solo**, v.35, p.559-566, 2011. DOI: <https://doi.org/10.1590/S0100-06832011000200025>.
- STYLES, D.; YESUFU, J.; BOWMAN, M.; WILLIAMS, A.P.; DUFFY, C.; LUYCKX, K. Climate mitigation efficacy of anaerobic digestion in a decarbonising economy. **Journal of Cleaner Production**, v.338, art.130441, 2022. DOI: <https://doi.org/10.1016/j.jclepro.2022.130441>.
- SUBBARAO, P.M.; D'SILVA, T.C.; ADLAK, K.; KUMAR, S.; CHANDRA, R.; VIJAY, V.K. Anaerobic digestion as a sustainable technology for efficiently utilizing biomass in the context of carbon neutrality and circular economy. **Environmental Research**, v.234, art.116286, 2023. DOI: <https://doi.org/10.1016/j.envres.2023.116286>.
- TELHADO, S.F.P.; CAPDEVILLE, G. de (Ed.). **Tecnologias poupa-terra 2021**. Brasília: Embrapa, 2021.
- THODAY-KENNEDY, E.; BANERJEE, B.; PANOZZO, J.; MAHARJAN, P.; HUDSON, D.; SPANGENBERG, G.; HAYDEN, M.; KANT, S. Dissecting physiological and agronomic diversity in safflower populations using proximal phenotyping. **Agriculture**, v.13, art.620, 2023. DOI: <https://doi.org/10.3390/agriculture13030620>.
- TOMM, G.O.; SMIDERLE, O.; RAPOSO, R.; DALMAGO, G.; SOARES, A.; FIGER, E.; LUFT, A.; BATISTELLA, E.; SCARANTI, V.; BENIN, F. Can canola be produced in tropical areas? In: INTERNATIONAL RAPESEED CONGRESS, 13., 2011, Prague. **Proceedings**. Prague: International Consultative Research Group on Rapeseed, 2011. p.2-5. Available at: <http://www.alice.cnptia.embrapa.br/alice/handle/doc/895037>. Accessed on: Sept. 3 2025.
- UNITED NATIONS. **Transforming our world: the 2030 Agenda** for sustainable development. New York, 2015.
- UNITED STATES. Department of Energy. **2023 Billion Ton Report: an assessment of U.S. renewable carbon resources**. Oak Ridge: Oak Ridge National Laboratory, 2025. (ORNL/SPR-2024/3103). Available at: <https://www1.eere.energy.gov/apps/bioenergy/bto-2023-billion-ton-report/assets/pdfs/beto-2023-billion-ton-report-1.pdf>. Accessed on: July 1 2025.
- USDA-ERS. United States Department of Agriculture. Economic Research Service. **Demand for domestic vegetable oils boosted by rising biofuels production**. 2024. (Chart of Note, Jan. 30 2024). Available at: <https://ers.usda.gov/data-products/charts-of-note/chart-detail?chartId=108363>. Accessed on: Feb. 13 2026.
- USDA-FAS. United States Department of Agriculture. Foreign Agricultural Service. **Biofuels annual: Brazil**. Report Number BR2024-0022. Aug. 31 2024. Available at: [https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Biofuels+Annual\\_Brasilia\\_Brazil\\_BR2024-0022.pdf](https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Biofuels+Annual_Brasilia_Brazil_BR2024-0022.pdf). Accessed on: Sept. 5 2025.
- USDA-FAS. United States Department of Agriculture. Foreign Agricultural Service. **Brazil's sunflower sector – opportunities and challenges in a growing market**. Report Number BR2025-0029. Aug. 28 2025a. Available at: [https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Brazil%27s+Sunflower+Sector+Opportunities+and+Challenge+in+a+Growing+Market\\_Brasilia\\_Brazil\\_BR2025-0029.pdf](https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Brazil%27s+Sunflower+Sector+Opportunities+and+Challenge+in+a+Growing+Market_Brasilia_Brazil_BR2025-0029.pdf). Accessed on: Sept. 5 2025.
- USDA-FAS. United States Department of Agriculture. Foreign Agricultural Service. **Commodity intelligence report: Brazil palm oil: potential and pitfalls**. Oct. 13 2022a. Available at: <https://ipad.fas.usda.gov/highlights/2022/10/Brazil/index.pdf>. Accessed on: Feb. 13 2026.
- USDA-FAS. United States Department of Agriculture. Foreign Agricultural Service. **Oilseed and products annual: Brazil**. Report Number BR2022-0026. Apr. 29 2022b. Available at: <https://apps.fas.usda.gov/newgainapi/api/Report/DownloadReportByFileName?fileName=Oilseeds+and+>

- Products+Annual\_Brasilia\_Brazil\_BR2022-0026>. Accessed on: Sept. 1 2025.
- USDA-FAS. United States Department of Agriculture. Foreign Agricultural Service. **Oilseeds: world markets and trade**. Sept. 2025b. Available at: <<https://apps.fas.usda.gov/psdonline/circulars/oilseeds.pdf>>. Accessed on: Sept. 1 2025.
- VALIN, H.; PETERS, D.; VAN DEN BERG, M.; FRANK, S.; HAVLIK, P.; FORSELL, N.; DIFULVIO, F. **The land use change impact of biofuels consumed in the EU: quantification of area and greenhouse gas impacts**. 2015. Available at: <[https://energy.ec.europa.eu/publications/land-use-change-impact-biofuels-consumed-eu\\_en](https://energy.ec.europa.eu/publications/land-use-change-impact-biofuels-consumed-eu_en)>. Accessed on: Sept. 1 2025.
- VAN LOO, E.N.; JONGSCHAAP, R.E.E. **JATROPT (*Jatropha curcas*: applied and technological research on plant traits)**: final report EU FP7 Grant 245236 (2010–2013). Wageningen: Plant Research International, 2014. Available at: <<https://research.wur.nl/en/publications/jatropt-jatropha-curcas-applied-and-technological-research-on-pla>>. Accessed on: Sept. 1 2025.
- VANDENBERGHE, L.P.S.; VALLADARES-DIESTRA, K.K.; BITTENCOURT, G.A.; TORRES, L.Z.; VIEIRA, S.; KARP, S.G.; SOCCOL, C.R. Beyond sugar and ethanol: the future of sugarcane biorefineries in Brazil. **Renewable and Sustainable Energy Reviews**, v.167, art.112721, 2022. DOI: <https://doi.org/10.1016/j.rser.2022.112721>.
- VARGAS-CARPINTERO, R.; HILGER, T.; MÖSSINGER, J.; SOUZA, R.F.; BARROSO ARMAS, J.C.; TIEDE, K.; LEWANDOWSKI, I. *Acrocomia spp.*: neglected crop, ballyhooed multipurpose palm or fit for the bioeconomy? A review. **Agronomy for Sustainable Development**, v.41, art.75, 2021. DOI: <https://doi.org/10.1007/s13593-021-00729-5>.
- VIEIRA, M.G.; BATISTA, L.R.; MUNIZ, A.S.; LICHTSTON, J.E.; OLIVEIRA, A.R.; CÉSAR-OLIVEIRA, M.A.; ANTONIOSI FILHO, N.R. Synthesis of antioxidant additive from safflower seed oil. **Journal of the Brazilian Chemical Society**, v.32, p.40-46, 2021. DOI: <https://doi.org/10.21577/0103-5053.20200151>.
- VON COSSEL, M.; SCORDIA, D.; ALTIERI, M.; GRESTA, M. Spotlight on agroecological cropping practices to improve the resilience of farming systems: a qualitative review of meta-analytic studies. **Frontiers in Agronomy**, v.7, art.1495846, 2025. DOI: <https://doi.org/10.3389/fagro.2025.1495846>.
- WATSON, M.J.; MACHADO, P.G.; SILVA, A.V. da; SALTAR, Y.; RIBEIRO, C.O.; NASCIMENTO, C.A.O.; DOWLING, A.W. Sustainable aviation fuel technologies, costs, emissions, policies, and markets: a critical review. **Journal of Cleaner Production**, v.449, art.141472, 2024. DOI: <https://doi.org/10.1016/j.jclepro.2024.141472>.
- WITT, T.W.; FLYNN, K.C.; ZOZ, T.; LEE, T.O.; MONTEIRO, J.E.B.A. A site suitability analysis for castor (*Ricinus communis* L.) production during Brazil's second harvest incorporating disease prediction. **Heliyon**, v.9, e18981, 2023. DOI: <https://doi.org/10.1016/j.heliyon.2023.e18981>.
- WORLD BIOENERGY ASSOCIATION. **Global bioenergy statistics report 2023**. 10<sup>th</sup> ed. 2023. Available at: <<https://www.worldbioenergy.org/uploads/231219%20GBS%20Report.pdf>>. Accessed on: Sept. 1 2025.
- ZHANG, H.; ZHOU, G.; VARSHNEY, R.K.; WAN, S. Advances in plant cultivation and physiology of oilseed crops. **Frontiers in Plant Science**, v.14, art.1280392, 2023. DOI: <https://doi.org/10.3389/fpls.2023.1280392>.
- ZHOU, Q.; LE, Q.V.; YANG, H.; GU, H.; YANG, Y.; SONNE, C.; TABATABAEI, M.; LAM, S.S.; LI, C.; CHEN, X.; PENG, W. Sustainable conversion of agricultural biomass into renewable energy products: a discussion. **BioResources**, v.17, p.3489-3508, 2022. DOI: <https://doi.org/10.15376/biores.17.2.Zhou>.

### Author contributions

**Mauricio Antonio Lopes:** conceptualization (lead), writing – original draft (lead), writing – review & editing (lead); **Bruno Galveas Laviola:** conceptualization (supporting), writing – original draft (supporting), writing – review & editing (supporting); **Simone Palma Favaro:** conceptualization (supporting), writing – original draft (supporting), writing – review & editing (supporting).

**Chief editor:** Edemar Corazza

**Edited by:** Mírian Baptista

### Data availability statement

No new data were generated or analyzed in this study. All information supporting the findings is contained within the article and in the cited references.

### Declaration of use of AI technologies

During the preparation of this manuscript, the authors used ChatGPT (developed by OpenAI) exclusively for support in English language grammar and stylistic refinement. The use of this tool did not influence the scientific content, interpretation,

or conclusions. All content was reviewed and approved by the authors, who take full responsibility for the manuscript.

### Conflict of interest statement

The authors declare no conflicts of interest.

### Acknowledgments

To Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) and to Ministério da Agricultura e Pecuária of Brazil (MAPA), for their financial support to the author's research on tropical canola and macaúba.

### Disclaimer/Publisher's note:

The statements, opinions, and data contained in all texts published in Pesquisa Agropecuária Brasileira (PAB) are solely those of the individual author(s) and not of the journal's publisher, editor, and editorial team, who disclaim responsibility for any injury to people or property resulting from any referred ideas, methods, instructions, or products.

The mention of specific chemical products, machines, and commercial equipment in the texts published in this journal does not imply their recommendation by the publisher.