



8TH INTERNATIONAL

MANGO SYMPOSIUM

SUN CITY, SOUTH AFRICA

5-10 FEBRUARY 2006

Mangoes - An Industry Committed to Quality

www.mangosa2006.co.za

634.44

I61p

2006

PC-2006.00102

ABSTRACT BOOK

SIYANAMUKELA E-AFRIKA! - "Welcome to Africa"



Effects of Postharvest Handling on Lenticel Spotting of 'Tommy Atkins' Mangoes from Northeast Brazil

Joston Simão de Assis

Embrapa Semi-Árido, C.P. 23, BR 428 km 152, 56300-970, Petrolina-PE, Brasil.

E-mail: joston@cpatsa.embrapa.br

Guy Self

Cirad-flhor, UR TROPIQUAL, TA50/16, avenue Jean François Breton, 34398 Montpellier Cedex 5, France. E-mail: guy.self@cirad.fr

Vanessa Cristina Caron

Escola Superior de Agricultura 'Luiz de Queiroz' (ESALQ), Universidade de São Paulo, Av. Padre Gaspar Bertoni, 567. BL T apto 234. Jd. Pacaembu, 13033-206, Campinas-SP, Brasil.

E-mail: vccaron@hotmail.com

Keywords: *Mangifera indica* L., lenticel discolouration, chilling injury

ABSTRACT

Both red and black lenticel spotting reduce the value of Tommy Atkins mangoes exported from northeast Brazil to North America and Europe. By applying none, some or all postharvest operations to early and mid-season fruit, it was shown that postharvest handling increases lenticel spotting in a cumulative way. Both types of lenticel spotting were caused principally by washing fruit in a calcium hydroxide solution to prevent sap burn. Following this, the USDA phytosanitary hot water treatment reduced the potential for subsequent red lenticel spotting during cold storage, but increased black lenticel spotting. The result of these reciprocal effects is that fruit for the EU market, which does not receive the hot water treatment, is affected principally by red spotting, and fruit for the US market by black spotting. It is suggested that black lenticel spotting occurs principally through physical processes involving the entry of water into the lenticels and the subsequent collapse and discolouration of sub-lenticellular cells. In contrast, red lenticel spotting is a physiological process involving anthocyanin production in response to low temperature in which water entry to the lenticels plays a role. The observation that lenticel spotting decreases during the season is consistent with the work of Joyce et al. (2001), and suggests that fruit become more physiologically resistant and that the properties of the lenticels change making them more resistant to water entry and damage as the season progresses. The nature of these changes remains to be determined.

BIOGRAPHY

Guy SELF is currently working for Cirad-flhor in Montpellier on the postharvest of tropical fruits. He has worked in Honduras on the postharvest of fruits and vegetables and in Brazil on the postharvest of mangoes.