

TERRA PRETA AND TERRA MULATA: Pre-Columbian Amazon kitchen middens and agricultural fields, their sustainability and their replication ⁽¹⁾

Wim Sombroek ⁽²⁾, Dirse Kern ⁽³⁾, Tarcício Rodrigues ⁽⁴⁾, Manoel da Silva Cravo ⁽⁴⁾, Tony Jarbas Cunha ⁽⁵⁾, William Woods ⁽⁶⁾ and Bruno Glaser ⁽⁷⁾

SP
4454

Abstract

Anthropogenic Black Earths (hortic Anthrosols) occur in many places of the Amazon region, in patches that vary from 5 to over 300 ha. They are locally known as Terra Preta (do Indio) and are apparently kitchen middens of pre-Columbian Amerindian settlements that have transformed the original soils (Ferralsols, Acrisols, a.o.) into black Fimic Anthrosols, with approximately double the amount of soil organic matter (SOM), higher amounts of Phosphorus and Calcium, and presence of ceramic artefacts.

Recent field studies indicate that these soils are often accompanied by bands of soils with dark brown rather than black colour and near absence of artefacts, but still with the same high amount of SOM. They are locally called Terra Mulata, and would appear to be the lasting result of intensive agricultural use by erstwhile Indian communities. The SOM of both the Terra Preta and the Terra Mulata is very stable and at the same time more re-active than the SOM of the original soils under primary forest cover.

An interdisciplinary and inter-institutional project is in the making, to establish the reasons for the sustainability of the qualities of both types of anthropogenically enriched soils; to obtain an inkling on the land use practises of the former Indian tribes that led to such a lasting enrichment, and to develop a replicating land management package for the benefit of more sustainable and more concentrated present-day small-holders settlement. A technical and socio-economically successful package would substantially contribute to sequestering of atmospheric CO₂ the managed soils, and at the same time leave more primary forest intact with its own CO₂ stock and sink function and biodiversity values.

Keywords: Amazon region; Anthrosols; pre-Columbian settlements; Soil organic matter; Carbon sequestration; Tropical forest conservation

1) Trabalho apresentado no Symposium 18 - Anthropogenic factors of soil formation – of the 17th World Congress of Soil Science – Bangkok, Thailand, 2002. CD-ROM, Paper 1.935

2) MPEG, Belém, Brazil and ISRIC, Wageningen, the Netherlands <sombroek@isric.nl>

3) Dept Ecologia, Museo Paraense Emilio Goeldi, Belém, Brazil

4) Dept Solos, Embrapa-CPATU, Belém, Brazil

5) Embrapa-CNPS, Rio de Janeiro, Brazil

6) Southern Illinois University, Edwardsville, USA

7) University of Bayreuth, Germany

INTRODUCTION

Anthropogenic black earths are of frequent occurrence in the Amazon region, especially the Brazilian part. They are locally known as *Terra Preta*, *Terra Preta-do-Indio*, or *Terra Preta arqueológica/antrópica* and are the lasting effect of intensive occupation by pre-Columbian Amer-indian tribal communities in the region. They have been mentioned and described by archeologists and anthropologists (Denevan 1992; Whitehead, 1996; Petersen et al, 2001), but also by ecologists, agronomists and soil scientists (Sombroek, 1966; Smith 1980; Moran, 1990; Pabst, 1993; Glaser, 1999; Woods, McCann and Meyer, 2000; Cleary, 2001; Kern et al, 2001). There are also two web-sites on the matter, one in Germany (english): www.geo.uni-bayreuth.de/bodenkunde/terra_preta/index.html, and one in Brasil (portuguese): www.mpeg.tpaindex.html>. Terra Preta soils contain an abundance of artefacts (ceramics, potsherds, stone tools, charcoal) , are black or very dark brown till at least 50 cm depth, have high levels of plant-available P levels (above 100 ppm) and are less acid than the surrounding soils. Nowadays, they are preferentially sought for agricultural settlement by traditional small-holder subsistence farming communities, who recognise them easily even when under high forest cover – the tree species assemblage being different. The soils occur usually in small patches averaging 20 ha, but 300-500 ha. sites have been reported as well, especially for the uplands (*terra firme*) in the neighbourhood of confluences of main rivers: Rio Negro – Rio Solimões near Manaus, Rio Madeira – Rio Amazonas near Itacoatiara, Rio Tapajos – Rio Amazonas near Santarem, and Marajó island in the mouth of the Amazonas –Tocantins system [composite maplet with all known sites and maplets of Belterra and Caxiuaná sites to be shown at oral presentation]. The smaller patches of Terra Preta soils are also found in upper catchment areas, all along the main tributaries of the Amazon, along rivulets (*igarapés*) feeding into these main tributaries such as former ring villages of the Upper Xingu (Heckenberger et al, 1999), and along age-old forest tracks between river systems (McCann, 1999). For south-eastern Amazonas state and north-eastern Rondonia, recent geo-pedological surveys for ecologic-economic zoning yielded estimates of TP sites every 5 km along the *igarapés*, and an over-all spatial coverage of one per 2 km² (verbal comm. Horbes, CPRM; Jarbas and Rodrigues, Embrapa-Soils; and Josuel Ravoni of SEDAM, respectively). Terra Preta soils occur also in the Amazon parts of Colombia (Araracuara area: Andrade, 1986; Eden et al, 1984; Mora, 1999), southern Venezuela and Peru, but apparently less frequently.

Archeologically, a number of different ceramic phases are distinguished, associated with former tribal chiefdoms (Hilbert, 1968), most of them having their peak of cultural development

between 500 and 1500 AD. Most researchers now estimate the total Indian population in Amazonia to have been between 5 and 10 million at the time of the Spanish and Portuguese conquest (Roosevelt, 1996; Denevan, 1998; Petersen et al, 2001) – soon to be decimated and then becoming by and large nomadic (Cleary, 2001).

In a number of cases the patches of Terra Preta are surrounded by a broad band of Terra Mulata soils. These are dark greyish brown rather than black in their upper horizons, have less elevated P levels, no or very few artefacts, but still elevated levels of soil organic matter in comparison with the undisturbed forest soils. They have been described by Sombroek (1966) for the Belterra area south of Santarém, by Woods and McCann (1999) for the Arapiuns area west of Santarém, by Kern (1996) for the Caxiuaná area east of the lower Xingú area, by Horbes (verbal comm.) along the Rio Juma, 100 km north of Apui on the TransAmazonica, and by Jarbas Cunha (in prep.) for the Humaitá area on the Madeira river. These authors consider such areas as pre-Columbian agricultural fields around the former Indian major villages, in-between the kitchen middens or homestead fields of the Terra Preta sites proper and the unaltered soils of the forests, where only foraging took place. Whereas the increase in soil organic matter and mineral fertility on the homesteads may have taken place accidentally, the Terra Mulata areas would be the result of intentional application of human or animal manure, green manure from forest or river, charcoal and ashes, fishing and hunting remains and calcium from pounded mollusc shells or root accretions (Roosmalen, pers. comm). This would allow a semi-permanent horticultural type of food production once a piece of forest-land had been cleared strenuously with stone hand-axes and subsequent burning.

Both Terra Preta and Terra Mulata soils are surrounded by natural soils under primary forest, where most of the organic matter is concentrated in the upper 20 cm. They may be Ferralsols (mainly xanthic, sometimes orthic or rhodic), Acrisols or else, have a sandy to clayey texture, variable stoniness, and nearly invariably with high amount of exchangeable Al (pH 4.5 or lower). It is nowadays generally accepted that the TP and TM are not the result of natural soil forming factors on rich parent material, but the lasting effect of anthropogenic influences on natural forest soils that happened to be located on strategic river bluffs, indigenous forest tracts between tribal communities, heads of rapids, etc. In fact, the deeper subsoils of the TP and TM soils have still the same clay mineralogy, colour, texture and structure as the nearby natural soils.

CHARACTERISTICS AND CLASSIFICATION CRITERIA

A geographic database is being compiled of all the sites where Terra Preta and Terra Mulata soils have been identified, accompanied by relevant site- and soil profile descriptions and

analytical data, and several representative soil monolith sequences. It is to be expected that there will be a substantial variation in detailed characteristics of both soil types, in dependence of the original soil, the length of the pre-Columbian occupation, the precise land use and management practises of the various Indian chiefdoms, and the type of water of the neighbouring river. The latter, in particular, will have caused major differences. The plant- and animal resources of clear waters (no sediment load, but neutral pH and presence of calcium: *agua clara* or *agua azul*) are different from the muddy river waters (an appreciable load of sediments and a variety of cations: *agua branca* or *barrenta*), or from the black waters (no sediments nor cations, but very acid due to the presence of humic acids: *agua preta*). The latter situation occurs most frequently in the area of podzolised soils in the upper and middle courses of the Rio Negro, where Terra Preta patches are rare and small, and Terra Mulata has not been reported thus far.

The data available thus far point to a much higher soil organic matter (SOM) content of the two soil types – often double the normal amount - which has moreover a high degree of permanency. The hetero-trophic respiration by microbial digestion is apparently low for the subsurface and subsoil layers, resulting in high percentages of stable or “recalcitrant“ SOM with C/N values of about 12. At the same time this SOM is apparently more re-active than the organic matter in the primary forest soils: the CEC-SOM values are 50-100% higher (Sombroek, 1995). [graph with additional curve points to be shown at oral presentation]. This means that the soils concerned can accept and subsequently release substantially more fresh plant nutrients.

Detailed studies on the specific composition of the SOM of TP and TM soils are few (Glaser, 1999). A specific microbial activity may be involved, with certain microbes acting as inoculate (Woods and McCann, 1999). Studies on bacterial functional redundancy” of the soils (Sparovek, in press) may elucidate this aspect.

The values of plant-available phosphorus are strikingly high in the TP variant (often over 150 ppm P-Bray in the subsurface layers), but only moderately high in the TM variant (50-150 ppm for the few profiles analysed), in comparison with normal forest soils (Sombroek, 1966; Smith, 1980). Detailed studies on fractional composition of the phosphorus in TP/TM are underway (Glaser; Cravo; Comerford and Falcão).

The amounts of exchangeable bases are higher than in the forest soils, but the base saturation is only moderately high: 20 to 70 %, compared to about 15 % in non-enriched soils, and associated with pH values of 4.5 –5.5, instead of 4.0-4.5. Calcium and Magnesium values are always higher, but their total amounts vary strongly – no doubt related with the calcium content of the nearby river water.

Agricultural occupation by present-day small-holder farmers without access to commercial fertilisers can be maintained for 10-15 years without appreciable diminution of the chemical

fertility status, but TP sites of longer occupation show a definite decline in agricultural productivity (Jarbas Cunha, in prep.).

Classification criteria

It may be obvious that of horthic Anthrosol is concerned. The data available thus far do not yet permit a field-relevant detailed pedologic classification scheme, but some suggested classification criteria are as follows (ref.: World Reference base for Soil Resources, FAO/ISRIC, 2001; US Soil Taxonomy, 1992 and EMBRAPA's new Classificação de Solos Brasileiros, 2000):

- Thickness of organic matter-rich layer: at least 50 cm (different for sandy vs. clayey?);
- Colour (in moist condition; Munsell terminology and notation): black or reddish black, very dark (reddish) brown, brownish black or dark (reddish) brown, i.e. value and chroma 3 or less; compare "mollic" and "umbric". For the soil profiles with abundance of ceramics - the Terra Preta proper - the colour values and chromas may be only 2 or less (reddish black, brownish black, or very dark reddish brown), comparable with the criterion "melanic";
- High organic-carbon content, averaged over 50 cm depth: 1.4% organic carbon or more (comparable with "humic" denominations for some Ferralsols and Nitisols);
- High value of CEC per percent organic carbon: more than 50% higher than in the adjacent normal soil (to be established by graphical method of Embrapa-Solos);
- High amount of easily extractable phosphorus content: more than 100 mg/kg Olsen-extractable P₂O₅ in the fine earth fraction, or more than 0.25% extractable in 1% citric acid, both averaged over the first 50 cm depth (critical values to vary with texture class?);
- Presence of artefacts/ceramics, averaged over 50 cm: higher or lower than 1% by volume, for TP and TM respectively;
- Abundance of "krotovinas", i.e. mesofauna channels below the blackish horizon, filled with organic matter-rich soil material as an additional criterion?;
- Presence of charcoal ("black" or "pyrogenic carbon") in the coarse-earth fraction of at least 1 % by volume, averaged over the first 50 cm, as another additional criterion?

Further subdivisions can be made on over-all texture (sandy vs clayey) and on pH or base saturation (<35, 35-50, 50-80 and > 80 %?). Some possible prefixes are: para-hortic, proto-hortic, hypo-hortic, hyper-hortic, molli-humic or umbri-humic. Tropo-plaggic could be applied if a clear thickening of the profile has taken place through solid earth additions, such as on burial mounds.

RE-DISCOVERING AND APPLYING THE PRE-COLUMBIAN INDIGENOUS TECHNIQUES FOR SUSTAINABLE LAND USE

Field research

For a number of the Terra Preta sites the enrichment may have been accidental as kitchen-midden formation. For others, especially where Terra Mulata bands of considerable extent are present, a special land management technique would have been involved, which resulted in a lasting soil organic matter increase and a substantially higher chemical soil fertility.

The present-day remnants of the Amer-indian population groups in the Amazon region have moved from their original sites to new areas (Cunha, 1992), with nomadic phases in-between, and often do not know how the TP/TM in their new living places happened to come about (Pabst, 1993). There are only a few verbal reports (Roosmalen; Mosimann da Silva) that in isolated upper catchment areas such as the upper Xingú, current Indian land use practises may still be reminiscent of pre-Columbian practises. Local interviews, patient on-the-spot recording of such practises on annual basis, and screening of fragmentary descriptions by early travellers will give some insight.

A number of research items can be formulated, as follows:

-Considering that relatively high contents of SOM, of phosphorus and of calcium/magnesium appear to be characteristic for TP/TM soils, how would former Amerindian groups have collected the required high amounts of the basic materials, in an environment of strongly weathered and acid soils: Were these materials forest litter or non-wooded undergrowth; aquatic grasses; shells, Ca-containing root exudates; animal bones; charcoal and ashes, or all together?

-What techniques were used to turn these basic materials into the characteristic stable and at the same time active SOM of the TP/TM soils: manual grinding and mixing, rotting and composting, or intentional animal manure production?

-If an animal digestion or trampling- and manure producing stage was essential (as in the case of the north-western European *plaggen* soils), then what type of animals were used in pens, corrals or stables, given the absence of cows and sheep? Could land- and /or aquatic giant turtles (*jabotis* or *tartarugas*) have been used for the purpose, as might be concluded from the Carvajal narrative (original in 1542; re-edited in 1934) on their abundance in erstwhile Amerindian villages?

Laboratory research

Advanced laboratory research on sample material of representative sequences of TP-TM-TC (the latter standing for the common, natural soil) may focus on the following questions:

- Are specific meso-fauna or microbial communities (worms, bacteria) involved in the creation and reproduction of the stable SOM in the TP/TM soils?

- What are the precise physico-chemical characteristics of the stable SOM in TP and TM soils respectively?
- Is the SOM stability the result of crypto-crystalline complexation with the predominant clay mineral such as kaolinite, through linkages with Ca and P compounds or elements, or is it the “black” or “pyrogenic” carbon of charcoal application on its own, or a combination of both?
- What is the reason for the relatively high activity per unit of SOM in the TP/TM, in comparison with to of non-enriched counterpart soils (TC), in the form of higher nutrient and moisture storage capacity?
- How long will these favourable properties of the TP/TM have been lasting after abandonment by the Amerindians (^{14}C and ^{13}C analysis), and to which degree are they degrading under present-day occupation by small-holding subsistence farmers who do not use indigenous soil improving techniques on permanent fields?

Application research

Once the above field and laboratory research items are elucidated, a number of application related items have to be solved:

- Can the favourable properties of the TP/TM soils be emulated/replicated on other sites, without depleting the surrounding countryside and within a time span of decades rather than centuries?
- What would be the most promising combination of methods to create new TP/TM, and can they be integrated in current agro-forestry practises and secondary forest (*capoeira*) management? If this is to include an animal husbandry stage, then which animals should be involved most effectively - from technical, economic and social points of view?
- Would the present-day small-holder rural population of the region be receptive to apply the required extra manual labour, and would they need incentives, financially or otherwise, such as an assured supply of lime and rock-phosphate?
- Which local institutions can best guide the experimental phase of a TP/TM replication programme, and how long should the testing last to become sure of the permanency of the newly added SOM?
- What is the quantitative potential, in the form of tonnes of Carbon sequestered and the reduction of hectares of deforestation, if the creation of new Terras Pretas and Terras Mulatas in the Amazon region would be undertaken systematically? How much international financial support per ha or farm household, through the Clean Development Mechanism of the Kyoto Protocol or else, would be needed annually?

- Finally, can and should the acquired insights and subsequent sustainable development packages be made beneficial to the remnants of the former Amer-indian tribes inside and outside their formal reserve areas, in the form of extension and/or financial compensation for the re-use of the traditional knowledge systems of their forefathers – considering the concept of communal intellectual property rights (cf. Posey, 2000)?

TOWARDS A TERRA-PRETA-NOVA PROJECT

Discussions of the above series of questions and hypotheses have led to the formulation of a multi-disciplinary and multi-institutional Brazilian research-cum-development project called Terra Preta Nova (TPN), with participation of scientists from other Latin American countries, USA and Europe. Its aims are: to contribute to the contracted and sustainable use of Amazon soils by small-holding farmers and the conservation of the primary forests, through the replication of the anthropogenic black earths of the pre-Columbian Indian tribal communities. It would also contribute to the carbon stock and sink functions of the Amazon forests and soils, and better preserve the Amer-indian Amazon cultural legacy.

The activities of the envisaged project are as follows:

- Region-wide surveys and local interviews on Terra Preta (TP) and associated Terra Mulata (TM) occurrences, genesis, and related ethno-biodiversity aspects;
- Anthropologic and archeologic site research and directed salvage activities in areas of rapid settlement (such as Apui, Humaitá, Itaituba, Novo Progresso; Araracuara in Colombia)
- Routine and specialised analysis on sample material, aiming to establish the reasons for the high stability and the high cation exchange capacity of the increased soil organic matter concerned;
- Laboratory experiments on the manufacturing of stable and active soil organic matter, using early indications of the surveys and the interviews;
- Directed agronomic and agro-forestry experiments in farmers' fields, to replicate TP/TM, in conjunction with existing projects on recuperation of degraded Amazonian lands, and on agro-forestry practises;
- Extension of the resulting package(s) of TPN soil management practises to priority areas of integrated environmental management - i.e. conservation of natural vegetative cover and sustainable small-holder settlement;
- Awareness promotion on the value of the indigenous patrimony, through state-level educational publications, reference collections on TP/TM material, etc.

It is hoped that the Global Environmental Facility (GEF, as an outcome of the Rio de Janeiro World Summit of the Environment in 1992), will be interested in co-financing of the project, through its focal area #13, entitled "Conservation and sustainable use of biological diversity important for agriculture".

REFERENCES.

- Andrade, A. 1986. Investigación arqueológica de los Anthrosoles de Araracuara, Amazonia. Fundación de Investigaciones Arqueológicas Nacionales, Banco da República, Bogotá. :19-64
- Carvajal, G. 1934. Discovery of the Orellana River. In: H.D. Heaton (ed). The Discovery of the Amazon According to the Account of Friar Gaspar de Carvajal and other Documents, with an Introduction by José Toribio Medina. American Geographical Society Publication 17. New York : 167-242.
- Cleary, D. 2001. Towards an environmental history of the Amazon; from pre-history to the nineteenth century. *Latin American Research Review* 36 (2): 65-95
- Cunha, M. C. (ed). 1992. História dos Índios no Brasil. Companhia das Letras da Secretaria Municipal da Cultura de São Paulo.
- Denevan, W. M. 1998. Comments on prehistoric agriculture in Amazonia. *Culture & Agriculture*, 20: 54-59.
- Denevan, W. M. 1992. The Aboriginal Population of Amazonia. In: The Native Population of the Americas in 1492, edited by W. Denevan, University of Wisconsin Press, Madison. : 205-234
- Eden, M. J., Bray, W., Herrera, L. and McEwan, C. 1984. Terra Preta soils and their archaeological context in the Caquetá basin of south-east Colombia. *American Antiquity*, 49: 125-140.
- Glaser, B. 1999. Eigenschaften und Stabilität des Humuskörpers der Indiasner Schwarzerden Amazoniens. *Bayreuther Bodenkundliche Berichte* 68. 195 pp
- Heckenberger, M. J. 1996. War and Peace in the Shadow of Empire: Socio-political change in the Upper Xingu of south-eastern Amazonia, A.D. 1400 -2000. In: University of Pittsburgh (Editor), 481 pp.
- Heckenberger, M.J., Petersen, J. B. and Neves, E. G. 1999. Village size and permanence in Amazonia: Two archaeological examples from Brazil. *Latin American Antiquity*, 10.
- Hilbert, P. 1968. Archäologische Untersuchungen am mittleren Amazonas, *Marburger Studien zur Volkerkunde*, Vol. 1, Verlag Dietrich Reimer, Berlin.
- Kern, D.C. 1996. Geoquímica e pedogeoquímica de sítios arqueológicos com Terra Preta na Floresta Nacional de Caxiuaná (Portel-Pará). Unpublished PhD thesis Univ. Federal de Belém, Brazil.
- Kern, D.C., Costa, M.L., and Frazão, F.J.L. 2001. Evolution of the scientific knowledge regarding Black Earth soil in the Amazon. *The Yearbook of the Conference of Latin American Geographers*, 26. (Conference in Benicassim, Spain, July 2001).
- McCann, J. M., Woods, W. I. and Meyer, D.W. 2000. Organic matter and Anthrosols in Amazonia: Interpreting the Amerindian Legacy. In: R.M. Rees et al (eds). *Sustainable Management of Soil Organic Matter*. CAB International, Wallingford. UK : 180-189.
- McCann, J.M. 1999. Before 1492. The making of the pre-Columbian Landscape. *Ecological Restoration* 17 (1&2) : 15-30
- Mora, S. 1999. Suelos negros y sociedad: un sistema agrícola de entonces, un sistema agrícola de ahora?. *Desarrollo sostenible en la Amazonía*. Bogotá : 32-45
- Moran, E.E. 1990. A ecologia humana das populações da Amazônia. Editora Voçes,

- Petrópolis, Brazil.
- Pabst, E. E. 1993. Terra Preta, ein Beitrag zur Genese-discussion auf der Basis von Geländearbeiten bei Tupi-Volkern Amazoniens. Unpublished Hab. thesis, Univ. Kassel.
- Petersen, J.B., Neves, E. and Heckenberger, M.J. 2001. Gift from the past: Terra Preta and prehistoric Amer- indian occupation in Amazonia. pp 86-107 .In: C. McEwan, C. Barreto and E. Neves (eds). *Unknown Amazon*. The British Museum Press, London. : 86-107
- Posey, D.A. 2000. Biodiversity, genetic resources and indigenous peoples in Amazonia: (re)discovering the wealth of traditional resources of native Amazonians. In: A. Hall (ed). *Amazonia at the crossroads*. Institute of Latin American Studies, London. : 188-204
- Roosevelt, A. 1996 "Paleo-indian Cave Dwellers in the Amazon: The Peopling of the Americas." *Science* 272 (19 Apr.): 373-84.
- Sombroek, W. G. 1966. *Amazon Soils: a reconnaissance of the soils of the Brazilian Amazon valley*. Pudoc, Wageningen. 292 pp
- Sombroek, W. G., Nachtergaele, F.O. and Hebel, A. 1993. Amounts, dynamics and sequestering of carbon in tropical and subtropical soils. *Ambio* 22: 417-426.
- Sombroek, W.G. 1995. Aspects of soil organic matter and nutrient cycling in relation to climate change and agricultural sustainability. In: *Proceedings Intern. Symposium on nuclear and related techniques in soil-plant studies on sustainable agriculture and environmental preservation*. IAEA/FAO Joint Division. Vienna. : 15-26
- Whitehead, N. 1996. Amazonian Archaeology: Searching for Paradise? A Review of Recent Literature and Fieldwork. *Journal of Archaeological Research* 4 (3): 241-64.
- Woods, W.I. and Mann, C.C. 2000. Earth movers of the Amazon. *Science* 287: 786-789.
- Woods, W.I. and McCann, J. M. 1999. The anthropogenic origin and persistence of Amazonian dark earths. *The Yearbook of the Conference of Latin American Geographers*, 25: 7-14.