

VARIABILITY IN INDIVIDUAL NODULE ACTIVITY OF SINGLE STRAINS OF *RHIZOBIUM ETLI* AND *R. TROPICI* IN SYMBIOSIS WITH *PHASEOLUS VULGARIS*¹

NADJA M. HORTA DE SÁ², IVANILDO EVÓDIO MARRIEL³ e AVÍLIO A. FRANCO⁴

ABSTRACT - A green-house experiment in Leonard jars was conducted to study the variability of nitrogenase activity (acetylene reduction) of individual nodules of *Phaseolus* beans inoculated separately with 19 strains of *Rhizobium tropici* and 6 strains of *R. etli*. Both species showed a wide range of activities with a C.V.=80% for *R. tropici* and C.V.=70% for *R. etli*. These data indicate that *Rhizobium* species with reiterated nif genes may present similar variability of single nodule activity as *Rhizobium* species containing a single copy of nif gene and that the use of either *R. tropici* or *R. etli* may be used for inoculant production with precaution for loss of effectiveness.

Index terms: *Phaseolus vulgaris*, leonard jars, nitrogenase, acetylene

VARIABILIDADE NA ATIVIDADE DE NÓDULOS INDIVIDUAIS FORMADOS POR ESTIRPES DE *RHIZOBIUM ETLI* E *R. TROPICI* EM SIMBIOSE COM O FEIJOEIRO

RESUMO - Foi conduzido, em casa de vegetação, em vasos-de-leonard, um experimento para estudar a variabilidade da atividade da nitrogenase (redução de acetileno) de nódulos individuais de feijão (*Phaseolus vulgaris*), infectados separadamente com 19 estirpes de *Rhizobium tropici* e 6 estirpes de *R. etli*. As 2 espécies mostraram grande variabilidade nas atividades, com um C.V.=80% para *R. tropici* e C.V.=70% para *R. etli*. Estes dados indicam que espécies de *Rhizobium* com "gens nif" repetidos podem apresentar variabilidade similar na atividade de nódulos individuais em relação às espécies de *Rhizobium* que contêm uma única cópia dos "gens nif"; indicam, também, que a utilização tanto de *R. tropici* como *R. etli* na produção de inoculante deve ser criteriosa, em face da perda de efetividade.

Termos para indexação: *Phaseolus vulgaris*, vasos-de-leonard, nitrogenase, acetileno.

INTRODUCTION

Biological nitrogen fixation in *Phaseolus* beans has frequently been used as an example of poor response of a legume crop to inoculation. The presence of large population of rhizobia able to nodulate and fix nitrogen with this crop, its short cycle, the sensitivity of the host to environmental stresses and the genetic instability of the symbiont may be responsible, under different cropping systems, to the lack of response of *Phaseolus* bean to inoculation (Franco, 1977).

Rhizobia genetic instability in laboratory manipulation, resulting in loss or decreased ability of the symbiont to nodulate or fix nitrogen in symbiosis with the host, have been registered for a long time. These alterations may be a result of mutagenic agents such as acridines, UV-light, SDS, several types of radiation, etc. (Zurkowski et al., 1973, Mathis et al., 1985 and Barbur & Elkan, 1989). The alterations have also been observed to occur under stress of high temperature (Djordjevic et al., 1983, Weaver & Wright, 1987) or even spontaneously during the routine sub-cultivation (Weaver & Frederick, 1982). Franco (1974) observed large variation in colony morphology and symbiotic effectiveness of several *Bradyrhizobium* spp. strains grown and stored in yeast mannitol agar under oil at room temperature. Herridge & Roughley (1975) tested 17 stock cul-

¹ Accepted for publication on September 20, 1994.

² Bióloga, Ph.D., Dep. Botânica - ICB/UFMG CEP 31270-901, Belo Horizonte, MG.

³ Eng.-Agr., M.Sc. CNPMS/EMBRAPA, CEP 35700-000, Sete Lagoas, MG.

⁴ Eng.-Agr., Ph.D., CNPMS/EMBRAPA, CEP 23851-970, Seropédica, RJ

tures of *Bradyrhizobium* sp. CB-756, obtained from several culture collections in different institutions and found great variation among cultures. Some of them have lost effectiveness to less than half to that with the highest effectiveness. Freeze drying or maintaining the culture in yeast mannitol agar medium may yield variants (Roughley, 1976). Peres et al. (1984) have shown great internal variation in the effectiveness of N₂ fixation in several strains of *Bradyrhizobium japonicum* and Weaver & Wright (1987) in *B. sp.* (Vigna).

The genetic instability seems to be more frequent in fast than in slow growing rhizobia. The three species of *Rhizobium* (*R. leguminosarum* bv. *phaseoli*, *R. tropici* and *R. etli*) that forms effective symbiosis with *Phaseolus* bean are fast growing. In these strains, the genes controlling biological nitrogen fixation (*nif* genes) are located in plasmids (Psym), in contrast with *Bradyrhizobium* that have the *nif* genes located in the chromosomes (Rosenberg et al., 1981, Kondorosi et al., 1982, Martinez et al., 1990). *R. leguminosarum* bv. *phaseoli* and *R. etli* strains present multicopies of the *nif* genes and nodulate only *Phaseolus* bean (Martinez et al., 1987 and Segóvia et al., 1993), while *R. tropici*, which is also able to nodulate and fix nitrogen in symbiosis with *Leucaena leucocephala*, has only one copy of the *nif* genes (Martinez et al., 1991). The Psym represent the molecular base of variability and instability in symbiotic properties found in rhizobia able to form effective symbiosis with *Phaseolus* beans. *R. tropici* strains have been found to be more heat tolerant and genetic stable than *R. etli* or *R. leguminosarum* bv. *phaseoli* (Mercante, 1993). Even though *R. etli* has only recently been described and there are not many studies on this species we may consider that most of bean specific rhizobia isolated from the Americas may be classified within this species (Segóvia et al., 1993).

The measurement of nitrogenase activity by acetylene reduction has been questioned by Minchin et al. (1983). They observed a sharp decline in ethylene production after a 10-minute exposure of nodules to acetylene. This inhibitory effect has been found to be variable with symbiotic systems, plant age, light intensity, pO₂, low temperature, water lodging, water stress and by differ-

ent ways of handling the material (Stralioto, 1990). In spite of those observations, Peres et al. (1984) have demonstrated that single nodule activities of several *Bradyrhizobium japonicum* strain were correlated with the effectiveness of their isolates with the host. Faria et al. (1984) have also observed in several legume trees good correlation between individual nodule activity and effectiveness of their isolates.

This study was aimed to compare, by measuring single nodule activity of 60 nodules of each strain, the genetic variability in the symbiotic effectiveness of 6 *R. etli* strains in comparison with 16 *R. tropici* strains.

MATERIALS AND METHODS

The experiment was conducted in a green-house using sterilized Leonard jars containing a 2:1 mixture sand: vermiculite (Vincent, 1970) and receiving 400 ml of nutrient solution without nitrogen (Norris, 1964). A complete randomized block design was used, with three repetitions, 6 strains of *Rhizobium etli* BR 365 (CNPAC 146), CPAC H 19 (Semia 476), CPAC H 30 (IPAGRO 1102), CPAC H 35 (IPAGRO 1378), CPAC H 23 (V23 RGS) and CPAC H 14; 13 strains of *R. tropici* BR 322 (CIAT 899), BR 10013 (Na 82), BR 817 (NGR 8), BR 818 (TAL 1145), BR 266 (Semia 492, CENA CO₅ II), BR 10014 (Car 22), BR 814 (DF 10), CPAC H 21 (UFP 491), CPAC H 20, CPAC H 36, CPAC H 38 (USA 1070), CPAC H 26 (IPAGRO 1020) and CFN 299. The rhizobia strains were tested in *Phaseolus vulgaris* L. cv. Negro Argel and at the same time in *Leucaena leucocephala* Witt cv. Peru to confirm to which species they belong. All strains, except BR 814, BR 817 and BR 818, were obtained originally from *Phaseolus* bean nodules.

The inoculum was prepared by growing each strain in yeast mannitol broth (Vincent, 1970) to the final logarithmic phase, standardizing to 10⁸ cells/ml and applying 2 ml of this suspension in each pot containing 2 plants.

The plants were harvested 28 days after emergence. Ten nodules of each plant of *Phaseolus* bean with fresh weight between 5 and 7 mg were detached from the root and placed into individual flasks, 10% of the air was replaced by acetylene, incubated for 10 minutes and the ethylene produced was measured as described by Peres et al. (1984).

In *Leucaena*, nodulation was considered positive when all plants of the three repetitions were nodulated.

RESULTS AND DISCUSSION

All 19 *Rhizobium* strains studied originated from the Americas were able to nodulate and fix nitrogen with *Phaseolus* bean, while 13 of them were also able to nodulate and fix nitrogen with *Leucaena*. As indicated by Segóvia et al. (1993), the first group may be classified as *R. etli* and the second group as *R. tropici* species. Even though only 3 strains were isolated from *Leucaena*, 13 of them were of the *R. tropici* species with a range of activity from 7 to 132 nmoles ethylene/h nodule⁻¹, similar to the range presented by the less promiscuous species *R. etli*, from 29 to 104 nmoles ethylene/h nodule⁻¹ (Sá et al., 1993).

Peres et al. (1984) had observed a good correlation between the activity of the individual nodules and the efficiency of their isolates in symbiosis with the host. The 19 rhizobial strains used in this study were chosen among the most efficient strains for inoculation for *Phaseolus* bean. A large

variation was found acetylene reduction of individual nodules within and amongst strains (Fig. 1, 2, 3, 4 and 5) with a range of coefficient of variation (C.V.) in each strain of 23 to 105 in *R. etli* and 32 to 165 in *R. tropici*. The mean C.V. for the 6 *R. etli* studied was 70% and for the 13 *R. tropici* was 80%. It is possible that amongst *R. tropici* there is a large range in effectiveness of the strains even though some of the strains are as/or more efficient than *R. etli* when in symbiosis with *Phaseolus* bean. The individual nodule activity of none of both *R. etli* or *R. tropici* fitted a normal distribution (Fig. 1, 2, 3, 4 and 5). When the activity of both species were pooled in the same figure (Fig. 6), the range of activity of individual nodules of both species was very similar, at the lowest (0-4 nmoles C₂H₄/h/nodule) up to the highest (100 nmoles 0-4 nmoles C₂H₄/h nodule⁻¹) activities.

Variability in symbiotic properties is a common characteristic in *Rhizobium* strains holding "nif gens" in plasmids (Flores et al., 1988). This fact can be more relevant among *R. etli* strains having more than one copy of "nif gens" in the plasmid (Psym), making them subject to a greater variabil-




FIG. 1. Variability in nitrogenase activity (acetylene reduction) in individual bean nodules inoculated with *R. etli* strains: (A) CPAC H35, (B) CPAC H14, (C) CPAC H30 and (D) CPAC H23.




FIG. 2. Variability in nitrogenase activity (acetylene reduction) in individual bean nodules inoculated with *R. etli* strains: (A) CPAC H19, (B) Br 365 and *R. tropici* strains: (C) Br 10013, (D) CPAC H26.




FIG. 3. Variability in nitrogenase activity (acetylene reduction) in individual bean nodules inoculated with *R. tropici* strains: (A) Br 322, (B) Br 814, (C) Br 266 and (D) Br 10014.




FIG. 4. Variability in nitrogenase activity (acetylene reduction) in individual bean nodules inoculated with *R. tropici* strains: (A) Br 817, (B) Br H36, (C) CPAC H20 and (D) CPAC H21.




FIG. 5. Variability in nitrogenase activity (acetylene reduction) in individual bean nodules inoculated with *R. tropici* strains: (A) CPAC H38, (B) CFN 299 and (C) Br 818.




FIG. 6. Variability in nitrogenase activity (acetylene reduction) in 360 bean nodules (60 / strain) inoculated with *R. etli* strains (Br 365, CPAC H19, CPAC H30, CPAC H35, CPAC H23, CPAC H14) and 780 bean nodules (60 / strain) inoculated with *R. tropici* strains (Br 322, CPAC H21, CPAC H20, Br 10013, Br 817, Br 818, Br 266, Br 10014, CPAC H36, CPAC H38, CPAC H26, Br 814, CFN 299).

* *R. etli*, C.V. 70%



R. tropici, C. V. 80%



ity in relation to the *R. tropici* strains which present only one copy of *nif* gens and are considered more stable (Flores et al., 1988). They are also tolerant to stress factors such as acidity (Vargas & Graham, 1988) and high temperatures (Karanja & Wood, 1988, Romero & Rosenblueth, 1990). However, the strains of both species herein studied, showed high variability and at similar levels. This was also observed when both species were grown at their highest temperature they would grow (Sá et al., 1993).

The variability of individual *phaseolus* beans nodule activity and the genetic variability of the symbiont must be further studied. The data presented in this study indicate however that the use of either *R. tropici* or *R. etli* may be used for inoculant production with precaution for loss of effectiveness.

CONCLUSIONS

Strains of *Rhizobium etli* and *R. tropici* showed a wide range of nitrogenase activity (acetylene reduction). That indicates that *Rhizobium* species with reiterated *nif* genes may present similar variability of single nodule activity as *Rhizobium* species containing a single copy of *nif* gene and that the use of either *R. tropici* or *R. etli* may be used for inoculant productions with precaution for loss of effectiveness.

ACKNOWLEDGEMENTS

The authors thank the laboratory technicians Paulo Figueiredo and Renato Simões for their collaboration; FINEP-PADCT, CNPq and EMBRAPA for their financial support.

"This work is part of the first author's Ph. D. thesis, taking place at the UFRJ Microbiology Institute".

REFERENCES

- BARBOUR, W. M.; ELKAN, G. H. Relationship of the presence and copy number of plasmids to exopolysaccharide production on symbiotic effectiveness in *Rhizobium fredii* USD A 206. **Applied and Environmental Microbiology**, v.55, p.813-818, 1989.
- DJORDJEVIC, M. A.; ZURKOWSKI, W.; SHINE, J.; ROLFE, B. Sym plasmid transfer to various symbiotic mutants of *Rhizobium trifolii*, *R. leguminosarum* and *R. meliloti*. **Journal of Bacteriology**, v.156, p.1035-1045, 1983.
- FARIA, S. M. de; MOREIRA, V. C. G.; FRANCO, A. A. Seleção de estirpes de *Rhizobium* para espécies leguminosas florestais. **Pesquisa Agropecuária Brasileira**, Brasília, v.19, s/n, p.175-179, 1984.
- FLORES, M.; GONZÁLEZ, V.; PARDO, M. A.; LEIJA, A.; MARTINEZ, E.; ROMERO, D.; PINERO, D.; D'ÁVILA, G.; PALACIOS, R. Genomic instability in *Rhizobium phaseoli*. **Journal of Bacteriology**, v.170. n.3, p.1191-1196, 1988.
- FRANCO, A. A. Competition amongst rhizobial strains of the colonization and nodulation of two tropical legumes. [S.l.]: University New South Wales, 1974. 183p. Tese de Mestrado.
- FRANCO, A. A. Nutritional restraints for tropical grain legume symbiosis. In: VINCENT, J. M.; WHITNEY, A.S.; BOSE, J. (Eds.). **Exploiting the legume *Rhizobium* symbiosis in tropical agriculture**. Hawaii: University of Hawaii. Niftal Project, 1977. p.237-252. (College Tropical Agriculture Miscellaneous Publication, 145).
- HERRIDGE, D. F.; ROUGHLEY, R. J. Variation in colony characteristics and symbiotic effectiveness of *Rhizobium*. **Journal of Applied Bacteriology**, v.38, p.19-27, 1975.
- KARANJA, N. K.; WOOD, M. Selecting *Rhizobium phaseoli* strains for use with beans (*Phaseolus vulgaris* L.) in Kenya: Tolerance of high soil temperature and antibiotic resistance. **Plant and Soil**, v.112, p.15-22, 1988.
- KONDOROSI, A.; KONDOROSI, E.; PANKHURST, C. E.; BROUGHTON, W. J.; BANFALVI, Z. Mobilization of a *Rhizobium meliloti* megaplasmid carrying nodulation and nitrogen fixation genes into other rhizobia and *Agrobacterium*. **Molecular and General Genetic**, v.188, p.433-439, 1982.
- MARTINEZ, E.; PALACIOS, R.; SÁNCHEZ, F. Nitrogen-fixing nodules induced by *Agrobacterium tumefaciens* harboring *Rhizobium phaseoli* plasmids. **Journal of Bacteriology**, v.169, n.6, p.2828-2834, 1987.
- MARTINEZ, E.; ROMERO, D.; PALACIOS, R. The *Rhizobium* genome. CRC. **Critical Review Plant Science**, v.9, p.59-93, 1990.
- MARTINEZ-ROMERO, E.; SEGOVIA, L.; MERCANTE, F. M.; FRANCO, A. A.; GRAHAM, P.; PARDO, M. A. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. **International Journal of Systematic Bacteriology**, v.41, n.3, p.417-426, 1991.
- MATHIS, J. N.; BARBOUR, W. M.; ELKAN, G. H. Effect of sym plasmid curing on symbiotic effectiveness in *Rhizobium fredii*. **Applied and Environmental Microbiology**, v.49, p.1385-1388, 1985.
- MERCANTE, F. M. **Uso de *Leucaena leucocephala* na obtenção de *Rhizobium* tolerante à temperatura elevada para inoculação do feijoeiro**. Itaguaí, Rio de Janeiro: Universidade Federal do Rio de Janeiro. Instituto de Agronomia, 1993. 126p. Dissertação de Mestrado.
- MINCHIN, F. R.; WITTY, J. F.; SHELHY, J. E.; MULLER, M. A. A major error in the acetylene reduction assay: decreases in nodular activity under assay conditions. **Journal of Experimental Botany**, v.34, p.641-649, 1983.
- NORRIS, D. O. Techniques used in work *Rhizobium*. In: NORRIS, D. O. (Ed.). **Some concepts and methods in sub-tropical pasture research**. London: Farmham Royal, Bucks, Commonwealth Agriculture Bureaux, 1964. p.186-198.
- PERES, J. R. R.; VARGAS, M. A. T.; SUHET, A. R. Variabilidade na eficiência em fixar nitrogênio entre isolados de uma mesma estirpe de *Rhizobium japonicum*. **Revista Brasileira de Ciência do Solo**, v.8, p.193-196, 1984.
- ROMERO, E. M.; ROSENBLUETH, M. Increased bean (*Phaseolus vulgaris* L.) nodulation competi-

- tiveness of genetically modified *Rhizobium* strains. **Applied and Environmental Microbiology**, v.56, n.8, p.2384-2388, 1990.
- ROSENBERG, G.; BOITARD, P.; DENARIE, J.; CASSE-DELBERT, F. Genes controlling early and late functions in symbiosis are located on a megaplasmid in *Rhizobium meliloti*. **Molecular and General Genetic**, v.184, p.326-333, 1981.
- ROUGHLEY, R. J. The production of high quality inoculants and their contribution to legume yield. In: NUTMAN, P. S. (Ed.). **Symbiotic nitrogen fixation in plants**. [S.l.]: Cambridge University Press, 1976. p.125-136.
- SÁ, N. M. H.; SCOTTI, M. R. M.; PAIVA, E.; FRANCO, A.; DÖBEREINER, J. Selection and characterization of *Rhizobium* spp. strains stable and capable in fixing nitrogen in bean (*Phaseolus vulgaris* L.). **Revista Brasileira de Microbiologia**, v.24, p.38-48, 1993.
- SEGOVIA, L.; YOUNG, J. P. W.; MARTINEZ-ROMERO, E. Reclassification of American *Rhizobium leguminosarum* biovar etli sp. nov. **International Journal of Systematic Bacteriology**, v.43, p.374-377, 1993.
- STRALIOTO, R. Custo energético da fixação de nitrogênio em soja inoculada com estirpes de *Bradyrhizobium japonicum* com diferentes capacidades de reciclar H_2 . Rio de Janeiro: Universidade Federal Rural do Rio de Janeiro. Instituto de Agronomia, 1990. Dissertação de Mestrado.
- VARGAS, A. A. T.; GRAHAM, P. H. *Phaseolus vulgaris* cultivar and *Rhizobium* strain variation in acid-pH tolerance and nodulation under acid condition. **Field Crops Research**, v.19, p.91-101, 1988.
- VINCENT, J. M. **A manual for the practical study at root-nodule bacteria**. Oxford: Blackwell, 1970. p.164.
- WEAVER, R. W.; FREDERICK, L. R. *Rhizobium*. In: PAGE, A. L.; MULLER, R. H.; KEENEY, D. R. (Eds.). **Methods of soil analysis**. Part 2. Chemical and microbiological properties. 2.ed. Madison: American Society of Agronomy, 1982. (Agronomy monograph, 9).
- WEAVER, R. W.; WRIGHT, S. F. Variability in effectiveness of Rhizobia during culture and in nodules. **Applied and Environmental Microbiology**, v. 53, n.12, p.2972-2974, 1987.
- ZURKOWSKI, W.; HOFFMAN, M.; LORKIEWICZ, Z. Effect of acriflavine and sodium dodecyl sulphate on infectiveness of *R. trifolii*. **Acta Microbiologica Polonica**, v.5, p.55-60, 1973.