



Genes associated with hypersensitive response (HR) in the citrus EST database (CitEST)

Simone Guidetti-Gonzalez¹, Juliana Freitas-Astúa^{2,4}, Alexandre Morais do Amaral^{3,4}, Natália F. Martins³, Angela Mehta³, Marília Santos Silva⁵ and Helaine Carrer¹

¹*Departamento de Ciências Biológicas, Escola Superior Luis de Queiróz, Universidade de São Paulo, Piracicaba, SP, Brazil.*

²*Embrapa Mandioca e Fruticultura Tropical, Cruz das Almas, BA, Brazil.*

³*Embrapa Recursos Genéticos e Biotecnologia, Brasília, DF, Brazil.*

⁴*Centro APTA Citros Sylvio Moreira, Instituto Agrônomo de Campinas, Cordeirópolis, SP, Brazil.*

⁵*Embrapa Cerrados, Planaltina, DF, Brazil.*

Abstract

Plants are continuously exposed to pathogen attack, but successful infection is rare because they protect themselves against pathogens using a wide range of response mechanisms. One of them is the hypersensitive response (HR), which is a form of cell death often associated with plant resistance to pathogen infection to prevent the spread of the pathogen from infected to uninfected tissues. Cell death is activated by recognition of pathogen-derived molecules by the resistance (*R*) gene products, and is associated with the massive accumulation of reactive oxygen species (ROS), salicylic acid (SA), and other pro-death signals such as nitric oxide (NO). The analysis of the citrus EST (CitEST) database revealed the presence of putative genes likely to be involved in HR through their products, like metacaspases, lipoxygenases, phospholipases, pathogenesis-related proteins, glutathione transferases/peroxidases, enzymes involved in the phenylpropanoid pathway and in the formation and detoxification of ROS, as well as those involved in the formation and regulation of ion channels, SA and NO. By analysis of the EST database of Citrus, it was possible to identify several putative genes that code for key enzymes involved in HR triggering and also in plant defense against biotic and abiotic stress.

Key words: biotic and abiotic stress response, plant-pathogen incompatible interaction, programmed cell death, plant disease resistance.

Received: July 21, 2006; Accepted: February 28, 2007.

Introduction

Plant-pathogen interactions involve complex, specific, and dynamic mechanisms. When a plant is resistant to a pathogen (incompatible interaction), its recognition often results in a hypersensitive reaction (HR), a rapid cell death of the plant tissues at the infection site that contributes to the limitation of growth and spread of the pathogen (Gabriel and Rolfe, 1990; Parker and Coleman, 1997; Lee and Hwang, 2005). It is generally believed that recognition of incompatible pathogens, via specific resistance genes encoded by the plant genome, leads to the activation of a number of different signaling mechanisms that initiate at the plasma membrane (Pontier *et al.*, 2002).

The initial events following the recognition of the pathogen elicitor by the plant are calcium influx, alkalization of the extracellular space, protein kinase activation, production of reactive oxygen species (ROS) and nitric oxide (NO), and transcriptional reprogramming (Dangl and Jones, 2001). The H₂O₂ produced by the oxidative burst in the dark does not play an important role in the execution of cell death, but an early and massive production of fatty acid hydroperoxides by lipoxygenases has a prominent role in cell death (Montillet *et al.*, 2005). NO and ROS could also contribute to rapid transcriptional activation of a battery of “defense genes” in and surrounding the infected cell. Some of these genes encode peroxidases, glutathione *S*-transferases, proteinase inhibitors, and various biosynthetic enzymes, such as phenylalanine ammonia lyase (PAL), which is the first enzyme in the phenylpropanoid pathway and is involved in the synthesis of low molecular weight, antimicrobial compounds known as phytoalexins

(Klessig *et al.*, 2000). Other defense mechanisms that are activated include biosynthesis of salicylic acid, induction of ethylene biosynthesis, cell-wall strengthening, and lignification (Scheel, 1998; Lam *et al.*, 2001). The oxidative burst leads to a HR and then to the establishment of systemic acquired resistance (SAR) (Rentel *et al.*, 2004).

HR has been a source of interest and controversy since its recognition in 1915; however, in recent years, the combination of molecular genetics, computer-enhanced microscopy, new cytochemical techniques and a better understanding of plant signal transduction and cell biology have helped to unravel this apparently universal plant response (Heath, 2000).

A functional genomics project has been initiated to study the molecular characterization of the main biological and agronomical traits of citrus (Forment *et al.*, 2005) and the present work also tries to help to uncover the genetic machinery of *Citrus* species. The objective of this work was to identify expressed sequence tags (ESTs) involved in the codification of key enzymes related to the HR in citrus plants. By searching the citrus EST database (CitEST), we describe putative pathways that may lead to HR in citrus plants.

Materials and Methods

In the Citrus EST Sequencing Project (Millennium Project-CNPq), various libraries of citrus cDNA were constructed with tissues from *Citrus sinensis*, *C. reticulata*, *C. limonia*, *C. aurantifolia*, *C. aurantium*, *C. limettiodes*, *C. latifolia*, *C. sunki* and *Poncirus trifoliata*, in different combinations of limiting factors (biotic and abiotic), physiological conditions (adult x juvenile) or tissue (leaf, bark, fruit, root, flower, and seed). The libraries description can be found in Targon *et al.* (this issue). The CitEST database contains over 181,000 sequenced ESTs (see CitEST: <http://biotecnologia.centrodecitricultura.br>), which were used in this work.

The identification of the genes related to HR was performed by using keywords (metacaspases, calcium and potassium channels, NAD(P)H oxidase, peroxidases, superoxide dismutase, nitrate reductase, chorismate mutase, phenylalanine ammonia-lyase, cinnamic acid 4-hydroxylase, chalcone synthase, isochorismate synthase, lipoxigenase, phospholipase, glutathione transferase, glutathione peroxidase, chitinase, osmotin, thaumatin, permeatin) and by BLASTX searches with related sequences present in other species, like *Arabidopsis thaliana*, *Lycopersicon esculentum* (tomato), *Oryza sativa* (rice), *Nicotiana tabacum* (tobacco) and others in the CitEST database. Reads with E value < 10⁻¹⁰ were individually inspected, excluding sequences unrelated to the target enzyme, whereas the remaining sequences were clustered using the CAP3 (Huang and Madan, 1999). In order to analyze phylogeny among CitEST contigs and coding region of the proteins identified in the NCBI database of known HR genes,

cladograms were made using ClustalW program (<http://www.ebi.ac.uk/clustalw/>).

Results and Discussion

The clusterization of the reads identified using keywords and BLASTX with known HR genes from other plant species in the CitEST database resulted in 227 contigs and 174 singletons. Table 1 shows 18 proteins involved in the HR process identified in the CitEST, including the total number of contigs and singletons obtained. The presence of these proteins permitted inference of a putative pathway that leads to HR in citric plants.

The NCBI database (<http://www.ncbi.nlm.nih.gov/>) has 184,881 and 28,822 nucleotides sequences of *Citrus* and *Poncirus* species, respectively. Searches in this database revealed the presence of two sequences related to glutathione peroxidase, one peroxidase, two superoxide dismutase, five phenylalanine ammonia-lyase, three cinnamic acid 4-hydroxylase, three chalcone synthase, three lipoxigenase, four chitinase and one osmotin in *Citrus* sequences, while for the *Poncirus* species there are just one superoxide dismutase, two nitrate reductase and one glutathione transferase. Some HR genes were not identified in this database, such as metacaspase, calcium and potassium channel related proteins, nitrate reductase, isochorismate synthase, chorismate mutase, phospholipase and glutathione transferase, however they were identified in the CitEST database. According to these data, this work would be con-

Table 1 - Classes of HR proteins identified in CitEST database.

HR Proteins	Total number of	
	Contigs	Singletons
Chalcone synthase	4	12
Chitinase	29	24
Chorismate mutase	5	4
Cinnamic acid 4-hydroxylase	3	2
Glutathione peroxidase	10	2
Glutathione transferase	36	4
Isochorismate synthase	2	1
Lipoxigenase	16	18
Metacaspase	3	4
Nitrate reductase	3	1
NADPH oxidase	3	3
PR-5-like proteins	10	4
Peroxidase	52	42
Phenylalanine ammonia-lyase	4	7
Phospholipase	9	12
Potassium channel	17	16
Superoxide dismutase	19	16
Two-pore calcium channel	2	2

sidered the first attempt to catalog the majority of genes related to HR in *Citrus* and *Poncirus* species.

The identification of resistance (*R*) genes that permits plants to perceive the pathogen invasion and trigger the ion flux, which culminates in HR, complements this work, and is reported in Guidetti-Gonzalez and Carrer (this issue). Several distinct MAPK (mitogen-activated protein kinases) cascades have been implicated in the regulation of plant disease resistance through signal transduction between perception of the elicitors and the activation of disease resistance genes. These MAPKs are reported in Mehta *et al.* (this issue). Besides, several pathogenesis related (PR) proteins are known to be expressed in the context of HR and usually act as antifungal factors at the end of the HR signaling cascade. Herein, the expression of two types of PR proteins typically presenting antifungal activity (chitinases and osmotins/thamatin/permeatins) were analyzed within the CiEST database. Campos *et al.* (this issue) are presenting a complete analysis of citrus PR proteins under biotic and abiotic stress.

The proteins involved in HR, found in the present work, are detailed below:

Metacaspases

Apoptosis in animals is characterized and commonly defined by the activation of a family of cysteine-dependent aspartate-specific proteases, or caspases (Pennell and Lamb, 1997). Caspases can proteolytically activate downstream caspases or cut various cellular substrates, leading to cell death (Utz and Anderson, 2000). Caspase homologues in plants have been reported and called metacaspase (Hoerberichts *et al.*, 2003; Vercammen *et al.*, 2004). In tomato, a metacaspase *LeMCA1*, is induced during infection of leaves with *Botrytis cinerea* (Hoerberichts *et al.*, 2003). Searching within the CiEST, we found 18 reads that match with metacaspases of *A. thaliana*, *L. esculentum* and *O. sativa*, which formed three contigs and four singletons (Table S1). One of these singletons, with similarity of 3e-84 to *L. esculentum*, came from *P. trifoliata* *Citrus tristeza virus*-infected leaf library, which suggests involvement in plant disease resistance, as observed in other species (De Jong *et al.*, 2000; Hoerberichts *et al.*, 2003).

Calcium and Potassium channels

Some of the earliest signaling events detected during the HR include ion fluxes, influx of calcium and H⁺, and efflux of K⁺ (Atkinson *et al.*, 1990).

Calcium, as the divalent cation (Ca²⁺), is an essential plant nutrient. It is required for structural roles in the cell wall and membranes, for inorganic and organic anion uptake in the vacuole, and as an intracellular messenger in the cytosol. Polarization or depolarization of the membrane voltage opens the gate subunits and allows calcium flow through the cell membrane. When calcium ions enter plant cells through Ca²⁺-permeable ion channels, tonoplast

and/or endoplasmic reticulum, cellular responses to a diverse range of developmental cues and environmental challenges are initiated (White, 2000; Sanders *et al.*, 2002).

The search for calcium channel sequences within the CiEST database revealed the presence of 15 reads (Table S1). The clustering identified as potential calcium channels, two contigs and two singletons, contig 1 (C1) was mainly composed of *C. sinensis* sequences with only one sequence from *P. trifoliata* and one occurrence of *C. reticulata*. Contig 2 (C2) was mostly composed of *P. trifoliata* sequences (two reads) and one read from *C. reticulata*. Singletons were found in *C. limonia* and *P. trifoliata*.

Contig 1 showed 75% identity of deduced protein sequence with the Two-pore channel 1 (TPC1) from *Arabidopsis thaliana* (Furuichi *et al.*, 2001). Contig 2 showed 61% amino acid identity with the Two-pore calcium channel TPCB1 from *N. tabacum*. One singleton is 74% identical to the *N. tabacum* TPC and the other is 50% identical to the tobacco TPC1B.

The wheat channel homologue to TPC1 from *A. thaliana* was cloned and expressed under abscisic acid (ABA) and various stress treatments indicating that these channels might be the first gate of control for several cell phenomena related to Ca²⁺ influx as cell signaling, homeostasis and stress response (Wang *et al.*, 2005).

Potassium is the predominant inorganic ion of plant cells, where it plays a major role in contributing to cellular hydrostatic (turgor) pressure, growth, and responses to the environment (Michard *et al.*, 2005).

The *HLM1* gene encodes a member of the CNGC (cyclic nucleotide - gated channel - CNGC4) ion channel family (Maser *et al.*, 2001). The study of this gene product showed that CNGC4 is permeable to both K⁺ and Na⁺, is activated by cGMP and cAMP, and its expression is induced in response to pathogen infection and some pathogen-related signals. So HLM1 may constitute a common downstream component of the signaling pathways leading to HR (Balague *et al.*, 2003). In our analyses, we found only one singleton similar to HLM1 (6e-96), the other 15 singletons and 17 contigs showed similarity to other members of the CNGC family (Table S1), possibly with potassium channel activity. Our comparative analysis of the potassium channel revealed its presence in the *C. reticulata*, *C. sinensis*, *C. latifolia*, *C. aurantifolia* and *P. trifoliata* libraries. These results were expected because potassium is involved in wide environmental responses and is possibly involved in the citrus HR as well.

Reactive oxygen species

The generation of reactive oxygen species (ROS), such as singlet oxygen, superoxide anion radicals, hydrogen peroxide, and hydroxyl radicals is one of the earliest responses during HR just after ion fluxes (Atkinson *et al.*, 1990). These ROS not only have direct antimicrobial activ-

ity (Lamb and Dixon, 1997), but also act as signaling molecules leading to up- or down-regulation of numerous genes involved in plant defense responses such as the induction of defense-related genes and initiation of programmed cell death (PCD) at the site of the attempted infection (Neill *et al.*, 2002; Apel and Hirt, 2004). This localized PCD generates a physical barrier restraining nutrient availability because of the rapid dehydration caused by tissue death (Parker and Coleman, 1997).

Plants have a number of ways of generating extracellular ROS (Bolwell and Wojtaszek, 1997). During the incompatible pathogen interaction, superoxides are produced enzymatically outside the cell and are rapidly converted to H₂O₂, which is able to cross the plasma membrane (Melillo *et al.*, 2006). Cell wall-bound peroxidase (POXs) and a plasma membrane-bound NADPH oxidase are considered the major sources for ROS production as defense mechanisms during biotic stresses (Heath, 2000; Kawano, 2003; Melillo *et al.*, 2006), even though there is evidence supporting a differential ROS generation for different stimuli as well (Bolwell, 1999).

The balance of production and detoxification of ROS is essential for aerobic life (Apel and Hirt, 2004) and the key enzymes or enzyme families involved in the production, as well as scavenging of ROS and induction of HR in plants are nicotinamide-adenine dinucleotide phosphate [NAD(P)H] oxidase, peroxidases and superoxide dismutases. Superoxide radicals generated by the one-electron reduction of molecular oxygen are rapidly converted within the chloroplast to hydrogen peroxide by CuZn-superoxide dismutase (Asada, 1999). The superoxide dismutases (SODs) are extremely effective antioxidant plant enzymes (Skinner and Baek, 2006), since they dismutate two superoxide anions into H₂O₂ and O₂ (Fridovich, 1991).

Three contigs and three singletons harboring NAD(P)H oxidase gene sequences were obtained from the CitEST database. The three contigs comprised 16, 4 and 5 reads, respectively. They were assembled with ESTs from both *C. sinensis* and *P. trifoliata* libraries. However contig 2 (C2) and contig 3 (C3) presented only reads from the leaf libraries, contig 1 (C1), with the largest number of reads, comprised sequences from fruit tissue as well. The three contigs contained partial gene sequences and exhibited similarity with NADPH oxidase protein from either *Nicotiana* sp. or *A. thaliana* as best BLASTX hits (Table S1).

Plants have multiple genes encoding peroxidases (Tognolli *et al.*, 2002; Apel and Hirt, 2004). Therefore, as expected, a large number of peroxidase transcripts were identified, 568 reads total, which clustered in 52 contigs and 42 singletons (Table S1).

Several interesting inferences could be drawn from these sequences. Considering that glutathione S-transferase/oxidase/peroxidase comprises a large number of reads, which are from all of the seven *Citrus* and *Poncirus* species and

also the six tissues available at the CitEST, we can infer that the expression of these transcripts are constitutive not only in the genus *Citrus* but, likely, in other related rutaceae.

However, in a few cases, the specificity of the contigs is evident, with some particular conditions leading to the clustering of similar sequences in well-defined contigs. One of the contigs (C63) was composed of three *C. limonia* sequences, all of them were from the library constructed from roots undergoing water stress, suggesting that this transcript may code for a peroxidase involved in drought response. Interestingly, three contigs (C22, C64 and C65) were composed by reads from a library constructed from *C. sinensis* symptomatic for citrus variegated chlorosis (CVC), caused by *Xylella fastidiosa*, one of the most devastating diseases of the crop, indicating that peroxidases may be involved in the response of citrus to this particular biotic stress.

The multicopy pattern of superoxide dismutase (*SOD*) gene observed in various hosts was also found within the CitEST database, which yielded 19 contigs and 16 singletons with similarity to translated SOD sequences (Table S1).

As in the peroxidases, at least one contig harbored 43 sequences from most of the species and tissues from where the libraries were constructed. Interestingly, no *SOD* gene was found to be expressed in the flower tissue, but that might be due to the relatively low number of reads of this library (4,330) compared to the others (Targon *et al.*, this issue). On the other hand, some contigs comprised reads from one or few species. While three contigs (C3, C18 and C19) were formed of only *Poncirus* sp. reads, the other 11 were formed of only *Citrus* sp. sequences. Within these latter, some harbored sequences from a particular species (two contigs from *C. reticulata* alone, five from *C. sinensis*, and one from *C. latifolia*), suggest high specificity of these transcripts.

Likewise the peroxidases, but on a lower level, it was possible to determine some tissue-specificity of the *SOD* transcripts. At least six contigs (C3, C4, C12, C13, C18, and C19) harbored sequences only from the leaf libraries, while the other three (contigs C7, C9, and C16) were formed solely by reads from the fruit libraries. Furthermore, few contigs comprised reads from a particular condition. One of them (contig C12), of great interest, was composed of 39 reads from both the *C. sinensis* and *C. reticulata* leaf libraries infected with *X. fastidiosa*. This transcript was similar to a Cu/Zn superoxide dismutase from *C. sinensis* (CAA03881.1) and, since the two species are susceptible and resistant to CVC, respectively, it is likely that such transcript is not involved in disease resistance, but in the response to this specific biotic stress.

Nitric oxide

Nitric oxide (NO) has well-known biological functions in mammals, but it was only recently recognized as a

signal compound in plants (Zhao *et al.*, 2005). In fact, it is now known that NO plays an important role in diverse physiological processes in plants (Grun *et al.*, 2006). The Avr factors from pathogens stimulate NO production, which promotes disease resistance in plants, collaborating with ROS in the oxidative burst (Delledonne *et al.*, 1998). The relative level of NO and H₂O₂ appears to be very important to HR-associated cell death in soybean cells (Delledonne *et al.*, 2001) and it was observed that only the simultaneous increase of NO and H₂O₂ in tobacco cells induced typical programmed cell death (de Pinto *et al.*, 2002).

Two potential enzymatic sources of NO in plants are NO synthase (NOS) and nitrate reductase (NR). NOS is a family of well characterized enzymes in mammalian cells (Desikan *et al.*, 2002) that was found in several plants (Neill *et al.*, 2002), and serves as a signal for plant growth, development, and defense (Delledonne *et al.*, 1998; Klessig *et al.*, 2000; Neill *et al.*, 2002). NR is a central enzyme of nitrogen assimilation in plants (Lea, 1999; Kaiser *et al.*, 2002), and is induced in potato tubers treated with either *Phytophthora infestans* or an elicitor derived from this oomycete pathogen (Yamamoto *et al.*, 2003).

Despite the large number of citrus ESTs sequenced in the CitEST Project, we did not find reads similar to NOS, which suggests that the NO synthesis in these plants can be preferentially performed by NR. Within the CitEST database, three contigs and one singleton similar to plant nitrate reductase were assembled (Table S1). One of these contigs (C10), similar to *Petunia x hybrida* and also *Tilia platyphyllos* (e-values = 0.0) nitrate reductase, seems to be expressed especially within non-infected leaf tissue from *C. reticulata* and *P. trifoliata*, while the unique read from bark *P. trifoliata* is from *Phytophthora*-infected library. These results suggest that this contig has species- and tissue-specific expression, showing that the bark tissue of *P. trifoliata* may be involved in the production of NO for plant defense, as with potato tubers infected with *P. infestans* (Yamamoto *et al.*, 2003).

Another contig (C11), similar to NR from *Tilia platyphyllos* (2e-152) presents reads solely from *C. reticulata* and a tendency of expression in *Xylella fastidiosa*-infected leaves (Figure 1a). This specific expression pattern suggests that this protein is important to plant defense specifically in *C. reticulata*. The third contig (C7) did not exhibit a species-specific expression, but comprised mainly sequences from the *C. reticulata* and *C. sinensis* leaf libraries infected with *X. fastidiosa*, indicating that this transcript may be involved in the response to this biotic stress. Figure 2 shows the estimate of a phylogeny among contigs C7, C10 and C11, with nitrate reductase from *Petunia x hybrida* (gi|294113), *Tilia platyphyllos* (gi|24474445) and *Ricinus communis* (gi|11119240). As expected, it was observed that Contigs C10 and C11 are more closely related to each other and are related with *Tilia platyphyllos*

and *Petunia x hybrida*, and C7 is more closely related to *Ricinus communis*.

Phenylpropanoid pathway

In plants, the phenylpropanoid pathway is responsible for the synthesis of a wide variety of secondary metabolic compounds, including lignins, salicylates, coumarins, hydroxycinnamic amides, pigments, and flavonoids (Hwang *et al.*, 2003).

In the plant phenylpropanoid pathway, chorismate mutase catalyses the reaction that forms phenylalanine, and the enzyme phenylalanine ammonia-lyase (PAL) deaminates phenylalanine to yield cinnamic acid (Gozzo, 2003; Kaneko *et al.*, 2003). Several other enzymes are present in the phenylpropanoid pathway, with cinnamate 4-hydroxylase (C4H)-the first cytochrome P450 upstream in this pathway - leading to the synthesis of a variety of compounds including lignin monomers, flavonoids, and furcoumarins (Morant *et al.*, 2003; Gravot *et al.*, 2004). Chalcone synthase (CHS) is an enzyme that catalyses the reaction that leads to the production of flavonoids and phytoalexins (Yang *et al.*, 2002). The final substance produced by this pathway (4,2', 4', 6'-tetrahydroxychalcone) serves as the backbone for the production of a wide array of flavonoid compounds that function in such diverse roles as flower pigmentation, UV protection, signaling, male fertility, and defense against microbial pathogens (Winkel-Shirley, 2001).

Within the CitEST database, we found five contigs and four singletons similar to chorismate mutase (Table S1), some of them showed specific patterns of expression. The contig 4 (C4) that is similar to *A. thaliana* chorismate mutase (accession number AAC73018) showed specific expression in the *C. sinensis* fruit, whereas the contig 6 (C6) (1e-67) (similar to accession number AAM61395) showed expression in *C. reticulata* fruits and in *C. sinensis* *X. fastidiosa*-infected leaves (Figure 1b). The other three contigs did not show an expression pattern, presenting reads from different libraries. Figure 3 shows a phylogeny analysis among contigs C4, C5, C6 and C7, and other chorismate mutase sequences from NCBI, showing coherence between contigs and the best hit in BLAST search (Table S1).

Four contigs and seven singletons were found in the search for phenylalanine ammonia-lyase (PAL) within the CitEST database (Table S1). The contig 10 presents high similarity (2e-107) with PAL from *Nerium oleander* and was expressed solely in *P. trifoliata* *Phytophthora*-infected bark (two reads), suggesting its importance in specific defense of this species. Another contig (C8), comprising sequences with similarity to that from *C. limon*, comprised 20 reads from different libraries. In *C. sinensis*, as in *C. reticulata*, reads showed a specific pattern of expression within *X. fastidiosa*-infected leaf libraries. However, PAL expression was also found in *C. reticulata* within the




Figure 1 - Transformed data representing the relative abundance of EST by library, expressed in 10^4 reads: (a) Nitrate reductase contig 11, (b) chorismate mutase contig 6, (c) chalcone synthase contig 1, (d) isochorismate synthase contig 2. Code of the libraries: CS: *Citrus sinensis*; CR: *Citrus reticulata*; C1: leaf cDNA; C3: fruit cDNA; 100: non-infected material; 101: infected with *Xylella fastidiosa*; 102: 30 days after *X. fastidiosa* infection; 700: stadium 1 of the fruit; 701: stadium 2 of the fruit; 703 stadium 4 of the fruit.




Figure 2 - Cladogram showing phylogeny relation among contigs 7, 10 and 11 with nitrate reductase from *Petunia hybrida* (gi|294113), *Tilia platyphyllos* (gi|24474445) and *Ricinus communis* (gi|11119240).

healthy fruit library. The contigs C3 (65 reads) and C9 (four reads) were similar to PAL from *Citrus clementina* x *Citrus reticulata* and *Jatropha curcas*, respectively, although they were highly similar to PAL (e-values = 0.0), they did not show any specific expression pattern, since they were formed by reads from a variety of libraries.

In a study with Valencia sweet orange, two cinnamic acid 4-hydroxylase (C4H) cDNAs were cloned and showed differential expression regulation (Betz *et al.*, 2001). One




Figure 3 - Cladogram showing phylogeny relation among contigs 4, 5, 6 and 7 with chorismate mutase from *Fagus sylvatica* (ABA54871), *Arabidopsis thaliana* (AAC73018) and *Arabidopsis thaliana* (AAM61395).

of the cDNAs, C4H2, appeared to be expressed constitutively and seemed to play the role of a 'housekeeping' gene in the phenylpropanoid pathway. Conversely, expression studies showed C4H1 to be strongly wound-inducible, but even in wounded tissue it reached much lower levels of mRNA than C4H2 (Betz *et al.*, 2001).

Within the CitEST, we found three contigs and two singletons related to C4H2 (Table S1). One of these contigs (C3) was assembled from 46 reads and showed 100% of identity with C4H2, for which the majority reads originated from *C. sinensis* and *C. reticulata* in different stages of fruit development. Contig 1 (C1) showed 91% of identity with C4H2 and was formed by 11 reads, mostly from *C. sinensis* (10 reads) in the early stages of fruit development and one read from the *P. trifoliata* *Citrus tristeza virus*-infected library. These results suggest a possible pattern of expression of this protein in the fruits of *C. sinensis* and *C. reticulata*. Moreover, a possible involvement in plant defense against pathogen in *P. trifoliata* cannot be excluded.

In contig 2 (C2), we observed 81% of amino acid identity with *Medicago sativa* C4H protein (Table S1), in agreement with its pattern of expression during fruit ripening (Fahrendorf and Dixon, 1993), since the two reads of this contig were originated from the *C. sinensis* fruit libraries. Figure 4 shows the phylogeny pattern among contigs C1, C2 and C3, and cinnamic acid 4-hydroxylase from *Citrus sinensis* (AAF66066) and *Medicago sativa* (gi|166372); coherence can be observed with the best hit in BLAST search and the phylogenetic relationship.

Two cDNA clones encoding chalcone synthase (CHS) were isolated (CitCHS1 and CitCHS2) from citrus (Moriguchi *et al.*, 1999). The accumulation of CitCHS2 mRNA was notably induced by embryogenesis but CitCHS1 mRNA was not. There was no detectable accumulation of flavonoid in the undifferentiated calli, but flavonoid accumulated after the morphological changes to embryoids. These results indicate that two CHS genes differentially expressed during citrus somatic embryogenesis and CitCHS2 may regulate the accumulation of flavonoids in citrus cell cultures (Moriguchi *et al.*, 1999). Within CitEST, we found four contigs and 12 singletons related to chalcone synthase; among them, three contigs and eight singletons related to CitCHS2. Contig 3 (C3) is formed by 75 reads that mostly came from *C. sinensis* fruits in the early stages of development and is 100% identical to the most homologous BLAST hit (*C. sinensis*). Contig 4 (C4) was composed of 54 reads mostly from *C. reticulata* in the early stages of the fruit development and shares 92% identity with *C. sinensis* CHS. Finally, another contig (C1) was formed by nine *C. sinensis* reads and presented a tendency of expression in early stages of *X. fastidiosa*-infected leaves




Figure 4 - Cladogram showing phylogeny relation among contigs 1, 2 and 3 with cinnamic acid 4-hydroxylase from *Citrus sinensis* (AAF66066) and *Medicago sativa* (gi|166372).

(Figure 1c), and was 79% identical to *C. sinensis* CHS from NCBI. The singletons related to CHS came especially from *C. sinensis* and *C. reticulata* fruit libraries. Figure 5 shows the expected phylogeny relationship between these contigs and chalcone synthase from *Citrus sinensis* (BAA81664) and *Betula pendula* (CAA71904).

These results indicate a possible presence of at least two CHS genes in *C. sinensis* that are expressed during fruit development as shown in other species (Aharoni and O'Connell, 2002; Kumar and Ellis, 2003). In addition, other CHSs may be involved in citrus response to pathogens. For more information of flavonoid biosynthesis, see Lucheta *et al.* (this issue).

Salicylic acid

The involvement of salicylic acid (SA) as a signal molecule for plant defense has been extensively studied (Shah and Klessig, 1999; Shah, 2003; Stout *et al.*, 2006). SA quickly accumulates at the site of infection during pathogen attack and plant HR, and it spreads to other parts of the plant to induce a wide range of defense responses (Zhao *et al.*, 2005). Besides inducing PR protein genes, SA is assumed to act before the divergence of the two metabolic branches leading to the phenylpropanoid pathway and the activation of the oxidative burst, respectively. In this way, SA amplifies its own synthesis, which, in addition, is stimulated by H₂O₂ (Gozzo, 2003; Zhao *et al.*, 2005).

Previous work has suggested that plants synthesize SA from phenylalanine utilizing chorismate mutase (Gozzo, 2003). However, evidence showed that, in the chloroplast of *Arabidopsis*, SA is synthesized from isochorismate catalysed by isochorismate synthase. This route, typical of bacteria, seems to be required even in other plants to produce SA for defense against pathogens (Métraux, 2002; Gozzo, 2003).

Although evidence from tobacco and potato would indicate that salicylic acid in plants is derived from phenylalanine via benzoic acid, catalysed by benzoic acid 2-hydroxylase (Yalpani *et al.*, 1993; Coquoz *et al.*, 1998) in citrus SA, formation seems to occur via the isochorismate route since the benzoic acid 2-hydroxylase was not found within the CitEST database.

Analysis within CitEST resulted in two contigs and one singleton similar to isochorismate synthases from *M.*




Figure 5 - Cladogram showing phylogeny relation among contigs 1, 3 and 4 with chalcone synthase from *Citrus sinensis* (BAA81664) and *Betula pendula* (CAA71904).

truncatula, *Capsicum annuum* and *Catharanthus roseus* (Table S1). Only one contig (C2), similar to *M. truncatula* isochorismate synthase, showed a specific pattern of expression, presenting a tendency to be expressed in *C. sinensis* leaves, in the early stage of *X. fastidiosa* infection (Figure 1d). This result may suggest that this contig was formed by a transcript that codes for a protein involved in the production of SA upon plant infection by a pathogen.

Lipoxygenases

Lipoxygenases (LOXs) constitute a family of enzymes that deoxygenate polyunsaturated fatty acids containing *cis*, *cis*-1,4-pentadiene structure. The hydroperoxides produced by the LOX reaction are converted into a series of compounds involved in several physiological processes including growth, fruit ripening and response to biotic and abiotic stresses (Hildebrand *et al.*, 1989; Gardner, 1991; Siedow, 1991; Brash, 1999). The most common substrates for plant LOXs are linoleic and linolenic acids, which are important constituents of plant membrane phospholipids. Plant LOXs are classified according to their positional specificity of linoleic or linolenic acid oxygenation, which occurs either at the C-9 or at the C-13 of the hydrocarbon backbone of the fatty acid. Therefore, these LOXs are often referred to as 9-LOXs or 13-LOXs.

Regarding sequence similarity, plant LOXs can be classified into two gene subfamilies. Enzymes with no transit peptide share a similarity higher than 75% with one another and are designated *type 1*-LOXs. The other subfamily, called *type 2*-LOX, contains a putative chloroplast transit peptide sequence and shares a sequence similarity of approximately 35% with each other (Feussner and Wastermack, 2002). In *C. jambhiri*, a LOX containing typical features of a transit peptide for chloroplastic targeting has been identified and expressed upon wounding and inoculation with nonpathogenic isolates of *Alternaria alternata* (Gomi *et al.*, 2002b).

The analysis of the CitEST database revealed 323 sequences related to lipoxygenases, which formed 16 contigs and 18 singletons. The best hit in a BLASTX search as well as the homologous organism are shown in Table S1. The data available from the CitEST database does not allow us to classify the LOXs found and further studies are necessary to characterize the different citrus LOXs. Moreover, most contigs and singletons were incomplete and therefore a classification of the LOXs encountered, based solely on partial sequence comparison, may be premature. The three largest contigs obtained (C3, C7 and C11) revealed deduced amino acid sequences of 921, 945 and 900 amino acids, respectively. The comparison of these sequences with those of the 8 other LOXs showed high similarity to the citrus LOXs (Table S1) revealed that the sequence of contig 7 presented an exclusive amino acid insertion sequence at po-

sitions 574 to 606. Contig 7 was formed by reads from healthy and *X. fastidiosa*-infected leaves of *C. reticulata* as well as *X. fastidiosa*-infected leaves of *C. sinensis* and showed 73% identity with *lox1* of *Sesbania rostrata*. Contig 3 showed 71% identity with a *lox2* of *Zantedeschia aethiopica* and contig 11 showed 78% identity with a *lox* of *C. jambhiri*. Contigs 3 and 11 represent complete sequences of the *lox* gene.

Some 9-LOXs have been associated to plant resistance and lipid peroxidation, which is involved in membrane damage and HR (Rance *et al.*, 1998; Kolomiets *et al.*, 2000; Jalloul *et al.*, 2002). The biosynthesis of oxygenated unsaturated fatty acids via LOX pathway may be important for plant defense against pathogen attack (Peng *et al.*, 1994). Most of the reads showing similarity to LOXs were obtained from pathogen-infected or fruit libraries. LOXs were highly expressed in the early stages of *X. fastidiosa*-infected libraries of sweet orange (*C. sinensis*) and mandarin (*C. reticulata*) (up to 30 days after inoculation-DAI). Some ESTs associated with lipoxygenase were also observed in *C. reticulata* up to 60 DAI. Several studies have reported that LOX activity in plants is higher and more rapid upon infection of avirulent pathogens as compared to virulent pathogens (Ocampo *et al.*, 1986; Yamamoto and Tani, 1986; Siedow, 1991; Melan *et al.*, 1993). *C. reticulata* is resistant to *X. fastidiosa* and, as expected, a higher expression of LOX was observed in *X. fastidiosa*-infected mandarin plants. LOX was also expressed in *C. sinensis* and *P. trifoliata* libraries infected with *Citrus leprosis virus* (CiLV) and *Citrus tristeza virus* (CTV), respectively. *P. trifoliata* is resistant to CTV and therefore the expression of LOX in this interaction was expected.

Several fruit ESTs associated with LOX were also identified within the CitEST database. The role of LOX in fruit development has been well documented (Griffiths *et al.*, 1999; Chen *et al.*, 2004a). LOXs play an important role in the production of volatile molecules that can influence flavor and aroma of many plant products (Robinson *et al.*, 1995). LOXs can also alter shelf life and organoleptical and nutritional characteristics of plant products (Robinson *et al.*, 1995). These are important features for the citrus industry, since fruit quality is essential for juice production as well as for the commercialization of fresh fruits.

Three *C. limonia* EST sequences related to LOX were also expressed during water stress. Membranes are main targets of degradative processes induced by drought and lipid peroxidation is one of the most important causes of cell deterioration, resulting in changes in the composition of fatty acids that affect the structural and functional properties of cell membranes (Smirnov, 1993; Asada, 1999). The expression of LOXs during drought has been reported (Gigon *et al.*, 2004; Sofu *et al.*, 2004); however, little is known about the relationship between drought stress and LOX activity. Further studies are necessary to better understand the function of citrus LOXs in drought response.

Phospholipases

Phospholipases (PLs) in plants are known to be involved in several cellular responses, including plant growth, development, and responses to stress and defense. PLs play a pivotal role in the plant response via lipids pathway. The superfamily is involved in processes such as cytoskeletal rearrangement, membrane trafficking and degradation (Meijer and Munnik, 2003; Wang, 2004). Several studies have associated PL activity to stress response and defense, and the involvement of PLs and other enzymes in cell death has also been reported (Spiteller, 2003; Wang, 2005).

Plant PLs are classified according to their site of hydrolysis on phospholipids as phospholipase A₁ (PLA₁), phospholipase A₂ (PLA₂), phospholipase C (PLC), and phospholipase D (PLD).

The analysis within the CitEST database revealed sequences related to PLs. The assembling of the reads (E-value < -10) provided 9 contigs and 12 singletons with E-value < -10. The best hit in a BLASTX search as well as the homologous organism are shown in Table S1.

The phospholipase A (PLA) activity increased rapidly and systemically in response to wounding by attack of herbivores in leaves of tomato plants (Narvaez-Vasquez *et al.*, 1999). Multiple secretory-phospholipase A₂ (sPLA₂) genes have been also identified in plants (Lee *et al.*, 2005) and within the CitEST database, four reads were identified with high similarity to PLA₂. The clusterization provided only one contig that was confirmed to be 75% identical to a rice PLA₂ form, and this contig was formed by reads from healthy *C. sinensis* and *C. aurantifolia* libraries. This PLA₂ contig contains twelve conserved cysteine residues and sequences that are likely to represent the Ca(2+)-binding loop and active-site motif, which are characteristic of animal PLA₂s. In the CitEST project, the PLA₂ identified was sequenced from bark tissue, probably indicating a secretory possibility and a defense response in plant to a variety of stimuli including biotic and abiotic stresses (Lee *et al.*, 2005).

Under the control of cell surface receptors, the phosphoinositide-specific PLC isozymes hydrolyze the highly phosphorylated lipid phosphatidylinositol 4,5-bisphosphate [PI(4,5)P₂], generating two intracellular products: inositol 1,4,5-trisphosphate (InsP₃), which is an universal calcium-mobilizing second messenger; and diacylglycerol, which is an activator of protein kinase C (Rebecchi and Pentylala, 2000). Evidence suggests that InsP₃ may be involved in the signal transduction pathway for the development of the HR of *C. lemon* against *A. alternata* (Ortega and Perez, 2001). A search within the CitEST identified reads homologous to PLC genes. Clustering of these reads resulted in five contigs and nine singletons. The similarity among the deduced PLC proteins found ranged from 59% to 73% identity to PLC from *A. thaliana*, *O. sativa*, *Nicotiana rustica*, and *Glycine max*. The most representa-

tive group for PLC was contig 1, which was formed by reads from the libraries, bark and leaves of healthy *C. sinensis* and from *C. sinensis X. fastidiosa*-infected leaves.

PLD mediates the hydrolysis of membrane lipids and diacylglycerol-kinase-mediated phosphorylation of diacylglycerol (Wang, 2002). The expression of several PLD genes has been extensively reported in plant defense induced by the elicitor treatment as well as pathogen infection (Laxalt *et al.*, 2001; De Torres *et al.*, 2002). The PLD group was present in 14 reads clustered into three contigs within CitEST. The PLD genes identified were classified as different isoforms related to the *A. thaliana* (AAP42742.1) PLD protein. The CitEST PLDs were 80 to 83% identical to their orthologues. The largest PLD was also formed by reads from healthy *C. sinensis*.

Analysis of the CitEST database revealed the presence of several citrus PLs; however, since most contigs and singletons are incomplete, it was not possible to unequivocally classify them within the family. Further studies are necessary to better understand the genetic and biochemical heterogeneity of PLs and to relate the function of citrus PLs in HR.

Glutathione transferases and Glutathione peroxidases

Plant tissues contain several ROS scavenging enzymes (like glutathione peroxidase) and detoxifying lipid peroxidation products (like glutathione S-transferases and phospholipid-hydroperoxide glutathione peroxidase) (Chen *et al.*, 2004b), which protect cells from ROS and damage from stress conditions.

By using an *in-silico* analysis of the CitEST database, we have identified 36 contigs and four singletons related to glutathione transferase (GST) genes (Table S1). Taking into consideration that the CitEST database was constructed based on a number of ESTs produced during different stages of the development and challenges, it is obvious that, as seen in other plant species (Sappl *et al.*, 2004), several mechanisms are acting to induce citrus GSTs.

By combining keyword and BLASTX searching within the CitEST database with the sequence from public database that showed the best match with citrus sequences, we found full-length genes with high homology to GST 22 from soybean (McGonigle *et al.*, 2000). Whether this putative gene plays a role in stress tolerance remains to be experimentally demonstrated in plant species, including citrus. In addition, we found full-length copies of LeGST-T3, a GST protein from tomato that was found to protect yeast cells from H₂O₂-induced oxidative stress (Kilili *et al.*, 2004). This indicates that the putative citrus genes could play a role in cell protection from pro-oxidant-induced cell death.

Glutathione peroxidases (GPX) are enzymes that reduce organic hydroperoxides produced during oxidative stress. The over-expression of a GPX inhibited oxidative

stress - induced PCD in tomato (Chen *et al.*, 2004b). It is strongly suggested that specific GPXs, such as Phospholipid Hydroperoxide GPX (PHGPX), play a specific role in ROS; however, their specific role in plants remains to be elucidated (Milla *et al.*, 2003; Chen *et al.*, 2004b). PHGPX functions, in the removal of phospholipid hydroperoxides, are generated as products of lipoxygenase catalysed oxygenation of fatty acids (Ursini *et al.*, 1985).

The analysis of the CitEST database also revealed sequences related to GPXs, which formed 10 contigs and 2 singletons (Table S1). The analysis of these sequences shows GPX and PHGPX similar to the ones found in orange, pea, *Momordica charantia*, chickpea, and *Arabidopsis*.

At least two copies of a gene found in this study code for the PHGPX that was experimentally found to be responsible for salt-stress signaling in citrus (Holland *et al.*, 1993; Avsian-Kretschmer *et al.*, 2004).

As compared to other stress-related genes found in this broad-scale search, an observation that emerged was the proportion of genes that code for GPX in citrus. On the basis of these results, it is suggested that GPX is an enzyme that shows a low number of isoforms and/or copies in the genome of citrus and, probably, plays a role in a condition not strongly induced in this study.

Pathogenesis-related proteins

The induction of PR proteins is triggered stereotypically in plants by infection with fungal, bacterial, and viral pathogens, and is part of a more general response to biotic and abiotic stress (Datta and Muthukrishnan, 1999). PR proteins comprise a group of 17 protein classes (Christensen *et al.*, 2002), most of which present antifungal activity, such as the PR-3, PR-4, PR-8 and PR-11 proteins (also known as chitinases) and the PR-5 proteins (also known as osmotins, permatins or thaumatins).

The chitinases, which are vacuolar PR proteins, act in concert to defend against a specific class of pathogens, fungi (Wirdnam *et al.*, 2004). Moreover, there is also an indication that chitinases may act as lysozymes, by degrading bacterial cell walls (Tiffin, 2004). This class of proteins can be induced by ethylene treatment and in association with HR (Leubner-Metzger and Meins, 1999).

The analysis of the CitEST database revealed several reads related to chitinases, which formed 29 contigs and 24 singletons (Table S1). The analysis of these sequences shows various acidic classes I and II chitinases that match with citrus libraries from elsewhere. In addition, we found a number of sequences that relate to chitinase, in general, from other plant species such as *A. thaliana*, *Tulipa bakeri*, *Hevea brasiliensis*, *Gossypium hirsutum*, and *N. tabacum*. In rough lemon (*C. jambhiri*), the expression of the two acidic chitinases (I and II) was not constitutive and accumulated in leaves two hours after induction by wounding or inoculation with non-pathogenic isolates of *A. alternata*

(Gomi *et al.*, 2002a). Our results reveal that a number of acidic class I and II chitinases are expressed in citrus. However, it remains unclear whether the citrus chitinases found in our study refer to isoforms of the same gene.

The PR-5 proteins are expressed in response to abscisic acid, ethylene, auxin, salinity, lack of water, cold, UV light, wounding, viral, fungal and oomycetal infection (Nelson *et al.*, 1992; Zhu *et al.*, 1995; Sato *et al.*, 1996). Within the CitEST database, we found 63 unique reads related to PR-5-like proteins, clustered into 10 contigs and four singletons (Table S1). These singletons encode a 219-amino acid ORF homologous to a sunflower (*H. annuus*) PR-5-like protein, a 190-amino acid ORF homologous to a turnip (*B. rapa*) PR-5-like protein, as well as a 182- and a 280-amino acid ORF, both homologous to a tobacco (*N. tabacum*) PR-5-like protein (Table S1). These singletons were originated from fruit libraries, from CTV-infected leaf libraries or from *X. fastidiosa*-infected leaf libraries, either from *C. reticulata*, *C. sinensis* or from *P. trifoliata* origin. The 10 PR-5-like contigs were assembled from two up to 13 reads each, mainly from the fruit libraries, but also from seed, bark, and flower libraries. Despite the fact that PR-5 proteins are expressed also in response to biotic stress, there were only five reads composing contigs which were originated from leaves infected either with *X. fastidiosa* or with CTV. Moreover, the majority of the PR-5-like reads forming contigs were from *C. sinensis* origin, secondarily from *C. reticulata* or *P. trifoliata* and seldomly from *C. aurantifolia* or *C. aurantium*. Two contigs homologous to tomato PR-5-like proteins, one homologous to an *Arabidopsis* PR-5-like protein (accession NP_173261), one homologous to a cultivated apple PR-5-like protein and one homologous to a grape PR-5-like protein presented complete coding sequences when aligned with the best hit in BLASTX search. As expected, a CitEST contig homologous to a *C. sinensis* PR-5-like protein (accession AAC02549; Table S1) presented high amino acid identity (97%). The 14 PR-5-like clusters found within the CitEST database, *i.e.*, 10 contigs and the four singletons, were considerably similar (from 73% to 97% similarity) to their corresponding most homologous amino acid sequences (Table S1).

To conclude, the CitEST database is considerably representative of citrus PR-5-like sequences, expressed in various biological situations, tissues and citrus species. Unexpectedly, within the CitEST database, no PR-5-like sequences associated with drought stress or with infection by *Phytophthora parasitica* were found. Surprisingly, there were only seven reads found, out of 63 reads in total, which originated either from *X. fastidiosa* infected-leaves or from CTV infected-leaves. Although most of the PR-5-like reads found within the CitEST database are associated with abiotic stress and plant development processes, these sequences potentially encode PR-5 proteins involved also in citrus HR. Therefore, the possible role of the CitEST PR-

5-like clusters in HR must be demonstrated experimentally. Further investigation is also necessary to clone complete citrus PR-5-like sequences, to validate their biological function and to better characterize their expression patterns under biotic and abiotic stress conditions.

Concluding Remarks

Despite a large number of potential pathogens that attempt to infect cultivated plants, disease is not the rule. Plants have an arsenal of mechanisms to defend themselves against pathogen invasion. One of these strategies is the expression of HR genes, elicited when resistant hosts are infected by incompatible races of a pathogen. It is characterized by localized cell death at the site of attempted infection, generating a physical barrier that restricts nutrient availability to the pathogen and prevents the dispersion of the pathogen to uninfected tissues. HR has been investigated extensively, and the increase of genomic sequences available has made it possible to indicate the existence of homologous genes in all plant species by *in silico* analysis. Moreover, the use of tools to search for sequence similarities at the amino acid level can reveal the overall patterns of gene network related to HR responses. In the present work, we tried to uncover the pathways that may lead to HR in citrus, based on the CitEST database. We found putative deduced proteins that can be important for HR and thus we have shown the putative pathways involved in this process. However, further studies are necessary to confirm the functions of these proteins and to better understand the genetic and biochemical network in citrus HR. To our knowledge, this is the first genome study that attempts to identify genes involved in HR in citrus plants. Functional characterization of them may contribute for the development of molecular tools to overcome economically important citrus diseases.

Acknowledgments

The authors would like to thank CNPq/ Millennium Institute (62.0054/01-8) for financially supporting this work and FAPESP for financial support of S. G.-G. (05/58117-4).

References

Aharoni A and O'Connell AP (2002) Gene expression analysis of strawberry achene and receptacle maturation using DNA microarrays. *J Exp Bot* 53:2073-2087.

Apel K and Hirt H (2004) Reactive oxygen species: Metabolism, oxidative stress, and signal transduction. *Annu Rev Plant Biol* 55:373-379.

Asada K (1999) The water-water cycle in chloroplasts: Scavenging of active oxygens and dissipation of excess photons. *Annu Rev Plant Physiol Plant Mol Biol* 50:601-639.

Atkinson MM, Keppler LD, Orlandi EW, Baker CJ and Mischke CF (1990) Involvement of plasma membrane calcium influx in bacterial induction of the K^+/H^+ and hypersensitive responses in tobacco. *Plant Physiol* 92:215-221.

Avsian-Kretchmer O, Gueta-Dahan Y, Lev-Yadun S, Gollop R and Ben-Hayyim G (2004) The salt-stress signal transduction pathway that activates the *gpx1* promoter is mediated by intracellular H_2O_2 , different from the pathway induced by extracellular H_2O_2 . *Plant Physiol* 135:1685-1696.

Balague C, Lin B, Alcon C, Flottes G, Malmstrom S, Kohler C, Neuhaus G, Pelletier G, Gaymard F and Roby D (2003) HLM1, an essential signaling component in the hypersensitive response, is a member of the cyclic nucleotide-gated channel ion channel family. *Plant Cell* 15:365-379.

Betz C, McCollum TG and Mayer RT (2001) Differential expression of two cinnamate 4-hydroxylase genes in 'Valencia' orange (*Citrus sinensis* Osbeck). *Plant Mol Biol* 46:741-748.

Bolwell GP (1999) Role of active oxygen species and NO in plant defence responses. *Curr Op Plant Biol* 2:287-294.

Bolwell GP and Wojtaszek P (1997) Mechanisms for the generation of reactive oxygen species in plant defence: A broad perspective. *Physiol Mol Plant Path* 51:347-366.

Brash AR (1999) Lipoxygenases: Occurrence, functions, catalysis and acquisition of substrate. *J Biol Chem* 274:23679-23682.

Chen G, Hackett R, Walker D, Taylor A, Lin Z and Grierson D (2004a) Identification of a specific isoform of tomato lipoxygenase (TomloxC) involved in the generation of fatty acid-derived flavor compounds. *Plant Physiol* 136:2641-2651.

Chen S, Vaghchhipawala Z, Li W, Asard H and Dickman MB (2004b) Tomato phospholipid hydroperoxide glutathione peroxidase inhibits cell death induced by bax and oxidative stresses in yeast and plants. *Plant Physiol* 135:1630-1641.

Christensen AB, Cho BHO, Næsby M, Gregersen PL, Brandt J, Madriz-Ordeñana K, Collinge DB and Thordal-Christensen H (2002) The molecular characterization of two barley proteins establishes the novel PR-17 family of pathogenesis-related proteins. *Mol Plant Pathol* 3:135-144.

Coquoz JL, Buchala A and Metraux JP (1998) The biosynthesis of salicylic acid in potato plants. *Plant Physiol* 117:1095-1101.

Dangl JL and Jones JD (2001) Plant pathogens and integrated defence responses to infection. *Nature* 411:826-833.

Datta SK and Muthukrishnan S (1999) Pathogenesis-Related Proteins in Plants. CRC Press LLC, Boca Raton, 290 pp.

De Jong AJ, Hoeberichts FA, Yakimova ET, Maximova E and Woltering EJ (2000) Chemical-induced apoptotic cell death in tomato cells: Involvement of caspase-like proteases. *Planta* 211:656-662.

Delledonne M, Xia Y, Dixon RA and Lamb C (1998) Nitric oxide functions as a signal in plant disease resistance. *Nature* 394:585-588.

Delledonne M, Zeier J, Marocco A and Lamb C (2001) Signal interactions between nitric oxide and reactive oxygen intermediates in the plant hypersensitive disease resistance response. *Proc Natl Acad Sci USA* 98:13454-13459.

de Pinto M, Tommasi F and De Gara L (2002) Changes in the antioxidant systems as part of the signaling pathway responsible for the programmed cell death activated by nitric oxide and reactive oxygen species in tobacco bright-yellow 2 cells. *Plant Physiol* 130:698-708.

Desikan R, Griffiths R, Hancock J and Neill S (2002) A new role for an old enzyme: Nitrate reductase-mediated nitric oxide generation is required for abscisic acid-induced stomatal

- closure in *Arabidopsis thaliana*. Proc Natl Acad Sci USA 99:16314-16318.
- De Torres Z, Fernandez-Delmond I, Niittyla T, Sanchez P and Grant M (2002) Differential expression of genes encoding *Arabidopsis* phospholipases after challenge with virulent or avirulent *Pseudomonas* isolates. Mol Plant Microbe Interact 15:808-816.
- Fahrendorf T and Dixon RA (1993) Stress responses in alfalfa (*Medicago sativa* L.). XVIII: Molecular cloning and expression of the elicitor-inducible cinnamic acid 4-hydroxylase cytochrome P450. Arch Biochem Biophys 305:509-515.
- Feussner I and Wasternack C (2002) The lipoxygenase pathway. Annu Rev Plant Biol 53:275-97.
- Forment J, Gadea J, Huerta L, Abizanda L, Agusti J, Alamar S, Alos E, Andres F, Arribas R, Beltran J *et al.* (2005) Development of a citrus genome-wide EST collection and cDNA microarray as resources for genomic studies. Plant Mol Biol 57:375-91.
- Fridovich I (1991) Molecular oxygen; friend and foe. In: Pell EJ, Steffen KL (eds) Active Oxygen/Oxidative Stress and Plant Metabolism. Amer Soc Plant Physiol, Rockville, pp 1-5.
- Furuichi T, Cunningham KW and Muto S (2001) A putative two pore channel AtTPC1 mediates Ca²⁺ flux in *Arabidopsis* leaf cells. Plant Cell Physiol 42:900-905.
- Gabriel DW and Rolfe BG (1990) Working models of specific recognition in plant - Microbe interactions. Annu Rev Phytopath 28:365-391.
- Gardner HW (1991) Recent investigations into the lipoxygenase pathway of plants. Biochim Biophys Acta 1084:221-239.
- Gigon A, Matos AR, Laffray D, Zuily-Fodil Y and Pham-Thi AT (2004) Effect of drought stress on lipid metabolism in the leaves of *Arabidopsis thaliana* (ecotype Columbia). Ann Bot 94:345-51.
- Gomi K, Itoh N, Yamamoto H and Akimitsu K (2002a) Characterization and functional analysis of class I and II acidic chitinase cDNA from rough lemon. J Gen Plant Pathol 68:191-199.
- Gomi K, Yamamoto H and Akimitsu K (2002b) Characterization of a lipoxygenase gene in rough lemon induced by *Alternaria alternata*. J Gen Plant Pathol 68:21-30.
- Gozzo F (2003) Systemic acquired resistance in crop protection: From nature to a chemical approach. J Agric Food Chem 51:4487-4503.
- Gravot A, Larbat R, Hehn A, Lievre K, Gontier E, Goergen JL and Bourgaud F (2004) Cinnamic acid 4-hydroxylase mechanism-based inactivation by psoralen derivatives: Cloning and characterization of a C4H from a psoralen producing plant-*Ruta graveolens*-exhibiting low sensitivity to psoralen inactivation. Arch Biochem Biophys 422:71-80.
- Griffiths A, Barry C, Alpuche-Soli AG and Grierson D (1999) Ethylene and developmental signals regulate expression of lipoxygenase genes during tomato fruit ripening. J Exp Bot 50:793-798.
- Grun S, Lindermayr C, Sell S and Durner J (2006) Nitric oxide and gene regulation in plants. J Exp Bot 57:507-516.
- Heath MC (2000) Hypersensitive response-related death. Plant Mol Biol 44:321-334.
- Hildebrand DF, Hamilton-Kemp TR, Legg CS and Bookjans G (1989) Plant lipoxygenases: Occurrence, properties and possible functions. Curr Top Plant Biochem Physiol 7:201-219.
- Hoerberichts FA, ten Have A and Woltering EJ (2003) A tomato metacaspase gene is upregulated during programmed cell death in *Botrytis cinerea*-infected leaves. Planta 217:517-522.
- Holland D, Ben-Hayyim G, Faltin Z, Camoin L, Strosberg AD and Eshdat Y (1993) Molecular characterization of salt-stress-associated protein in citrus: Protein and cDNA sequence homology to mammalian glutathione peroxidases. Plant Mol Biol 21:923-927.
- Huang X and Madan A (1999) CAP3: A DNA sequence assembly program. Genome Res 9:868-877.
- Hwang EI, Kaneko M, Ohnishi Y and Horinouchi S (2003) Production of plant-specific flavanones by *Escherichia coli* containing an artificial gene cluster. Appl Environ Microbiol 69:2699-2706.
- Jalloul A, Montillet JL, Assigbetsé K, Agnel JP, Delannoy E, Triantaphylides C, Daniel JF, Marmey P, Geiger JP and Nicole M (2002) Lipid peroxidation in cotton: *Xanthomonas* interactions and the role of lipoxygenases during the hypersensitive reaction. Plant J 32:1-12.
- Kaiser WM, Weiner H, Kandlbinder A, Tsai C-B, Rockel P, Sonoda M and Planchet E (2002) Modulation of nitrate reductase: Some new insights, an unusual case and a potentially important side reaction. J Exp Bot 53:875-882.
- Kaneko M, Hwang EI, Ohnishi Y and Horinouchi S (2003) Heterologous production of flavanones in *Escherichia coli*: Potential for combinatorial biosynthesis of flavonoids in bacteria. J Ind Microbiol Biotechnol 30:456-461.
- Kawano T (2003) Roles of the reactive oxygen species-generating peroxidase reactions in plant defense and growth induction. Plant Cell Rep 21:829-837.
- Kilili KG, Atanassova N, Vardanyan A, Clatot N, Al-Sabarna K, Kanellopoulos PN, Makris AM and Kampranis SC (2004) Differential roles of Tau class glutathione S-transferases in oxidative stress. J Biol Chem 279:24540-24551.
- Klessig DF, Durner J, Noad R, Navarre DA, Wendehenne D, Kumar D, Zhou JM, Shah J, Zhang S, Kachroo P, *et al.* (2000) Nitric oxide and salicylic acid signaling in plant defenses. Proc Natl Acad Sci USA 97:8849-8855.
- Kolomiets MV, Chen H, Gladon RJ, Braun EJ, and Hannapel DJ (2000) A leaf lipoxygenase of potato induced specifically by pathogen infection. Plant Physiol 124:1121-1130.
- Kumar A and Ellis BE (2003) A family of polyketide synthase genes expressed in ripening *Rubus* fruits. Phytochemistry 62:513-526.
- Lam E, Kato N and Lawton M (2001) Programmed cell death, mitochondria and the plant hypersensitive response. Nature 411:848-853.
- Lamb C and Dixon RA (1997) The oxidative burst in plant disease resistance. Annu Rev Plant Physiol Plant Mol Biol 48:251-275.
- Laxalt AM, ter Riet B, Verdonk JC, Parigi L, Tameling WI, Vossen J, Haring M, Musgrave A and Munnik T (2001) Characterization of five tomato phospholipase D cDNAs: Rapid and specific expression of LePLDbeta1 on elicitation with xylanase. Plant J 26:237-247.
- Lea PJ (1999) Nitrogen metabolism. In: Lea PJ and Leegood RC (eds) Plant Biochemistry and Molecular Biology. 2nd edition. Wiley, Chichester, pp 163-192.

- Lee HY, Bahnb SC, Shinb JS, Hwanga I, Backc K, Doellingd JH and Ryu SB (2005) Multiple forms of secretory phospholipase A2 in plants. *Prog Lipid Res* 44:52-67.
- Lee SC and Hwang BK (2005) Induction of some defense-related genes and oxidative burst is required for the establishment of systemic acquired resistance in *Capsicum annuum*. *Planta* 221:790-800.
- Leubner-Metzger G and Meins FJr (1999) Function and regulation of plant b-1,3-glucanases (PR-2). In: Datta SK and Muthukrishnan S (eds) *Pathogenesis Related Proteins in Plants*, CRC Press, Boca Raton, pp 49-76.
- Maser P, Thomine S, Schroeder JI, Ward JM, Hirschi K, Sze H, Talke IN, Amtmann A, Maathuis FJM, Sanders D, *et al.* (2001) Phylogenetic relationships within cation transporter families of *Arabidopsis*. *Plant Physiol* 126:1646-1667.
- McGonigle B, Keeler SJ, Lau SM, Koeppel MK and O'Keefe DP (2000) A genomics approach to the comprehensive analysis of the glutathione S-transferase gene family in soybean and maize. *Plant Physiol* 124:1105-1120.
- Meijer HJ and Munnik T (2003) Phospholipid-based signaling in plants. *Annu Rev Plant Biol* 54:265-306.
- Melan MA, Dong X, Endara ME, Davis KR, Ausubel FM and Peterman TK (1993) An *Arabidopsis thaliana* lipoxygenase gene can be induced by pathogens, abscisic acid, and methyl jasmonate. *Plant Physiol* 101:441-50.
- Melillo MT, Leonetti P, Bongiovanni M, Castagnone-Sereno P and Bleve-Zacheo T (2006) Modulation of reactive oxygen species activities and H₂O₂ accumulation during compatible and incompatible tomato - Root-knot nematode interactions. *New Phytol* 170:501-512.
- Métraux J-P (2002) Recent breakthroughs in the study of salicylic acid biosynthesis. *Trends Plant Sci* 7:332-334.
- Michard E, Lacombe B, Poree F, Mueller-Roeber B, Sentenac H, Thibaud JB and Dreyer I (2005) A unique voltage sensor sensitizes the potassium channel AKT2 to phosphoregulation. *J Gen Physiol* 126:605-617.
- Milla MAR, Maurer A, Rodriguez Huete A and Gustafson JP (2003) Glutathione peroxidase genes in *Arabidopsis* are ubiquitous and regulated by abiotic stresses through diverse signaling pathways. *Plant J* 36:602-615.
- Montillet JL, Chamnongpol S, Rusterucci C, Dat J, van de Cotte B, Agnel JP, Battesti C, Inze D, Van Breusegem F and Triantaphylides C (2005) Fatty acid hydroperoxides and H₂O₂ in the execution of hypersensitive cell death in tobacco leaves. *Plant Physiol* 138:1516-1526.
- Morant M, Bak S, Moller BL and Werck-Reichhart D (2003) Plant cytochromes P450: Tools for pharmacology, plant protection and phytoremediation. *Curr Opin Biotechnol* 14:151-162.
- Moriguchi T, Kita M, Tomono Y, EndoInagaki T and Omura M (1999) One type of chalcone synthase gene expressed during embryogenesis regulates the flavonoid accumulation in citrus cell cultures. *Plant Cell Physiol* 40:651-655.
- Narvaez-Vasquez J, Florin-Christensen J and Ryan CA (1999) Positional specificity of a phospholipase A activity induced by wounding, systemin, and oligosaccharide elicitors in tomato leaves. *Plant Cell* 11:2249-2260.
- Neill SJ, Desikan R, Clarke A, Hurst RD and Hancock JT (2002) Hydrogen peroxide and nitric oxide as signaling molecules in plants. *J Exp Bot* 53:1237-1247.
- Nelson DE, Raghothama KG, Singh NK, Hasegawa PM and Bressan RA (1992) Analysis of structure and transcriptional activation of an osmotin gene. *Plant Mol Biol* 19:577-588.
- Ocampo CA, Moerschbacher B and Grambow HJ (1986) Increase lipoxygenase activity is involved in the hypersensitive response of wheat leaf cells infected with avirulent rust fungi or treated with fungal elicitor. *Z Naturforsch* 41c:558-563.
- Ortega X and Perez LM (2001) Participation of the phosphoinositide metabolism in the hypersensitive response of *Citrus limon* against *Alternaria alternata*. *Biol Res* 34:43-50.
- Parker JE and Colemann MJ (1997) Molecular intimacy between proteins specifying plant-pathogen recognition. *Trends Biochem Sci* 22:291-296.
- Peng YL, Shirano Y, Ohta H, Hibino T, Tanaka K and Shibata D (1994) A novel lipoxygenase from rice. Primary structure and specific expression upon incompatible infection with rice blast fungus. *J Biol Chem* 269:3755-3761.
- Pennell RI and Lamb C (1997) Programmed cell death in plants. *Plant Cell* 9:1157-1168.
- Pontier D, Mittler R and Lam E (2002) Mechanism of cell death and disease resistance induction by transgenic expression of bacterio-opsin. *Plant J* 30:499-509.
- Rance I, Fournier J and Esquerre-Tugaye MT (1998) The incompatible interaction between *Phytophthora parasitica* var. *nicotianae* race O and tobacco is suppressed in transgenic plants expressing antisense lipoxygenase sequences. *Proc Natl Acad Sci USA* 95:6554-6559.
- Rebecchi MJ and Pentylala SN (2000) Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol Rev* 80:1291-1335.
- Rentel MC, Lecourieux D, Ouaked F, Usher S, Petersen L, Okamoto H, Knight H, Peck SC, Grierson CS, Hirt H, *et al.* (2004) OXII kinase is necessary for oxidative burst-mediated signaling in *Arabidopsis*. *Nature* 427:858-861.
- Robinson DS, Wu Z, Domoney C and Casey R (1995) Lipoxygenase and quality of foods. *Food Chem* 54:33-43.
- Sanders D, Pelloux J, Brownlee C and Harper JF (2002) Calcium at crossroads of signaling. *Plant Cell* 14 (supplement):S401-S417.
- Sappl PG, Onate-Sanchez L, Singh KB and Millar AH (2004) Proteomic analysis of glutathione S-transferases of *Arabidopsis thaliana* reveals differential salicylic acid-induced expression of the plant-specific phi and tau classes. *Plant Mol Biol* 54:205-219.
- Sato F, Kitajima S, Koyama T and Yamada Y (1996) Ethylene-induced gene expression of osmotin-like protein, a neutral isoform of tobacco PR-5, is mediated by the AGCCGCC *cis*-sequence. *Plant Cell Physiol* 37:249-255.
- Scheel D (1998) Resistance response physiology and signal transduction. *Curr Opin Plant Biol* 1:305-310.
- Shah J (2003) The salicylic acid loop in plant defense. *Curr Opin Plant Biol* 6:365-371.
- Shah J and Klessig DF (1999) Salicylic acid: Signal perception and transduction. In: Libbenga K, Hall M and Hooykaas PJJ (eds) *Biochemistry and Molecular Biology of Plant Hormones*. Elsevier, London, v 33, pp 513-541.
- Siedow JN (1991) Plant lipoxygenases: Structure and function. *Annu Rev Plant Physiol Plant Mol Biol* 42:145-188.
- Skinner DZ and Baek K-H (2006) Differential expression of manganese superoxide dismutase sequence variants in near iso-

- genic lines of wheat during cold acclimation. *Plant Cell Rep* 25:223-230.
- Smirnoff N (1993) The role of active oxygen in the response to water deficit and desiccation. *New Phytol* 125:27-58.
- Sofa A, Dichio B, Xiloyannis C and Masia A (2004) Lipoxygenase activity and proline accumulation in leaves and roots of olive trees in response to drought stress. *Physiol Plant* 121:58-65.
- Spiteller G (2003) The relationship between changes in the cell wall, lipid peroxidation, proliferation, senescence and cell death. *Physiol Plant* 119:5-155.
- Stout MJ, Thaler JS and Thomma BPHJ (2006) Plant-mediated interactions between pathogenic microorganisms and herbivorous arthropods. *Annu Rev Entomol* 51:663-689.
- Tiffin P (2004) Comparative evolutionary histories of chitinase genes in the genus *Zea* and family Poaceae. *Genetics* 167:1331-1340.
- Tognolli M, Penel C, Greppin H and Simon P (2002) Analysis and expression of the class III peroxidase large gene family in *Arabidopsis thaliana*. *Gene* 288:129-138.
- Ursini F, Maiorino M and Gregolin C (1985) The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochim Biophys Acta* 839:62-70.
- Utz PJ and Anderson P (2000) Life and death decisions: Regulation of apoptosis by proteolysis of signaling molecules. *Cell Death Differ* 7:589-602.
- Vercammen D, van de Cotte B, De Jaeger G, Eeckhout D, Cas-teels P, Vandepoele K, Vandenberghe I, Van Beeumen J, Inze D and Van Breusegem F (2004) Type II metacaspases Atmc4 and Atmc9 of *Arabidopsis thaliana* cleave substrates after arginine and lysine. *J Biol Chem* 279:45329-45336.
- Wang X (2002) Phospholipase D in hormonal and stress signaling. *Curr Opin Plant Biol* 5:408-414.
- Wang X (2004) Lipid signaling. *Curr Opin Plant Biol* 7:329-336.
- Wang X (2005) Regulatory functions of phospholipase D and phosphatidic acid in plant growth, development, and stress responses. *Plant Physiol* 139:566-573.
- Wang YJ, Yu JN, Chen T, Zhang ZG, Hao YJ, Zhang JS and Chen SY (2005) Functional analysis of a putative Ca²⁺ channel gene *TaTPC1* from wheat. *Jour Exp Bot* 56:3051-3060.
- White PJ (2000) Calcium channels in higher plants. *Biochim Biophys Acta* 1465:171-189.
- Winkel-Shirley B (2001) Flavonoid biosynthesis. A colorful model for genetics, biochemistry, cell biology, and biotechnology. *Plant Physiol* 126:485-493.
- Wirdnam C, Motoyama A, Arn-Bouidoires E, van Eeden S, Iglesias A and Meins F Jr (2004) Altered expression of an ankyrin-repeat protein results in leaf abnormalities, necrotic lesions, and the elaboration of a systemic signal. *Plant Mol Biol* 56:717-730.
- Yalpani N, Leon J, Lawton MA and Raskin I (1993) Pathway of salicylic acid biosynthesis in healthy and virus-inoculated tobacco. *Plant Physiol* 103:315-321.
- Yamamoto A, Katou S, Yoshioka H, Doke N and Kawakita H (2003) Nitrate reductase, a nitric oxide-producing enzyme: Induction by pathogen signals. *J Gen Plant Pathol* 69:218-229.
- Yamamoto H and Tani T (1986) Possible involvement of lipoxygenase in the mechanism of resistance of oats *Avena sativa* to *Puccinia coronata* pv. *avenae*. *J Phytopathol* 116:329-337.
- Yang J, Huang J, Gu H, Zhong Y and Yang Z (2002) Duplication and adaptive evolution of the chalcone synthase genes of *Dendranthema* (Asteraceae). *Mol Biol Evol* 19:1752-1759.
- Zhao J, Davis LC and Verpoorte R (2005) Elicitor signal transduction leading to production of plant secondary metabolites. *Biotechnol Adv* 23:283-333.
- Zhu B, Chen THH and Li PH (1995) Activation of two osmotin-like protein genes by abiotic stimuli and fungal pathogen in transgenic potato plants. *Plant Physiol* 108:929-937.

Internet Resources

- CitEST (Citrus ESTs database), <http://biotecnologia.centrodecitricultura.br> (June 25, 2006).
- ClustalW program, <http://www.ebi.ac.uk/clustalw/> (November 13, 2006).
- NCBI, <http://www.ncbi.nlm.nih.gov/> (November 6, 2006).

Supplementary Material

The following online material is available for this article:

Table S1

This material is available as part of the online article from <http://www.scielo.br/gmb>.

Associate Editor: Raquel Luciana Boscarol-Camargo

Table S1 - Description of citrus ESTs encoding proteins involved in HR.

Gene product and organism	Accession Number	Best <i>e</i> -value	Best Similarity	Number of reads	Number Contigs*	Number Singletons
Calcium channels						
two-pore calcium channel[A. <i>thaliana</i>]	gi14349162	0.0	371/432	10	1 (C1)	0
two-pore calcium channel [N. <i>tabacum</i>]	gi46275794	9e-52	133/170	5	1 (C2)	2
Chalcone synthase						
chalcone synthase [<i>C. sinensis</i>]	BAA81664	0.0	391/391	146	3 (C1; C3; C4)	8
chalcone synthase [<i>Betula pendula</i>]	CAA71904	2e-40	89/108	3	1	1
chalcone synthase [<i>Aubrieta deltoidea</i>]	AAG43406	1e-25	86/146	1	0	1
chalcone synthase 3 [<i>Ruta graveolens</i>]	Q9FSB7	3e-17	106/239	1	0	1
chalcone synthase [<i>Chrysanthemum indicum</i>]	AAM46963	7e-19	43/46	1	0	1
Chitinase						
Acidic class II chitinase [<i>C. jambhiri</i>]	BAC20285	e-169	282/293	185	6	11
Acidic class I chitinase [<i>C. jambhiri</i>]	BAC20284	e-177	289/300	154	4	6
Bulb chitinase-1 [<i>Tulipa bakeri</i>]	BAA88408	e-110	230/279	7	1	0

Chitinase [<i>C. sinensis</i>]	CAA93847	e-178	292/292	64	3	1
<u>Chitinase [<i>Hevea brasiliensis</i>]</u>	CAA09110	e-139	269/302	6	1	0
Chitinase CHI1 [<i>C. sinensis</i>]	AAC35981	4e-75	166/232	13	4	1
Chitinase-like protein [<i>G.hirsutum</i>]	AAQ56598	e-164	291/322	18	3	2
Chitinase/lysozyme [<i>N.tabacum</i>]	CAA55128	4e-48	126/195	4	1	2
Putative chitinase [<i>A. thaliana</i>]	CAA19698	9e-45	145/220	3	1	1
Putative class I chitinase [<i>A. thaliana</i>]	AAG48821	e-149	277/321	115	3	0
Receptor-like kinase CHRK1 [<i>N. tabacum</i>]	AAD52097	6e-76	299/484	9	2	0

Chorismate mutase

putative chorismate mutase [<i>Fagus sylvatica</i>]	ABA54871	8e-122	265/326	7	1 (C5)	0
putative chorismate mutase [<i>A. thaliana</i>]	AAC73018	5e-100	219/290	4	1 (C4)	1
chorismate mutase ATCM2 [<i>A. thaliana</i>].	NP_196648	4e-65	145/182	2	1	0
chorimate mutase [<i>L. esculentum</i>]	AAD48923	4e-15	49/60	1	0	1
chorismate mutase-T [<i>A. thaliana</i>]	BAB10786	1e-35	97/137	1	0	1
chorismate mutase [<i>A. thaliana</i>]	AAM61395	3e-107	270/370	10	2 (C6; C7)	1

Cinnamic acid 4-hydroxylase

cinnamate 4-hydroxylase CYP73 [<i>C. sinensis</i>]	AAF66066	0.0	485/485	59	2 (C1; C3)	2
Cinnamic acid 4-hydroxylase [<i>M.</i>]	P37114	4e-74	108/121	2	1 (C2)	0

sativa]

Glutathione peroxidases

Phospholipid hydroperoxide glutathione peroxidase [<i>C. sinensis</i>]	CAE46896	1e-91	167/167	13	1	0
CIT-SAP GPX4_CITS Probable phospholipid hydroperoxide glutathione peroxidase (PHGPx) (Salt-associated protein) [<i>C. sinensis</i>]	CAA47018	9e-92	167/167	49	1	2
Phospholipid glutathione peroxidase [<i>Pisum sativum</i>]	CAA04142	2e-89	193/246	28	2	0
Phospholipid hydroperoxide glutathione peroxidase [<i>Momordica charantia</i>]	AAL40914	4e-76	148/164	14	2	0
Putative phospholipid hydroperoxide glutathione peroxidase [<i>Cicer arietinum</i>]	CAD31839	8e-24	83/137	3	1	0
Glutathione peroxidase, putative [<i>A. thaliana</i>]	NP_564813	2e-69	139/153	5	1	0
Putative glutathione peroxidase [<i>A. thaliana</i>]	AAM67012	5e-68	145/169	8	1	0
Glutathione peroxidase, putative [<i>A. thaliana</i>]	NP_194915	6e-86	168/184	6	1	0

Glutathione transferases

Glutathione S-transferase GST 22 [<i>G. max</i>]	AAG34812	3e-89	181/209	51	6	1
Glutathione S-transferase GST 14 [<i>G. max</i>]	AAG34804	2e-66	145/220	29	3	1
Glutathione S-transferase GST 23 [<i>G. max</i>]	AAG34813	5e-35	79/85	2	1	0
Glutathione S-transferase T3 LeGST-T3 [<i>L. esculentum</i>]	AAG16758	4e-82	173/223	29	5	1
Glutathione S-transferase [<i>Pisum sativum</i>]	BAC81649	8e-97	196/236	10	3	0
Glutathione transferase [<i>Carica papaya</i>]	CAA04391	2e-82	159/180	6	2	0
Glutathione S-transferase [<i>Euphorbia esula</i>]	AAF65767	3e-79	166/212	45	2	0
Glutathione S-transferase [<i>E. esula</i>]	AAF72197	6e-89	188/210	15	1	0
Glutathione S-transferase 2 [<i>Papaver somniferum</i>]	AAF22518	2e-85	175/215	37	2	0
Glutathione S- transferase/peroxidase [<i>Capsicum chinense</i>]	CAI48072	e-100	193/220	13	2	0
Glutathione S-transferase [<i>Thlaspi caerulescens</i>]	AY917145	5e-174	204/240	13	1	0
Glutathione transferase AtGST 10	BAB09723	4e-99	208/237	3	1	1

[*A. thaliana*]

Glutathione S-transferase [*A. thaliana*] AAG30140 6e-77 164/215 6 1 0

Putative glutathione S-transferase AAM64426 5e-62 153/205 5 1 0

[*A. thaliana*]

Putative glutathione S-transferase XP_470193 3e-71 163/218 14 1 0

[*O. sativa*]

Putative glutathione transferase [*O. sativa*] BAD28950 7e-22 52/60 2 1 0

Glutathione S-transferase GST 9 AAG34817 3e-31 85/110 5 1 0

[*Zea mays*]

Putative glutathione S-transferase AAM34480 4e-13 42/48 2 1 0

[*Phaseolus acutifolius*]

Glutathione S-transferase BAC21263 5e-66 153/189 4 1 0

[*Cucurbita maxima*]

Isochorismate synthase

isochorismate synthases, putative ABE91019 1e-81 185/202 4 1 (C2) 0
[*M. truncatula*].

isochorismate synthase [*C. annuum*] AAW66457 3e-65 175/244 2 1 0

Isochorismate synthase, chloroplast precursor [*Catharanthus roseus*] Q9ZPC0 1e-58 163/246 1 0 1

Lipoxygenase

Lipoxygenase [*Fragaria x ananassa*] CAE17327 e-126 219/255 4 1 1

Lipoxygenase [<i>Corylus avellana</i>]	CAD10740	4e-76	158/187	1	0	1
Lipoxygenase [<i>Solanum tuberosum</i>]	CAA65268	0.0	622/803	29	3	0
Lipoxygenase [<i>C. jambhiri</i>]	BAB84352	0.0	762/904	226	6 (C11)	10
Lipoxygenase 1 [<i>Sesbania rostrata</i>]	CAC43237	0.0	803/950	9	1 (C7)	0
Lipoxygenase 1 [<i>Prunus dulcis</i>]	CAD10779	8e-05	44/74	2	0	2
Lipoxygenase 2 [<i>Zantedeschia aethiopica</i>]	AAG18376	0.0	690/819	27	2 (C3)	0
Lipoxygenase 2 [<i>Nicotiana attenuata</i>]	AAP83137	4e-83	207/289	25	3	4

Metacaspases

metacaspase 1 [<i>A. thaliana</i>]	AAP84706	e-104	202/243	9	2	0
metacaspase 3 [<i>A. thaliana</i>]	AAP44516	6e-38	125/226	1	0	1
metacaspase 7 [<i>A. thaliana</i>]	AAP84710	4e-88	206/267	5	1	0
metacaspase 1 [<i>L. esculentum</i>]	AAM51555	3e-84	181/234	2	0	2
putative metacaspase [<i>O. sativa</i>]	AAR06365	4e-20	81/150	1	0	1

NADPH

NADPH oxidase [<i>N. tabacum</i>]	AAM28891	0.0	639/720	16	1 (C1)	0
FRO2-like protein; NADPH oxidase-like [<i>A. thaliana</i>]	BAA98161	e-104	235/310	4	1 (C2)	0
ferric reductase-like transmembrane component family protein [<i>A. thaliana</i>]	NP_199785	e-152	317/401	5	1 (C3)	1
FRO2 homolog [<i>A. thaliana</i>]	BAB08722	1e-79	178/272	1	0	1

respiratory burst oxidase homolog [<i>N. benthamiana</i>]	BAC56865	3e-69	81/119	1	0	1
--	----------	-------	--------	---	---	---

Nitrate reductase

nitrate reductase [NADH] (NR) [<i>Petunia x hybrida</i>]	P36859	0.0	495/567	11	1 (C10)	0
---	--------	-----	---------	----	---------	---

nitrate reductase [<i>Tilia platyphyllos</i>]	AAN15927	2e-152	294/337	3	1 (C11)	0
---	----------	--------	---------	---	---------	---

nitrate reductase [<i>Ricinus communis</i>]	AAG30576	1e-85	179/211	6	1 (C7)	0
---	----------	-------	---------	---	--------	---

nitrate reductase [<i>Brassica napus</i>]	BAA07395	4e-23	60/89	1	0	1
---	----------	-------	-------	---	---	---

PR-5-like protein (Osmotin, Permeatin or Thaumatin)

PR-5-like protein [<i>A. thaliana</i>]	AAM44961	1e-103	218/296	2	1	0
--	----------	--------	---------	---	---	---

PR-5-like protein [<i>A. thaliana</i>]	NP_173261	1e-109	208/243	13	1	0
--	-----------	--------	---------	----	---	---

PR-5-like protein [<i>A. thaliana</i>]	BAB11214	5e-75	158/203	3	1	0
--	----------	-------	---------	---	---	---

PR-5-like protein [<i>A. thaliana</i>]	AAM64698	1e-101	193/231	8	1	0
--	----------	--------	---------	---	---	---

PR-5-like protein [<i>Citrus sinensis</i>]	AAC02549	2e-63	111/114	3	1	0
--	----------	-------	---------	---	---	---

PR-5-like protein OLP1 [<i>L. esculentum</i>]	Q41350	1e-117	204/235	25	3	0
---	--------	--------	---------	----	---	---

PR-5-like protein Mdt11 [<i>Malus x domestica</i>]	AAC36740	4e-98	191/235	3	1	0
--	----------	-------	---------	---	---	---

PR-5-like protein VVTL1 [<i>Vitis vinifera</i>]	AAB61590	9e-89	177/227	2	1	0
---	----------	-------	---------	---	---	---

PR-5-like protein [<i>Brassica rapa</i>]	T14428	6e-75	152/188	1	0	1
--	--------	-------	---------	---	---	---

PR-5-like protein [<i>Helianthus</i>]	AAM21199	9e-84	150/188	1	0	1
---	----------	-------	---------	---	---	---

annuus]

PR-5-like protein SE39b [*N.*

BAA74546

1e-83

161/210

2

0

2

tabacum]

Peroxidases

thioredoxin peroxidase [*N.*

CAC84143

e-113

234/278

45

3

2

tabacum]

apoplastic anionic guaiacol

AAL92037

e-129

282/352

47

4

1

peroxidase [*G. hirsutum*]

peroxidase [*Spinacia oleracea*]

CAA71492

e-111

246/314

23

2

0

ascorbate peroxidase [*Ipomoea*

AAP42501

e-126

231/250

41

4

0

batatas]

peroxidase [*N. tabacum*]

AAK52084

e-125

271/354

16

1

0

peroxidase precursor [*Quercus*

AAR31106

e-92

189/218

12

1

0

suber]

putative peroxidase [*A. thaliana*]

AAO50583

e-117

233/250

20

1

0

cationic peroxidase isozyme 40K

BAA07664

e-98

240/321

53

7

0

precursor [*N. tabacum*]

class III peroxidase [*G. hirsutum*]

AAP76387

e-152

291/318

11

2

0

putative peroxidase [*A. thaliana*]

AAM20043

e-129

254/294

28

1

0

bacterial-induced peroxidase

AAD43561

e-136

272/320

22

1

1

precursor [*G. hirsutum*]

peroxidase [*Populus balsamifera*

CAA66037

e-120

258/343

13

3

0

subsp. Trichocarpa]

peroxidase ATP2a [*A. thaliana*]

CAA66863

e-126

244/281

21

2

0

putative L-ascorbate peroxidase [*A.*

BAC42431

9e-53

133/170

64

1

1

thaliana]

peroxidase precursor [<i>G. max</i>]	AAD11482	e-127	267/316	10	1	1
putative ascorbate peroxidase [<i>A. thaliana</i>]	AAF23294	6e-98	183/202	7	1	0
putative trypanothione-dependent peroxidase [<i>O. sativa</i> (japonica cultivar-group)]	AAP50932	1e-86	248/406	3	1 (C22)	0
L-ascorbate peroxidase 1b (APX1b) [<i>A. thaliana</i>]	NP_187575	e-125	230/248	21	1	0
ascorbate peroxidase pirl T09845 L-ascorbate peroxidase (EC 1.11.1.11), glyoxysomal - upland cotton	AAB52954	e-119	217/223	3	1 (C65)	0
secretory peroxidase [<i>N. tabacum</i>]	AAD33072	e-133	246/257	16	1	5
chloroplast stromal ascorbate peroxidase [<i>Vigna unguiculata</i>]	AAS55853	e-165	321/369	10	1	0
Peroxidase pirl T10790 peroxidase (EC 1.11.1.7) - upland cotton	AAA99868	e-168	314/332	13	1	4
At2g29670/T27A16.23 [<i>A. thaliana</i>]	AAN18208	6e-88	212/278	13	1	0
thylakoid-bound ascorbate peroxidase [<i>N. tabacum</i>]	BAA78552	e-100	218/264	4	1	0
peroxidase precursor [<i>Populus kitakamiensis</i>]	BAA07240	e-127	263/327	7	1	0
putative ascorbate peroxidase	CAB64343	e-138	297/347	5	2	0

(TL29) [*L. esculentum*]

putative peroxidase [<i>A. thaliana</i>]	AAP40436	5e-91	181/198	4	2	0
putative peroxidase ATP2a [<i>A. thaliana</i>]	AAM65003	3e-31	71/80	6	2	1
peroxidase ATP17a like protein [<i>A. thaliana</i>]	BAD44575	1e-17	51/67	2	1	0
peroxidase [<i>P. nigra</i>]	BAA11853	e-120	253/325	2	1	0
peroxidase-like protein [<i>A. thaliana</i>]	BAB08730	1e-25	62/67	1	0	1
peroxisomal ascorbate peroxidase [<i>Cucurbita</i> cv. Kurokawa Amakuri]	BAB64351	1e-18	52/66	1	0	1
glutathione peroxidase, putative [<i>A. thaliana</i>]	NP_564813	1e-15	41/43	1	0	1
peroxidase C2 precursor-like protein [<i>A. thaliana</i>]	AAN15499	3e-14	42/75	1	0	1
glutathione transferase [<i>Carica papaya</i>]	CAA04391	8e-62	130/160	1	0	1
korean-radish isoperoxidase [<i>Raphanus sativus</i>]	CAA62597	1e-42	130/233	1	0	1
putative glutathione peroxidase [<i>A. thaliana</i>]	AAL34198	4e-42	112/174	1	0	1
peroxidase [<i>Linum usitatissimum</i>]	AAB02926	1e-72	162/213	1	0	1
putative peroxidase [<i>A. thaliana</i>]	CAB39666	3e-20	61/150	1	0	1
glutathione peroxidase (EC	S33618	5e-30	42/44	1	0	1

1.11.1.9) - sweet orange

ascorbate peroxidase [<i>Pimpinella brachycarpa</i>]	AAF22246	e-101	196/250	1	0	1
PER7_ARATH Peroxidase 7 precursor (Atperox P7)	Q9SY33	4e-58	149/214	1	0	1
cytosolic ascorbate peroxidase [<i>Fragaria x ananassa</i>]	AAB94574	2e-28	54/98	1	0	1
putative thioredoxin peroxidase [<i>O. sativa</i> (japonica cultivar-group)]	BAD28826	2e-26	64/73	1	0	1
peroxidase ATP1a [<i>A. thaliana</i>]	CAA66862	2e-29	112/199	1	0	1
peroxidase (EC 1.11.1.7), anionic, precursor - wood tobacco	B56555	3e-59	139/190	1	0	1
peroxidase [<i>A. thaliana</i>]	BAB10239	e-70	176/260	1	0	1
thylakoid-bound L-ascorbate peroxidase precursor [<i>Mesembryanthemum crystallinum</i>]	AAC19393	e-34	70/94	1	0	1
thylakoid-bound ascorbate peroxidase [<i>Cucurbita cv. Kurokawa Amakuri</i>]	BAA12029	e-60	130/146	1	0	1
thioredoxin peroxidase [<i>Secale cereale</i>]	AAC78473	4e-19	71/142	1	0	1
peroxidase [<i>A. thaliana</i>]	CAA66957	2e-43	89/140	1	0	1
putative peroxidase [<i>O. sativa</i> (japonica cultivar-group)]	BAB39281	2e-63	150/216	1	0	1

ascorbate peroxidase [<i>O. sativa</i> (japonica cultivar-group)]	BAB17666	2e-22	67/142	1	0	1
putative ascorbate peroxidase APX4 [<i>A. thaliana</i>]	AAP72143	3e-30	67/75	1	0	1
peroxidase [<i>N. tabacum</i>]	BAA82306	8e-71	155/235	1	0	1
PE48_ARATH Putative Peroxidase 48 (Atperox P48)	O81755	6e-20	61/77	1	0	1

Phenylalanine ammonia-lyase

phenylalanine-ammonia lyase [<i>C. clementina</i> x <i>C. reticulata</i>]	CAB42794	0.0	708/713	69	1 (C3)	2
phenylalanine ammonia-lyase [<i>C. limon</i>]	Q42667	0.0	715/722	23	1 (C8)	3
phenylalanine ammonia-lyase [<i>Jatropha curcas</i>]	ABI33979	0.0	356/380	4	1 (C9)	0
phenylalanine ammonia-lyase [<i>Nerium oleander</i>]	AAX18625	2e-107	200/208	2	1 (C10)	0
phenylalanine ammonia-lyase 1 [<i>Manihot esculenta</i>]	AAK62030	e-116	231/270	1	0	1
phenylalanine ammonia-lyase [<i>Populus trichocarpa</i>]	P45730	9e-55	110/145	1	0	1

Phospholipases

putative phospholipase A2 [<i>O. sativa</i>]	XP_468549	6e-47	98/120	4	1	0
phosphoinositide-specific	AAB03258	0.0	512/602	20	1	0

phospholipase C P13 [<i>G. max</i>]						
putative phospholipase C [<i>A. thaliana</i>]	AAC32238	e-127	229/276	2	1	0
phospholipase C [<i>O. sativa</i>]	CAC81703	e-101	94/118	4	0	4
phospholipase C [<i>A. thaliana</i>]	NP_175685	e-108	200/222	1	0	1
1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase [<i>N. rustica</i>]	CAA72681	e-147	273/306	1	1	0
phosphoinositide-specific phospholipase C P12 [<i>G. max</i>]	gi7435171	5e-33	67/85	1	0	1
phosphoinositide-specific phospholipase C [<i>N. rustica</i>]	CAA65127	e-112	236/286	3	2	0
PLC [<i>A. thaliana</i>]	AAL16172	9e-92	178/204	1	0	1
ATPLC2 phospholipase C (ATPLC2) [<i>A. thaliana</i>]	NP_187464	2e-70	153/191	3	0	3
phospholipase C2 [<i>N. tabacum</i>]	AAF33824	3e-73	164/214	1	0	1
PLD [<i>A. thaliana</i>]	AAP42742	e-160	289/312	14	3	1

Potassium channels

CNGC1 [<i>A. thaliana</i>]	NP_200125	0.0	366/386	48	11	7
CNGC2 [<i>G. hirsutum</i>]	AAX18166	0.0	337/359	19	3	0
CNGC3 [<i>A. thaliana</i>]	NP_566075	e-100	204/245	2	0	2
CNGC4/ HLM1 [<i>A. thaliana</i>]	Q94AS9	6e-96	175/194	1	0	1
CNGC9 [<i>A. thaliana</i>]	NP_194785	1e-81	179/234	1	0	1
CNGC10 [<i>A. thaliana</i>]	NP_563625	1e-12	83/160	2	1	0
CNGC14 [<i>A. thaliana</i>]	NP_850056	2e-84	121/158	1	0	1

CNGC15 [<i>A. thaliana</i>]	NP_180393	3e-60	135/182	2	0	2
CNGC17 [<i>A. thaliana</i>]	NP_194765	3e-179	347/394	5	1	2
CNGC19 [<i>A. thaliana</i>]	NP_188396	2e-118	254/311	6	1	0

Superoxide dismutase

SODM_HEVBR Superoxide dismutase [Mn], mitochondrial precursor S39492 superoxide dismutase (EC 1.15.1.1) (Mn)	P35017	e-105	199/231	43	1	0
Fe-superoxide dismutase precursor [<i>M. sativa</i>]	AAL32441	e-105	216/260	58	3 (C4)	0
copper-zinc superoxide dismutase [<i>Nelumbo nucifera</i>]	AAW80437	2e-55	107/112	3	1	0
Fe-superoxide dismutase precursor [<i>A. thaliana</i>]	AAM61633	3e-86	177/205	3	1 (C3)	0
iron superoxide dismutase 3 [<i>A. thaliana</i>]	AAM63713	3e-64	139/178	4	1 (C7)	0
superoxidase dismutase [<i>L. esculentum</i>]	AAQ09007	3e-77	75/83	4	1(C9)	0
superoxide dismutase [Fe] [<i>L. esculentum</i>]	CAE22480	3e-76	179/252	2	1(C18)	0
Cu/Zn-superoxide dismutase copper chaperone precursor [<i>G. max</i>]	AAK01931	3e-99	199/217	7	2 (C16)	0
copper/zinc superoxide dismutase [<i>C. limon</i>]	AAQ14591	7e-85	151/152	30	4 (C19)	1

Cu/Zn superoxide dismutase [<i>C. sinensis</i>]	CAA03881	5e-67	125/126	59	4 (C12)	5
MnSOD [<i>H. brasiliensis</i>]	CAB53458	3e-16	40/48	1	0	1
manganese superoxide dismutase [<i>Capsicum annuum</i>]	AAB88870	1e-50	101/110	1	0	1
putative CuZn-superoxide dismutase [<i>Populus tremula</i> x <i>P. tremuloides</i>]	CAC33847	3e-69	116/127	2	0	2
iron superoxide dismutase 3 [<i>A. thaliana</i>]	AAC24834	2e-59	142/200	2	0	2
iron-superoxide dismutase [<i>G. max</i>]	AAQ13492	3e-17	69/120	1	0	1
copper/zinc superoxide dismutase [<i>Bruguiera gymnorrhiza</i>]	BAB78597	3e-13	38/40	1	0	1
superoxide dismutase (EC 1.15.1.1) (Mn) precursor - curled-leaved tobacco	S03639	e-62	128/153	1	0	1
iron superoxide dismutase 3 [<i>A. thaliana</i>]	BAB11186	2e-47	65/106	1	0	1

*Between parentheses are represented the contigs (C) commented in the Results and Discussion. As search for different proteins had been made in separate archives, different contigs (representing different proteins) can have the same number.