## Theobroma grandiflorum (WILLD. ex SPRENG.) SCHUM.: Production factors in agro-ecosystems

Reisdorff, Christoph (1), Marino, Wolfgang (1), Claret de Souza, Aparecida das Graças (2), Schroth, Götz (1) and Schmidt, Petra (1)

One long term objective of the field experiments conducted on a former terra firme rain forest site near Manaus is to evaluate the feasibility of sustainable agroforestry in Amazonia, paying special attention to the potential to recultivate degraded areas. For this purpose different mixed cultivation systems are being tested and compared to monocultures designed according to common practices. Study groups of various SHIFT-projects are working on this experimental site, each group dealing with particular aspects of the different agro-ecosystems.

There are manifold aspects which need to be considered, one of these being the cultivated species itself. It needs to be studied with respect to its behavior within particular field environments in order to come to a kind of agroecological profile of the species and its varieties. In addition, the economic potential of a crop plant is important as well, considering that an ecologically sound cultivation system will only be put into practice if it promises to be profitable.

One of the species being part of the field experiments is *Theobrona grandiflorum*. This species is attracting particular interest because of the following facts:

- It is native in the Amazon region;
- The livestock of cultivated plants represents a primary selection with a high degree of variability, thus, still offering the opportunity to combine economic criteria with agro-ecological criteria for selection and breeding;
- T. grandiflorum. is economically attractive because of the high market value of the fruit flesh;
- The species is closely related to cocoa, hence offering an use option in addition to pulp-production, namely the use of the seeds for the preparation of a chocolate like product.

Our poster sketches some main lines of research with respect to the species *Theobroma grandiflorum* which have been done within the frame of the SHIFT experimental site at the EMBRAPA Amazônia Ocidental near Manaus. This includes aspects of eco-physiology, biotic interactions, plantation management, infra-specific variability and the utilization of the seeds.

<sup>(1)</sup> Universität Hamburg, Hamburg, Germany. (2) EMBRAPA Amazônia Ocidental, Manaus, Brazil.