

P2-223

# MEMOGAIN; SUPRA-BIOAVAILABLE NICOTINIC ENHANCER FOR THE TREATMENT OF ALZHEIMER'S DISEASE

Alfred Maelicke, Galantos Pharma GmbH, Mainz, Germany.  
Contact e-mail: alfred.maelicke@galantos.com

**Background:** Galantamine (Reminyl, Razadyne) is a potent allosterically potentiating ligand (APL) of nicotinic receptors and a moderate inhibitor of cholinesterases. In addition to its cognition-enhancing properties and mediated by its action on  $\alpha 7$  nicotinic receptors, galantamine reduces amyloid toxicity and neuronal death. Manifestation of these properties in patients is limited by low dosing necessitated by need to control the usual gastrointestinal side effects (nausea, vomiting, diarrhoea) of cholinesterase inhibitors. We have developed a derivative of galantamine ("Memogain") for the following purposes: increased bioavailability in the brain and reduced, if not negligible, GI side effects at efficacious doses. **Methods:** Animal model studies to test for enhanced bioavailability in the brain, reduced dosing schemes, absence of gastrointestinal side effects, potency in overcoming drug-induced cognitive impairment, tissue distribution of pro-drug and active moiety, pharmacokinetics and toxicity. **Results:** Memogain is an inactive pro-drug of galantamine having more than 15fold higher bioavailability in the brain than the same doses of galantamine. Memogain is enzymatically cleaved in the brain to release its active moiety galantamine. In animal models of dementia, Memogain produced several fold larger cognitive improvement than galantamine, in the absence of any GI side effects. In the ferret dramatically reduced emetic responses were observed when Memogain was administered instead of galantamine. **Conclusions:** Should the advantageous properties of Memogain seen in animal models hold up to its action in man, we expect a dramatically improved side effects profile, no requirement for up-titration of dose to achieve satisfactory compliance, improved potency because of better delivery to the brain, and neuroprotective activity in addition to symptomatic cognitive enhancement. Memogain may become a second generation antidegenerative drug of combined symptomatic and disease-modifying properties.

P2-224

# ACTIVITY OF PHOSPHOLIPASE A<sub>2</sub> (PLA<sub>2</sub>) SUBTYPES IN RAT BRAIN IS ALTERED BY FEEDING CONJUGATED LINOLEIC ACID (CLA) AND LINSEED OIL

Nádia R. Barbosa<sup>1</sup>, Marco A. S. Gama<sup>2</sup>, Fernando C. F. Lopes<sup>2</sup>, P. P. Defillipo<sup>3</sup>, Fábio B. Mury<sup>3</sup>, Rafael Trevisani<sup>4</sup>, Dimas E. Oliveira<sup>5</sup>, Wagner F. Gattaz<sup>3</sup>, <sup>1</sup>Faculty of Pharmacy and Biochemistry, Universidade Federal de Juiz de Fora, Laboratory of Neuroscience, IPQ, FMUSP, Juiz de Fora/São Paulo, Brazil; <sup>2</sup>National Dairy Cattle Research Centre, Embrapa, Juiz de Fora, Brazil; <sup>3</sup>Psychiatry Institute, Faculty of Medicine, University of São Paulo, São Paulo, São Paulo, Brazil; <sup>4</sup>Faculty of Pharmacy and Biochemistry, Universidade Federal de Juiz de Fora, Juiz de Fora, Brazil; <sup>5</sup>Universidade do Estado de Santa Catarina, UDESC, Chapecó, Brazil.  
Contact e-mail: nadiafox@gmail.com

**Background:** There are growing evidences that pathogenesis of neurological disorders are associated with an altered inflammatory response, where PLA<sub>2</sub> enzyme plays a key critical role. However, different PLA<sub>2</sub> subtypes have been identified in mammals which can differ in several aspects such as substrate specificity. Noteworthy, recent studies found reduced iPLA<sub>2</sub> activity in blood samples and postmortem brain tissues of Alzheimer patients. Since certain CLA isomers and n-3 polyunsaturated fatty acids (n-3 PUFA) have been shown to modulate selectively PLA<sub>2</sub> activity, dietary supplementation with these FA could influence Alzheimer's disease progression. **Methods:** This study was designed to evaluate the effects of CLA and linseed oil on activity of PLA<sub>2</sub> subtypes (iPLA<sub>2</sub>, sPLA<sub>2</sub> and cPLA<sub>2</sub>) in rat brain. Forty-eight Wistar rats (75±10 days-old) received by gavage 1.1 ml/day of the following treatments, for 30 days: 1) Control (saline solution); 2) CLA supplement; 3) sunflower oil (SO) and 4) linseed oil (LO). The CLA supplement contained 60% of total CLA (30% of trans-10 cis-12 and 30% of cis-9 trans-11) as methyl esters. SO and LO contained 65% of linoleic acid and 50% of  $\alpha$ -linolenic acid and were used as a source of n-6 and n-3 PUFA, respectively. After 30 days, animals were decapitated and samples of hippocampus and frontal cortex were collected to perform PLA<sub>2</sub> subtypes activity essays. Results

were analyzed by ANOVA followed by Tukey-test and differences were declared significant at  $P < 0.05$ . **Results:** Activity of iPLA<sub>2</sub> in the cortex increased by 160 and 145% ( $P < 0.05$ ) in LO when compared with SO and control, respectively. However, activities of sPLA<sub>2</sub> and cPLA<sub>2</sub> were unaltered by treatments in the cortex. In the hippocampus, there were no differences among treatments for both iPLA<sub>2</sub> and sPLA<sub>2</sub>, but activity of cPLA<sub>2</sub> increased by 262, 210 and 400% ( $P < 0.01$ ) in CLA when compared with LO, SO and Control (0.16 vs. 0.061, 0.076 and 0.040 pMol/mg prot/min, respectively). **Conclusions:** Our results suggest a nutraceutical potential of linseed oil and CLA for prevention and/or inhibition of Alzheimer's disease progression.

P2-225

# CHARACTERIZATION OF INOSITOL TRANSPORTERS AS A METHOD FOR DRUG DELIVERY TO THE CNS

Daniela Fenili<sup>1,2</sup>, Mark Nitz<sup>3</sup>, JoAnne McLaurin<sup>1,2</sup>, <sup>1</sup>Department of Laboratory Medicine and Pathobiology, University of Toronto, Toronto, ON, Canada; <sup>2</sup>Centre for Research in Neurodegenerative Diseases, Toronto, ON, Canada; <sup>3</sup>Department of Chemistry, University of Toronto, Toronto, ON, Canada. Contact e-mail: daniela.fenili@utoronto.ca

**Background:** One challenge to designing drugs for targeting CNS diseases is to develop compounds that when given orally, cross the blood-brain barrier to reach their target sites in the brain. One design strategy is to develop drugs that can cross into the brain using a known transport system. myo-Inositol is able to cross into the brain using the inositol transporters, of which there are three: one hydrogen/myo-inositol cotransporter (HMIT) and two sodium/myo-inositol cotransporters (SMIT1, SMIT2). Our laboratory has previously shown that scyllo-inositol, ELND005, is able to both prevent and reverse Alzheimer's disease pathology in the TgCRND8 mouse model of Alzheimer's disease. We have also shown, using gas chromatography/mass spectrometry, that ad libitum treatment of mice with scyllo-inositol results in a significant increase in the brain levels of scyllo-inositol in both TgCRND8 and non-Tg littermates (Fenili et al, 2007, J Mol Med). **Methods:** Competitive transport assays have been developed to examine the minimal binding requirements for transport by the three known inositol transporters in astrocytic and neural cell lines. Inositol, simple sugars and related compounds, 20 in total, were used to define the structural features required for active transport. Further the relative expression of the transporters as a function of age and disease progression was analyzed in the cortex, hippocampus, septum and cerebellum using QPCR. **Results:** Across all brain regions analyzed the expression level of the transporters were HMIT > SMIT1 > SMIT2. This result did not vary with age or disease phenotype. **Conclusions:** All three inositol transporters are expressed in the brain, with a similar pattern of expression being observed regardless of brain region.

P2-226

# PUBLIC SUPPORT FOR AN ANNUAL TAX INCREASE TO PROVIDE ACCESS TO AN ALZHEIMER'S DISEASE MEDICATION IN CANADA

Natasha Clayton, Mark Oremus, Jean-Eric Tarride, Parminder Raina, McMaster University, Hamilton, ON, Canada. Contact e-mail: clayton@mcmaster.ca

**Background:** In Canada, the cost of Alzheimer's disease (AD) medications is mainly borne by patients and caregivers. The primary objective of this study was to measure the Canadian general public's support for a program of unrestricted access to AD medications. The secondary objective was to identify determinants of support for the program. **Methods:** A national sample of 500 Canadians, aged 18 years or older, was randomly recruited to participate in a computer-assisted telephone interview. The sample was stratified by income. Participants were presented with a set of randomly-ordered scenarios describing a hypothetical, new AD medication. The efficacy of the medication was varied by scenario: the medication was alternately described as modifying the symptoms of cognitive decline or actually halting disease progression. The adverse effects profile was also varied in the scenarios: no adverse effects or a 30% chance of some adverse effects. For each scenario, participants were asked whether they supported an annual increase in personal income taxes to fund unrestricted access to the AD medication.



ScienceDirect - Alzheimer's & Dementia, Volume 5, Issue 4, Supplement 1, Pages P1-P552 (July 20 - Windows Internet Explorer

http://www.sciencedirect.com/science?\_ob=PublicationURL&\_tockey=%23TOC%2326546%232009%23999949995.6996%231239092%23FLA%238\_c...

Arquivo Editar Exibir Favoritos Ferramentas Ajuda

Emissão de Parecer Ad hoc... ScienceDirect - Alzheimer...

ScienceDirect

Login: Register

Home Browse Search My Settings Alerts Help

Quick Search All fields Author search tips Journal/book title --This Journal/Book-- Volume Issue Page Clear Go Advanced Search

Alzheimer's & Dementia

Copyright © 2009 The Alzheimer's Association. All rights reserved.

Sample Issue Online | About this Journal | Submit your Article | Shortcut link to this Title

New Article Feed Alert me about new Volumes / Issues

Add to collab Add to Favorites

Alzheimers Association

ISTAART: Join now!

Font Size: + -

Full-text available  
Abstract only  
Articles in Press

Volume 5 (2009)

Volume 5, Issue 4  
pp. 287-368 (July 2009)

Volume 5, Issue 4, Supplement 1  
pp. P1-P552 (July 2009)

Alzheimer's Association International Conference on Alzheimer's Disease, Alzheimer's Association International Conference on Alzheimer's Disease  
Vienna, Austria  
11-16 July 2009

Volume 5, Issue 3  
pp. 199-286 (May 2009)

Volume 5, Issue 2  
pp. 81-198 (March 2009)

Volume 5, Issue 1  
pp. 1-80 (January 2009)

Volume 5, Issue 4, Supplement 1, Pages P1-P552 (July 2009)

Alzheimer's Association International Conference on Alzheimer's Disease, Alzheimer's Association International Conference on Alzheimer's Disease  
Vienna, Austria  
11-16 July 2009

Previous vol/iss Next vol/iss

Add to my Quick Links

No prev art. 1 - 100 of 1,660 Next

Email Articles Export Citations Download PDFs Open All Previews

1. Subscriber information page  
Page A1  
Preview PDF (50 K) | Related Articles

2. ICAD Scientific Program Committee  
Page A2  
Preview PDF (37 K) | Related Articles

3. Contents  
Pages A3-A5  
Preview PDF (75 K) | Related Articles

Editorial Board

Internet | Modo Protegido: Ativado

Windows Vista

09:08

Iniciar PTBIC-2009 ScienceDirect - ... http://efomento.c... Encaminhamento-F... Edital PTBIC - CNPq... Sem título - Paint