



Association mapping for yield and grain quality traits in rice (*Oryza sativa* L.)

Tereza Cristina de Oliveira Borba¹, Rosana Pereira Vianello Brondani¹, Flávio Breseghello¹, Alexandre Siqueira Guedes Coelho², João Antônio Mendonça¹, Paulo Hideo Nakano Rangel¹ and Claudio Brondani¹

¹*Embrapa Arroz e Feijão, Santo Antônio de Goiás, GO, Brazil.*

²*Setor de Melhoramento de Plantas, Escola de Agronomia, Campus-II Samambaia, Universidade Federal de Goiás, Goiânia, GO, Brazil.*

Abstract

Association analysis was applied to a panel of accessions of Embrapa Rice Core Collection (ERiCC) with 86 SSR and field data from two experiments. A clear subdivision between lowland and upland accessions was apparent, thereby indicating the presence of population structure. Thirty-two accessions with admixed ancestry were identified through structure analysis, these being discarded from association analysis, thus leaving 210 accessions subdivided into two panels. The association of yield and grain-quality traits with SSR was undertaken with a mixed linear model, with markers and subpopulation as fixed factors, and kinship matrix as a random factor. Eight markers from the two appraised panels showed significant association with four different traits, although only one (RM190) maintained the marker-trait association across years and cultivation. The significant association detected between amylose content and RM190 was in agreement with previous QTL analyses in the literature. Herein, the feasibility of undertaking association analysis in conjunction with germplasm characterization was demonstrated, even when considering low marker density. The high linkage disequilibrium expected in rice lines and cultivars facilitates the detection of marker-trait associations for implementing marker assisted selection, and the mining of alleles related to important traits in germplasm.

Key words: association analysis, core collection, genetic structure.

Received: March 16, 2009; Accepted: February 23, 2010.

Introduction

Association analysis, or linkage disequilibrium mapping, is a notable strategy used for identifying genes controlling important traits. It is already being successfully applied for identifying genes related to human diseases. Research in humans has turned to association analysis, since linkage analysis has not been successful in the fine-scale mapping of disease loci, due to the impossibility of undertaking controlled-breeding crosses (Flint-Garcia *et al.*, 2003). Unlike humans, in most plant species, the identification of those genomic regions which contribute to important characteristics has been mostly achieved through linkage analysis within segregating populations, the result of crosses between genitors with contrasting phenotypes and genotypes (Buntjer *et al.*, 2005; Skot *et al.*, 2005).

In breeding and pre-breeding programs, QTL detection is an important tool for the identification of favorable alleles and for identifying and validating molecular assisted

selection (MAS). The identification of genomic regions related to quantitative traits in plants was largely achieved through QTL mapping (Skot *et al.*, 2005). However, there are some inherent limitations to QTL analysis. First, in linkage studies, the segregating population usually presents only two segregating alleles per locus, which means that in any given cross, the sampled genetic diversity may be limited. In addition, the potential use of the genetic diversity available in species is restricted (Peleman and van der Voort, 2003). Another disadvantage as regards the substantiation of a low number of traits per cross, is through the difficulty in identifying parents with contrasting genotypes and phenotypes for all those traits of interest (Buntjer *et al.*, 2005). Furthermore, the high resolution desired for MAS or cloning candidate genes requires developing large segregating populations, possibly difficult in some species (Skot *et al.*, 2005).

According to Zondervan and Cardon (2004), the main purpose in linkage analysis, as in association mapping, is the detection of correlations between phenotypic variation and genotypes through linkage disequilibrium. However, association analysis has the advantage of contemplating all the meiotic and recombination events that may occur in the

evaluated population (Ferreira and Grattapaglia, 2006). Furthermore, this form is highly dependent on the extent of linkage disequilibrium (LD), a higher degree implying the use of less markers per chromosome, without the loss of genetic resolution for marker assisted selection (MAS) (Rostoks *et al.*, 2006). As rice is a self-pollinating species, it is expected to present high linkage disequilibrium (Flint-Garcia *et al.*, 2003), thereby requiring fewer markers. In addition, the recent bottlenecks encountered since the beginning of rice breeding have given rise to high linkage disequilibrium blocks, thereby facilitating association studies (Patron *et al.*, 2002).

One of the great advantages of association mapping lies in the fact that no mapping population needs to be developed, as the sampling of non-related individuals represents a series of advantages towards developing and validating MAS in breeding programs (Jannink *et al.*, 2001), as well as an opportunity for increasing the exploitation of germplasm accessions in the search for advantageous allele combinations. Such a strategy, unlike traditional linkage analysis, facilitates the search for functional variation in a much broader germplasm context (Zhu *et al.*, 2008). Thus, experimental populations may constitute a representative sample of a larger population for which inferences are sought (Breseghello and Sorrels, 2006b). In panels with highly divergent individuals and assumed random mating, only polymorphisms with extremely tight linkage to a locus with desirable phenotypic effects are likely to be significantly associated with a given trait (Remington *et al.*, 2001). Furthermore, association analysis can benefit by including data collected over years of experimental analysis with genotypes of breeding programs, with the additional possibility of analyzing several traits simultaneously.

Improving grain yield and quality are important challenges in rice breeding, thus priorities for the international market (Fan *et al.*, 2005). Although quality assumes many aspects and is highly related to preference in diverse cultures, its characteristics are mainly defined by milling properties, grain size and shape, cooking and eating characteristics, and nutritional qualities (He *et al.*, 1999). According to He *et al.* (1999), of these the most relevant are appearance and cooking quality, reported to be directly related to amylose content, gel consistency and gelatinization temperature (Fan *et al.*, 2005).

The aims of this work were to analyze and identify the association of simple sequence repeat (SSR) markers with yield and grain quality traits in a panel of accessions from the Embrapa Rice Core Collection (ERiCC), represented by breeding material from Brazil and other countries.

Material and Methods

Plant material and genomic DNA extraction

The evaluated panel of 242 accessions from ERiCC (Abadie *et al.*, 2005) was composed of: a) 94 accessions of

inbred lines and cultivars developed by rice breeding programs in Brazil (57 upland and 37 lowland accessions); and b) 148 accessions of inbred lines and cultivars developed by breeding programs worldwide (76 upland and 72 lowland accessions) (Table S1). Each accession was evaluated in a four-plant bulk, the total genomic DNA being extracted as described by Brondani *et al.* (2002).

Phenotypic data

The phenotypic evaluation of 242 ERiCC accessions was carried out in Santo Antônio de Goiás, the state of Goiás, Brazil (altitude 749 m; 16°40'43" S; 49°15'14" W), in 2004 and 2005, under irrigated conditions, following an augmented block design with plots of 4 rows x 5 m, at a density of 20 plants m⁻¹. Data were taken from the two middle rows and the 4 central meters of each. In 2004, the following traits were evaluated: 1) YLD - grain yield, in kg ha⁻¹; 2) TILN - tiller number per plant; and 3) PANN - panicle number per plant. In 2005, two traits were evaluated: YLD and the yield from ratooning (RYLD), this being the yield from the second harvest, approximately 40 days post-main. In both years, grain quality data were evaluated for cooking and milling traits, measured as to amylose content (AC) and head-milled rice (MR), respectively. Amylose content was determined according to Juliano (1979), whereas head-milled rice was measured as the proportion of the weight of whole kernels over the weight of paddy rice. The descriptive statistics of phenotypic data were computed using the Genes 4.1 program (Cruz, 1997).

SSR characterization of ERiCC

The 86 SSR fluorescent markers were dispersed in all the 12 rice chromosomes, an average of seven markers per chromosome (ranging from a maximum of eight markers and a minimum of five per chromosome). The markers were labeled with the fluorescent dyes HEX (hexachlorine - 6 carboxyfluorescein) and 6-FAM (fluorochrome 6-carboxyfluorescein) (Table S2). PCR was carried out in a final volume of 15 µL containing a customized concentration of primers (forward and reverse) (Table S2), 1X reaction buffer (50 mM of KCl, 10 mM Tris-HCl pH 8.4, 0.1% Triton X-100 and 1.5 mM of MgCl₂), 0.22 mM of dNTP, 15 ng of template DNA, and one unit of the Taq DNA Polymerase enzyme. Thermocycling was carried out in a GeneAmp PCR System 9700 (Applied Biosystems) and the amplification conditions were 94 °C for 5 min followed by 40 cycles of 94 °C for 1 min, specific annealing temperature for 1 min, and 72 °C for 1 min, and a final extension of 72 °C for 7 min. PCR products were analyzed in an ABI 3100 DNA sequencer (Applied Biosystems) and the alleles were scored with GeneMapper 3.5 software (Applied Biosystems). The size standard used was obtained according to Brondani and Grattapaglia (2001).

Statistical analysis

Allele diversity and genetic structure: The 242 inbred lines and cultivars of ERiCC were analyzed in a pooled DNA sample, composed of four individual plants. From SSR analysis, heterogeneity (the presence of individual heterozygous or homozygous plants with different alleles in the bulk) was evident in certain accessions. In order to proceed with association analysis, the accessions were treated as pure lines, under the definition of working alleles, and represented by the most common allele. Rare alleles (frequency below 5%) were treated as missing data in population structure analysis, and as null alleles in association analysis, according to the strategy described by Breseghello and Sorrells (2006a). The hypothesis of division from one to four subpopulations was tested with structure software (Pritchard *et al.*, 2000), allowing for admixture and correlated allele frequencies, with a burn-in of 10,000 and a run-length of 100,000. The *Fst* parameter (software FSTAT 2.9.3.2; Goudet, 2002) and factorial correspondence analysis (FCA) (software Genetix 4.03; Belkhir *et al.*, 2004) were also applied for investigating accession genetic structure.

Association analysis: The association between markers and phenotypic traits was done using the Mixed Linear Model (MLM), an available option in Tassel version 1.9.6 software, where markers tested and subpopulation data (Q matrix) were considered as fixed-effect factors, whereas the kinship matrix was considered as a random-effect factor. The kinship matrix was obtained from SPAGeDi version 1.2 software (Hardy and Vekemans, 2002). To confirm the significance of associations between loci and traits, a correction for multiple testing was applied using the false discovery rate (FDR) method with Qvalue version 1.0 software (Storey, 2002). The FDR level was set at 0.05, and the π_0 method for bootstrap analysis. The FDR method, expressed as a *q*-value, is defined as the expected proportion

of true null hypotheses within the class of rejected null hypotheses (Kraakman *et al.*, 2004).

The significance of differences between allele effects was obtained from Kruskal-Wallis nonparametric rank testing with *R* program (*R* Development Core Team). Non-parametrical multiple test procedures, regarding amylose content and each pair of alleles, were carried out as described by Campos (1983).

Results

Phenotypic data

Experimental field data were distributed normally, except for traits related to grain quality (AC and MR). ERiCC accessions revealed wide variation in performance of all the evaluated traits (Table 1). In the 2004 experiment, the firmest correlation detected was between TILN and PANN (0.88; $p < 0.01$), and in the 2005, between YLD and MR (0.38; $p < 0.01$) (Table 2). No correlation was detected in yield data from 2004 and 2005 experiments.

Allele diversity and population structure

A total of 1,066 alleles were detected with the set of 86 SSR markers on a panel of 242 accessions. The average number of alleles per locus was 12.4, ranging from three (RM484) to 32 (RM204). Most loci presented one or more alleles with a frequency below 5%. These rare alleles represented, approximately, 48% of the total allele number, and to avoid an increase in variance errors in association analysis, they were not considered. The remaining alleles (554), referred to as common alleles, ranged from two to eight per locus. In lowland accessions, the mean was 4.80 alleles per locus, with gene diversity of 0.64, whereas in upland accessions this was 4.76 alleles per locus and gene diversity 0.56. Common SSR alleles were used to check the structure of ERiCC genetic variation. The model-based clustering method resulted in the highest likelihood from data, *i.e.* the

Table 1 - Descriptive statistics for yield (YLD), tiller number (TILN), panicle number (PANN), yield from ratooning (RYLD), amylose content (AC) and head-milled rice (MR).

	2004 Experiment					2005 Experiment			
	YLD (kg/ha)	TILN	PANN	AC (%)	MR (%)	YLD (kg/ha)	RYLD (kg/ha)	AC (%)	MR (%)
Average	4685.0	166.43	143.3	24.3	60.4	4298.1	1403.4	24.1	43.1
Minimum	900	71	32	4	42.0	435	0	8	3.57
Maximum	8844	281	236	31	70.9	8130	3020	31	66.1
Standard deviation	41.6	6.7	6.2	2.5	2.5	39.0	24.4	2.2	3.8
BR IRGA 409 [#]	5993.5	154	128	27	65.1	5372.3	1425.7	26	46.9
CAIAP0 [#]	2948.7	114	102	26	-	5720.2	1125.0	24	61.9
METICA 1 [#]	6243.5	197	175	25	-	3282.9	1884.2	25	43.17
COLOSSO [#]	3911.1	127	108	25	-	4785.0	1350.0	24	65.4

[#] Controls of field experiments for both years.

Table 2 - Pearson correlation coefficients among the phenotypic traits: yield (YLD), tiller number (TILN), panicle number (PANN), yield from ratooning (RYLD), amylose content (AC) and head-milled rice (MR).

		2004 Experiment					2005 Experiment			
		YLD (kg/ha)	TILN	PANN	AC (%)	MR (%)	YLD (kg/ha)	RYLD (kg/ha)	AC (%)	MR (%)
2004 Experiment	YLD	-								
	TILN	0.20**								
	PANN	0.31**	0.88**							
	AC	0.19**	0.26**	0.25**						
	MR	0.31**	-	-	-					
2005 Experiment	YLD	-	-	-	-	-				
	RYLD	-	-	-	-	-	-			
	AC	-	-	-	0.82**	-	0.20**	0.26**		
	MR	-	-	-	-	0.14*	0.38**	-	-	-

Only significant values are shown (* $p < 0.05$; ** $p < 0.01$).

probability that a given individual originated from a certain population, when the number of subpopulations (k) was set at 2, thereby indicating a subdivision among accessions caused by the cultivation system (lowland or upland rice accessions). No population structure was detected due to the origin of accessions (Brazilian or worldwide breeding programs).

Thirty-two accessions (13%) were predicted to have admixed ancestry, this meaning that their origin could not be attributed exclusively to one of the two inferred subpopulations. Consequently, they were discarded from the analysis. Association analysis was then undertaken with 210 ERiCC accessions (92 lowland and 118 upland). The overall F_{st} statistics across subpopulations was 0.775, and F_{st} values for lowland and upland groups were 0.135 and 0.205, respectively, thereby indicating high differentiation among subpopulations, and low to intermediate levels of

differentiation within accessions from the same subpopulation. FCA was applied in order to visualize the subdivision among accessions, whereby it was possible to confirm the division into two subpopulations, based on the cultivation system (Figure 1).

Association analysis

Three different accession panels, viz., a complete panel of 210 accessions, and from this, 92 lowland accessions and 118 upland, were analyzed (association analysis). 23 SSR markers (27%) in the complete panel were identified as being significantly associated to at least one of the evaluated traits (data not shown). On considering the low genomic coverage of the SSR set, and the complexity of the traits evaluated, the number of associated markers was high, thereby implying the presence of spurious marker-trait associations. On considering the system of cultivation




Figure 1 - Spatial distribution of genetic variability in the 210 selected accessions from ERiCC, based on factorial correspondence analysis (FCA). The white dots represent lowland accessions and the gray, upland ones.

as the main generator of genetic structuring among all the 210 accessions, and using subpopulations as a basis for re-analysis, two accession panels were defined, with a lower number of marker-trait associations. From the 86 SSRs, eight markers were associated with four traits in lowland, and only one with amylose content in upland accessions (Table 3). In lowland accessions, the marker OG60 was associated with PANN in the 2004 experiment and RM190 with AC in both the 2004 and 2005. In 2005, two markers (RM1 and 4653) were associated with MR, and four (RM264, RM267, RM125 and RM38) with YLD. In upland accessions and in both years, there was a significant association of only RM190 with AC. There was no association with any marker in the case of RYLD and TILN. Despite the significant associations between marker-trait, the only consistent association over years and cropping systems was between the RM190 marker and AC. This association was also significant in the analysis of all the 210 accessions (data not shown).

On considering the 210 accessions, six alleles (frequencies over 5%) were distinguished for RM190, with four common in both accession panels (alleles 105, 107, 121 and 125 bp). However, two alleles (117 and 119 bp) were identified (frequencies over 5%) one each only in upland and lowland accession panels, respectively (Figure 2). From Kruskal-Wallis testing, it was possible to identify a significant difference in allele effects across lowland and upland panel accessions. From a multiple test based on a non-parametrical approach it was possible to identify the relationships of the six RM190 alleles with diverse classes of amylose content, as well as positive and significant correlation, for both years, in upland and lowland panel accessions (0.58 with $p < 0.01$, and 0.54 with $p < 0.01$, respectively).

In the lowland accession panel, the 105 and 107 bp (base pair) alleles presented significantly different effects from the remainder. According to 2004 and 2005 experi-

mental data, both were correlated to higher amylose content than the others, although in 2004, there was a significant difference between the two themselves, with the 105 bp revealing higher AC than the 107 bp (Table 4). Despite statistical differences between AC and the respective alleles, no clear pattern in allele effects could be identified in lowland accessions. In the upland accession panel, both in 2004 and 2005, 121 bp allele AC was lower than in 105, 107 and 125 bp, whereas in 2005, 117 bp allele AC was lower than in 105 and 107 bp. As with lowland accessions, no clear pattern in allele effects was identified.

Discussion

SSR allele diversity and population structure

A set of 86 highly informative SSR markers was used in genotyping ERiCC inbred lines and cultivars. The number of common alleles detected (frequency $\geq 5\%$) was similar to that previously identified for inbred rice lines and cultivars (Lu *et al.*, 2005). In the present work, rare alleles were not integrated into analysis, as low frequency alleles inflate variance estimates of linkage disequilibrium (Remington *et al.*, 2001). Additionally, rare alleles are more susceptible to bias caused by covariance between markers and population structure, thus increasing the chance of type I error (Brescghello and Sorrels, 2006a). The presence of admixture also contributes to overestimating linkage disequilibrium, due to causes not related to physical connections on a chromosome (Flint-Garcia *et al.*, 2003).

According to Brescghello and Sorrels (2006a), the selection process of a minimum sample with maximum variation results in a normalizing effect that is expected to minimize population structure, thereby creating a favorable situation for association analysis. Structure-presence specification is a prior requirement in core collections, since certain procedures adopted to build these collections may lead

Table 3 - Association of SSR markers with phenotypic traits. The statistics shown refer to the coefficient of determination (R^2).

	Marker	Chromosome	Experiment 2004		Experiment 2005		
			PANN	AC	MR	YLD	AC
Lowland accessions	RM1	1	0.019	0.031	0.039* (<i>q</i>)	0.000	0.000
	RM38	8	0.083	0.049	0.021	0.040* (<i>q</i>)	0.074
	RM125	7	0.039	0.040	0.037	0.002* (<i>q</i>)	0.000
	RM190	6	0.062	0.425* (<i>q</i>)	0.012	0.049	0.36* (<i>q</i>)
	RM264	8	0.000	0.170	0.027	0.001* (<i>q</i>)	0.110
	RM267	5	0.020	0.019	0.020	0.011* (<i>q</i>)	0.035
	4653	12	0.137	0.068	0.004* (<i>q</i>)	0.143	0.070
	OG60	4	0.352* (<i>q</i>)	0.023	0.053	0.000	0.204
Upland accessions	RM190	6	0.000	0.390** (<i>q</i>)	0.000	0.090	0.490** (<i>q</i>)

Panicle number (PANN); amylose content (AC); head-milled rice (MR); yield (YLD).

Only SSR markers with significant marker-trait association are given. The *q* indicates the false discovery rate control value set to 0.05. * $p < 0.005$; ** $p < 0.0001$.

to a structured accession panel. The stratification of accessions into meaningful groups, with the maximum variation between groups and the minimum within, as suggested by van Hintum *et al.* (2000), may result in accession panels with low potential for detecting gene effects through association analysis, since most variance is attributed to population structure. ERiCC inbred lines and cultivars were origi-

nally selected according to the origin of accessions (Brazil or worldwide breeding programs), and their system of cultivation (lowland or upland) (Abadie *et al.*, 2005). The ERiCC structure based on a cultivation system is probably due to most lowland accessions being from *indica* subspecies, whereas upland accessions are mostly from *japonica* (Khush, 1997).




Figure 2 - Empirical distribution of amylose content (y-axis) among alleles identified for RM190 SSR (x-axis). The subdivisions in amylose content data refer to quartile division, and the lines in boxes are the median of amylose content in each allele. The A and B boxes refer to amylose content data on lowland accessions from the 2004 and 2005 experiments, respectively. The C and D boxes refer to amylose content data on upland accessions from the 2004 and 2005 experiments, respectively. The pool of rare and missing alleles is represented by the PMR denominated allele.

Table 4 - Pairwise statistical differences in average amylose content values of each identified allele of the RM190 marker in both accession panels and over experimental years.

	105	107	117	119	121	125	PMR
105	-						
107	L/04	-					
117	U/05	U/05	-				
119	L/04, L/05	L/04, L/05	-	-			
121	L/04, U/04, U/05	L/04, L/05, U/04, U/05	-	-	-		
125	L/04	L/05	-	-	U/04, U/05	-	
PMR	-	-	-	-	U/04	-	-

L - Lowland accession panel. U - Upland accession panel. 04 - Data from 2004 experiment. 05 - Data from 2005 experiment. PMR - pool of missing and rare alleles for the RM190 marker.

A very substantial differentiation between lowland and upland accessions was identified, after the removal of rare alleles and admixed accessions, with *Fst* increasing from 0.11 to 0.77. In Garris *et al.* (2005), *Fst* values for *indica* and *temperate japonica* and *indica* and *tropical japonica* accessions were, respectively, 0.43 and 0.36, with the proportion of admixed ancestry estimated at 10%. The higher *Fst* value identified between lowland and upland accession panels, even when compared to the divergent *indica* and *japonica* accessions found by Garris *et al.* (2005), may be due to the elimination of rare alleles and accessions with mixed ancestry, thereby increasing divergence between the two.

Association analysis

Despite the broad genome coverage provided by the set of 86 SSR markers used in the analysis, the whole genome scan designed for association analysis was limited by the low density of markers. However, due to numerous bottlenecks in the history of rice domestication, the level of linkage disequilibrium in some regions is estimated to be greater than the 250 kb found in *Arabidopsis* (Garris *et al.*, 2003). Considering this situation, a target marker can be associated to a candidate gene responsible for certain traits, even though thousands of base pairs apart, thereby favoring marker-assisted selection procedures.

Association analysis with MLM combines information from kinship and population structure, thereby reducing type I errors (Yu *et al.*, 2006), that is the error of rejecting a null hypothesis when it is actually true. Here, 23 SSR markers, associated to yield and grain quality traits, were identified, when such an approach was applied to the panel of 210 accessions. Agrama *et al.* (2007), on using the same model in association analysis of yields and their components in 103 rice germplasm accessions with 123 SSR markers, identified 25 marker-trait associations with 21 SSRs. The main subpopulations identified from their data set corresponded to the geographic origin of accessions. However, from our results there was every indication that the subdivision of the evaluated accessions could be attributed to differences in the cultivation system. In the case of Agrama *et al.* (2007), the difference could be due to bias in favor of *japonica* as against *indica* accessions, for, from the 103 rice accessions, 49 corresponded to *japonica* accessions, whereas 25 were classified as of an admixed ancestry, mainly between those with an *indica* and *temperate japonica* background.

In a comparative analysis of yield components, with data from 2004 and 2005, a difference was noted, from one year to the next, in the marker-trait association of yield-related genes, as detected by SSR markers. Quantitative agronomic traits, such as yield, are especially affected by the interaction of both genotype and environment, an understanding of how this interaction is controlled being a basis

for defining breeding strategies that would improve genetic gains within these traits.

Two markers (RM38 and RM267) were previously detected by QTL mapping. RM38 was related to yield when assaying 190 lines in an *indica* x *japonica* double-haploid population, genotyped with 179 SSR markers for agronomic traits in a two-year replicated-field experiment (Jiang *et al.*, 2004). This finding was corroborated by Marri *et al.* (2005) who also found the RM38 marker related to a yield QTL (R^2 of 7.99), by using an interspecific population of 251 advanced backcross families (*O. rufipogon* x *O. sativa*), genotyped with 80 SSR markers. Cho *et al.* (2003) identified the association of RM267 with a yield QTL detected in an interspecific backcross F_2 population (*O. rufipogon* x *O. sativa*). A recent study in rice, dealt with the association between markers previously identified as linked at QTLs (Agrama *et al.*, 2007). With this base, QTL analysis can be considered a pre-requisite for distinguishing markers related to important traits, by constituting, together with complementary association studies using highly divergent accession panels from core collections, a means of identifying the different alleles of these markers and attributing phenotypic weights to each.

No difference in allelic effects on yield was identified through association analysis, when considering markers previously associated to this trait in rice. This may be due to the complex inheritance of the YLD trait, as not only a single allele, but also epistatic alleles are involved. Haplotypes related to highest performance may be identified through association analysis, and, an additional advantage over QTL mapping analysis, values can be attributed to alleles present in a given panel of individuals, whereby the most favorable combinations can be traced by breeders in all elite lines and cultivars. Consequentially, there is an increase in the capacity of novel sets in detecting marker-trait associations, even of alleles with minor or modest phenotypic effects (Risch and Zhang, 1996). In QTL linkage analysis, on the contrary, only a maximum of two alleles per locus are involved (diploid individual), with detection being restricted to the size of the effect and the presence of contrasts between genitor alleles in the desired trait.

As to grain quality traits, RM1 was associated to MR in the 2005 field experiment, and to yield QTL in previous works (Yu *et al.*, 1997; Brondani *et al.*, 2002; Septiningsih *et al.*, 2003), thereby indicating its location in a genomic region, and thus requiring detailed analysis, in order to identify genes and alleles of agronomic interest. In Brazil, rice breeders consider MR to be a very important trait, since cultivars, wherein the percentage of intact grains falls below 60%, are considered to be economically of less value. As to RM190 (*Waxy* gene), which is related to expression of the granule-bound starch-synthase enzyme, and is largely responsible for amylose content in rice grains (Ayres *et al.*, 1997), association to AC was reported in seven different segregating populations when applying QTL linkage

analysis (He *et al.*, 1999; Tan *et al.*, 1999; Lanceras *et al.*, 2000; Septiningsih *et al.*, 2003; Zhou *et al.*, 2003; Aluko *et al.*, 2004; Fan *et al.*, 2005). AC was significantly related to RM190 through association analysis in both panels of upland and lowland accessions, throughout the two years.

The confirmation of an association between the RM190 marker and AC, when using core collection accessions, is additional evidence of efficiency in applying association analysis to gene identification. The significant results from Kruskal-Wallis testing, comprising experimental years and accession panels, implies that at least one of the alleles, exerted a discriminating effect on amylose content. Nevertheless, in spite of the lack of complete correspondence between RM190 alleles and amylose content, as to experimental-year and accession panel, some indication of a correlation for alleles 105 and 107 bp to intermediate amylose content, in the panel of lowland accessions and by experimental year, was found. On evaluating a panel of 89 non-waxy accessions, Ayres *et al.* (1997) identified seven alleles from a SSR marker in the *Wx* gene, whence four were correlated with different patterns of amylose content (high, intermediate and low), thus defining approximately 83% of AC variation. Bao *et al.* (2006) when examining a panel of 499 non-waxy accessions with RM190, identified ten alleles that together accounted for nearly 90% of AC variation. The difference of allelic correlation to amylose content classes recorded in the present work, Ayres *et al.* (1997) and Bao *et al.* (2006), could be due to the different composition of accession panels, since Ayres *et al.* (1997) analyzed a set of rice accessions from the U.S.A. with a narrower genetic base when compared to the ERiCC accession panel. Bao *et al.* (2006), when comparing the ERiCC accession panel, analyzed a panel of germplasm accessions with low diversity for the *Waxy* SSR marker, since from the ten alleles identified, only two accounted for a frequency of approximately 83%. The evaluation of germplasm accessions with a narrower genetic basis may be reflected in a low variation in modifier genes that could influence AC, particularly at the *Waxy* locus. This low variation in rice genomes, even in traits controlled by epistatic alleles, may restrict phenotypic trait variability. On the other hand, a wider genetic basis, as found in ERiCC accessions, could increase AC phenotypic variation, due to the allelic variability in genes located upstream of the *Waxy* gene in the starch synthesis route. This would reduce the power of a marker assisted selection in pre-breeding and breeding programs, if based solely on *Waxy* gene alleles, thereby requiring, as mandatory, the study of allelic variation in transcripts from the starch synthesis route in rice grains. As with Ayres *et al.* (1997), RM190 alleles with no relation to any specific class of amylose content were observed in the present work.

Starch makes up for 90% of polished grain in rice, and AC is recognized as one of the most important components in rice grain products (Bao *et al.*, 2006). The use of marker

assisted selection for this trait would be of help in breeding programs, since accessions could be genotyped at an early stage for a trait that is normally evaluated after harvesting. However, the lack of correspondence in RM190 alleles to specific AC classes, as well as rice accessions with similar AC values, maybe showing differences in rice eating and textural qualities (Bao *et al.*, 2006), indicate the need for further studies before MAS implementation. Currently in the Embrapa breeding program, AC analysis is an annual, routine procedure in hundreds of inbred lines. In advanced elite inbred lines, the result of this analysis is being correlated with the direct determination of gel consistency and pasting viscosity by a panel of trained panelists, by using a small quantity of grains cooked in Petri dishes, thereby providing precise standards for defining grain quality.

This study demonstrated the feasibility of conducting association analysis together with germplasm characterization of a rice core collection using SSR markers. It also facilitated the identification of markers related to yield, panicle number, milled-head rice and amylose content, in a panel of genetically unrelated ERiCC accessions. Quantitative traits, such as yield, have complex gene and allele interactions, and studies to dissect this trait may start from QTL analysis, due to consolidated statistics and higher resolution potential. The expected high linkage disequilibrium of rice inbred lines and cultivars, although facilitating the detection of marker-trait associations, makes gene identification more difficult, since LD spans many thousands of base pairs. However, for breeding purposes the correlations detected by association analysis may be sufficient for marker assisted selection and mining alleles related to important traits in germplasm collections.

Acknowledgments

We gratefully acknowledge the Ph.D. scholarship for Tereza Borba and financial support from CNPq.

References

- Abadie T, Cordeiro CMT, Fonseca JR, Alves RBN, Burle ML, Brondani C, Rangel PHN, Castro EM, Silva HT, Freire MS, *et al.* (2005) Constructing a rice core collection for Brazil. *Pesq Agropec Bras* 40:129-136.
- Agrama HA, Eizenga GC and Yan W (2007) Association mapping of yield and its components in rice cultivars. *Mol Breed* 19:341-356.
- Aluko G, Martinez C, Tohme J, Castano C, Bergman C and Oard JH (2004) QTL mapping of grain quality traits from the interspecific cross *Oryza sativa* x *O. glaberrima*. *Theor Appl Genet* 109:630-639.
- Ayres NM, McClung A, Larkin PD, Bligh HFJ, Jones CA and Park WD (1997) Microsatellites and a single nucleotide polymorphism differentiate apparent amylose classes in an extended pedigree of US rice germplasm. *Theor Appl Genet* 94:773-781.
- Bao JS, Corke H and Sun M (2006) Microsatellites, single nucleotide polymorphisms and a sequence tagged site in starch-

- synthesizing genes in relation to starch physicochemical properties in nonwaxy rice (*Oryza sativa* L.). *Theor Appl Genet* 113:1185-1196.
- Breseghele F and Sorrells ME (2006a) Association mapping of kernel size and milling quality in wheat (*Triticum aestivum* L.) cultivars. *Genetics* 172:1165-1177.
- Breseghele F and Sorrells ME (2006b) Association analysis as a strategy for improvement of qualitative traits in plants. *Crop Sci* 46:1323-1330.
- Brondani C, Rangel PHN, Brondani RPV and Ferreira ME (2002) QTL mapping and introgression of yield related traits from *Oryza glumaepatula* to cultivated rice (*Oryza sativa*) using microsatellite markers. *Theor Appl Genet* 104:1192-1203.
- Brondani RPV and Grattapaglia D (2001) Cost-effective method to synthesize a fluorescent internal DNA standard for automated fragment sizing. *Biotechniques* 31:793-800.
- Buntjer JB, Sorensen AP and Peleman JD (2005) Haplotype diversity: The link between statistical and biological association. *Trends Plant Sci* 10:466-471.
- Campos H (1983) *Estatística Experimental Não-Paramétrica*. 4ª edição. ESALQ, Piracicaba, 349 pp.
- Cho YC, Suh JP, Choi IS, Hong HC, Baek MK, Kang KH, Kim YG, Ahn SN, Choi HC, Hwang HG, *et al.* (2003) QTLs analysis of yield and its related traits in wild rice relative *Oryza rufipogon*. *Treat Crop Res* 4:19-29.
- Cruz CD (1997) *Programa Genes: Aplicativo Computacional em Genética e Estatística*. 1ª edição. Editora UFV, Viçosa, 442 pp.
- Fan CC, Yu XQ, Xing YZ, Xu CG, Luo LJ and Zhang Q (2005) The main effects, epistatic effects and environmental interactions of QTLs on the cooking and eating quality of rice in a doubled-haploid line population. *Theor Appl Genet* 110:1445-1452.
- Ferreira ME and Grattapaglia D (2006) *Genética de associação em plantas*. In: Borém A and Caixeta, ET (eds) *Marcadores Moleculares*. Editora UFV, Viçosa, pp 273-306.
- Flint-García AS, Thornsberry JM and Buckler ES (2003) Structure of linkage disequilibrium in plants. *Annu Rev Plant Biol* 54:357-374.
- Garris AJ, McCouch SR and Kresovich S (2003) Population structure and its effect on haplotype diversity and linkage disequilibrium surrounding the *xa5* locus of rice (*Oryza sativa* L.). *Genetics* 165:759-769.
- Garris AJ, Tai TH, Coburn J, Kresovich S and McCouch S (2005) Genetic structure and diversity in *Oryza sativa* L. *Genetics* 169:1631-1638.
- Hardy OJ and Vekemans X (2002) SPAGeDi: A versatile computer program to analyse spatial genetic structure at the individual or population levels. *Mol Ecol Notes* 2:618-620.
- He P, Li SG, Qian Q, Ma YQ, Li JZ, Wang WM, Chen Y and Zhu LH (1999) Genetic analysis of rice grain quality. *Theor Appl Genet* 98:502-508.
- Jannink JL, Bink CAMM and Jansen RC (2001) Using complex plant pedigrees to map valuable genes. *Trends Plant Sci* 6:337-342.
- Jiang GH, Xu CG, Li XH and He YQ (2004) Characterization of the genetic basis for yield and its component traits of rice revealed by doubled haploid population. *Yi Chuan Xue Bao* 31:63-72.
- Juliano B (1979) Amylose analysis - A review. In: *Chemical Aspects of Rice Grain Quality*. IRRI, Los Banos, pp 251-260.
- Khush GS (1997) Origin, dispersal, cultivation and variation of rice. *Plant Mol Biol* 35:25-34.
- Kraakman ATW, Niks RF, van den Berg PMMM, Stam P and van Eeuwijk FA (2004) Linkage disequilibrium of yield and yield stability in modern spring barley cultivars. *Genetics* 168:435-446.
- Lanceras JC, Huang ZL, Naivikul O, Vanavichit A, Ruanjaichon V and Tragoonrun S (2000) Mapping of genes for cooking and eating qualities in Thai jasmine rice (KDML105). *DNA Res* 7:93-101.
- Lu H, Redus MA, Coburn JR, Rutger JN, McCouch SR and Tai TH (2005) Population structure and breeding patterns of 145 U.S. rice cultivars based on SSR marker analysis. *Crop Sci* 45:66-76.
- Marri PR, Sarla N, Reddy LV and Siddiq EA (2005) Identification and mapping of yield and yield related QTLs from an Indian accession of *Oryza rufipogon*. *BMC Genet* 6:1-14.
- Patron NJ, Smith AM, Fahy BF, Hylton CM, Naldrett MJ, Rosnagel BG and Denyer K (2002) The altered pattern of amylose accumulation in the endosperm of low-amylose barley cultivars is attributable to a single mutant allele of granule-bound starch synthase I with a deletion in the 5'-non-coding region. *Plant Physiol* 130:190-198.
- Peleman JD and van der Voort JR (2003) Breeding by design. *Trends Plant Sci* 8:330-334.
- Pritchard JK, Stephens M and Donnelly P (2000) Inference of population structure using multilocus genotype data. *Genetics* 155:945-959.
- Remington DL, Thornsberry JM, Matsuoka Y, Wilson LM, Whitt SR, Doebley J, Kresovich S, Goodman MM and Buckler ES (2001) Structure of linkage disequilibrium and phenotypic associations in the maize genome. *Proc Natl Acad Sci USA* 98:11479-11484.
- Risch N and Zhang H (1996) Mapping quantitative trait loci with extreme discordant pairs, sampling considerations. *Am J Hum Genet* 58:836-843.
- Rostoks NT, Ramsay L, Mackenzie K, Cardle L, Bhat PR, Roose ML, Svensson JT, Stein N, Varshney RK, Marshall DF, *et al.* (2006). Recent history of artificial outcrossing facilitates whole-genome association mapping in elite inbred crop varieties. *Proc Natl Acad Sci USA* 103:18656-18661.
- Septiningsih EM, Prasetyono J, Lubis E, Tai TH, Tjubaryat T, Moeljopawiro S and McCouch S (2003) Identification of quantitative trait loci for yield and yield components in an advanced backcross population derived from the *Oryza sativa* variety IR64 and the wild relative *O. rufipogon*. *Theor Appl Genet* 107:1419-1432.
- Skot L, Humphreys MO and Armstead I (2005) An association mapping approach to identify flowering time genes in natural populations of *Lolium perenne* (L.). *Mol Breed* 15:233-245.
- Storey JD (2002) A direct approach to false discovery rates. *J R Stat Soc Ser B* 64:479-498.
- Tan YF, Li JX, Yu SB, Xing YZ, Xu CG and Zhang Q (1999) The three important traits for cooking and eating quality of rice grains are controlled by a single locus in a elite rice hybrid, Shanyou 63. *Theor Appl Genet* 99:642-648.
- Yu JG, Pressoir WH, Briggs I, Bi IV, Yamasaki M, Doebley JF, McMullen MD, Gaut BS, Nielsen DM, Holland JB, *et al.* (2006) A unified mixed-model method for association map-

ping that accounts for multiple levels of relatedness. *Nat Genet* 38:203-208.

Yu SB, Li JX, Xu CG, Tan YF, Gao YJ, Li XH, Zhang Q and Maroof MAS (1997) Importance of epistasis as the genetic basis of heterosis in an elite rice hybrid. *Proc Natl Acad Sci USA* 94:9226-9231.

Zhou PH, Tan YF, He YQ, Xu C and Zhang Q (2003) Simultaneous improvement for four quality traits of Zhenshan 97, an elite parent of hybrid rice, by molecular marker-assisted selection. *Theor Appl Genet* 106:326-331.

Zhu C, Gore M, Buckler ES and Yu J (2008) Status and prospects of association mapping in plants. *Plant Genome* 1:5-20.

Zondervan KT and Cardon LR (2004) The complex interplay among factors that influence allelic association. *Nat Rev Genet* 5:89-100.

Internet Resources

Belkhir K, Borsa P and Chikhi L (2001) *Genetix* v. 4.05.2. Université de Montpellier.
<http://www.univ-montp2.fr/~genetix/genetix/genetix.htm> (May 5, 2004).

Goudet J (2002) *FSTAT*, v. 2.9.3.2.

<http://www.unil.ch/izea/software/fstat.html> (February 5, 2002).

R Development Core Team. <http://www.r-project.org> (January 4, 2006).

Tassel software, v. 1.9.6. <http://www.maizegenetics.net> (December 2, 2006).

van Hintum TJJ, Brown AHD, Spillane C and Hodgkin T (2000) Core collections of plant genetic resources. http://www.bioversityinternational.org/publications/pubfile.asp?ID_PUB=614 (December 2, 2006).

Supplementary Material

The following online material is available for this article:

Table S1. Identification of ERiCC accessions, their origin and common name.

Table S2. Identification and information on SSR Markers used for ERiCC evaluation

This material is made available as part of the online article from <http://www.scielo.br.gmb>.

Associate Editor: Everaldo Gonçalves de Barros

License information: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Supplemental Table 1. Identification of the ERiCC accessions, their origin and common name.

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
1	CNA0000082	L	Suriname	AWINI
2	CNA0000122	L	India	ARC-10666
3	CNA0000586	L	Sri Lanka	BG 090-2
4	CNA0000692	L	India	BKN 6820-6-3-2
5	CNA0000754	L	Peru	CHANCAY
6	CNA0000798	L	Colombia	CICA 4
7	CNA0000923	L	Mexico	C 79-272-4-1-2-3-10
8	CNA0000950	L	India	IET 0355
9	CNA0000952	L	Pakistan	CR 36-148
10	CNA0001419	L	Colombia	COLOMBIA 1
11	CNA0001420	L	Colombia	CARREON
12	CNA0001423	L	Colombia	TAPURIPA-161
13	CNA0001467	L	India	IET 2881
14	CNA0002222	L	Lao	KH.YOUANE (V.T.A 13)
15	CNA0002246	L	Lao	KH.NGANH TAMAY (SVA 33)
16	CNA0002253	L	Lao	KH.KHAO BAY (VT-A 25)
17	CNA0002258	L	Thailand	KU 56-3
18	CNA0002293	L	Thailand	KU 94-2
19	CNA0002416	L	Liberia	LAC 12
20	CNA0002437	L	Liberia	LAC 28
21	CNA0002480	L	Cameroon	M 40
22	CNA0002482	L	Cameroon	M 44
23	CNA0002529	L	Philippines	MEHR
24	CNA0002672	L	Thailand	NAHNG PAYA 132
25	CNA0002871	L	Portugal	RIZZOTO 159
26	CNA0003195	L	Egypt	GZ 944-5-2-2
27	CNA0003196	L	Egypt	GZ 809-4-1-2
28	CNA0003005	U	India	110281
29	CNA0003241	L	India	PAU 41-306-2-1-PR 405
30	CNA0003411	L	Colombia	CICA 8
31	CNA0003417	L	Japan	PI 294351
32	CNA0003446	L	Philippines	IR 34
33	CNA0003452	L	-	CHIANUNG SEN 25

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
34	CNA0003569	L	India	KAU 2110
35	CNA0003591	L	India	TNAU 2686-1
36	CNA0003602	L	India	UPR 79-23
37	CNA0003665	L	Taiwan	KAOHSIUNG SEN YU 104
38	CNA0003668	L	Philippines	MTU 7029
39	CNA0004308	L	USA	CAROLINA SP 407
40	CNA0004552	L	Colombia	CR 1113
41	CNA0004566	L	Colombia	METICA 1
42	CNA0004576	L	Colombia	ZENIT
43	CNA0004579	L	Colombia	CAMPONI
44	CNA0004625	L	Colombia	TETEP
45	CNA0004629	L	Philippines	IR 50
46	CNA0005014	L	China	WU 10 B
47	CNA0005015	L	China	YAR AI ZHAO B
48	CNA0005016	L	China	ZHENSHAN 97 A
49	CNA0005477	L	China	K ASKHAM 36/14
50	CNA0005478	L	China	SZU MAIO
51	CNA0005853	L	Russia	WIR 5621
52	CNA0006910	L	Indonesia	MANINJAU
53	CNA0006943	L	Philippines	IR 54 R
54	CNA0006955	L	Italy	KORAL
55	CNA0006961	L	Italy	VITRO
56	CNA0007408	L	Colombia	WC 0144
57	-	L	Suriname	ELONI
58	-	L	Suriname	CIWINI
59	-	L	Suriname	CESWONI
60	-	L	Philippines	IR8
61	-	L	Philippines	IR36
62	-	L	USA	LEBONNET
63	-	L	India	BASMATI 370
64	-	L	Japan	NOURIN MOCHI
65	-	L	Japan	MINAMI HATA MOCHI
66	-	L	Japan	TOMOE MOCHI
67	-	L	Japan	MOGAMI CHIKANARI
68	-	L	Colombia	ORYZICA LHANOS4
69	-	L	Colombia	ORYZICA 1
70	-	L	Colombia	CICA 7
71	-	L	Colombia	CICA 9
72	-	L	Philippines	HUAN-SEN-GO
73	-	L	-	5287
74	-	L	-	RAMTULASI
75	CNA0000482	U	USA	BLUEBONNET
76	CNA0001006	U	USA	DAWN
77	CNA0002524	U	France	MOROBEREKAN
78	CNA0003287	U	France	IREM 123-2-1
79	CNA0003288	U	France	IREM 293-B

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
80	CNA0003289	U	France	IREM 247
81	CNA0003362	U	France	IRAT 142
82	CNA0003375	U	France	IRAT 13
83	CNA0003395	U	France	IRAT 141
84	CNA0003397	U	France	IRAT 144
85	CNA0003403	U	Nigeria	TOX 490-3-108-D1-B-B
86	CNA0004193	U	France	IREM 656
87	CNA0004428	U	France	N.7384 [RPL X DANIELA]
88	CNA0004463	U	France	N.7441 [CA 435 X TAINUNG CHUEN 2]
89	CNA0004480	U	France	IRAT 124
90	CNA0004487	U	France	MAKOUTA
91	CNA0004543	U	Nigeria	TOX 1012-12-3-1
92	CNA0004617	U	Nigeria	TOX 1011-4-2
93	CNA0004640	U	Nigeria	TOX 1785-19-18
94	CNA0004697	U	France	N.2583
95	CNA0004752	U	France	IRAT 122
96	CNA0004759	U	Nigeria	TOX 514-16-101-1
97	CNA0004788	U	Nigeria	TOX 503-4-115-B-B
98	CNA0004796	U	Nigeria	TOX 516-28-10B-D2-B-B
99	CNA0005277	U	Nigeria	TOX 1858-114
100	CNA0005326	U	Nigeria	TOX 1780-8-5
101	CNA0005334	U	Nigeria	TOX 1871-29
102	CNA0005358	U	Nigeria	TOX 1858-101
103	CNA0005970	U	Nigeria	FAROX 299
104	CNA0005972	U	Nigeria	FAROX 301
105	CNA0005994	U	Nigeria	TOX 995-208-1-101
106	CNA0006034	U	Nigeria	ITA 150
107	CNA0006035	U	Nigeria	ITA 225
108	CNA0006572	U	France	IREM 195
109	CNA0006574	U	France	IRAT 112
110	CNA0006940	U	USA	LEMONT
111	CNA0006941	U	USA	NEW BONNET
112	CNA0008092	U	France	L 141
113	CNA0008093	U	France	L 285
114	CNA0008411	U	USA	BLUE BELLE
115	CNA0008412	U	USA	BLUEBONNET 50
116	CNA0008432	U	USA	LACASSINE
117	CNA0008545	U	Colombia	CT 11216-10-12-B-BRM-10
118	CNA0009102	U	Colombia	CT10006-7-2-M-5-1P-3
119	CNA0009113	U	Colombia	CT10037-9-4-M-1-1P-2-M
120	CNA0009115	U	Colombia	CT11632-3-3-M
121	CNA0009123	U	Colombia	CT11891-3-3-3-M
122	CNA0009124	U	Colombia	CT13364-7-1
123	CNA0009139	U	Colombia	CT13366-15-4
124	CNA0009154	U	Colombia	CT13370-2-M

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
125	CNA0009197	U	Colombia	CT13377-8-4
126	CNA0009199	U	Colombia	CT13381-1-3
127	CNA0009223	U	Colombia	CT13569-5-7
128	CNA0009227	U	Colombia	CT13570-3-2
129	CNA0009240	U	Colombia	CT13572-6-2
130	CNA0009280	U	Colombia	CT13573-11-2
131	CNA0009319	U	Colombia	CT13579-3-4
132	CNA0009364	U	Colombia	CT13581-5-2
133	CNA0009415	U	Colombia	CT13582-11-4
134	CNA0009561	U	Colombia	CT13584-12-9
135	CNA0009591	U	Colombia	CT13585-12-3
136		U	-	B6149F-MR-19
137	-	U	Philippines	B8503-TB-19-B-3
138	-	U	-	BR4742-B-19-23
139	-	U	India	CUTACK 4
140	-	U	Philippines	IR65907-188-1-B
141	-	U	France	IRAT 10
142	-	U	Philippines	KATAKTARA
143	-	U	Philippines	TB154E-TB-2
144	-	U	Philippines	TB47H-MR-11-51-3
145	-	U	Philippines	YN1905-UUL-62
146	-	U	Philippines	YN906-UUL 65
147	-	U	China	YUNLU N 1
148	-	U	China	YUNLU N 7
149	-	L	EMBRAPA-CNPAF/ Brazil	RS16PL12-35-1-B
150	-	L	EMBRAPA-CNPAF/ Brazil	RS16PL1-34-4-B
151	-	L	EMBRAPA-CNPAF/ Brazil	RS16PL5-12-6-B
152	-	L	EMBRAPA-CNPAF/ Brazil	RS16PL12-10-1-B
153	CNA0001416	L	EMBRAPA-CNPAF/ Brazil	IPSL 0574
154	CNA0001117	L	EEPG/ Brazil	EEPG-1-169
155	-	L	EMBRAPA-CPACT/ Brazil	BRS AGRISUL
156	-	L	EMBRAPA-CPACT/ Brazil	BRS BOJURU
157	-	L	EMBRAPA-CNPAF/ Brazil	DIAMANTE
158	-	L	EMBRAPA-CNPAF/ Brazil	BRS FORMOSO
159	-	L	EMBRAPA-CNPAF/ Brazil	MARAJÓ
160	-	L	EMBRAPA-CNPAF/ Brazil	BRS JABURU
161	-	L	EMBRAPA-CNPAF/ Brazil	BRS BIGUA
162	-	L	EMBRAPA-CNPAF/ Brazil	RIO GRANDE
163	-	L	EPAGRI/ Brazil	EPAGRI 107
164	-	L	EPAGRI/ Brazil	EPAGRI 108
165	-	L	EPAGRI/ Brazil	SCS 111
166	-	L	EPAGRI/ Brazil	SCS 112
167	CNA0001337	L	IPEACO/ Brazil	IPEACO-SL 1969
168	CNA0001339	L	IPEACO/ Brazil	IPEACO-SL 0769
169	CNA0001344	L	IPEACO/ Brazil	IPEACO-SL 1469

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
170	CNA0001407	L	IPEACO/ Brazil	IPSL 2070
171	CNA0001413	L	IPEACO/ Brazil	IPSL 0970
172	CNA0001414	L	IPEACO/ Brazil	IPSL 0570
173	CNA0001106	L	IRGA/ Brazil	EEA 405
174	CNA0001107	L	IRGA/ Brazil	EEA 401
175	CNA0001108	L	IRGA/ Brazil	EEA 404
176	CNA0001109	L	IRGA/ Brazil	EEA 406
177	-	L	IRGA/ Brazil	IRGA 409
178	-	L	IRGA/ Brazil	IRGA 413
179	-	L	IRGA/ Brazil	IRGA 416
180	-	L	IRGA/ Brazil	IRGA 417
181	-	L	IRGA/ Brazil	IRGA 418
182	-	L	IRGA/ Brazil	IRGA 419
183	-	L	IRGA/ Brazil	IRGA 420
184	CNA0006129	L	EMPASC/ Brazil	EMPASC 103
185	CNA0006130	L	EMPASC/ Brazil	EMPASC 104
186	CNA0001118	U	EEPG/ Brazil	EEPG-1-269-FURNAS
187	CNA0000963	U	IPEACO/ Brazil	AMARELAO X GUEDES
188	CNA0000969	U	IPEACO/ Brazil	HONDURAS X MATAO
189	CNA0000976	U	IPEACO/ Brazil	SATURNO X PRATAO PRECOCE
190	CNA0000994	U	IPEACO/ Brazil	ESAV X MATAO
191	CNA0001347	U	IPEACO/ Brazil	IPEACO-SL 2270
192	CNA0001350	U	IPEACO/ Brazil	IPEACO-SL 1970
193	CNA0004098	U	EMBRAPA-CNPAF/ Brazil	XINGU
194	CNA0004120	U	EMBRAPA-CNPAF/ Brazil	RIO PARAGUAY
195	CNA0004121	U	EMBRAPA-CNPAF/ Brazil	GUARANI
196	CNA0004141	U	EMBRAPA-CNPAF/ Brazil	TRIUNFO
197	CNA0004172	U	EMBRAPA-CNPAF/ Brazil	GUAPORE
198	CNA0004206	U	EMBRAPA-CNPAF/ Brazil	ARAGUAIA
199	CNA0004748	U	EMBRAPA-CNPAF/ Brazil	CUIABANA
200	CNA0005180	U	EMBRAPA-CNPAF/ Brazil	TANGARA
201	CNA0006187	U	EMBRAPA-CNPAF/ Brazil	CAIAPO
202	CNA0006701	U	EMBRAPA-CNPAF/ Brazil	CARAJAS
203	CNA0007024	U	EMBRAPA-CNPAF/ Brazil	CNAX 1503-12-9-4-B
204	CNA0007119	U	EMBRAPA-CNPAF/ Brazil	AIMORE
205	CNA0007706	U	EMBRAPA-CNPAF/ Brazil	CONFIANCA
206	CNA0008070	U	EMBRAPA-CNPAF/ Brazil	PRIMAVERA
207	CNA0008711	U	EMBRAPA-CNPAF/ Brazil	SOBERANA
208	CNA0005673	U	IAC/ Brazil	IAC 81-176
209	CNA0005901	U	IAC/ Brazil	URUCUI
210	CNA0006170	U	IAC/ Brazil	LS 85-125
211	CNA0006174	U	IAC/ Brazil	LS 85-158
212	CNA0007799	U	IAC/ Brazil	IAC 1191
213	CNA0006406	U	IAC/ Brazil	LS 86-68
214	CNA0004168	U	IAPAR/ Brazil	L 80-68

	CNPAF Germplasm Bank Identification Number	Cultivation System*	Origin**	Common Name
215	CNA0005672	U	IAPAR/ Brazil	L 82-192
216	CNA0006666	U	IAPAR/ Brazil	A12-286-1-1
217	CNA0006672	U	IAPAR/ Brazil	A8-204-1-1
218	CNA0008309	U	IAPAR/ Brazil	L 92-342
219	CNA0006413	U	IAPAR/ Brazil	L 85-20
220	CNA0006422	U	IAPAR/ Brazil	IAPAR L 99-98
221	CNA0005650	U	IPEACO/ Brazil	IPEACO 11-P
222	CNA0005660	U	IPEACO/ Brazil	IPEACO 77-P
223	CNA0003281	U	Brazil	IRAT 177
224	CNA0005342	U	Brazil	RIO VERDE
225	CNA0002123	U	EEPG/ Brazil	JAPONES X PRAIANA
226	-	U	IAC/ Brazil	IAC 201
227	-	U	IAC/ Brazil	IAC 202
228	-	U	IAC/ Brazil	IAC 165
229	-	U	IAPAR/ Brazil	IAPAR 9
230	-	U	IAPAR/ Brazil	IAPAR 62
231	-	U	IAPAR/ Brazil	IAPAR 63
232	CNA0003490	U	Brazil	MEARIN
233	CNA0004078	U	Brazil	-
234	CNA0004243	U	Brazil	-
235	CNA0005975	U	Brazil	-
236	CNA0006030	U	EMBRAPA-CNPAF/ Brazil	-
237	CNA0007425	U	EMBRAPA-CNPAF/ Brazil	CANASTRA
238	CNA0007937	U	EMBRAPA-CNPAF/ Brazil	PROGRESSO
239	CNA0008172	U	EMBRAPA-CNPAF/ Brazil	BONANCA
240	CNA0008305	U	EMBRAPA-CNPAF/ Brazil	CARISMA
241	CNA0008533	U	EMBRAPA-CNPAF/ Brazil	MARAVILHA
242	CNA0008540	U	EMBRAPA-CNPAF/ Brazil	TALENTO

* L = lowland accessions ; U = upland accessions. For comparison with other studies, the lowland accessions used in this study were considered as indica and the upland accessions, as japonica [Khush GS (1997) Origin, dispersal, cultivation and variation of rice. Plant Molecular Biology 35:25-34].

** EMBRAPA – CNPAF = Empresa Brasileira de Pesquisa Agropecuária – Centro Nacional de Pesquisa (Brazil); IAC = Instituto Agrônomo de Campinas (Brazil); IPEACO = Instituto de Pesquisa Agropecuária do Centro Oeste (Brazil); IAPAR = Instituto Agrônomo do Paraná (Brazil); EMPASC = Empresa de Pesquisa Agropecuária de Santa Catarina (Brazil); EEPG = Estação Experimental de Ponta Grossa (Brazil); IRGA = Instituto Rio Grandense do Arroz (Brazil); EPAGRI = Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina (Brazil)

Supplemental Table 2. SSR Markers used for the evaluation of the ERiCC accessions (Data on “Marker concentration for multiplexed PCR”, “Missing data”, “Number of alleles” and “PIC” were obtained in this study).

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
1	RM229	129	5.9	HEX	106–131	11	0.83	(TC) ₁₁ (CT) ₅ C ₃ (CT) ₅	11	Chen et al. (1997)
	RM207	129	3.1	FAM	84–158	23	0.86	(CT) ₂₅	2	Chen et al. (1997)
2	RM222*	129	10	FAM	199–215	12	0.72	(CT) ₁₈	10	Chen et al. (1997)
	RM1	71.67	0.8	FAM	76–119	23	0.75	(AG) ₂₆	1	Panaud et al. (1996)
	RM248	150	4.1	HEX	80–104	11	0.81	(CT) ₂₅	7	Chen et al. (1997)
	RM38	71.67	5.7	HEX	246–278	18	0.86	(GA) ₁₆	8	Chen et al. (1997)
3	RM5	71.67	1.8	FAM	109–133	9	0.73	(GA) ₁₄	1	Panaud et al. (1996)
	RM253	71.67	9.8	HEX	89–119	11	0.75	(GA) ₂₅	6	Chen et al. (1997)
4	RM11	71.67	3.6	HEX	123–143	11	0.78	(GA) ₁₇	7	Panaud et al. (1996)
	RM13*	129	4.6	FAM	129–164	15	0.72	(GA) ₆ -GA ₁₆	5	Panaud et al. (1996)
5	RM252	129	11	FAM	193–277	23	0.62	(CT) ₁₉	4	Chen et al. (1997)
	RM277	129	1.8	FAM	110–126	8	0.58	(GA) ₁₁	12	Temnykh et al. (2000)
	RM204	129	6.9	HEX	106–194	32	0.87	(CT) ₄₄	6	Chen et al. (1997)
6	RM231	129	10.5	FAM	170–196	10	0.78	(CT) ₁₆	3	Chen et al. (1997)
	RM304	129	10	HEX	82–198	21	0.84	(GT) ₂ (AT) ₁₀ (GT) ₃₃	10	Temnykh et al. (2000)

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
7	OG17	71.67	2.3	FAM	114-180	24	0.77	(CT) ₂₄	2	Brondani et al. (2001)
	OG44*	71.67	2.8	HEX	152-182	15	0.80	(CT) ₄ -23bp-(CT) ₂₂ -(GT) ₄ -(GC) ₆	3	Brondani et al. (2001)
8	OG106*	129	10	FAM	200-250	22	0.90	(CT) ₂₇	9	Brondani et al. (2001)
	4879	129	9.5	HEX	108-150	10	0.80	(AAT) ₈	4	Rangel (2005)
9	4653*	129	3.9	FAM	80-170	19	0.85	(AAG) ₂₅	12	Rangel (2005)
	RM224	129	4.1	HEX	124-158	10	0.81	(AAG) ₈ -(AG) ₁₃	11	Chen et al. (1997)
10	RM223*	129	3.6	FAM	139-163	15	0.64	(CT) ₂₅	8	Chen et al. (1997)
	RM257	71.67	4.1	HEX	130-192	22	0.90	(CT) ₂₄	9	Chen et al. (1997)
11	OG61*	129	2.3	FAM	96-152	28	0.88	(CT) ₁₈	5	Brondani et al. (2001)
	RM234	129	0.8	HEX	126-156	13	0.78	(CT) ₂₅	7	Chen et al. (1997)
12	RM103	198	3.0	FAM	300-350	4	0.64	(GAA) ₅	6	Temnykh et al. (2000)
	RM334	258	3.3	FAM	120-210	14	0.86	(CTT) ₂₀	5	Temnykh et al. (2000)
	RM309	198	1.9	HEX	150-180	14	0.79	(GT) ₁₃	12	Temnykh et al. (2000)
13	RM286	138	2.1	FAM	80-140	14	0.84	(GA) ₁₆	11	Temnykh et al. (2000)
	RM276	180	3.9	HEX	75-160	17	0.84	(AG) ₈ -A ₃ -(GA) ₃₃	6	Temnykh et al. (2000)

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
14	RM125	129	0.3	FAM	100-150	13	0.58	(GCT) ₈	7	Temnykh et al. (2000)
	RM171	71.67	7.5	FAM	310-344	8	0.59	(GATG) ₅	10	Akagi et al. (1996)
	RM484	71.67	7.5	HEX	290-300	3	0.53	(AT) ₉	10	Temnykh et al. (2001)
	RM341	71.67	1.0	HEX	110-180	19	0.76	(CTT) ₂₀	2	Temnykh et al. (2000)
15	RM169*	258	0.3	FAM	150-200	14	0.75	(GA) ₁₂	5	Temnykh et al. (2000)
	RM216	129	0.5	HEX	120-155	10	0.81	(CT) ₁₈	10	Chen et al. (1997)
16	RM152	258	0.3	FAM	150-180	10	0.58	(GGC) ₁₀	8	Akagi et al. (1996)
	RM267	129	0.3	HEX	125-175	11	0.49	(GA) ₂₁	5	Temnykh et al. (2000)
17	RM264	180	3.3	FAM	125-195	22	0.83	(GA) ₂₇	8	Temnykh et al. (2000)
	RM263	138	0.8	HEX	164-220	13	0.80	(CT) ₃₄	2	Chen et al. (1997)
18	RM335	180	3.3	FAM	100-170	26	0.90	(CTT) ₂₅	4	Temnykh et al. (2000)
	RM287	138	8.0	HEX	90-140	12	0.75	(GA) ₂₁	11	Temnykh et al. (2000)
19	RM151	258	2.3	FAM	197-220	7	0.74	(TA) ₂₃	1	Akagi et al. (1996)
	RM271	138	4.3	FAM	75-125	6	0.80	(GA) ₁₅	10	Temnykh et al. (2000)
	RM336*	129	3.3	HEX	100-200	6	0.79	(CTT) ₁₈	7	Temnykh et al. (2000)

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing Data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
20	RM51	270	14.0	FAM	120-150	9	0.77	(GA) ₁₃	7	Chen et al. (1997)
	RM332	129	12.0	HEX	140-190	6	0.62	(CTT) ₅ -(CTT) ₁₄	11	Temnykh et al. (2000)
21	RM310	116.67	3.3	HEX	80-130	9	0.84	(GT) ₁₉	8	Temnykh et al. (2000)
	RM9	100	4.5	FAM	120-200	14	0.80	(GA) ₁₅ GT(GA) ₂	1	Panaud et al. (1996)
	RM22*	132	1.3	HEX	150-230	7	0.55	(GA) ₂₂	3	Panaud et al. (1996)
22	RM55	150	3.2	HEX	200-245	18	0.89	(GA) ₁₇	3	Chen et al. (1997)
	RM266	210	8.8	FAM	100-150	13	0.79	(GA) ₁₉	2	Temnykh et al. (2000)
	RM278	258	1.3	HEX	120-160	17	0.89	(GA) ₁₇	9	Temnykh et al. (2000)
23	RM85	180	2.3	FAM	80-120	8	0.79	(TGG) ₅ (TCT) ₁₂	3	Temnykh et al. (2000)
	RM117	258	4.2	FAM	190-220	12	0.71	(AG) ₇	12	Temnykh et al. (2000)
	RM302	180	1.3	HEX	70-210	8	0.82	(GT) ₃₀ (AT) ₈	1	Temnykh et al. (2000)
24	RM206	258	7.8	FAM	115-210	12	0.80	(CT) ₂₁	11	Chen et al. (1997)
	RM159	258	3.1	HEX	220-280	9	0.75	(GA) ₁₉	5	Temnykh et al. (2000)
25	OG99	71.67	2.3	FAM	110-160	7	0.19	(CT) ₄ TT(CT) ₄ CC(CT) ₃ – 16pb – (CT) ₈	3	Brondani et al. (2001)
	OG60	71.67	0.5	HEX	100-160	14	0.74	(CT) ₂₁ (TCC) ₂	4	Brondani et al. (2001)

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing Data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
26	RM261	138	4.5	FAM	100-130	15	0.59	C ₉ (CT) ₈	4	Chen et al. (1997)
	RM3	210	3.2	HEX	115-160	4	0.70	(GA) ₂ GG(GA) ₂₅	6	Panaud et al. (1996)
27	RM220	150	2.4	FAM	90-135	9	0.69	(CT) ₁₇	1	Chen et al. (1997)
	RM280	258	4.1	HEX	140-190	6	0.88	(GA) ₁₆	4	Temnykh et al. (2000)
	RM7	210	3.0	FAM	160-195	15	0.85	(GA) ₁₉	3	Panaud et al. (1996)
28	RM144*	258	9.5	FAM	200-250	15	0.75	(ATT) ₁₁	11	Temnykh et al. (2000)
	RM243	129	4.2	FAM	85-130	10	0.75	(CT) ₁₈	1	Chen et al. (1997)
	RM317	258	5.0	HEX	120-180	9	0.76	(GC) ₄ (GT) ₁₈	4	Temnykh et al. (2000)
29	RM190*	150	0.5	FAM	100-130	11	0.75	(CT) ₁₁	6	Akagi et al. (1996)
	RM205	210	0.5	HEX	115-170	14	0.78	(CT) ₂₅	9	Chen et al. (1997)
	RM70	258	11.5	FAM	150-215	11	0.80	(ATT) ₃₃	7	Chen et al. (1997)
30	RM14	270	10.0	HEX	160-120	9	0.81	(GA) ₁₈	1	Panaud et al. (1996)
	5371*	258	8.4	FAM	150-210	5	0.83	(GAA) ₁₄	12	Rangel (2005)
	RM178	258	4.3	HEX	110-140	8	0.69	(GA) ₅ (AG) ₈	5	Temnykh et al. (2000)
31	RM210	129	1.3	FAM	115-175	15	0.87	(CT) ₂₃	8	Chen et al. (1997)
	RM240	129	0.3	HEX	120-160	5	0.47	(CT) ₂₁	2	Chen et al. (1997)

Panel	SSR Marker	Marker concentration for multiplexed PCR (nM)	Missing Data (%)	Fluorochrome	Allele size range (bp)	Number of alleles	PIC	Motif	Chromosome	References**
32	RM161	198	4.4	HEX	150-200	8	0.59	(AG) ₂₀	5	Temnykh et al. (2000)
	RM135	198	1.3	FAM	100-140	5	0.55	(CGG) ₁₀	3	Akagi et al. (1996)
	OG10*	258	0.0	HEX	40-130	21	0.81	(CT) ₂₉	9	Brondani et al. (2001)
33	RM30	270	4.9	HEX	70-120	15	0.64	(AG) ₉ A(GA) ₁₂	6	Panaud et al. (1996)
	RM154	258	5.3	HEX	140-210	14	0.91	(GA) ₂₁	2	Akagi et al. (1996)
	RM119	258	7.0	FAM	150-185	6	0.75	(GTC) ₆	4	Temnykh et al. (2000)
34	RM311	270	5.0	HEX	160-200	8	0.74	(GT) ₃ (GTAT) ₈ (GT) ₅	10	Temnykh et al. (2000)
	RM53*	258	10.0	FAM	168-208	7	0.76	(GA) ₁₄	2	Chen et al. (1997)
35	RM12	129	2.1	FAM	150-190	7	0.41	(GA) ₂₁	12	Panaud et al. (1996)
	RM19	258	11.0	HEX	200-270	11	0.72	(ATC) ₁₀	12	Panaud et al. (1996)
Average	-	-	-	-	-	12.4	0.75	-	-	-

* Markers with presence of PCR artifacts (especially stutters)

**References

- Akagi H, Yokozeki Y, Inagaki A, Fujimura T (1996) Microsatellite DNA markers for rice chromosomes. *Theor Appl Genet* 93: 1071-1077.
- Brondani C, Brondani RPV, Rangel PHN, Ferreira ME (2001) Development and mapping of *Oryza glumaepatula* derived microsatellite markers in the interspecific cross *O. glumaepatula* x *O. sativa*. *Hereditas* 134: 59-71.
- Chen X, Temnykh S, Xu Y, Cho YG, McCouch SR (1997) Development of a microsatellite framework map providing genome-wide coverage in rice (*Oryza sativa* L.). *Theor Appl Genet* 95: 553-567
- Panaud O, Chen X, McCouch SR (1996) Development of microsatellite markers and characterization of simple sequence length polymorphism (SSLP) in rice (*Oryza sativa* L.). *Mol Gen Genet* 252: 597-607
- Rangel, PN. Construção de mapa de ligação e mapeamento de QTLs utilizando marcadores SSRs, ESTs e SNPs em cruzamento interespecífico *Oryza glumaepatula* X *Oryza sativa*. Thesis, Universidade Federal de Goiás, 111 pp.
- Temnykh S, Park WD, Ayres NM, Cartinhour S, Hauck N, Lipovich L, Cho YG, Ishii T, McCouch SR (2000) Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theor Appl Genet* 100: 697-712