

Characterization of organic matter's quality evaluated through ^{13}C NMR of the Serra do Sudeste's soils in Rio Grande do Sul

Roberta Jeske Kunde^{a*}; Daiane Carvalho dos Santos^b; Otávio dos Anjos Leal^b; Wilson Tadeu Lopes da Silva^c; Marcos Flávio Silva Corba^d; Clenio Nailto Pillon^e

^aUCPel, Rua Félix da Cunha, 412, CEP 96010-000 Pelotas, RS, Brasil; ^bUFPel, Campus Universitário s/nº, Caixa Postal 354, CEP 96010-900, Capão do Leão, RS, Brasil; ^cEmbrapa Instrumentação Agropecuária Rua XV de Novembro 1452, CEP 13560-97, São Carlos, SP, Brasil; ^dEmbrapa Pecuária Sul, BR 153 Km 603, Caixa Postal, 242, CEP 96401-970, Bagé, RS, Brasil; ^eEmbrapa Clima Temperado, Rodovia BR 392, km 78, Caixa Postal, 403, CEP 96001-970, Pelotas, RS, Brasil.

*E-mail: roberta_kunde@hotmail.com

Keywords: density physical fractionation, natural field

Introduction

Due to the absence of significant research work on the characterization of soil organic matter (SOM) in the region of Serra do Sudeste do Rio Grande do Sul, the purpose of this work is to evaluate qualitatively the physical fractions of a Lithic Leptosol (RL), Haplic Planosol (SX) and Haplic Vertisol (VX) under a native grassland in Torrinhás, 2nd District of Pinheiro Machado, RS.

The soil sampling was performed on the soil's layer from 0.000 to 0.025 m and part of the undisturbed samples were used for the density physical fractionation of the SOM [1], in which a solution 2.0 mg m⁻³ of sodium polytungstate was used. The energy dispersion per ultrasound was of 357 J mL⁻¹ for RL, 374 J mL⁻¹ for the SX and 461 J mL⁻¹ for VX.

The samples of the free light fraction (FLF) and occluded light fraction (OLF) were treated with aqueous solution of HF 10%, to perform the analysis on the ^{13}C nuclear magnetic resonance (NMR).

Results and Discussion

The spectra were similar for the different soil types analyzed (Figure 1). In the region between 25-35 ppm, the C alkyl signal is from methylene, derived from long chain aliphatic. In the region of C O-alkyl (60-110 ppm), one can observe two distinct peaks; the one at 72-75 ppm is assigned to cellulose and the other at 105 ppm, derived from hemicellulose and other carbohydrates. The signal between 160-230 ppm corresponds, in part, to the carboxyl groups of organic acids [2].

In the analyzes of different soil types, there was a predominance of substituted C alkyl groups (C O-alkyl/C di-O-alkyl + C N-alkyl/C methoxyl), whose proportion varied from 36% to

37%, followed by C alkyl groups (26% to 36%). The aromatic structures (C aryl + C phenyl) contributed approximately 13% of the composition of the samples, the carboxyl groups with 8% of and carbonyl groups with 6% (Table 1).

Considering that the substituted C alkyl groups indicate the presence of polysaccharides and proteins structures in the SOM. These compounds are easily decomposed by microorganisms, therefore it can be inferred that the presence of these structures in the OLF indicates the process of physical protection of SOM by occlusion into stable aggregates.

Analyzing the chemical composition of FLF of the SOM, the proportions of substituted C alkyl, varied from 37% to 51% for RL and VX, respectively. For C alkyl, these ratios ranged 26-32% for VX and RL, respectively. Possibly the greatest abundance of substituted C alkyl in the FLF of VX can be explained by differences in chemical composition (recalcitrance) of crop residues and its relationship with the environment where decomposition occurs.

Considering the aromatic structures (C aryl + C phenyl), the highest percentage was found in RL (17%). These higher proportions, found in substituted C alkyl groups in relation to the aromatic structures, show that carbohydrates are the main organic constituents of the soil, although they have low molecular recalcitrance. This fact associated with the aromatic compounds that correspond to smaller proportions is an indication that the colloidal and physical protection of SOM are overlapping the magnitude of protection by recalcitrance [3].

In OLF, similar proportion was observed among the functional groups of carbon in comparison to FLF. However, the average of the different types of soils were decreased by

3% in FLF compared to the OLF for the substituted C alkyl groups of and the C alkyl.

These variations indicate an increase in the degree of decomposition of OLF [4].




Figure 1. ^{13}C NMR spectroscopy of free light fraction (FLF) and occluded light fraction (OLF) of organic matter in a Lithic Leptosol (RL), Haplic Planosol (SX) and Haplic Vertisol (VX), under natural field in the layer of 0.000 to 0.025 m.

Table 1. Percentage distribution of the C's functional groups determined by ^{13}C NMR spectroscopy in the free light fraction (FLF) and occluded light fraction (OLF) of organic matter in a Lithic Leptosol (RL), Haplic Planosol (SX) and Haplic Vertisol (VX), under field natural in the layer of 0.000 to 0.025 m.

Soils *	Distribution of types of C / chemical shifts (ppm)						
	C alkyl 0 - 45	C N-alkyl C Methoxyl 45 - 60	C O-alkyl C di-O-alkyl 60 - 110	C-aryl 110 - 140	C phenolic 140 - 160	C carboxyl 160 - 185	C carbonyl 185 - 230
				FLL			
RL	32	10	27	13	4	8	6
SX	30	10	33	8	5	8	5
VX	26	9	42	7	3	7	6
				FLO			
RL	36	10	27	8	4	8	7
SX	33	9	29	9	5	9	6
VX	27	15	32	8	4	8	6

*RL - Lithic Leptosol; SX - Haplic Planosol e VX - Haplic Vertisol.

Conclusions

The occluded light fraction has a higher proportion of more recalcitrant compounds in relation to the free light fraction, suggesting a more advanced stage of humification, regardless of the soil's type.

In the fractions analyzed there is not evidence of charred material, due to the absence of fires in this study area.

Acknowledgements

We thank Temperate Climate Embrapa (CPACT) and Agricultural Instrumentation Embrapa (CNPDIA) for supporting the research and for laboratory evaluations.

¹Conceição, P.C.; Boeni, M.; Dieckow, J.; Bayer, C.; Mielniczuk, J. 2008. *Rev. Bras. Ci. Solo*, 32,541-549.

²Kögel-Knabner, I. 2002. *Soil Biol. Bioch.*, 34,139-162.

³Diekow, J. *Estoque e qualidade da matéria orgânica do solo em função de sistemas de culturas e adubação nitrogenada no sistema plantio direto*. 2003. 164f. Tese (Doutorado) - Universidade Federal do Rio Grande do Sul, Porto Alegre.

⁴Baldock, J.A. et al. 1992. *Biogeochemistry*, Netherlands, 16,1- 42.