

Yacare Caiman *Caiman yacare*

Zilca Campos¹, Alfonso Q. Llobet², Carlos I. Piña³ and William E. Magnusson⁴

¹Embrapa Pantanal, Caixa Postal 109, Corumbá, MS, Brazil, 79320-900 (zilca@cpap.embrapa.br);

²Calle Chochi #2755, Condominio Guembe, Bloque 5, Depto. 1B Barrio San Luis, Santa Cruz, Bolivia (allobet@cotas.com.bo);

³CIC y TTP-CONICET, Dr. Matteri y España Diamante CP 3105, Argentina (cidcarlos@infoaire.com.ar);

⁴Instituto Nacional de Pesquisas da Amazonia/CPEC, Caixa Postal 478, Manaus, Amazonas 69011-970, Brazil (bill@inpa.gov.br)

Common Names: Yacaré, Jacaré, Lagarto, Yacaré negro, Yacaré tinga, jacaré-do-Pantanal, Pantanal Caiman

Important Synonyms: *Caiman crocodilus yacare* (see Velasco and Ayarzagüena 2009)

Range: Argentina, Bolivia, Brazil, Paraguay



Figure 1. Distribution of *Caiman yacare*.

Conservation Overview

CITES: Appendix II

CSG Action Plan:

Availability of survey data: Adequate

Need for wild population recovery: Generally Low.
Moderate in some places.

Potential for sustainable management: Highest

2009 IUCN Red List: LRLc (Lower Risk, least concern). Widespread and numerous populations throughout its distribution, although in some places still in process of recovery; IUCN 2009) (last assessed in 1996).

Principal threats: Habitat destruction, hydroelectric dam, siltation, illegal hunting.

Ecology and Natural History

The Yacare caiman is found in the lowlands of northern and eastern Bolivia and western Brazil, from the Amazon southwards through the Guaporé/Madeira and Paraguay/Paraná River systems and into northern Argentina. Morphologically and ecologically, this species is similar to the common caiman (*Caiman crocodilus crocodilus*), and integrates with that subspecies along a large area (probably more than 1000 km) of the Madeira River in Amazonia. As with the common caiman, *C. yacare* is found in a wide spectrum of habitat types. Most ecological studies have been carried out in the Pantanal region of southern Brazil (Crawshaw and Schaller 1980; Schaller and Crawshaw 1982; Cintra 1986, 1988; Campos 1993, 2003; Catto and Amaro 1994; Campos and Magnusson 1995; Campos *et al.* 1995, 2003, 2005, 2006a, 2006b; Santos *et al.* 1996; Pinheiro *et al.* 1997; Rebelo *et al.* 1997).

More recent studies in Bolivia included assessing the distribution and population status of the species, monitoring harvests, and gathering additional data to improve the national program of management (Pacheco 1993; Godshalk 1994; Rebolledo-Garin and Tapia-Arauz 1994; Llobet 1996; Llobet and Goitia 1997; Llobet and Aparicio 1999; Rios 2003; Cisneros 2005; Llobet and CIPTA 2006; Peters 2006). In Brazil, aerial surveys of caimans and nests have been carried out in the Pantanal since 1991 (Mourão *et al.* 1994, 2000; Coutinho and Campos 1996). Nesting habitats were also monitored during incubation and hatching periods

(Borges and Filho 1993; Campos and Magnusson 1995), and the effect of water level on nest production and hatchling survivorship (Coutinho and Campos 1996; Coutinho *et al.* 2001).



Figure 2. Adult *C. yacare*. Photograph: Jemeema Carrigan.

Terrestrial movement in groups was described for caimans in the Pantanal, in response to disturbance (Campos *et al.* 2003). This behavior may make caimans vulnerable if intensive hunting were it to resume (Campos 2003).

Resolution of the systematic relationships within the very widespread *C. crocodilus* complex is needed. Extensive surveys and specimen collection in southern Brazil, northern Bolivia and Paraguay suggest a very complex gradient of morphological features between *C. c. crocodilus* and the *C. yacare*. As there are no fixed differences between the two taxa, they have to be defined geographically, rather than morphologically, which obviously poses legal difficulties if an individual crosses the hypothetical line and changes “species”.

Morphological and genetic analyses to resolve the relationships between these taxa are incomplete and inconclusive, mainly because data from the zone of hybridization between the two taxa was not included (Brazaitis *et al.* 1998; Busack and Pandya 2001). Researchers from the Federal University of Amazonas, Instituto Nacional de Pesquisa da Amazônia and Embrapa Pantanal have returned to the Pantanal-Gauporé-Madeira region to quantify the variation and define the geographical boundaries of the postulated species and their variants (Vasconcelos and Campos 2007; Hrbek *et al.* 2008). This study on morphological and genetic variation of the caimans should resolve the issue by the end of 2009.

Caiman yacare is still, and always has been, listed technically as a full species (Medem 1983). Some authors do not recognize it as being distinct from the common caiman and use the subspecies *C. c. yacare*. Recent molecular data (Godshalk 2006, 2008a,b) provide evidence for long-term biological separation of large populations of caimans in Central and South America, which may require a new

approach to the systematics of the caiman group. The historic forces, driven largely by plate tectonics, have subdivided this neotropical species complex into at least four evolutionarily significant units (ESUs) that are reproductively isolated. The Andes, in conjunction with a rugged desert coastline, have effectively isolated *C. c. fuscus* to the west and north through Central America to Honduras. Two similar but reproductively separate basin populations of *C. c. crocodilus* have evolved in the Orinoco and Amazon River drainages, and *C. yacare* populates the southern Amazon and Paraguay-Parana basins (Godshalk 2008a).

More work is needed to map the boundaries of the various caiman populations and to characterize their phylogeography. Contact zones between the populations may prove particularly valuable in resolving some taxonomic issues, not only between the four major source populations, but also the fringe populations along the Atlantic coast from Guyana to Brazil.

Yacare caiman are mound nesters, with egg-laying usually peaking in the middle of the wet season, from December to February. Clutch size typically consists of 22-35 eggs, with a maximum of 42 eggs. It seems that habitat type could influence clutch size (Campos 1993; Zambrana *et al.* 2008). The natural tendency of females to guard their nests is apparently influenced by human hunting pressure (Crawshaw 1987), which results in decreased nest attendance and a lower hatching success.



Figure 3. Hatchling *C. yacare*. Photograph: Charlie Manolis.

Conservation and Status

Most of what is written about the conservation of the common caiman applies equally well to *C. yacare*. Basic survey information is available for this species in all countries where it occurs. This information has resulted principally from programs using the species since late 1990s (Bolivia), or new programs whose goal is to utilize the species in near future (Argentina and Brazil). Some basic information for these programs resulted from a series of CITES-sponsored surveys in southern Brazil, Bolivia, and Paraguay, as well as surveys sponsored by local governments, scientific institutions and

NGOs working within these countries. During the early 1990s, populations of Yacare caiman were considered to be somewhat depleted in all four Range States, principally due to widespread illegal hunting during the 1970s and 1980s. Poaching is no longer a problem throughout much of this species' range, but habitat destruction, construction of hydroelectric dams, and siltation of rivers continue to affect caiman populations, particularly in Brazil.

Due to its small size at maturity, ability to adapt to a wide variety of habitat types, and learned wariness, *C. yacare* is particularly resilient to hunting pressure. Studies in Brazil suggest that illegal hunting did not seriously impact populations (Mourão *et al.* 1996). Surveys in Paraguay (Scott *et al.* 1990; King *et al.* 1994) and Argentina (Siroski 2004; Piña *et al.* 2008) indicate that extensive populations remain or have recovered from previous exploitation, since the closure of uncontrolled hunting and implementation of export controls.

Commercially-oriented management programs are in place in three of the four Range States for *C. yacare*. Paraguay suspended its national program in 2003. Cropping is permitted in Bolivia, where hides are currently exported under a CITES quota (50,000 skins).

The conservation and use program for *C. yacare* in Bolivia began in 1997 with few evaluations of its population, and a small experimental harvest. During the following years, between 30,000 (1999) and 59,000 (2003) individuals were harvested annually through a management model based on eco-regional population sizes. From 2004, a large group of institutions and specialists participated in the re-design of the program to adapt it to the national reality, and strengthen the regulatory and administrative framework by means of a monitoring plan. In addition, standardized sampling protocols were designed; monitoring of harvests commenced, and a new classification of water bodies was made. Additionally, a new basis of technical data (counts, harvests, habitats) and administrative data (users, farms, quotas history, contraventions) was developed, and geographically assessed together through a GIS. This database is continuously updated with new information. Parallel to this activity, management plans were suggested for development in Indigenous Lands (TCOs = Spanish acronym) and Protected Areas to increase local community participation in Yacare caiman use, and carrying out a group of activities directed to strengthen local communal structures.

Currently, there are two management plans in operation, and another 12 are under evaluation, with the intention of covering some 8,503,400 ha of the *C. yacare* distribution in Bolivia (Llobet *et al.* 2004; Llobet and Bello 2008). Between 1999 and 2007, more than 298,000 individuals were harvested. The initial national export quota (1999) was established at 36,500 individuals, but it fluctuated from 45,000 to 50,000 animals per year between 2001 and 2007 (Llobet and Bello 2008).

In Brazil, hunting of wild animals is not permitted. However, ranching and farming are allowed (Coutinho *et al.* 1998), and

new legislation for the management of the Pantanal caiman - including a head-starting program - is under evaluation by the Federal and State Governments. The aim is to implement an adult caiman harvesting quota in the Brazilian Pantanal. This program is based principally on proposals by Coutinho (2000).

In Argentina, four ranching programs are utilising *C. yacare* (Larriera *et al.* 2008); two in Formosa Province, one in Corrientes Province, and one in Chaco Province. Together, they produce around 15,000 skins per year. Recently, a harvesting program was approved by the Federal and State Governments, to collect wild adult caimans from Bañado la Estrella, Formosa, northern Argentina.



Figure 4. Captive adult *C. yacare* in Bolivia. Photograph: Charlie Manolis.



Figure 5. Captive adult *C. yacare* in Bolivia. Photograph: Charlie Manolis.

Priority Projects

High priority

1. **Strengthen the monitoring of wild populations and harvesting in Bolivia:** The development of a series of management plans has already produced more reliable

data on the species. However, it is important to ensure a comprehensive monitoring program to follow the implementation of these management plans, and to make corrections through an adaptive management approach.

2. **Reassessment and implementation of a management program in Paraguay:** Following the self-imposition of a CITES moratorium on Paraguay in 2003 a reassessment of the Paraguay National Program and the design of a reliable monitoring program for harvesting of wild *C. yacare* are considered important to ensure the sustainable use of the species, specially now that the partial lifting of the moratorium is ready to start. Even though Paraguayan *C. yacare* populations do not depend on a management program for their conservation, a weak management program could affect those populations and other programs for the species in the region.
3. **Study of *Caiman* systematics:** The aim is to clarify the complex taxonomic situation of *C. yacare*, develop a map with boundaries for the various caiman populations and to characterize their phylogeography. If an arbitrary geographic definition is maintained, a zone with a low density of caimans should be selected as the limit to reduce the problem of individuals changing taxa as they cross the line. For enforcement purposes, it would be useful to be able to distinguish the taxon via observable meristic characteristics that can be seen in commercial skins, even if this results in two taxa co-existing in the same population in some areas. Studies of morphological and genetic variation in Brazil and others countries should clarify the issues in the near future.

Moderate priority

4. **Control of illegal trade:** Management national programs for the species need to improve law enforcement and enhance the controls in border areas to avoid illegal trade between countries.
5. **Quantification of *C. yacare* distribution on the border between Peru and Bolivia:** The border between Peru and Bolivia appears to be the limit of Yacare caiman distribution from the south and also the limit for *C. crocodilus* distribution from the north. No studies have been carried out to assess the northwestern distribution limits for *C. yacare*, or the existence of a sympatric area for both species in this region between Peru and Bolivia.
6. **Long-term ecological studies in the Pantanal, Brazil:** Like the llanos of Venezuela and Colombia, the Pantanal is a large seasonally inundated savanna that offers excellent research opportunities for the study of caiman population dynamics. Monitoring programs of the populations via aerial survey and radio-telemetry have been made in the Pantanal region, and habitat destruction, such as deforestation, siltation, hydroelectric dam, agriculture, and mining have been evaluated in the medium-term. This part of the research and conservation of caiman in Brazil should be continued and implemented in other countries

(Bolivia, Paraguay and Argentina). The analysis of the effects of global climate change on the biology of the caimans in the Pantanal will be made in the near future. The large population of caiman in the area is a function of the quality of the wetlands habitat, and changes to the flooding regime will affect the distribution and abundance of the species.

7. **Implementation of ranching programs in Bolivia, Paraguay and Brazil:** Brazil has undertaken a very rapid development of captive breeding (farming) of caiman similar to that in Colombia. A number of facilities are registered with the Government and CITES as captive breeding facilities, and are producing skins for export. However, since ranching offers greater positive impact than farming for effective conservation of both, species and habitats, it is recommended that a ranching model be implemented, initially on an experimental basis. The produced information would serve to assess the feasibility of starting ranching programs on a national scale.

References

- Borges, G.A. and Arruda Filho, A. (1993). Avaliocio do potencial de ninhos de *Caiman crocodilus yacare* numa sub-regiao do Pantanal de Pocone-MT. Pp. 55-117 in Zoocria de los Crocodylia. Memorias de la I Reunion Regional de Especialistas en Cocodrilos de la IUCN. IUCN: Gland.
- Busack, S.D., and Pandya, S. (2001). Geographic variation in *Caiman crocodilus* and *Caiman yacare* (Crocodylia: Alligatoridae): Systematic and legal implications. *Herpetologica* 57(3): 294-312.
- Brazaitis, P., Rebelo, G. and Yamashita, C. (1998). The distribution of *Caiman crocodilus crocodilus* and *Caiman yacare* populations in Brazil. *Amphibia-Reptilia* 19: 193-201.
- Campos, Z. (1993). Effect of habitat on survival of eggs and sex ratio of hatchlings of *Caiman crocodilus yacare* in the Pantanal, Brazil. *J. Herpetol.* 27(2): 127-132.
- Campos, Z. (2003). *Caiman crocodilus yacare*. Food-related movement. *Herp. Rev.* 34: 141.
- Campos, Z., Coutinho, M. and Magnusson, W. (2003). Terrestrial activity of caiman in the Pantanal, Brazil. *Copeia* 2003: 628-634.
- Campos, Z., Coutinho, M. and Magnusson, W. (2005). Field body temperatures of caimans in the Pantanal, Brazil. *Herpetol. J.* 15: 97-106.
- Campos, Z., Coutinho, M. and Magnusson, W. (2006b). *Caiman crocodilus yacare* (Pantanal Caiman). *Aestivation. Herp. Rev.* 37: 343-344.
- Campos, Z., Coutinho, M., Mourão, G., Bayliss, P. and

- Magnusson, W. (2006a). Long distance movement by *Caiman crocodilus yacare*: implications for management of the species in the Brazilian Pantanal. *Herp. J.* 16: 123-132.
- Campos, Z. and Magnusson, W. (1995). Relationships between rainfall, nesting habitat and fecundity of *Caiman crocodilus yacare* in the Pantanal, Brazil. *J. Trop. Ecol.* 11: 351-358.
- Campos, Z. and Mourão, G. (1995). *Caiman latirostris* (Broad-snouted caiman) nesting. *Herp. Rev.* 26:203-204.
- Campos, Z., Mourão, G., Coutinho, M. and Abercrombie, C. (1995). Night-light counts, size structures and sex ratios in wild populations of *Caiman crocodilus yacare* in the Brazilian Pantanal. *Vida Silvestre Neotropical* 4(1): 46-50.
- Catto, J.B. and Amato, F.R. (1994). Helminth community structure of the Caiman *Caiman crocodilus yacare* (Crocodylia, Alligatoridae) in the Brazilian Pantanal. *Brazil. J. Vet. Parasitol.* 3: 109-118.
- Cintra, R. (1986). Nascimento de filhotes de *Caiman yacare* (Daudin, 1802) em condicoes semi-naturais no pantanal Matogrossense. *Papeis Avulsos Zool., S. Paulo.* 36: 91-101.
- Cintra, R. (1988). Nesting ecology of the Paraguayan Caiman (*Caiman yacare*) in the Brazilian Pantanal. *J. Herpetol.* 22: 219-222.
- Cisneros, F. (2005). Estado de la población y uso de microhábitat del lagarto (*Caiman yacare*) y caimán negro (*Melanosuchus niger*) en lagunas fluviales en la cuenca baja del río Ichoa del Territorio Indígena y Parque Nacional Isiboro Sécura (Cochabamba, Bolivia). Tesis de Licenciatura, Universidad Mayor de San Simón, Cochabamba, Bolivia.
- Coutinho, M. (2000). Population ecology and the conservation and management of *Caiman yacare* in the Pantanal Brazil. Unpublished PhD Thesis, University of Queensland, Australia.
- Coutinho, M., Campos, Z., Bampi, I. and Dal'Ava, F. (1998). Preliminary report for the management system of Yacare caiman in the Pantanal: a proposal for future research and the development of a monitoring system for wild population subjected to nest harvests. *Ciência e Cultura* 50: 60-64.
- Coutinho, M., Campos, Z., Cardoso, F., Massara, P. and Castro, A. (2001). Reproductive biology and its implication for management of Caiman *caiman yacare* in the pantanal wetland, Brazil. Pp. 229-243 *in* *Biology and Evolution of Crocodylians*. Vol. 2, ed. by G. Grigg, F. Seebacher and C. Franklin. Surrey Beatty and Sons: Sydney.
- Coutinho, M. and Campos, Z. (1996). Effect of habitat and seasonality on the density of caiman (*Caiman crocodilus yacare*) in the southern Pantanal. *J. Trop. Ecol.* 12: 741-747.
- Crawshaw Jr., P.G. (1987). Nesting ecology of the Paraguayan caiman (*Caiman yacare*) in the Pantanal of Mato Grosso, Brazil. Unpublished MSc Thesis, University of Florida, Gainesville, Florida, USA.
- Crawshaw Jr., P.G. and Schaller, G. (1980). Nesting of the Paraguayan caiman (*Caiman yacare*) in Brazil. *Pap. Avulsos de Zoologia* 33: 283-292.
- Godshalk, R.E. (1994). Feasibility of a conservation and sustainable yield plan for the Yacare Caiman, *Caiman crocodilus yacare*, in Bolivia. Unpublished MSc Thesis, University of Florida, Gainesville, Florida, USA.
- Godshalk, R.E. (2006). Phylogeography and conservation genetics of the Yacare caiman (*Caiman yacare*) of South America. Unpublished PhD Thesis, University of Florida, Gainesville, Florida, USA.
- Godshalk, R.E. (2008a). The invasion and subsequent radiation of Caimaninae in the Neotropics, and the relationships of the extant species, with special reference to the Caiman complex. Pp. 352-367 *in* *Crocodyles*. Proceedings of the 19th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Godshalk, R.E. (2008b). The phylogeography of the Yacare caiman, *Caiman yacare*, of central South America. pp. 137-152 *in* *Crocodyles*. Proceedings of the 19th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Hrbek, T., Vasconcelos, W.R., Rebelo, G. and Farias, I.P. (2008). Phylogenetic relationships of south American alligatorids and the Caiman of Madeira River. *J. Exp. Zool.* 309A(10): 588-599.
- IUCN (2009). IUCN Red List of Threatened Species. Ver. 2009.1 (www.iucnredlist.org; viewed 30 September 2009).
- King, F.W., Aquino, A.L., Scott, N. and Palacios, R. (1994). Survey of caimans in Paraguay. Pp. 162-198 *in* *Crocodyles*: Proceedings of the 12th Working Meeting of the IUCN-SSC Crocodile Specialist Group IUCN: Gland.
- Larriera, A., Imhof, A. and Siroski, P. (2008). Estado actual de los programas de conservación y manejo del género Caiman en Argentina. Pp. 141-180 *in* *Contribución al Conocimiento del género Caiman de Sudamérica*, ed. By J. Castroviejo, J. Ayarzagüena and A. Velasco. Publ. Asoc. Amigos de Doñana 18: Seville.
- Llobet, A. (1996). Distribución y abundancia de *Caiman yacare* en la cuenca del río Ichilo. Tesis de Licenciatura.

- Universidad Mayor de San Simón. Cochabamba, Bolivia.
- Llobet, A. and Aparicio, J.K. (1999). Abundancia, estructura poblacional y perspectivas de aprovechamiento de *Caiman yacare* en cinco estancias del Departamento del Beni. Pp. 285-293 in Manejo y Conservación de Fauna Silvestre en América Latina, ed. by T.G. Fang, O.L. Montenegro and R.E. Bodmer. Instituto de Ecología: La Paz, Bolivia.
- Llobet, A. and Bello, A.K. (2008). Programa de conservación y aprovechamiento del yacaré o lagarto (*Caiman yacare*) en Bolivia. Lecciones aprendidas. Pp. 83-118 in Contribución al Conocimiento del Género Caiman de Suramérica, ed. by J. Castroviejo, J. Ayarzagüena and A. Velasco. Publ. Asoc. Amigos de Doñana No. 18: Seville.
- Llobet, A. and CIPTA. (2006). Propuesta de Plan de Manejo: "Aprovechamiento y Manejo Sostenible del Lagarto (*Caiman yacare*) en la Tierra Comunitaria de Origen Tacana, Provincia Iturrealde, Departamento de La Paz". Documento técnico. Wildlife Conservation Society: La Paz, Bolivia.
- Llobet, A. and Goitia, E. (1997). Abundancia y estructura poblacional de *Caiman yacare* en lagunas de la llanura inundable de los ríos Ichilo y Chapare (Bolivia). Revista Boliviana Ecología Conservación Ambiental 2: 39-47.
- Llobet, A., Pacheco, L.F. and Aparicio, J.K. (2004). Analysis of the program of conservation and use of the Spectacled Caiman (*Caiman yacare*) in Bolivia, and recommendations to improve it. Pp. 257-267 in Crocodiles. Proceedings of the 17th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Medem, F. (1983). Los Crocodylia de Sur América. Vol. 2. Univ. Nac. Colombia y Fondo "Francisco José de Caldas": Bogotá.
- Mourão, G., Coutinho, M., Mauro, R., Campos, Z., Tomás, T. and Magnusson, W. (2000). Aerial survey of caiman, marsh deer, and pampas deer in the Pantanal Wetland of Brazil. Biol. Conserv. 92: 175-183.
- Mourão, G. M., Campos, Z., Coutinho, M. and Abercrombie, C. (1996). Biological size structure of illegal harvested and surviving caiman (*Caiman crocodilus yacare*) in the Pantanal - Brazil. Conserv. Biol. 75: 261-265.
- Mourão, G., Campos, Z. and Coutinho, M. (1994). Test of an aerial survey for caiman and other wildlife in the Pantanal, Brazil. Wildl. Soc. Bull. 22: 50-56.
- Pacheco, L.F. (1993). Abundance, distribution, and habitat use of crocodylians in Beni, Bolivia. Unpublished MSc Thesis, University of Florida, Gainesville, Florida, USA.
- Peters, G. (2006). Estado poblacional y lineamientos de manejo para *Caiman yacare* en la Laguna Bolivia y el canal de conexión con el río Ichoa, dentro del Territorio Indígena y Parque Nacional Isiboro Sécure (TIPNIS), Beni, Bolivia. Tesis de Licenciatura, Universidad Católica Boliviana, Cochabamba, Bolivia.
- Pinheiro, M., Mourão, G., Campos, Z. and Coutinho, M. (1997). Influência da temperatura de incubação na determinação do sexo do jacaré (*Caiman crocodilus yacare*). Revista Brasileira Biologia 57: 383-391.
- Piña, C.I., Siroski, P., Simoncini, M. and Príncipe, G. (2008). Status of *Caiman latirostris* and *Caiman yacare* populations in north Argentina. Pp. 219 in Crocodiles. Proceedings of the 19th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Rebello, G.H., Borges, G.A.N., Yamashita, C. and Filho, A.G. (1997). Growth, sex ratio, population structure, and hunting mortality of *Caiman yacare* in the Pantanal, Brazil. Vida Silvestre Neotropical 6: 29-36.
- Rebolledo-Garin, P.P. and Tapia-Arauz, M.C. (1994). La herpetofauna de la Reserva de Vida Silvestre de los Ríos Blanco y Negro: Diversidad, distribución, pautas para su conservación y uso sostenible. Informe preliminar. Fundación Amigos de la Naturaleza (FAN): Santa Cruz, Bolivia.
- Rios, J.N. (2003). Estado de conservación de *Caiman yacare* en áreas de uso de recursos naturales del Territorio Comunitario de Origen (TCO) Tacana, Prov. Iturrealde, Departamento de La Paz. Tesis de Licenciatura, Universidad Mayor de San Andrés. La Paz, Bolivia.
- Schaller, G.B. and Crawshaw, P.G. (1982). Fishing behaviour of the Paraguayan caiman. Copeia 1982: 66-72.
- Scott Jr., N.J., Aquino, A. and Fitzgerald, L.A. (1990). Distribution, habitats and conservation of the caiman (Alligatoridae) of Paraguay. Vida Silvestre Neotropical 2: 43-51.
- Siroski, P. (2004). *Caiman latirostris* and *Caiman yacare* population surveys in Formosa Province, Argentina. Pp. 443-446 in Crocodiles. Proceedings of the 17th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.
- Vasconcelos, W. and Campos, Z. (2007). Geographic variation between Pantanal Caiman (*Caiman crocodilus yacare*) and Amazonian Caiman (*Caiman crocodilus crocodilus*): First Phase. Crocodile Specialist Group Newsletter 26(4): 6-7.
- Zambrana, M.D., Severiche, J. and Llobet, A. (2008). Nesting habits of spectacled caiman (*Caiman yacare*) in Natural Area of Integrated Management (ANMI) San Matías. Pp. 49-60 in Crocodiles. Proceedings of the 19th Working Meeting of the IUCN-SSC Crocodile Specialist Group. IUCN: Gland.