



A new polymorphism in the *Growth and Differentiation Factor 9 (GDF9)* gene is associated with increased ovulation rate and prolificacy in homozygous sheep

B. D. M. Silva*, E. A. Castro[†], C. J. H. Souza[‡], S. R. Paiva[†], R. Sartori[§], M. M. Franco^{†,*}, H. C. Azevedo[¶], T. A. S. N. Silva*, A. M. C. Vieira*, J. P. Neves* and E. O. Melo^{†,*}

*Universidade de Brasília, Brasília, Brazil. [†]Embrapa Recursos Genéticos e Biotecnologia, Brasília, Brazil. [‡]Embrapa Pecuária Sul, Bagé, Brazil. [§]Universidade de São Paulo, São Paulo, Brazil. [¶]Embrapa Tabuleiros Costeiros, Aracaju, Brazil

Summary

Brazilian Santa Inês (SI) sheep are very well-adapted to the tropical conditions of Brazil and are an important source of animal protein. A high rate of twin births was reported in some SI flocks. *Growth and Differentiation Factor 9 (GDF9)* and *Bone Morphogenetic Protein 15 (BMP15)* are the first two genes expressed by the oocyte to be associated with an increased ovulation rate in sheep. All *GDF9* and *BMP15* variants characterized, until now, present the same phenotype: the heterozygote ewes have an increased ovulation rate and the mutated homozygotes are sterile. In this study, we have found a new allele of *GDF9*, named *FecG^E* (Embrapa), which leads to a substitution of a phenylalanine with a cysteine in a conservative position of the mature peptide. Homozygote ewes presenting the *FecG^E* allele have shown an increase in their ovulation rate (82%) and prolificacy (58%). This new phenotype can be very useful in better understanding the genetic control of follicular development; the mechanisms involved in the control of ovulation rate in mammals; and for the improvement of sheep production.

Keywords growth factor, *Ovis aries*, prolificacy.

Some breeds of sheep are naturally prolific and they are very informative for the study of reproductive genetics and physiology. It is postulated that *GDF9* and *BMP15* may form non-covalent homo and heterodimers *in vivo* and, in a species-specific way, modulate the ovulation rate in mammals (Moore *et al.* 2004). The TGF β -family member *BMP15* was the first gene to be associated with increased ovulation rate in Inverdale (*FecX^I* polymorphism) and Hanna (*FecX^H*) sheep (Galloway *et al.* 2000). Soon after, the Booroola variant (*FecB^B*) was found in the *BMP1B* gene of Merino Booroola (Mulsant *et al.* 2001; Souza *et al.* 2001; Wilson *et al.* 2001). The *BMP1B*, together with its partner *BMPRII*, is responsible for the *BMP15* signalling in the ovarian follicles (Moore *et al.* 2003). The last major gene found to be associated with prolific phenotype was *GDF9*, in which a polymorphism (*FecG^H*) found in Cambridge and F700-Belclare sheep is responsible for an increased ovulation rate in heterozygotes and sterility in homozygotes, in a way very similar to all *BMP15* variants (Hanrahan *et al.*

2004; Bodin *et al.* 2007; Martinez-Royo *et al.* 2008; Monteagudo *et al.* 2008).

Besides the *GDF9* and *BMP15* activities during cumulus expansion, oocyte maturation and ovulation (Elvin *et al.* 1999, 2000; Gui & Joyce 2005; Yoshino *et al.* 2006), these two paracrine factors play important roles during many steps of follicular development. They influence follicle growth (Dong *et al.* 1996; Nilsson & Skinner 2002), cumulus and granulosa cell proliferation (Hayashi *et al.* 1999; Gilchrist *et al.* 2006; Spicer *et al.* 2006), cell-survival signalling (Hussein *et al.* 2005; Orisaka *et al.* 2006) and act as modulators of many other growth-factors and endocrine hormones (Juengel *et al.* 2004). As a result of their role in the folliculogenesis, the availability of *GDF9* and *BMP15* polymorphisms can be very useful in the study of animal reproductive genetics and physiology. In this study we have shown, for the first time, a polymorphism in the *GDF9* gene that increases the ovulation rate in homozygotes, without sterility, in sheep.

In this study, 23 ewes (*Ovis aries*) from a Santa Inês (SI) population with a history of multiple births (twin and triplet births) were investigated for SNPs in the *BMP15* and *GDF9* genes. The ewes were genotyped for the Booroola SNP (*FecB^B*) as previously described (Wilson *et al.* 2001). Subsequently, exon 2 of the *GDF9* and *BMP15* genes were

Address for correspondence

E. O. Melo, Embrapa Recursos Genéticos e Biotecnologia PBI sala 7B, PqEB Final W5 Norte, Asa Norte, Brasília 70770-9001, Brazil.
E-mail: eom@cenargen.embrapa.br

Accepted for publication 15 April 2010

screened for SNPs by DNA sequencing of PCR amplicons using the following primers: *GDF9* (forward 5'-GGAGAAAAGGGACAGAAGC; reverse 5'-ACGACAGGTACTACTAGT); and *BMP15* (forward 5'-GGCTGCTTGTCAGTTTGTAC; reverse 5'-GAGCACTTTCAGATTTAA) (see Appendix S1 for details). Seven (GI to GVII) single nucleotide polymorphisms (SNPs) were found in *GDF9* (Table A1, additional data). Only the GVII polymorphism is a non-conservative change in position 345 (phenylalanine to cysteine), which was detected in 43% of the sequenced animals, and it is in the mature peptide of *GDF9*. This polymorphism provokes a change in a residue which is 100% conserved in the sequence of four representative mammalian species (Figure A1 in Appendix S1), and was named *FecG^E* (GenBank FJ429111) according to the previous nomenclature for the high fertility *GDF9* allele (*FecG^H*) (Hanrahan et al. 2004).

To find the frequency of the *FecG^E* polymorphism, a total of 334 animals from a separate flock (Appendix S1) that had not been selected for prolificacy have had their genotypes identified by a PCR-RFLP strategy (Appendix S1). All data about the parturition of these ewes during the period of 2002 to 2008 were collected, and the association between the number of lamb births and genotypes was tested. The genotype distribution and allele frequency were analysed by the Chi-square test. A difference ($P < 0.001$) in the frequency of *FecG^E* and *FecG⁺* alleles, as well as in the genotype distribution, was observed between the randomly selected and the prolificacy-selected flocks (Table 1).

To investigate the association between the *FecG^E* genotypes (E/E, +/E, and +/+) and the ovulation rate, 39 ewes (15 *FecG^{+/+}*, 15 *FecG^{+/E}* and 9 *FecG^{E/E}*) were selected from the genotyped flocks and submitted to oestrus synchronization. The animals were oestrus synchronized twice, with eCG and PGF2alpha-based protocols in a cross-over design (Appendix S1). Eleven days after the last oestrus detection, laparoscopy was performed as previously described (Killen & Caffery 1982) to infer ovulation rate by counting corpora lutea (CL). At the end of the breeding season, pregnancy status was evaluated by ultrasound. All animals submitted to laparoscopy had their *GDF9* and *BMP15* exon 2 sequenced to confirm the *FecG^E* genotyping and to verify that there was no other characterized polymorphism

associated with ovulation rate. The animals were handled in accordance with pertinent Brazilian legislation and following Embrapa's procedures for animal care.

The CL number and the number of lamb births were fitted to the GLM (Generalized Linear Model), where the Poisson distribution was attributed to the ovulation rate, pregnancy and lambing data. In this analysis, the lamb count was considered as response variable, measured for each animal in seven different breeding seasons from 2002 to 2008 (time variable). To measure the influence of genotype over the offspring number through the time, a generalized linear mixed model (GLMM, SAS software) was applied. The offspring number y_{ij} of the i^{th} animal at the j^{th} time was modelled as the per following model:

$$y_{ij}|\gamma_i \sim \text{Poisson}(\mu_{ij})$$

$$\gamma_i \sim N(0, \sigma^2)$$

$$\mu_{ij} = \exp(\beta_0 + \beta_1 X1_i + \beta_2 X2_i + \beta_3 Z_{ij} + \gamma_i)$$

$$\text{Var}(y_{ij}|\gamma_i) = \sigma^2 \mu_{ij},$$

where y_{ij} follows the Poisson distribution conditioned to the random effect for animal γ_i , which was assumed to be normally distributed with variance σ^2 . The expected mean μ_{ij} is a non-linear function of the effects of genotype (β_1 and β_2), time when the counting was made (β_3) and the random effect because of each animal. The variance for y_{ij} irrespective of the random effect γ_i is $\text{Var}(y_{ij}) = \sigma^2 \mu_{ij}$, where the extra (or sub) variation is taken into account. The estimate for σ^2 is 0.1696 (standard error = 0.0087), indicating a strong under-dispersion, but this is correctly modelled by GLMM (see Appendix S1, SAS output in additional data).

The parturition data of the 334 genotyped ewes showed a difference ($P < 0.0001$) in the prolificacy amongst the groups (Table 2). Regarding the ovulation rate, it was greater ($P < 0.001$) in the homozygote (E/E) group, which showed an 82% increase in CL average (2.22 ± 0.12 , Fig. 1a), as well as the highest frequency (96.3%) of multiple-ovulating ewes (Fig. 1b), when compared with +/E and +/+ groups. The heterozygote group (+/E) presented no difference ($P = 0.612$) in CL average (1.34 ± 0.08) or in the frequency (31.8%) of ewes with multiple ovulations (Fig. 1a and b), when compared with the wild-type ewes (1.22 ± 0.11 and 14.6%

Table 1 Genotypic and allelic frequencies of *FecG^E* in Santa Inês flocks.

SI Flock	Genotype	Frequency (N)	Allele	Frequency (N)
Prolific-selected ^a	+/+	0.174 (4)	<i>FecG⁺</i>	0.478 (22)
	+/E	0.609 (14)	<i>FecG^E</i>	0.522 (24)
	E/E	0.217 (5)	–	–
Randomly selected ^b	+/+	0.656 (219)	<i>FecG⁺</i>	0.808 (540)
	+/E	0.305 (102)	<i>FecG^E</i>	0.192 (128)
	E/E	0.0389 (13)	–	–

Distinct letters are different ($P < 0.001$) for genotype distribution.

Table 2 The effect of of *FecG^E* in Santa Inês prolificacy.

SI Flock (F1)	Genotype	Prolificacy of F1 (mean; [95% CI])
Randomly selected	+/+	1.13; [1.11, 1.16] ^a
	+/E	1.44; [1.41, 1.48] ^b
	E/E	1.78; [1.69, 1.87] ^c

Distinct letters are different ($P < 0.001$).

Non-selected SI; $N = 334$ ewes (219 +/+; 102 +/E and 13 E/E), called F1.

Prolificacy = mean of 764 offspring (F2) from the 334 genotyped ewes (F1); separated according to their genotype category.




Figure 1 The effect of genotypes on ovulation. (a) The average number of corpora lutea (CL) per ewe in each genotype: E/E (*FecG^E* in homozygosis) $N = 9$, +/E (*FecG^E* in hereozygosis) $N = 15$ and +/+ (without *FecG^E* allele) $N = 15$. The CL data are presented as mean \pm SE. (b) The frequency of multiple-ovulating ewes (≥ 2 CL) in each genotype as described in (a). (c) The frequency of twinning scored by ultrasonography at the 45th day of gestation in each genotype: E/E ($N = 9$), +/E ($N = 14$) and +/+ ($N = 14$). Groups with different letters differ ($P < 0.001$; Figures 1a and 1b) or ($P = 0.0136$; Figure 1c).

respectively). We observed a genotype effect on the number of twins per ewe ($P = 0.0136$); E/E ewes showed 44% of twin-pregnancy, while no twin-pregnancy was observed in

+/+ ewes (Fig. 1c). Moreover, the E/E ewes presented no observable effect of *FecG^E* other than the increased ovulation rate and twinning.

It has been suggested that increasing multiple births may be an efficient way to improve meat production per ewe, and an increase of 50% in total weight weaned per ewe lambing twins has been reported (Rajab *et al.* 1992). The increase of one extra CL and 58% more lambs born observed in E/E ewes compared with +/+ was a strong evidence of the *FecG^E* effect on ovulation rate control and prolificacy, and represents a new phenotype for *GDF9* in sheep. Our parturition data point to an additive effect for the *FecG^E* allele, despite no difference being observed in the ovulation rate between +/+ and E/+ ewes. However, the allele interactions of *FecG^E* are certainly distinct from the over-dominant behaviour observed in *FecG^H* and all *FecX* alleles described until now. The E/E pregnancy and parturition data confirm that their oocytes were viable and fertile; which correlate with the increased prolificacy (number of lambs/ewe) observed amongst these animals. In this study, for the first time, a new SNP that increased the ovulation rate and prolificacy of homozygote sheep was documented for the *GDF9* gene. This new genetic variant, together with the other documented variants in *GDF9* and *BMP15*, can be very useful to obtain a better understanding of the genetic control of ovulation rate in mammals. Moreover, this major gene variant can be applied in breeding programmes by gene-assisted selection (GAS), aiming towards the improvement of sheep reproductive potential and production. However, further investigation is necessary to shed light on the allelic interactions of the *FecG^E* variant.

Acknowledgements

We thank Dr Margot A. N. Dode for her help in manuscript revision; Alexandre Lim, Rogério Tokarsky and Concepta M. Pimentel for their assistance with the animals; Joseane P. da Silva for statistical assistance. This work was supported by IFS, Sweden (Grant B/4317-1) and CNPq, Brazil (Grant 474619/2007-9).

References

- Bodin L., Di Pasquale E., Fabre S., Bontoux M., Monget P., Persani L. & Mulsant P. (2007) A novel mutation in the *bone morphogenetic protein 15* gene causing defective protein secretion is associated with both increased ovulation rate and sterility in Lacaune sheep. *Endocrinology* **148**, 393–400.
- Dong J., Albertini D.F., Nishimori K., Kumar T.R., Lu N. & Matzuk M.M. (1996) Growth differentiation factor-9 is required during early ovarian folliculogenesis. *Nature* **383**, 531–5.
- Elvin J.A., Clark A.T., Wang P., Wolfman N.M. & Matzuk M.M. (1999) Paracrine actions of growth differentiation factor-9 in the mammalian ovary. *Molecular Endocrinology* **13**, 1035–48.
- Elvin J.A., Yan C. & Matzuk M.M. (2000) Growth differentiation factor-9 stimulates progesterone synthesis in granulosa cells via a

- prostaglandin E2/EP2 receptor pathway. *Proceedings of the National Academy of Sciences of the United States of America* **97**, 10288–93.
- Galloway S.M., McNatty K.P., Cambridge L.M. *et al.* (2000) Mutations in an oocyte-derived growth factor gene (*BMP15*) cause increased ovulation rate and infertility in a dosage-sensitive manner. *Nature Genetics* **25**, 279–83.
- Gilchrist R.B., Ritter L.J., Myllymaa S., Kaivo-Oja N., Dragovic R.A., Hickey T.E., Ritvos O. & Mottershead D.G. (2006) Molecular basis of oocyte-paracrine signalling that promotes granulosa cell proliferation. *Journal of Cell Science* **119**, 3811–21.
- Gui L.M. & Joyce I.M. (2005) RNA interference evidence that growth differentiation factor-9 mediates oocyte regulation of cumulus expansion in mice. *Biology of Reproduction* **72**, 195–9.
- Hanrahan J.P., Gregan S.M., Mulsant P., Mullen M., Davis G.H., Powell R. & Galloway S.M. (2004) Mutations in the genes for oocyte-derived growth factors *GDF9* and *BMP15* are associated with both increased ovulation rate and sterility in Cambridge and Belclare sheep (*Ovis aries*). *Biology of Reproduction* **70**, 900–9.
- Hayashi M., McGee E.A., Min G., Klein C., Rose U.M., van Duin M. & Hsueh A.J. (1999) Recombinant growth differentiation factor-9 (*GDF-9*) enhances growth and differentiation of cultured early ovarian follicles. *Endocrinology* **140**, 1236–44.
- Hussein T.S., Froiland D.A., Amato F., Thompson J.G. & Gilchrist R.B. (2005) Oocytes prevent cumulus cell apoptosis by maintaining a morphogenic paracrine gradient of bone morphogenetic proteins. *Journal of Cell Science* **118**, 5257–68.
- Juengel J.L., Bodensteiner K.J., Heath D.A., Hudson N.L., Moeller C.L., Smith P., Galloway S.M., Davis G.H., Sawyer H.R. & McNatty K.P. (2004) Physiology of *GDF9* and *BMP15* signalling molecules. *Animal Reproduction Science* **82–83**, 447–60.
- Killen I.D. & Caffery G.J. (1982) Uterine insemination of ewes with the aid of a laparoscope. *Australian Veterinary Journal* **59**, 95.
- Martinez-Royo A., Jurado J.J., Smulders J.P. *et al.* (2008) A deletion in the *bone morphogenetic protein 15* gene causes sterility and increased prolificacy in Rasa Aragonesa sheep. *Animal Genetics* **39**, 294–7.
- Monteagudo L.V., Ponz R., Tejedor M.T., Lavina A. & Sierra I. (2008) A 17 bp deletion in the *Bone Morphogenetic Protein 15* (*BMP15*) gene is associated to increased prolificacy in the Rasa Aragonesa sheep breed. *Animal Reproduction Science* **110**, 139–46.
- Moore R.K., Otsuka F. & Shimasaki S. (2003) Molecular basis of bone morphogenetic protein-15 signaling in granulosa cells. *Journal of Biological Chemistry* **278**, 304–10.
- Moore R.K., Erickson G.F. & Shimasaki S. (2004) Are *BMP-15* and *GDF-9* primary determinants of ovulation quota in mammals? *TRENDS in Endocrinology and Metabolism* **15**, 356–61.
- Mulsant P., Lecerf F., Fabre S. *et al.* (2001) Mutation in bone morphogenetic protein receptor-IB is associated with increased ovulation rate in Booroola Merino ewes. *Proceedings of the National Academy of Sciences of the United States of America* **98**, 5104–9.
- Nilsson E.E. & Skinner M.K. (2002) Growth and differentiation factor-9 stimulates progression of early primary but not primordial rat ovarian follicle development. *Biology of Reproduction* **67**, 1018–24.
- Orisaka M., Orisaka S., Jiang J.Y., Craig J., Wang Y., Kotsuji F. & Tsang B.K. (2006) Growth differentiation factor 9 is antiapoptotic during follicular development from preantral to early antral stage. *Molecular Endocrinology* **20**, 2456–68.
- Rajab M.H., Cartwright T.C., Dahm P.F. & Figueiredo E.A. (1992) Performance of three tropical hair sheep breeds. *Journal of Animal Science* **70**, 3351–9.
- Souza C.J., MacDougall C., Campbell B.K., McNeilly A.S. & Baird D.T. (2001) The Booroola (*FecB*) phenotype is associated with a mutation in the *bone morphogenetic receptor type 1 B* (*BMPRI1B*) gene. *Journal of Endocrinology* **169**, R1–6.
- Spicer L.J., Aad P.Y., Allen D., Mazerbourg S. & Hsueh A.J. (2006) Growth differentiation factor-9 has divergent effects on proliferation and steroidogenesis of bovine granulosa cells. *Journal of Endocrinology* **189**, 329–39.
- Wilson T., Wu X.Y., Juengel J.L. *et al.* (2001) Highly prolific Booroola sheep have a mutation in the intracellular kinase domain of bone morphogenetic protein IB receptor (*ALK-6*) that is expressed in both oocytes and granulosa cells. *Biology of Reproduction* **64**, 1225–35.
- Yoshino O., McMahon H.E., Sharma S. & Shimasaki S. (2006) A unique preovulatory expression pattern plays a key role in the physiological functions of *BMP-15* in the mouse. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 10678–83.

Supporting Information

Additional supporting information may be found in the online version of this article.

Appendix S1 Additional data, materials and methods.

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.