

Butterflies and moths (Insecta: Lepidoptera) associated with *erva-mate*, the South American Holly (*Ilex paraguariensis* St. Hil.), in Rio Grande do Sul, Brazil

Edegar Fronza^{1,2*}, Alexandre Specht³ and Elio Corseuil⁴

1 Pontifícia Universidade Católica do Rio Grande do Sul, Faculdade de Biociências. Avenida Ipiranga, n° 6681, Partenon. CEP 90619-900. Porto Alegre, RS, Brasil.

2 Universidade de Caxias do Sul, Campus da Região dos Vinhedos, Laboratório de Biologia, Departamento de Ciências Exatas e da Natureza. Alameda João Dal Sasso, n° 800, Universitário, Caixa Postal 32. CEP 95700-000. Bento Gonçalves, RS, Brasil.

3 Embrapa Cerrados. BR 020, Km 18. Caixa Postal 08223. CEP 73310-970. Planaltina, DF, Brasil.

4 Rua Cecimbra Jaques, 111. CEP 90870-150. Porto Alegre, RS, Brasil.

* Corresponding author. E-mail: fronzabio@yahoo.com.br

ABSTRACT: The result of an inventory on the lepidopteran fauna associated with the South American Holly (*erva-mate*) is described. This inventory was based on a bibliographic review, the examination of specimens deposited in scientific collections from the state of Rio Grande do Sul, the systematic collection within commercial *erva-mate* farms located in Anta Gorda, RS, and on the occasional collections from native trees within the Região dos Vinhedos campus of the Universidade de Caxias do Sul. Every fifteen days between 20 August 2005 and 19 August 2006, 150 trees were visually inspected and all Lepidoptera were collected. A list of 75 lepidopteran taxa belonging to 16 families is presented, including 27 new occurrences for the *erva-mate* crop, of which 14 species represent new records for the state.

INTRODUCTION

Among the plants explored in the southern region of Brazil, the *erva-mate* (*Ilex paraguariensis* – Aquifoliaceae, South American Holly) is a very evident crop with significant socioeconomic, environmental and cultural importance that forms one of the most characteristic agroforestry systems (Bozzetto 1995; Tormen 1995). As a tree that is naturally ombrophilous, it becomes predisposed to pest attacks after being subjected to physiological stress, when cultivated in open areas, and especially as a monoculture (Carpanezzi 1995; Borges *et al.* 2003). The simplification of the ecosystem to a monoculture tends to generate impacts that favor some phytophagous insect species. The observed incidence of pests is less abundant in natural *erva-mate* stands that include other forest species (Borges *et al.* 2003).

Almost one hundred phytophagous insects are associated with *erva-mate* plants, especially among the Lepidoptera and Coleoptera (Diaz 1997). With respect to Lepidoptera, species which harm *erva-mate* plants have been mentioned since the beginning of the last century (Jorgensen 1921; Ronna 1934; Parseval 1937; Bourquin 1944). The number of harmful species has increased over time, where, Penteado (1995) listed 20, Azevedo and Corseuil (1996) 30, and Diaz (1997) 41 lepidopteran taxa. This increasing tendency in the number of associated species is mainly due to most previous research only recording species with large economic impact, namely *Hedypathes betulinus* (Coleoptera: Cerambycidae) (d'Avila *et al.* 2006) and *Gyropsylla spagazziniana* (Hemiptera: Psyllidae) (Chiaradia *et al.* 2002), and the failure to consider the ecosystem as a whole. Most Lepidoptera occur at low population levels, without causing expressive economic damage to the *erva-mate* crop, resulting in very

restricted knowledge of their species diversity, biology, and population dynamics. Nevertheless, together, these insects can limit production, making periodical monitoring fundamental, considering that ecological imbalances, including those occurring due to the monoculture itself, can elevate one or more species to pest condition (Penteado 1995).

Given that specific knowledge can provide grounds for alternative cultural management and, at the same time, contribute to the knowledge of the insect fauna associated with regional native plants, the objective of this study is to contribute an updated list of the Lepidoptera associated with the *erva-mate* plant.

MATERIALS AND METHODS

The inventory data of the Lepidoptera associated to the *erva-mate* crop was obtained from a bibliographic review, from the examination of specimens deposited in scientific collections within the state of Rio Grande do Sul (RS), from the systematic collection within commercial *erva-mate* crops, and from occasional collecting.

The bibliographic research encompassed all literature on the *erva-mate* plant, from Brazil, Paraguay, Argentina and Uruguay, registering the first author to record a specific Lepidoptera species on the crop. The records with only generic level information were considered when associated specimens were present in collections.

Specimen examination included those from scientific collections of the following institutions: the Museu de Ciências e Tecnologia of the Pontifícia Universidade Católica do Rio Grande do Sul, Porto Alegre (MCTP); the Museu Ramiro Gomes Costa, Fundação Estadual de Pesquisa Agropecuária, Porto Alegre (MRGC); the Museu Anchieta de Ciências Naturais, Porto Alegre (MAPA); the

Museu de Ciências Naturais of the Fundação Zoobotânica do Rio Grande do Sul, Porto Alegre (MCNZ); the Museu Entomológico Ceslau Biezanko, Faculdade de Agronomia Eliseu Maciel, Universidade Federal de Pelotas, Pelotas (MECB); the Coleção Andrej Menchov Bertels of the Centro de Pesquisa Agropecuária de Clima Temperado, EMBRAPA, Pelotas (CAMB); the Museu de Ciências Naturais, Universidade Católica de Pelotas, Pelotas (MUCP); the Setor de Entomologia, Faculdade de Agronomia da Universidade Federal do Rio Grande do Sul, Porto Alegre (FASE); and the collection of the Laboratório de Biologia do Departamento de Ciências Exatas e da Natureza of the Universidade de Caxias do Sul (CUCS).

The systematic collecting occurred every fifteen days, between 20 August, 2005 and 19 August, 2006, for a total of 27 collections, in the municipality of Anta Gorda, of the Encosta Inferior do Nordeste region, state of RS. Three commercial *erva-mate* farms with a minimum regrowth of one year and without the use of insecticides were selected and fifty randomly chosen trees were visually examined in each farm. The study areas are located within a quadrilateral between the coordinates 28°56'40" S, 52°01'12" W; 28°57'11" S, 52°01'13" W; 28°56'56" S, 52°00'56" W and 28°56'44" S, 52°01'00" W.

The occasional collecting occurred on several native *erva-mate* plants found in Bento Gonçalves, within the Região dos Vinhedos campus of the Universidade de Caxias do Sul, CARVI-UCS, from which leaves were removed for the feeding of reared larvae.

After collection, larvae were reared in the CARVI Laboratório de Biologia, under a temperature of 25 ± 1°C, RH 70 ± 10%, with a 14 hour photophase, and fed with *erva-mate* leaves. Upon emergence, adults were spread and dried and eventually incorporated in the CUCS and MCTP collections as voucher material. The specimens were identified using specific bibliographic resources (e.g.

D'Abbrera 1995; 1986; Kitching and Cadiou 2000; Willmott 2003; Lamas 2004) and sent to specialists when necessary.

The nomenclature was updated following Davis (1964) for Psychidae; Heppner (1995) for Dalceridae, Limacodidae, Megalopygidae and Tortricidae; Heppner (1996) for Apatelodidae, Lasiocampidae and Saturniidae; Kitching and Cadiou (2000) for Sphingidae; Lamas (2004), for Hesperidae, Nymphalidae and Riodinidae; Pogue (2002) for *Spodoptera*; Poole (1989) for the remaining Noctuidae; Scoble (1999) for Geometridae and Teston and Corseuil (2004) for Arctiinae.

RESULTS AND DISCUSSION

Table 1 lists the Lepidoptera in alphabetical order, by genus and species, also including the first bibliographical reference which cites the *erva-mate* as a host plant and the occurrence of the taxon in the state of Rio Grande do Sul. Additionally, when deposited material exists, the collection and/or field capture data is indicated.

A list of 75 lepidopteran taxa belonging to 16 families is presented, including 27 new occurrences for the *erva-mate* crop whose adults are shown in Figures 1-27. The number of lepidopteran taxa from published records, the specimens found in collections, and those obtained by collecting are presented in Figure 28.

A bibliographic review, with the updated nomenclature and the exclusion of Lepidoptera cited as "sp." when voucher material was not available, resulted in 48 taxa. During the systematic collecting in farms and/or occasional collections in CARVI-UCS, thirty seven Lepidoptera were obtained, including 26 new occurrences for the *erva-mate* crop, of which 15 species represent new records for the state of Rio Grande do Sul. Six other taxa were found after examining collections, of which two are new records for the *erva-mate* crop, although one was also found during collecting.

TABLE 1. Lepidoptera occurring on *erva-mate* plants, indicating the presence in the sampling (collecting in three commercial *erva-mate* farms within the municipality of Anta Gorda, between August 20, 2005 and August 19, 2006, and occasional samples), Museum and/or literature records, with first author to record the species for the crop and for the state of Rio Grande do Sul.

Number of individuals collected. * First records for the *erva-mate* crop. ** First records for the state of Rio Grande do Sul. Literature records: 1- Baucke 1960a; 2- Baucke 1960b; 3- Bertels and Bauke 1966; 4- Biezanko 1949; 5- Biezanko 1961; 6- Biezanko 1982; 7- Biezanko 1986; 8- Biezanko and Freitas 1938; 9- Biezanko et al. 1974; 10- Bourquin 1943; 11- Bourquin 1944; 12- Bouvier 1930; 13- Corseuil et al. 2002; 14- De coll and Saini 1992; 15- Hayward 1969; 16- Iede and Machado 1989; 17- Jorgensen 1921; 18- Jorgensen 1932; 19- Kober and Vargas 1960; 20- Koehler 1934; 21- Koehler 1935; 22- Lima 1947; 23- Mabilde 1896; 24- Mattos 1982; 25- Muelo 1946; 26- Oliveira et al. 1999; 27- Parseval 1937; 28- Pastrana 2004; 29- Ronna 1923; 30- Ronna 1933; 31- Ronna 1934; 32- Silva et al. 1968; 33- Specht and Corseuil 1996; 34- Specht et al. 2005; 35- Teston and Corseuil 2004; 36- Trujillo 1995; 37- Weimer 1894; 38- Weimer 1907.

SPECIES	FAMILY	SAMPLING #	MUSEUM RECORDS	LITERATURE RECORDS	
				MATE	RS
<i>Acharya barbara</i> (Dyar, 1905)	Limacodidae			1	
<i>Acharya nesea</i> (Stoll, 1780)	Limacodidae			5	30
<i>Acharya trimacula</i> (Sepp, [1848])	Limacodidae			5	23
<i>Acraga moorei</i> Dyar, 1898	Dalceridae: Acraginae			14	
<i>Adelpha calliphane</i> Fruhstorfer, 1915	Nymphalidae: Limenitidinae			15	32
<i>Adelpha hyas hyas</i> (Doyère, [1840])	Nymphalidae: Limenitidinae	30		32	2
<i>Adelpha serpa serpa</i> (Boisduval, 1836)	Nymphalidae: Limenitidinae			3	3
<i>Adelpha thessalia indefecta</i> Fruhstorfer, 1913	Nymphalidae: Limenitidinae	172		17	4
<i>Adelpha thoasa gerona</i> (Hewitson, 1867)	Nymphalidae: Limenitidinae			15	
<i>Adelpha zea</i> (Hewitson, 1850)	Nymphalidae: Limenitidinae			9	4
<i>Alabama argillacea</i> (Hubner, 1823)	Noctuidae: Calpinae			31	31

TABLE 1. CONTINUED.

SPECIES	FAMILY	SAMPLING #	MUSEUM RECORDS	LITERATURE RECORDS	
				MATE	RS
<i>Apatelodes martia</i> (Stoll, 1782)	Apatelodidae: Apatelodinae			9	7
<i>Apatelodes tropea</i> (Schaus, 1896)	Apatelodidae: Apatelodinae			9	
<i>Argyrotaenia fletcheriella</i> (Köhler, 1939)	Tortricidae: Tortricinae			28	
<i>Argyrotaenia sphaleropa</i> (Meyrick, 1909)	Tortricidae: Tortricinae			14	5
<i>Astrartes fulgurator fulgurator</i> (Walch, 1775)	Hesperidae: Pyrginae			17	32
<i>Automeris illustris</i> (Walker, 1855)	Saturniidae: Hemileucinae	8		21	23
<i>Automeris naranja</i> Schaus, 1898	Saturniidae: Hemileucinae	18		10	38
<i>Bertholdia soror</i> Dyar, 1901	Noctuidae: Arctiinae	5		*	35
<i>Carales astur</i> (Cramer, 1777)	Noctuidae: Arctiinae	11		*	23
<i>Cerodirphia opis</i> (Schaus, 1892)	Saturniidae: Hemileucinae			34	12
<i>Citheronia brissotii brissotii</i> (Boisduval, 1868)	Saturniidae: Ceratocampinae	7		25	23
<i>Citheronia brissotii meridionalis</i> Bouvier, 1927	Saturniidae: Ceratocampinae			15	3
<i>Citheronia laocoon laocoon</i> (Cramer, 1777)	Saturniidae: Ceratocampinae			32	32
<i>Chalodeta theodora</i> (C. and R. Felder, 1862)	Riodinidae: Riodininae	1		*	23
<i>Dalcerina tijucana</i> (Schaus, 1892)	Dalceridae: Dalcerinae	1		24	5
<i>Dysschema sacrifica</i> (Hübner, [1831])	Noctuidae: Arctiinae	3		*	37
<i>Eacles imperialis magnifica</i> (Drury, 1773)	Saturniidae: Ceratocampinae	4		*	8
<i>Emesis mandana mandana</i> (Cramer, 1780)	Riodinidae: Riodininae			17	23
<i>Hylesia nigricans</i> (Berg, 1875)	Saturniidae: Hemileucinae	26		11	23
<i>Hylesia oratex</i> Dyar, 1913	Saturniidae: Hemileucinae			22	13
<i>Iridopsis aviceps</i> Prout, 1932	Geometridae: Ennominae	1		*	**
<i>Leucanella janeira</i> (Westwood, [1854])	Saturniidae: Hemileucinae			34	13
<i>Leucanella viridescens</i> (Walter, 1855)	Saturniidae: Hemileucinae			2	23
<i>Lophocampa citrina</i> (Sepp, [1843])	Noctuidae: Arctiinae			16	32
<i>Lumacra</i> sp.	Psychidae	5		*	**
<i>Manduca albiplaga</i> (Walker, 1856)	Sphingidae: Sphinginae			20	6
<i>Megalopyge urens</i> Berg, 1882	Megalopygidae	6		3	5
<i>Melanolophia</i> sp.	Geometridae: Ennominae	1		*	**
<i>Microgonia perfulvata</i> Dognin, 1916	Geometridae: Ennominae	1		*	**
<i>Nyceryx continua continua</i> Walker, 1856	Sphingidae: Macroglossinae	2	MRGC	*	26
<i>Oenoptila mixtata</i> Gueneé, [1858]	Geometridae: Ennominae	18		*	**
<i>Oxydia apidania</i> Cramer, [1779]	Geometridae: Ennominae			17	
<i>Oxydia distichata</i> Gueneé, [1858]	Geometridae: Ennominae			36	23
<i>Oxydia mundata</i> Guenee, [1858]	Geometridae: Ennominae			28	
<i>Oxydia sociata</i> Warren, 1895	Geometridae: Ennominae	2		*	**
<i>Oyketicus geyeri</i> (Berg, 1877)	Psychidae			27	30
<i>Oyketicus kirbyi</i> (Lands-Guilding, 1827)	Psychidae			27	29
<i>Paracles fusca</i> (Walker, 1856)	Noctuidae: Arctiinae	22		*	30
<i>Paracles variegata</i> (Schaus, 1896)	Noctuidae: Arctiinae	2		*	35
<i>Parasa cucumenica</i> Dyar, 1926	Limacodidae	1		*	**
<i>Pelochita cinerea</i> (Walker, 1855)	Noctuidae: Arctiinae	3		*	35
<i>Periga circumstans</i> Walker, 1855	Saturniidae: Hemileucinae	14		*	7
<i>Perigonia ilus</i> Boisduval, 1870	Sphingidae: Macroglossinae			17	
<i>Perigonia lusca lusca</i> (Fabricius, 1777)	Sphingidae: Macroglossinae			20	23
<i>Perigonia passerina</i> Boisduval, [1875]	Sphingidae: Macroglossinae	3		*	**
<i>Pherotesia subjecta</i> Warren, 1905	Geometridae: Ennominae	1		*	**
<i>Phobetron hipparchia</i> (Cramer, 1777)	Limacodidae	2		14	23
<i>Pleuopruca asthenaria</i> Walker, 1861	Geometridae: Ennominae	1		*	**
<i>Polia marea</i> (Schaus, 1894)	Noctuidae: Hadeninae		MRGC	*	33

TABLE 1. CONTINUED.

SPECIES	FAMILY	SAMPLING #	MUSEUM RECORDS	LITERATURE RECORDS	
				MATE	RS
<i>Rothschildia aurota aurota</i> (Cramer, 1775)	Saturniidae: Saturniinae			17	23
<i>Rothschildia aurota speculifera</i> (Walker, 1855)	Saturniidae: Saturniinae	8		18	7
<i>Rothschildia jacobaeae</i> (Walker, 1855)	Saturniidae: Saturniinae			9	37
<i>Sabulodes</i> sp.	Geometridae: Ennominae	3		*	**
<i>Spodoptera cosmioides</i> (Walker, 1858)	Noctuidae: Xyleninae	1		*	23
<i>Stenalcidia dukinfieldia</i> Schaus, 1897	Geometridae: Ennominae	1		*	**
<i>Stenoma armata</i> Zeller, 1877	Oecophoridae			28	
<i>Stenoma decora</i> Zeller, 1854	Oecophoridae			14	
<i>Tachuda pachydexius</i> Forbes, 1939	Notodontidae	1		*	**
<i>Talima</i> sp.	Limacodidae	1		*	**
<i>Thelosia camina</i> Schaus, 1896	Apatelodidae: Apatelodinae			19	19
<i>Thyrintheina arnobia arnobia</i> Stoll, 1782	Geometridae: Ennominae	2		9	**
<i>Titya proxima</i> (Burmeister, 1878)	Lasiocampidae: Macromphalinae	3		*	7
<i>Urbanus proteus proteus</i> (Linnaeus, 1758)	Hesperidae: Pyrginae			25	23
<i>Utethesia ornatrix</i> (Linnaeus, 1758)	Noctuidae: Arctiinae			28	37

In the current list, Geometridae was the family with the greatest increase in number of taxa, going from four to 13, which is reinforced by observations from local farmers who report an increase in the occurrence of “looper/spanworm” caterpillars. Within this group, *Thyrintheina arnobia arnobia*, recorded in the *erva-mate* crop from Argentina (Chiarelli 1943; De Coll and Saini 1992; Trujillo 1995), and referred to as the eucalyptus pest in Brazil (Peres Filho and Berti Filho 2003; Holtz et al. 2003), presents its first distributional record for the state of Rio Grande do Sul.

Among the *Adelpha*, identified using Willmott (2003) and Lamas (2004), only *A. hyas hyas* and *A. thessalia indefecta* were recorded, already cited in the bibliography although under names considered synonyms. The records of *A. serpa hyas*, published by Bertels and Baucke 1966 and Azevedo and Corseuil 1996 and based on Mattos (1982) citing *A. hyas*, were eliminated as they have *A. serpa hyas* as a synonym.

The *erva-mate* caterpillar *Thelosia camina* (Kober and Vargas 1960), whose presence in crops had already been detected in 1931 and 1934 (Parseval 1937), was not found in collections nor was it collected during this study. This species is described in the literature as one of the main pests for this crop, due to the intensive damage it caused during past decades. However, the attack by larvae of this species has not been further recorded in the literature, although adults have been captured in light traps, reflected by recently deposited specimens in the MCTP (under numbers 15744 and 15745).

Saturniidae presented the highest number of taxa (Figure 29), with the collection of *Periga circumstans* representing a new record. The larvae with the highest defoliating potential are found within this group, be it due to the size of the specimens, which require a larger quantity of food for their development, or due to their gregarious behavior, observed mostly among the Hemileucinae, which can completely defoliate trees.

Furthermore, larvae belonging to this subfamily contain urticating hairs (Lemaire 2002) and can cause problems and inconveniences among laborers. The occurrence of *Leucanella janeira* and *Cerodirphia opis* was recently recorded (Specht et al. 2005), although the literature has two citations of *Cerodirphia* sp. by Iede and Machado (1989) and Penteado (1995).

Perigonia passerina, identified using the image in D'Abbrera (1986) and the systematic proposal of Kitching and Cadiou (2000), and *Nyceryx continua continua*, both new records for the crop, were Sphingids collected during field activities. The taxa registered in the literature as *Pengonia lusca* F. forma *ilus* [sic] (Jorgensen 1921) and *Perigonia lusca ilus* (Muelo 1946) are herein considered as being *P. ilus*, following the systematic proposal of Kitching and Cadiou (2000). Likewise, *Perigonia lusca*, considered as one of the main pests of this crop in Argentina (De coll and Saini 1992; Trujillo 1995) and also registered in Paraná (Alves et al. 2001), is considered as being *P. lusca lusca*.

The Noctuidae, previously represented by three species, now has eleven, including the Arctiinae (*sensu* Lafontaine and Fibiger 2006). *Spodoptera cosmioides* is one of the new records among the Lepidoptera sampled during systematic collecting. Although the literature already has a citation of *Spodoptera* sp., by Diaz (1997), referring to Trujillo (1995), this taxon does not appear in the publication. This is possibly based on De Coll and Saini (1992), who present the image of a caterpillar of this genus which corresponds to the description of Zenker et al. (2007) for *S. cosmioides*. Another new record is of *Polia marea*, resulting from the examination of specimens in collections with an occurrence record for Rio Grande do Sul (Specht and Corseuil 1996).

Two families are included in the list: Lasiocampidae and Notodontidae, represented by the occurrence of *Titya proxima* and *Tachuda pachydexius*, respectively. Several other collected specimens are not included here because

the larvae, clearly different from the listed species, died due to pathogens and parasitoids, preventing the emergence of adults.


The examination of the previously deposited material in the State's scientific collections resulted in two new records for the *erva-mate* crop: *Nyceryx continua continua* and *Polia marea*, of which the former was also collected during the present study. These two species belong to the MRGC, whose collection is characterized by providing host plant information, which, with rare exceptions, does not occur in other collections. The absence of information in other collections only permitted the occurrence record by locality and season. This indicates the importance of also including biological information for specimens which are deposited in collections. With this information it is possible to assemble precise data banks on species occurrence, pest status, host plant range, and would assist in the detection of long-term environmental and ecological changes.

The final list excluded some taxa from the literature due to identification problems and incongruences. *Biblis hyperia* cited by De Coll and Saini (1992) was excluded because the published image corresponds to the caterpillar of *Adelpha hyas hyas*. *Hylesia paulex* and *H. remex* cited by Borges et al. (2003) were not included, as the record refers only to the presence of adults, captured in light traps, which does not necessarily indicate that their larvae are


hosted by the crop. Likewise, *H. nanus* was not included because the citation by Santos et al. (1988) was based on Vellozo (1951) who mentions only *Hylesia* sp. without a specific name.

Leucanella memusae registered by Pastrana (2004) was not included because it is a synonym of *L. viridescens*. For the same reason, although previously registered by Silva et al. (1968), this citation should not be considered because it refers to *L. viridescens* cited by Baucke (1960b). Specht et al. (2005) also registered *L. memusae*, taking the name adopted by Silva et al. (1968) into consideration. Furthermore, according to Lemaire (2002), the geographical distribution of *L. memusae* does not include Argentina.


The diversity revealed in this study is well above that which had been registered for this crop (Figure 28), although the majority of the species and subspecies collected are represented by few specimens. This means that greater attention should be given to the Lepidoptera and other insects associated with this crop, in virtue of both their ecological function and management practices. Rapid population growth of many insect species is observed as a result of increases in crop area, the formation of artificially selected genetically similar populations, and the management of native plants into monocultural systems (Andow 1983; 1990; Iede and Machado 1989; Penteadó 1995).




FIGURES 1-4. New records for the *erva-mate* crop: 1- *Bertholdia soror*; 2- *Carales astur*; 3- *Chalodeta theodora*; 4- *Dysschema sacrificia*. Photos: Edegar Fronza.



FIGURES 5-14. New records for the erva-mate crop: 5- *Eacles imperialis magnifica*; 6- *Iridopsis aviceps*; 7- *Lumacra* sp.; 8- *Melanolophia* sp.; 9- *Microgonia perfulvata*; 10- *Nyceryx continua continua*; 11- *Oenoptila mixtata*; 12- *Oxydia sociata*; 13- *Paracles fusca*; 14- *Paracles variegata*. Photos: Edgar Fronza.



FIGURES 15-24. New records for the *erva-mate* crop: 15- *Parasa cucumenica* 16- *Pelochita cinerea*; 17- *Periga circumstans*; 18- *Perigonia passerina*; 19- *Pherotesia subjecta*; 20- *Pleuroprucha asthenaria*; 21- *Polia marea*; 22- *Sabulodes* sp.; 23- *Spodoptera cosmioides*; 24- *Stenalcidia dukinfieldia*. Photos: Edegar Fronza.



FIGURES 25-27. New records for the *erva-mate* crop: 25- *Tachuda pachydexius*; 26- *Talima* sp.; 27- *Titya proxima*. Photos: Edgar Fronza.




FIGURE 28. Number of Lepidoptera taxa associated with *erva-mate* plants in the State of Rio Grande do Sul, resulting from the bibliographic review, examination of scientific collections and sampling (field collecting in three commercial *erva-mate* farms within the municipality of Anta Gorda, between August 20, 2005 and August 19, 2006, and occasional samples). There is a new record (*Nyceryx continua continua*) both from the collections and the collecting.




FIGURE 29. Number of Lepidoptera taxa per family occurring on *erva-mate* plants in the state of Rio Grande do Sul.

ACKNOWLEDGMENTS: To the curators Eduardo J. E. Silva (MECB), Elvia E. S. Vianna (MUCP), Felipe do Canto Quadros (MCTP), Fernando R. Meyer (MAPA), Luciano A. Moura (MCNZ), Mirtes Mello (CAMB), Vera R. S. Wolf (MRGC) and their staff, for permitting access and support during specimen examination. To José A. Teston and Alfred Moser, for the identification of the Arctiinae and Geometridae, respectively. To Professor Gervásio Carvalho for his advice during the finalization of the Masters degree. To the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - CAPES, for the fellowship.

LITERATURE CITED

Alves, L.F.A., D.L.Q Santana and R.M.C. Brancalhão. 2001. Ocorrência de *Perigonia lusca* (Fabr.) (Lep., Sphingidae) em *erva-mate* (*Ilex paraguariensis*) no Brasil. *Neotropical Entomology* 30(2): 339-340.

Andow, D. 1983. The extent of monoculture and its effects on insect pest populations with particular reference to wheat and cotton. *Agriculture, Ecosystems and Environment* 9: 25-35.

Andow, D.A. 1990. Population dynamics of insect herbivore in simple and diverse habitats. *Ecology* 71(3): 1006-1017.

Azevedo, E.C.G. and E. Corseuil. 1996. Insetos ocorrentes em *erva-mate*; p. 35-45 In Prefeitura Municipal de Venâncio Aires (org.). *Erva-mate: diagnósticos e perspectivas de desenvolvimento*. Venâncio Aires: Prefeitura Municipal.

Baucke, O. 1960a. Notas entomológicas II: *Sibine bárbara* Dyar, 1905 (Lep. Eucleidae) em *erva-mate* no Paraná. *Iheringia, Série Zoologia* 12: 7-9.

Baucke, O. 1960b. Notas entomológicas III: O gênero *Automeris* Hübner, 1819 (Lepidoptera, Hemileucidae) no Rio Grande do Sul. *Iheringia, Série Zoologia* 12: 11-19.

Bertels, A.M. and O. Baucke. 1966. Segunda relação das pragas das plantas cultivadas no Rio Grande do Sul. *Pesquisa agropecuária brasileira* 1: 17-46.

Biezanko, C.M. 1949. *Acraeidae, Heliconidae et Nymphalidae de Pelotas e seus arredores*. Pelotas: Ed. do autor. 16 p.

Biezanko, C.M. 1961. Castniidae, Zygaenidae, Dalceridae, Eucleidae, Megalopygidae, Cossidae et Hepialidae da Zona Missioneira do Rio Grande do Sul. *Arquivos de Entomologia, Série B*, 14:1-12.

Biezanko, C.M. 1982. Sphingidae da região sueste do Rio Grande do Sul. *Revista Centro de Ciências Rurais* 12(1): 59-75.

Biezanko, C.M. 1986. Adelocephalidae, Saturniidae, Mimallonidae, Lasiocampidae, Eupterotidae e Lymantridae da Região Missioneira do Rio Grande do Sul. *Revista Centro de Ciências Rurais* 16(2): 89-112.

Biezanko C. M. and R.G. de Freitas. 1938. Catálogo dos insetos encontrados na cidade de Pelotas e seus arredores. Fascículo 1 – Lepidopteros. *Boletim Escola Agronomia "Eliseu Maciel"* 25: 1-32.

Biezanko, C.M., A. Ruffinelli and D. Link. 1974. Plantas y otras sustancias alimenticias de las orugas de los lepidopteros uruguayos. *Revista Centro Ciências Rurais* 4(2): 107-148.

Borges, L.R., S.M.N. Lazzari and F.A. Lazzari. 2003. Comparação dos sistemas de cultivo nativo e adensado de *erva-mate*, *Ilex paraguariensis* St. Hil., quanto à ocorrência e flutuação populacional de insetos. *Revista Brasileira de Entomologia* 47(4): 563-568.

Bourquin, F. 1943 Observaciones sobre *Automeris aurantiaca*, Weyen., 1907 (Lep. Saturniidae). *Revista Sociedade Entomologica Argentina, Comunicaciones* 12 (2): 149.

- Bourquin, F. 1944. Observaciones sobre *Hylesia nigricans* (Berg, 1875); p. 45-47 In F. Bourquin (ed.). *Mariposas Argentinas Vida, desarrollo, costumbres y hechos curiosos de algunos lepidopteros argentinos*. Buenos Aires: El Ateneo.
- Bouvier, E.L. 1930. Sur la systématique des Saturnioides de la famille des Hémileucidés. *Comptes Rendus Hebdomadaires des Séances de l'Académie des Sciences* 191: 743-746.
- Bozzetto, D.J. 1995. Aspectos econômicos e sociais da cultura da *erva-mate* na região alta do Vale do Alto Taquari, Rio Grande do Sul, Brasil; p. 207-213 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi. *Erva-mate: Biologia e cultura no Cone Sul*. Porto Alegre: Ed. UFRGS.
- Carpanezi, A.A. 1995. Cultura da *erva-mate* no Brasil: conflitos e lacunas; p.43-46 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi. *Erva-mate: Biologia e cultura no Cone Sul*. Porto Alegre: Ed. UFRGS.
- Chiaradia, L.A., J.M. Milanez and A. Zidko. 2002. Estimativas das gerações anuais de *Gyropsylla spegazziniana* (Lizer, 1917) em função de sua exigência térmica. *Ciência Rural* 32(3): 385-391.
- Chiarelli, A. 1943. Um geometrido prejudicial a la yerba mate, *Thyrintheina arnobia*. *Revista Argentina de Agronomía* 10(3): 250-255.
- Corseuil, E., A. Specht and C. Lang. 2002. Saturniídeos (Lepidoptera, Saturniidae) registrados para o Rio Grande do Sul, Brasil. I. Hemileucinae. *Biociências* 10(2): 147-155.
- D'Abbrera, B. 1986. *Sphingidae Mundi. Hawk moths of the world*. Faringdon: E.W. Classey. 225 p.
- D'Abbrera, B. 1995. *Saturniidae Mundi – saturniid moths of the world*. Keltorn: Automeris Press, 178 p.
- Davis, D.R. 1964. Bagworm Moths of the Western Hemisphere. *Smithsonian Institution United States National Museum Bulletin* 244: 1-232.
- D'Ávila, M., E.C. Lúcio and J.V.C. Guedes. 2006 Bioecologia e manejo da broca-da-erva-mate *Hedypathes betulinus* Klug (1825) (Coleoptera: Cerambycidae). *Ciência Florestal* 16(2): 233-241.
- De Coll, O.R. and E.D. Saini. 1992. *Insectos y acaros perjudiciales al cultivo de la yerba mate en la Republica Argentina*. Montecarlo: Instituto Nacional de Tecnología Agropecuária. 48 p.
- Diaz, C.Y.F. 1997. Perspectivas del manejo integrado de plagas em yerba mate; p. 371-390 In I Congresso Sul-Americano da Erva-mate e II Reunião Técnica do Cone Sul Sobre a Cultura da Erva-mate. Curitiba: Anais.
- Hayward, K.J. 1969. Datos para el estudio de la ontogenia de lepidópteros argentinos. *Miscelánea del Instituto Miguel Lillo* 31: 1-142.
- Heppner, J.B. 1995. *Atlas of Neotropical Lepidoptera. Checklist Part 2: Hyblanoidea – Pyraloidea – Tortricoidea*. Gainesville: Association for Tropical Lepidoptera. 243 p.
- Heppner, J.B. 1996. *Atlas of Neotropical Lepidoptera. Checklist Part 4: Drepanoidea – Bombycoidea – Sphingoidea*. Gainesville: Association for Tropical Lepidoptera. 88 p.
- Holtz, A.M., H.G. Oliveira, A. Pallini, M. Venzon, J.C. Zanuncio, J.S. Marinho and M.C. Rosado. 2003. Desempenho de *Thyrintheina arnobia* Stoll (Lepidoptera: Geometridae) em Eucalipto e Goiaba: o Hospedeiro Nativo Não é um Bom Hospedeiro? *Neotropical Entomology* 32(3): 427-431.
- Iede, E.T. and D.C. Machado. 1989. Pragas da *erva-mate* (*Ilex paraguariensis* St. Hill.) e seu controle. *Boletim de Pesquisa Florestal* 18/19: 51-60.
- Jorgensen, P. 1921. Sobre algunos nuevos enemigos de la yerba-mate *Ilex paraguariensis*. *Revista de la Sociedad Científica del Paraguay* 1(1): 27-30.
- Jorgensen, P. 1932. Lepidopterologisches aus Sudamerika. *Deutsche Entomologische Zeitschrift* 46: 37-66.
- Kitching, I.J. and J.M. Cadiou. 2000. *Hawkmoths of the world: an annotated and illustrated revisionary checklist*. Ithaca: Cornell University Press, 227 p.
- Kober, E. and E. Vargas. 1960. Biologia e controle da "lagarta da erva-mate" (*Thelosia camina* Schaus, 1920; Lepidoptera – Eupterotidae). *Revista Escola Agronomia Veterinária UFRGS* 3(1): 7-35.
- Koehler, P. 1934. Catálogo preliminar de los lepidópteros argentinos dañinos. *Boletín Ministerio Agricultura de la Nación* 36(1): 25-45.
- Koehler, P. 1935. Notas sobre Automeriæ (Lep. Saturn.) Argentinos. *Revista Sociedade Entomologica Argentina* 7: 79-91.
- Lafontaine, J.D. and M. Fibiger. 2006. Revised higher classification of the Noctuoidea (Lepidoptera). *Entomological Society of Canada* 138: 610-635.
- Lamas, G. 2004. Checklist: part 4A. Hesperioidea – Papilionoidea; p. 1-439 In G. Lamas (ed.). *Atlas of neotropical lepidoptera*. Gainesville: Association for Tropical Lepidoptera.
- Lemaire, C. 2002. *The Saturniidae of America - Hemileucinae*. Keltorn: Goecke & Evers. 1388 p.
- Lima, A.D.F. 1947 [1945]. Insetos fitófagos de Santa Catarina. *Boletim Fitossanitário* 2(3-4): 233-251.
- Mabilde, A. P. 1896. *Guia practica para os principiantes colecionadores de insectos, contendo a descrição fiel de perto de 1000 borboletas com 180 figuras lithographadas em tamanho, formas e desenhos conforme o natural. Estudo sobre a caça, classificação e conservação de uma coleção mais ou menos regular*. Porto Alegre: Gundlach & Schuldt, 238 p.
- Mattos, N.F. 1982. *Estudos preliminares sobre pragas da erva-mate*. Porto Alegre: Instituto de Pesquisas de Recursos Naturais Renováveis. 18 p. [Publicação IPRNR n° 9].
- Muelo, A.C. 1946. *Yerba mate: su cultivo y explotacion*. Buenos Aires: Sudamericana, 217 p.
- Oliveira, R.B., A. Specht and E. Corseuil. 1999. Esfingídeos (Lepidoptera, Sphingidae) ocorrentes no Rio Grande do Sul. *Biociências* 7(1): 167-177.
- Parseval, M. von. 1937. *As moléstias e pragas mais comuns da erva-mate no Rio Grande do Sul*. Porto Alegre: Secretaria do Estado dos Negócios, da Agricultura, Indústria e Comércio. 30 p. [Boletim n° 55].
- Pastrana, J.A. (Coord.) 2004. *Los lepidópteros argentinos: sus plantas hospedadoras y otros sustratos alimentícios*. 1ª ed. Buenos Aires: Sociedad Entomológica Argentina, 350 p.
- Penteado, S.R.C. 1995. Principais pragas da *erva-mate* e medidas alternativas para o seu controle; p. 109-120 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi. *Erva-mate: Biologia e cultura no Cone Sul*. Porto Alegre: Ed. UFRGS.
- Peres Filho, O. and E. Berti Filho. 2003. Exigências térmicas de *Thyrintheina arnobia* (Stoll, 1782) e efeitos da temperatura na sua biologia. *Ciência Florestal* 13 (2): 143-151.
- Pogue, G.M. 2002. A world revision of the genus *Spodoptera* Guenée (Lepidoptera: Noctuidae). *Memoirs of the American Entomological Society* 43: 1-202.
- Poole, R.W. 1989. Noctuidae; p. 501-1013. In J.B. Heppner (ed.). *Lepidopterorum Catalogus*. New York: Brill.
- Ronna, E. 1923. Apotamentos de microfauna Rio-grandense. *Egatea - I Série: Lepidópteros* 3(4): 253-257.
- Ronna, E. 1933. Catálogo dos insetos até hoje encontrados nas plantas do Rio Grande do Sul. *Egatea* 18(1/2): 47-53; (4): 197-202.
- Ronna, E. 1934. Catálogo dos insetos até hoje encontrados nas plantas do Rio Grande do Sul. *Egatea* 19(1): 15-20.
- Santos, G.P., N. Anjos and J.C. Zanuncio. 1988. Biologia de *Hylesia nanus* (Walker, 1855) (Lepidoptera: Attacidae) desfoliadora de Citeiira (*Joannesia priceps*: Euphorbiaceae). *Revista Ceres* 35(201): 479-485.
- Scoble M.J. (1999). *Geometrid moths of the world: a catalogue (Lepidoptera, Geometridae)*. London: CSIRO Publishing, Natural History Museum. 1016 p.
- Silva, A.G.D'A., C.R. Gonçalves, D.M. Galvão, A.J.L. Gonçalves, J. Gomes, M.M. Silva and L. Simoni. 1968. *Quarto catálogo dos insetos que vivem nas plantas do Brasil, seus parasitos e predadores. Parte II – 1º Tomo*. Rio de Janeiro: Ministério da Agricultura. 622 p.
- Specht, A. and E. Corseuil. 1996. Lista documentada dos noctuídeos (Lepidoptera: Noctuidae) ocorrentes no Rio Grande do Sul, Brasil. *Biociências* 4(2): 131-170.
- Specht, A., E. Corseuil and A.C. Formentini. 2005. Lepidópteros de importância médica ocorrentes no Rio Grande do Sul. III. Saturniidae - Hemileucinae. *Biociências* 13(2): 149-162.
- Teston, J.A. and E. Corseuil. 2004. Diversidade de Arctiinae (Lepidoptera, Arctiidae) capturados com armadilha luminosa, em seis comunidades no Rio Grande do Sul, Brasil. *Revista Brasileira de Entomologia* 48(1): 77-90.
- Tormen, M.J. 1995. Economia ervateira brasileira; p. 27-40 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi. *Erva-mate: Biologia e cultura no Cone Sul*. Porto Alegre: Ed. UFRGS.
- Trujillo, M.R. 1995. Agroecosistema yerbatero de alta densidad: plagas y enemigos naturales; p. 129-134 In H. Winge, A.G. Ferreira, J.E.A. Mariath and L.C. Tarasconi. *Erva-mate: Biologia e cultura no Cone Sul*. Porto Alegre: Ed. UFRGS.
- Vellozo, L.G.C. 1951. Sobre a ocorrência de *Hylesia* sp nos ervais do Paraná. *Ciência e Cultura* 3(1): 33.
- Weymer, G. 1894. Exotische Lepidopteren. VII. Beitrag zur Lepidopterenfauna von Rio Grande do Sul. *Stettiner Entomologische Zeitung* 55(10-12): 311-333.
- Weymer, G. 1907. Exotische lepidopteren. *Deutsche Entomologische Zeitschrift* 20: 1-51.
- Willmott, K.R. 2003. *The genus Adelpha: its systematics, biology and biogeography (Lepidoptera: Nymphalidae: Limenitidini)*. Gainesville: Association for Tropical Lepidoptera. 323 p.
- Zenker, M.M., A. Specht and E. Corseuil. 2007. Estágios imaturos de *Spodoptera cosmíodes* (Walker) (Lepidoptera, Noctuidae). *Revista Brasileira de Zoologia* 24(1): 99-107.

RECEIVED: April 2011

LAST REVISED: June 2011

ACCEPTED: July 2011

PUBLISHED ONLINE: July 2011

EDITORIAL RESPONSIBILITY: Matthew Smart