

ISSN 2040-4700

SEPTEMBER 2011

VOLUME 2 PART 2



Advances in Animal Biosciences

Proceedings of the 8th International
Symposium on the Nutrition of Herbivores
(ISNH8)



advances in
animal
biosciences

CAMBRIDGE
UNIVERSITY PRESS

The effect of lipid sources on the methane emission of beef cattle at pasture using the SF6 tracer technique

Isabela P.C. Carvalho, Telma T. Berchielli, Alexandre Berndt, Rosa T. S. Frighetto, Giovanni Fiorentini

Construction and installation of six modular open-circuit chambers for monitoring gas emissions from large ruminants

Nico Peiren, Bart Sonck, Joaquin Castro Montoya, Daniël De Brabander, Veerle Fievez, Sam De Campeneere

Molecular analysis of methanogenic archaea in the forestomach of the alpaca (*Vicugna pacos*) reveals differences in population structure between individual hosts

Benoit St-Pierre, André-Denis G. Wright

Optimization of Roughage to Concentrate Ratio in Sweet Sorghum Bagasse Based Complete Ration for Efficient Microbial Biomass Production in Sheep Using In Vitro Gas Technique

N. Nalini Kumari, Y. Ramana Reddy, M. Blummel, D. Nagalakshmi, T. Monika, M. Pavani, M. Sudhakar Reddy, Ch. Ravinder Reddy, B.V.S. Reddy

The effect of incremental levels of Sandalwood oil on methane emissions and animal performance in sheep

Jolien Veneman, John Newbold, Hink Perdok, Walter Gerrits, Jan Dijkstra, Sander van Zijderveld

Potential methane production using different feed formulation by in vitro rumen fermentation

Seon-Ho Kim, Sang-Suk Lee, Che-Ok Jeon, Young-Kyoon Oh, Md. Jahangir Alam, Lovelia Mamuad, Chang-Dae Jeong

Effects of stearidonic acid-rich Echium oil on methane production in a RUSITEC system

P. Amaro, M.R.G. Maia, R.J. Dewhurst, A.J.M. Fonseca, A.R.J. Cabrita

The effect of differential physical forms of rapeseed as fat supplement on rumen NDF digestion and methane emission in dairy cows

Maike Johannes, Peter Lund, Martin Riis Weisbjerg, Anne Louise Hellwing, Torben Hvelplund

Effect of decreasing phosphorus intake and structure value on intake, milk production and phosphorus status of dairy cows

Wilfried van Straalen, Bart Tas¹, Ivonne Kok, Thomas Schonewille

Methane emissions using tanniferous native plants in ruminal fermentation in vitro

Emmanuel Andrade-Rivero, Octavio Alonso Castelán-Ortega, Julieta Gertrudis Estrada-Flores

The role of genetic improvement for productivity traits in reducing enteric methane relative to the amount of lamb carcass produced from the Welsh national sheep flock

Catherine Nakielny, Janet Roden, Jamie Newbold

Evaluation of volatile fatty acid stoichiometry models to predict enteric methane production in lactating Holstein cows

Jan Dijkstra, Andre Bannink, Jennifer Ellis, James France, Ermias Kebreab, Yael Morvay

The effect of lipid sources on the methane emission of beef cattle at pasture using the SF6 tracer technique

I P C Carvalho¹, T T Berchielli^{1,2}, A Berndt³, R T S Friguetto⁴, G Fiorentini¹

¹UNESP, Jaboticabal, Brazil, ²INCT/CA Member, Brazil, Brazil, ³EMBRAPA, São Carlos, Brazil, ⁴EMBRAPA, Jaguariúna, Brazil

Email: carvalho_isabela@yahoo.com.br

Introduction Methane is known to be a potent greenhouse gas and its release to the atmosphere contributes to global warming. The losses of dietary energy average about 5-12% in methane (Van Soest, 1991) and it contributes to 68 % of the Brazilian emission of this gas. The methane production in the rumen is intrinsic to the rumen fermentation, acting as a hydrogen (H₂) drain, so the reduction on methane emission requires an alternative route for the H₂, avoiding its accumulation. Lipid sources can reduce methanogens and protozoa through a toxic effect and also act as an H₂ drain due to the biohydrogenation process.

Materials and methods Twenty Nellore steers (initial average body weight of 440kg) were assigned to five treatments of a completely randomized design. The animals were divided in to 10 paddocks (2 paddocks per treatment) of *Brachiaria brizantha* cv. Xaraés. Linseed oil, palm oil, soy bean grain and by-pass fat (Lactoplus®) were added to a supplement offered to the animals once a day (an amount of 1,0% of the body weight). The control treatment was composed of an energy-protein supplement with no additional fat. All the concentrate containing 20% CP and 10% EE (except the control supplement, which contained 3% EE). The animals were weighed every 28 days and slaughtered at 495.6 kg. The methane emissions were evaluated using the SF6 tracer gas technique (Johnson *et al.*, 1994; Westberg *et al.*, 1998, adapted by Primavesi *et al.* 2004). Permeation tubes with known release rates for SF6 were placed in the rumen, 72 hours prior to the start of gas sampling. Each tube was charged with 500-600 mg of SF6 and incubated at 39°C for calibration. Release rates of SF6 were determined by measuring the weight loss of tubes for 6 wk to establish a steady pre-determined rate. Release rates of the permeation tubes used in this study ranged from 930 to 1600 ng min⁻¹. Gases exhaled from the nose and mouth were drawn into pre-evacuated collection tubes, through a capillary tubing. The collection system was designed to deliver half its volume during a 24-hr collection, ensuring a uniform collection rate. Five consecutive 24-hr gas samples were collected from each animal. The fixed effect was the treatment. The paddocks were used as the experimental unit. The methane emission and average daily gain were analysed using the tukey test, with a 5% probability.

Results There was no effect ($P>0.05$) of lipid sources on the average daily gain (ADG) of beef steers at pasture fed with 1% body weight supplement. The methane emissions were expressed in kilograms of methane emitted per year (kg CH₄.yr⁻¹), gram of methane emitted per day (g CH₄.day⁻¹), milligram of methane emitted per day per kilogram of body weight (mg CH₄.day.BW⁻¹), gram of methane emitted per day per kilogram of metabolic body weight (g CH₄.day.MBW⁻¹) and kilograms of methane emitted per kilogram of carcass produced (kg CH₄.kg CAR⁻¹). The methane emission was also not different ($P>0.05$) among the treatments in any unit expressed.

Table 1 Average daily gain and methane emission of animals receiving lipid sources on supplement

| Variables | Treatments | | | | | CV (%) |
|--|------------|----------|-------------|-------------|----------|--------|
| | Control | Palm oil | Linseed oil | By pass fat | Soy bean | |
| ADG | 0.564 | 0.567 | 0.661 | 0.556 | 0.561 | 12.123 |
| kg CH ₄ .yr ⁻¹ | 41.515 | 41.115 | 25.615 | 37.190 | 30.100 | 19.443 |
| g CH ₄ .day ⁻¹ | 113.735 | 112.645 | 70.180 | 101.880 | 82.470 | 19.446 |
| mg CH ₄ .day.BW ⁻¹ | 238.651 | 228.516 | 147.044 | 208.547 | 180.847 | 11.770 |
| g CH ₄ .day.MBW ⁻¹ | 1.115 | 1.077 | 0.687 | 0.980 | 0.836 | 13.627 |
| kg CH ₄ .kg CAR ⁻¹ | 0.236 | 0.245 | 0.144 | 0.227 | 0.180 | 22.604 |

Means within each variable followed by different letters differ by tukey test ($P < 0.05$), $n=2$.

ADG = average daily gain; BW = body weight; MBW = metabolic body weight; CAR = carcass

Conclusions Lipid sources such as palm oil, linseed oil, soy bean and by pass fat were not effective in mitigating methane emissions in Nellore steers, when included in a supplement with 10% EE, offered at a quantity of 1,0% BW to animals at tropical pasture.

Acknowledgements Bellman® Animal Nutrition; FAPESP

References

- Johnson, K.A. & Johnson, D.E. Measurement. 1994. Environmental Science and Technology. 28, 359-362.
- Primavesi, o., Friguetto, R. T. S., Pedreira, M. S., Lima, M. A., Berchielli, T. T., Demarchi, J. J. A. A., Manella, M. Q., Barbosa, P. F., Johnson, K. A., Westberg, H.H. Técnica do Gás traçador SF6 para medição de campo do metano ruminal em bovinos: adaptações para o Brasil. São Carlos: EMBRAPA Pecuária Sudeste, 74 p. (Documentos, 39).
- Van Soest, P.J. 1994. Nutritional ecology of the ruminant. 2. ed. Ithaca: Comstock. 476p.
- Westberg, H., Johnson, K.A., Cossalman, M.W., Michal, J.J. 1998. A SF6 tracer technique: methane measurement from ruminants. Washington State University, Pullman, Washington.: 40p.