




16th WORLD FERTILIZER CONGRESS OF CIEC

TECHNOLOGICAL INNOVATION FOR A
SUSTAINABLE TROPICAL AGRICULTURE

PROCEEDINGS



International Scientific Centre of Fertilizers (CIEC)



16th WORLD FERTILIZER CONGRESS OF CIEC

**TECHNOLOGICAL INNOVATION FOR A
SUSTAINABLE TROPICAL AGRICULTURE**

October 20-24, 2014
Rio de Janeiro, RJ - Brazil

PROCEEDINGS

*Vinicius de Melo Benites
Adilson de Oliveira Junior
Paulo Sergio Pavinato
Paulo César Teixeira
Milton Ferreira Moraes
Regina Maria Villas Bôas de Campos Leite
Ronaldo Pereira de Oliveira*
Editors

Rio de Janeiro, RJ
2014

CHARCOAL (BIOCHAR) AS SOIL CONDITIONER TO ENHANCE FERTILIZERS AND WATER USE EFFICIENCY IN AGRICULTURE IN ACID TROPICAL SOILS IN THE CENTRAL AMAZON BASIN

WENCESLAU G. TEIXEIRA¹, CHRISTOPH STEINER², GILVAN C. MARTINS³, MURILO R. DE ARRUDA⁴

¹ Embrapa Soil, Rua Jardim Botânico, 1024, Rio de Janeiro, 22460-000, Brazil (wenceslau.teixeira@embrapa.br); ² University of Kassel, Steinstr. 13, 37213, Witzenhausen, Germany; ^{3,4} Embrapa Amazônia Ocidental, km 30 – Rod. AM 010 – km 30, Manaus, Brazil (gilvan.coimbra@embrapa.br; murilo.arruda@embrapa.br)

Introduction

Biochar is a name for charred organic material (charcoal - carvão vegetal in Portuguese) when it is used as a soil amendment. Different methods of pyrolysis is used to make charcoal. In the Amazon the traditional way is the called "*caiera*", that consist in earth kiln is digged and filled with vegetal debris (mainly trunks of trees) its is covered with soil. Some opening vents in the soil allow to control the flux of oxygen into the kiln. Another traditional technique is the called hot tail (*rabo quente*) which is still a simple technique using a kiln made with bricks. After the kiln is filled a small fire is ignited at the kiln entrance then the door is sealed with soil and the vegetal residues is left to undergo incomplete combustion, creating charcoal and some amount of ash (Swami et al. 2009). Charcoal as soil amendment is traditionally used by the population in the Amazon to enhance soils to cultivate some horticultural crops as medicines and spices. Moreover in the Amazon ash from burned vegetation is the lime and the fertilizers used in the traditional slash and burn or shifting cultivation, a system of land clearing practiced by several thousand of small farmers in the Amazon. In spite of potential agronomic benefits to enhance soil quality (Teixeira et al., 2010) biochar has also the potential to reduce carbon emissions (Lehmann et al., 2005). Many publications sad that biochar can increase soil fertility, increase agricultural productivity, and provide protection against some foliar and soil-borne diseases. Furthermore, adding biochar can prolongate the time of cultivation it may reduces pressure on forests areas to agriculture as happens in Amazonian Dark Eath sites (German, 2003).

The Amazon soils comprise a large spectrum of soil with different potential to be used in the agriculture, in the Central Amazon the soils of

the upland areas are mainly dystrofic Ferralsols and Acrisols (Teixeira et al., 2011). However in the floodplains of many rivers, rich in sediments, the predominant soils are eutrofic Gleysols and Fluvisols. In the upland area in the Central Amazon, the only fertile soil (eutrofic) were the pretic Anthrosols (Amazonian Dark Earths), called regionally as Terra Preta de Índio, this type of soil has many good agronomic qualities (Teixeira et al., 2010) and can be considered the source of inspiration to apply biochar in soils (Benitez et al., 2010; Steiner et al., 2010; Steiner et al., 2009). The TPI shows normally a high concentration of P, Ca, Mn and large stocks of carbon. Moreover a large amount of the carbon in this type of soil seems to be of pyrogenic origin (Glaser et al. 2001) and has large amount of charge (CEC) (Liang et al., 2010). This article summarize result of many experiments conducted in 15 years of research with biochar in nursery and field experiments in the Central Amazon.

Material and methods

The material and methods used in the experiments were detailed described in the publications listed in the reference section.

Results and discussion

The results summarize the main results of many experiments conducted by the authors et al during the last decade

Fields experiments

In Central Amazon many fields experiments were conducted testing different levels of biochar in combination with different treatments (organic and mineral fertilizers). Also the response in productivity in annual crops (rice and sorghum) were studied (Steiner et al., 2007). In experiments with perennial crops, Banana (*Musa sp*) and Guarana (*Paulinia*

cupana) the treatments with addition of biochar reconfirm the previous results of potential enhance in the productivity for a longer period and to improve the CEC, respectively the studies of Steiner et al., 2007, Steiner et al., 2008 and Arruda and Teixeira, 2010 and Arruda et al. 2012. In an field experiment, with application of charcoal pieces in planting holes for a banana plantation (Steiner et al., 2009) addition of charcoal increased the water hold at potential of 100 kPa that indicate a that charcoal ammendment enhance soil pores with radius of ~ 0.015 mm. Either biochar promoted the aggregation with soil pores with pores of this size or biochar's porosity itself was responsible for this increase. The experiments with application of biochars in the Central Amazon show that only mineral fertilizer can also not guarantee higher productivity for a long period. Is that necessary to apply some source of organic matter in combination with the fertilizers and biochar is an option. In spite of the difficult to isolate single mechanism to explain some improvement in productivity and soil quality in acid tropical soils, the results are clearly related with reduction of leaching rates and enhancing efficiency of fertilization management.

Nursery experiments

Experiments were conducted using charcoal as component o growing media in nurseries for different plant species: Guaraná (*Paulinia cupana*); Crajiru (*Arabidae chica*) and Brazil Nut (*Bertholletia excelsa*) the results are published respectively in Arruda et al., (2007); Souza et al., (2007) and Nunes and Teixeira, (2010). The main conclusion of these experiments is that charcoal pieces can be a component of growing media in substitution to A horizon and commercial components (as vermiculite, montmorillonite) with similar results and the advantage to be cheaper and regionally available. The water holding capacity in lower tension of soil with adition of charcoal can be similar or higher that provide by addition of active clays.

In the future more sophisticated technologies may produce biochar by specific demand combining more stable charred material to reduce emission or to delivery nutrients and to enhance cation exchange capacity (CEC). Many benefits of charcoal application related in the literature are probably

related with the liming effect of ash that "contaminated" many biochar. Application of biochar with large amount of ash in alkaline soils some advantages may change to disadvantages. The most biochar can not be commercialized in the Brazil as fertilizer as do not attend the minimum required by legislation by the most part of nutrients and it should be refered as soil ammendment or conditioner.

Conclusions

Biochar has a potential when combined with fertilizers (mineral and organic) to enhance productivity in the dystrofic Ferrralsols in the Central Amazon.

Biochar has a potential to replace the A horizon in the nursery as a component of the growing media

Biochar from wood residues done in the traditional ways in the central Amazon (caieras and rabo quente) is a mixed of charred material and ash on liming, fertilizing and enhance water hold capacity.

Keywords: Biochar, tropical soils, CEC, Latossolo Amarelo

Acknowledgements

To Embrapa Amazônia Ocidental and our colleagues there (Gilvan Coimbra Martins, Murilo Arruda) to University of Bayreuth that carried for many years a cooperation with Embrapa in Brazil concerning Terra Preta de Índio and Biochar application (Wolfgang Zech, Bruno Glaser, Johannes Lehmann). To Embrapa, University Federal do Amazon (UFAM), CNPq and Capes for funding the studies and and scholarships for students and authors.

References

- ARRUDA M, MOREIRA A, TEIXEIRA W, SOUZA W, ATROCH A AND NASCIMENTO FILHO FD. 2012. Yield, soil fertility, and nutritional status of guarana clones cultivated with organic source of nutrients. *Revista de Ciências Agrárias / Amazonian Journal of Agricultural and Environmental Sciences* 55: 311-317.
- ARRUDA MRD AND TEIXEIRA WG 2010. Utilização de resíduos de carvão vegetal associado a fontes orgânicas de nutrientes no manejo sustentável do solo e do guaranazeiro (*Paullinia cupana* var. *sorbilis* (Mart.) Ducke) na Amazônia Central.

- As Terras Pretas de Índio da Amazônia: sua caracterização e uso deste conhecimento na criação de novas áreas, Manaus: Edua / Embrapa, p. 307-314.
- ARRUDA M.R.D., PEREIRA J.C.R.R., MOREIRA A.N. & TEIXEIRA W.G. Survival rate of guarana herbaceous cuttings in different growing media. *Ciência Agrotécnica*, 31:236-241, 2007
- BENITES VDM, TEIXEIRA WG, REZENDE ME AND PIMENTA AS 2009. Utilização de Carvão e Subprodutos da Carbonização Vegetal na Agricultura: Aprendendo com as Terras Pretas de Índio. In: TEIXEIRA, WG, KERN, DC, MADARI, BE, LIMA, HN AND WOODS, WI (Eds.) *As Terras Pretas de Índio da Amazônia: sua caracterização e uso deste conhecimento na criação de novas áreas*, Manaus: Embrapa Amazônia Ocidental, p. 286.
- GERMAN L 2003. Historical contingencies in the coevolution of environment and livelihood: contributions to the debate on Amazonian Black Earth. *Geoderma*, p. 307-331.
- GLASER B, HAUMAIER L, GUGGENBERGER G AND ZECH W. 2001. The "terra preta" phenomenon: a model for sustainable agriculture in the humid tropics. *Naturwissenschaften* 88: 37 - 41.
- LEHMANN J, GAUNT J AND RONDON M. 2005. Bio-char sequestration in terrestrial ecosystems - a review. *Mitigation and Adaptation Strategies for Global Change* 11: 403-427.
- LIANG, B.; LEHMANN, J.; SOLOMON, D.; KINYANGI, J.; GROSSMAN, J.; O'NEILL, B.; SKJEMSTAD, O.; THIES, J.; LUIZÃO, F. J.; PETERSEN, J.; NEVES, E. G. Black carbon increases cation exchange capacity in soils. *Soil Science Society of American Journal*, Madison, v. 70, p. 1719, 2006.
- NUNES M.M. & TEIXEIRA W.G. Crescimento de mudas de Castanheiras do Brasil (*Bertholletia excelsa*) em função de doses de carvão vegetal como componente de substrato. In: III Reunião Científica da Rede CTPETRO Amazônia, 2010, Manaus.
- SOUZA G., TEIXEIRA W., REIS R., CHAVES F. & XAVIER J. Growth of cajuru (*arrabidaea chica* verlot.) on different growing media. *Revista Brasileira de Plantas Mediciniais*, Botucatu, 8:62-65, 2006
- STEINER C., DE ARRUDA M.R., TEIXEIRA W.G. & ZECH W. Soil respiration curves as soil fertility indicators in perennial central amazonian plantations treated with charcoal, and mineral or organic fertilizers. *Tropical Science*, 47:218-230, 2007
- STEINER C., GARCIA M. & ZECH W. Effects of charcoal as slow release nutrient carrier on n-p-k dynamics and soil microbial population: Pot experiments with ferralsol substrate. In: W. I. Woods, W. G. Teixeira, J. Lehmann, C. Steiner and A. Winkler-Prins, eds. *Book Title* Berlin, Springer, 2009. p. 325.
- STEINER C., GLASER B., GERALDES TEIXEIRA W., LEHMANN J., BLUM W.E. & ZECH W. Nitrogen retention and plant uptake on a highly weathered central amazonian ferralsol amended with compost and charcoal. *Journal of Plant Nutrition and Soil Science*, 171:893-899, 2008
- STEINER, C.; TEIXEIRA, W.; ZECH, W. The Effect of Charcoal in Banana (*Musa Sp.*) Planting Holes: an on-farm study in Central Amazonia, Brazil. In: (Ed.) *Amazonian Dark Earths: Wim Sombroek's Vision*: Springer, 2009. p.423-432.
- STEINER C, GLASER B, TEIXEIRA WG, LEHMANN J, BLUM WEH AND ZECH W. 2008. Nitrogen retention and plant uptake on a highly weathered Central Amazonian Ferralsol amended with compost and charcoal. *Journal of Plant Nutrition and Soil Science* 47: 218.
- STEINER, C.; TEIXEIRA, W.; WOODS, W.; ZECH, W. Indigenous knowledge about terra preta formation. In: (Ed.) *Amazonian Dark Earths: Wim Sombroek's Vision*: Springer, 2009. p.193-204.
- STEINER, C.; TEIXEIRA, W.; ZECH, W. The Effect of Charcoal in Banana (*Musa Sp.*) Planting Holes: an on-farm study in Central Amazonia, Brazil. In: (Ed.) *Amazonian Dark Earths: Wim Sombroek's Vision*: Springer, 2009. p.423-432
- STEINER C., TEIXEIRA W.G., LEHMANN J., NEHLS T., BLUM W. & ZECH W. Long term effects of manure, charcoal and mineral fertilization on crop production and fertility on a highly weathered central amazonian upland soil. *Plant and Soil*, 291:275, 2007

STEINER, C.; TEIXEIRA, W. G.; NARAYAN, S.; MAJOR, J.; LEHMANN, J.; ZECH, W.; BLUM, W. E. H. Corte e carbonização como uma alternativa ao corte e queima: estudos na Amazônia. In: TEIXEIRA, W. G.; KERN, D. C.; MADARI, B. E.; LIMA, H. N.; WOODS, I. W. (Ed.). *As terras pretas de índio da Amazônia: sua caracterização e uso deste conhecimento na criação de novas áreas*. Manaus: EDUA/Embrapa Amazônia Ocidental, 2010. p. 298-306. ed., 2010.

SWAMI SN, STEINER C, TEIXEIRA W AND LEHMANN J 2009. Charcoal making in the Brazilian Amazon: economic aspects of production and carbon conversion efficiencies of kilns. *Amazonian Dark Earths: Wim Sombroek's Vision*: Springer, p. 411-422.

TEIXEIRA, W. G.; ARRUDA, W.; SHINZATO, E.; MACEDO, R. S.; MARTINS, G. C.; LIMA, H. N.; RODRIGUES, T. E. Solos do Estado do Amazonas. In: MAIA, M. A. M. e MARMOS, J. L. (Ed.). *Geodiversidade do estado do Amazonas*. 1. Manaus: CPRM, 2010. p.71 - 86.

TEIXEIRA W.G., KERN D.C., MADARI B.E., LIMA H.N. & WOODS W. *As terras pretas de índio da Amazônia: Sua caracterização e uso deste conhecimento na criação de novas áreas*. Manaus, Editora da Universidade Federal do Amazonas (EDUA) / Embrapa Amazônia Ocidental, 2010. 420 p.



Figure 1. Traditional use of charcoal pieces (biochar) to enhance soil to cultivated horticultural crops in the Amazon